

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

Forecasted deep stratospheric intrusions over Central Europe: case studies and climatologies

T. Trickl¹, H. Feldmann^{2,*}, H.-J. Kanter¹, H.-E. Scheel¹, M. Sprenger³, A. Stohl⁴, and H. Wernli^{3,**}

¹Forschungszentrum Karlsruhe, Institut für Meteorologie und Klimaforschung (IMK-IFU), Kreuzeckbahnstr. 19, 82467 Garmisch-Partenkirchen, Germany

²Rheinisches Institut für Umweltforschung, Univ. zu Köln, Aachener Str. 201–209, 50931 Köln, Germany

³Eidgenössische Technische Hochschule (ETH) Zürich, Institut für Atmosphäre und Klima, Universitätstr. 16, 8092 Zürich, Switzerland

⁴Norwegian Institute for Air Research, P.O. Box 100, Instituttveien 18, 2027 Kjeller, Norway

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

* now at: Forschungszentrum Karlsruhe, Institut für Meteorologie und Klimaforschung (IMK-TRO), Postfach 3640, 76021 Karlsruhe, Germany

** now at: Johannes-Gutenberg-Universität Mainz, Institut für Physik der Atmosphäre, 55099 Mainz, Germany

Received: 8 September 2008 – Accepted: 22 December 2008 – Published: 26 January 2009

Correspondence to: T. Trickl (thomas.trickl@imk.fzk.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

Based on daily predictions of stratospheric air intrusions, obtained from trajectory calculations by ETH Zürich with wind fields from ECMWF forecasts, a high number of measurements with the ozone lidar at IMK-IFU (Garmisch-Partenkirchen, Germany) were carried out in 2001. The lidar measurements show a large variety of rather different cases. In part, tropopause folds could be fully captured. The frequency of intrusion cases forecasted and verified by vertical sounding or in the in-situ data recorded at the nearby Zugspitze summit (2962 m a.s.l.) exceed that in previous work by more than a factor of two. Three cases mapped with the lidar were selected to validate the results for the corresponding time periods extracted from a one-year run with the new hemispheric version of the chemistry-transport model EURAD. Due to the high spatial resolution chosen for these simulations the agreement with the lidar measurements is satisfactory. The Zugspitze ozone data from 1978 to 2004 were recently filtered by applying different criteria for stratospheric air, based on the ^7Be and humidity measurements. Here, by using the daily model forecasts during the time period 2001–2005, we examine three criteria and determine how well they represent the stratospheric air intrusions reaching the mountain site. Seasonal cycles for the period 2001–2005 were derived for the forecasts as well as the intrusion frequency per month for the forecasted intrusions and each of the criteria, distinguishing eight different characteristic transport pathways. In most cases a winter maximum and a summer minimum was obtained, but in the case of cyclonic arrival of intrusions starting over Greenland a late-spring maximum is seen. Two of the filtering criteria examined, based on combining a relative-humidity (RH) threshold of 60% with either a ^7Be threshold of 5.5 mBq m^{-3} or the requirement for $\text{RH} \leq 30\%$ within $\pm 6 \text{ h}$, rather reliably predict periods of deep intrusions reaching the Zugspitze station. An “or” combination of both these criteria yields slightly more cases and covers 77.9% of the intrusions identified. The lack of observations in the complementary 22.1% are mostly explained by overpasses. In this way the long-term trend of stratospheric ozone observed at this site as well as the correspond-

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ing ozone budget may be derived on the basis of measurements only. This effort will be the subject of a subsequent publication.

1 Introduction

With the advent of quasi-continuous lidar remote sensing the investigation of stratosphere-to-troposphere transport (STT) has strongly revived (e.g., Browell et al., 1987; Ancellet et al., 1991; Langford et al., 1996; Ravetta et al., 1999; Eisele et al., 1999; Stohl and Trickl, 1999; Baray et al., 2000; Grant et al., 2000; Cho et al., 2001; Zanis et al., 2003; Trickl et al., 2003; Galani et al., 2003; Flentje et al., 2005; Papayannis et al., 2005). The ability of the lidar method to generate vertically highly resolved dense time series yields complete or partial mappings of stratospheric air intrusions previously not that easily accessible. In this way also a valuable basis for the quality assurance of numerical models is established. In particular in the frame of the European (EU) projects VOTALP (Vertical Ozone Transport in the ALPS; VOTALP, 2000a, b) and STACCATO (Influence of Stratosphere-Troposphere Exchange in a Changing Climate on Atmospheric Transport and Oxidation Capacity; STACCATO, 2003, <http://www.forst.tu-muenchen.de/EXT/LST/METEO/staccato/>) the lidar measurements and the continuous measurements at high-altitude Alpine summit stations have stimulated major efforts in model development, application and validation (e.g., Stohl and Trickl, 1999; Stohl et al., 2000; Zanis et al., 2003; Cristofanelli et al., 2003; Meloen et al., 2003; Roelofs et al., 2003). Many models are, still, far from being quantitative. This may, in part, be ascribed to insufficient spatial resolution, combined in Eulerian models with numerical diffusion, and problems in properly describing the free-tropospheric mixing. The meteorological fields provided, e.g., by the European Centre for Medium-Range Weather Forecasts (ECMWF) underlying many of the simulations, in principle, capture stratospheric intrusions rather accurately (Zanis et al., 2003). However, vertical displacements with respect to the observed ozone structures in part visible in model results for the northern rim of the Alps suggest to improve the model orography, i.e.,

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the horizontal resolution.

The vertical sounding activities within STACCATO (2000–2002) and the German follow-up project ATMOFAST (Atmospheric Long-range Transport and its Impact on the Trace-gas Composition in the Free Troposphere over Central Europe; ATMOFAST, 2005) were based on four-day trajectory forecasts by ETH (Eidgenössische Technische Hochschule) Zürich (Zanis et al., 2003). In this way, the measurements could be called for and synchronized in a highly systematic way. In Garmisch-Partenkirchen (Germany), particularly frequent lidar measurements were carried out during the period between February and August 2001, covering 20 of the 33 predicted STT events. This selection was mainly due to limiting the warnings at that time to sufficiently large direct intrusions reaching the majority of the partner stations. The measurements were, whenever possible, extended over several days which yielded a rich data set containing information on both STT and trans-Atlantic transport. A first case study was published by Zanis et al. (2003). In this study and in a companion paper on long-range advection of stratospheric air (Trickl et al., 2009a) we give a more complete report of the results obtained in 2001 which cover a large variety of rather different situations.

Lidar measurements based on intrusion forecasts were started in Garmisch-Partenkirchen in 1996 (Eisele et al., 1999), initially using a “classical” approach previously derived for the intrusion forecast for the nearby mountain-top stations Zugspitze (2962 m a.s.l.) and Wank (1780 m a.s.l.) since the 1970s (e.g., Sladkovic and Munzert, 1990). This forecast method implied criteria such as the position and an anti-cyclonic bending of the jet stream and a potential-temperature range typically between 295 and 305 K. A 98-% probability of detecting at least one stratospheric air tongue at the Zugspitze summit during a predicted intrusion period was found from a comparison with the lidar measurements for the 23 months of investigation (1996–1997). The occasional discrepancies found in the 1990s were mostly limited to the warm season.

Although this approach is not expected to yield a coverage of intrusion events as complete as the more recently used trajectory calculations by ETH Zürich based upon ECMWF forecasts this demonstrates that the continuous data series available for these

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



two summit stations since the 1970s could be highly suitable for quantifying STT reaching the respective altitudes in this area from measurements alone. The ETH results are now also used to review the data-selection criteria.

There have been numerous attempts to quantify STT from experimental data (e.g., Danielsen and Mohnen, 1977; Viezee et al., 1983; Beekmann, et al., 1997; more information may be obtained from the references in these papers and in the general overview papers by Holton et al., 1996 and Stohl et al., 2003). Results from case studies or a large number of vertical profile measurements have been used and extrapolated to the Northern Hemisphere or the entire globe by using assumptions such as model-derived geographical distributions of the tropopause folding activity. Beekmann et al. (1997) analysed sonde data and selected relative-humidity (RH) thresholds of 20 or 25% together with ozone increases by at least 20% with respect to a climatological mean profile for their analysis of tropopause folds in the entire free troposphere. In addition, meteorological criteria such as the existence of a nearby upper-tropospheric jet stream, a pronounced potential temperature gradient or elevated potential vorticity were included. This effort, extended to up to 26 years of sounding data, yielded an annual coverage by STT of 3 to 5% for the free troposphere above western European stations as well as a rather constant seasonal cycle. A cross-tropopause ozone flux of 5.7×10^{10} molecules $\text{cm}^{-2} \text{s}^{-1}$ was estimated for the Northern Hemisphere and 1.2×10^{11} molecules $\text{cm}^{-2} \text{s}^{-1}$ for western Europe.

A number of studies to assess the ozone budget for intrusions reaching 3000 m over Central Europe have already been carried out based on filtering Zugspitze data. For this station almost continuous measurements (24-h averages) of the isotope ^7Be , formed in the stratosphere to about 67% (Lal and Peters, 1967), have been made since 1970 (e.g., Pötzl and Carnuth, 1966; Reiter et al., 1971). A threshold for the integrated specific activity of ^7Be of 8 mBq m^{-3} for unambiguous intrusions was recommended by Sladkovic and Munzert (1990), while Eisele et al. (1999) state that 5 mBq m^{-3} is already strongly indicative of a deep intrusion event. Other filtering criteria, further narrowing the STT time intervals, are based on significant drops in relative humidity

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

and a simultaneous rise in ozone. Elbern et al. (1997) carried out a statistical analysis of the Zugspitze data with variable thresholds for the ten-year period 1984–1993. As a first step the correlated behaviour of ^7Be , ozone and relative humidity was examined. In a second step the statistical significance of the identified potential intrusion events was calculated for an eleven-day period around each event and insignificant cases were excluded. For ^7Be and O_3 thresholds of 2.9 and 2.1 times the standard deviation of the values with respect to the running monthly mean were chosen, respectively. In this way, 195 stratospheric intrusions could be unambiguously identified for the Zugspitze summit, 85 of which were also observed at the lower-lying Wank station. A seasonal cycle of the stratospheric fraction of the Zugspitze ozone with a winter maximum and a summer minimum was determined.

In a rather thorough study Stohl et al. (2000), again, analysed the Zugspitze data (1991–1997) as well as those from the VOTALP partner stations Jungfraujoch (3580 m a.s.l.), Sonnblick (3106 m a.s.l.) and Monte Cimone (2165 m a.s.l.) for a more confined period of two to three years. For the Zugspitze data the results are strongly depending on the thresholds for the data filtering when going from the selection criteria $[^7\text{Be}] > 10 \text{ mBq m}^{-3}$ and $\text{RH} < 30\%$ to $[^7\text{Be}] > 6 \text{ mBq m}^{-3}$ and $\text{RH} < 30\%$. No proof of the predominantly stratospheric nature of the air corresponding to the less critical thresholds was given, which is an important topic of the investigations presented here. In addition to the filtering of the station data, a three-year run (January 1995–September 1997) with the FLEXPART tracer model was carried out which yielded rather good agreement with the analyses of the station results as well as the influence of the tropospheric age of the stratospheric air tongues on the seasonal cycle. The results for a calculation subsequently extended to the years 1995–1999 are shown in Fig. 1. The seasonal cycle published by Elbern et al. (1997) for the Zugspitze summit is qualitatively confirmed, but both analyses for the late 1990s (station and model), exhibit a more pronounced summer minimum. The summer minimum is less pronounced at the higher-lying Jungfraujoch than at the other three stations, and, thus, the seasonal cycle resembles more that obtained by Beekmann et al. (1997) for the entire free tro-

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

posphere.

The FLEXPART results for the Zugspitze summit suggest an annual average of about 4.2 ppb (about 8.4% of the annual average ozone) of ozone originating in direct deep intrusions reaching 3000 m in that area. This was approximately confirmed in more recent re-analyses of the measurements by Scheel (Scheel, 2003; ATMOfAST, 2005) yielding 8.8% of stratospheric ozone contribution per year, however, based on a restrictive threshold for ^7Be (85th percentile, which coincides with 8 mBq m^{-3} in the late 1990s; see also (Cristofanelli et al., 2006) for a similar approach for the partner station Mte.-Cimone in Italy). Furthermore, a positive trend of STT ozone reaching the Zugspitze summit was found. The ^7Be data suggest a start of that increase in the mid-seventies.

In this paper, we give a final summary of the observations of STT events within STACCATO and ATMOfAST, in relation to the forecasts. In particular, we discuss (as mentioned) the most interesting examples from the very successful lidar sounding period 2001, including additional analyses with the FLEXPART and EURAD models for further clarification and for validation of the new hemispheric version of EURAD. In addition, based on the forecast data, we address the important question of how many of the predicted stratospheric air intrusions descend to the Alpine summit levels here represented by the Zugspitze station (2962 m a.s.l.). This effort, in turn, also confirms the excellent quality of the forecasts. Furthermore, the forecasts serve as a referee to judge the existing and new approaches for identifying the presence of stratospheric air based on correlating complementary station data. We also give some statistical analysis (2001–2005) for the most important transport pathways and their relative impact on the observations at this Alpine site.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2 Methods

2.1 Ozone lidar

The tropospheric ozone lidar at IMK-IFU (47°28'37" N, 11°3'52" E, 740 m a.s.l.) was completed in its first version in 1990 (Kempfer et al., 1994) and upgraded in 1994 and 1995 (Eisele and Trickl, 1996; 2005). It has a unique vertical range between 0.2 km and roughly 15 km above the ground, features low uncertainties of accuracies of about ± 3 ppb in the lower troposphere and ± 6 ppb (under optimum conditions) in the upper troposphere. The upper-tropospheric performance may be degraded in the presence of high lower-tropospheric ozone concentrations absorbing a lot of the ultraviolet laser emission and by enhanced sky light in summer, in particular in the presence of clouds. The vertical resolution is dynamically varied, roughly between 50 m and a few hundred metres, depending on the signal-to-noise ratio that decreases with altitude. The lidar has been used in numerous investigations mostly focussing on atmospheric transport (e.g., Eisele et al., 1999; Stohl and Trickl, 1999; Seibert et al., 2000; Carnuth et al., 2002; Trickl et al., 2003; Zanis et al., 2003).

2.2 Zugspitze series

For the identification of stratospheric intrusions at the Zugspitze station (summit: 2962 m a.s.l.) half-hourly averages of ozone, carbon monoxide and relative humidity (RH) as well as 24-h averages of the isotope ^7Be have been used. The ^7Be series at the Zugspitze is the longest and goes back to 1970 (e.g., Pöttl and Carnuth, 1966; Reiter et al., 1971; Scheel, 2002; Zanis et al., 2003a; Tositti et al., 2004). Ozone has been measured since 1978 (e.g., Reiter et al., 1987; Scheel et al., 1997; Oltmans et al., 2006). A chemiluminescence instrument (Bendix Corporation, model 8002) was in operation until 1999. Thereafter, only ultraviolet absorption instruments have been employed (Thermo Electron, TE49 analysers). Carbon monoxide has been measured since 1990, using instrumentation based on gas chromatography with a mercury re-

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

duction detector (Trace Analytical, RGD2) or gas filter correlation infrared absorption (Thermo Electron, TE48S) as well as vacuum resonance fluorescence (AL5001, Aero-Laser, Germany), with two different systems running in parallel for most of the time. RH is measured with a dew-point mirror (Meteorolabor, Thygan VTP6) with an uncertainty below 5% RH.

2.3 Forecasts

The stratospheric intrusion forecasts, based on operational ECMWF forecasts, have aided the STACCATO measurement groups with the set-up and planning of the observations. Every twelve hours four-day forward trajectories were calculated with the Lagrangian Analysis Tool (LAGRANTO; Wernli and Davies, 1997), starting in the entire region covering the Atlantic Ocean and western Europe (20° E to 80° W and 40° to 80° N) between 250 and 600 mbar. From this large set of trajectories those initially residing in the stratosphere (potential vorticity larger than 2 pvu) and descending during the following four days by more than 300 mbar into the troposphere were selected as “stratospheric intrusion trajectories”. The same selection criterion was used in a previous case study (Wernli, 1997) to study an intrusion associated with a major Atlantic cyclone. A first case study based on the forecast was described by Zanis et al. (2003).

These forecasts, daily distributed by automatically generated e-mail messages, have been made available almost continuously since November 2000, with only less than 3% missing in the five years selected for our analysis (2001–2005). Since June 2001 so-called “intrusion hit tables” were additionally distributed giving a crude estimate of the time-height development of stratospheric air above the four partner stations Jungfraujoch, Zugspitze, Monte Cimone and Thessaloniki over several days.

For the analysis of a specific intrusion event the entire sequence of forecast plots showing the relevant intrusion(s) were examined. On each day, forecasts are generated for the start times 00:00 and 12:00 UTC, as well as for 00:00 and 12:00 UTC on the following day. There are differences between the trajectory plots for the same start time, but generated on subsequent days, in particular variations in the positions of

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the source regions. However, the sequences of trajectory plots very reliably predict the observations of ozone peaks with the lidar or at the Zugspitze summit. However, due to the coarse vertical and temporal resolution of the graphical representations the forecasts could only be used to identify the presence of stratospheric air above the Garmisch-Partenkirchen area. For more refined analyses additional trajectory or model calculations were made (see below and Sect. 2.4).

For the analysis of the Zugspitze series (2001–2005, see Sect. 4) extensive trajectory calculations were carried out with the HYSPLIT model whenever the measurements indicated an intrusion not clearly identified by the forecasts. The advantage of this approach is that HYSPLIT can be promptly operated via the internet on a server of the National Oceanic and Atmospheric Administration (<http://www.arl.noaa.gov/ready/hysplit4.html>; Draxler, 1998), which helped us to save a lot of time in this considerable effort. The trajectories were calculated 315 h backward in time (maximum time span allowed) for several height levels (3000 m and higher) since 3000 m a.s.l. was converted into the wrong relative height above the ground, very likely due to a crude orography of the underlying data field. Also, in many cases trajectories were calculated for times adjacent to a pronounced humidity dip in order to make sure that there are no temporal shifts. A case was accepted as stratospheric when at least one HYSPLIT trajectory reached or exceeded an altitude of roughly 8 km at sufficiently high latitudes (and depending on the season) where one could expect the tropopause. It is important to note that most contributions from outside the ETH model domain come from the colder seasons during which a lower tropopause height may be assumed. Some cases in the ETH forecasts with trajectories not reaching Garmisch-Partenkirchen, but linked to STT observations at the Zugspitze summit by HYSPLIT runs even more than 10 days backward in time were a posteriori accepted as successful forecasts. Although comparisons with successful ETH forecasts indicate that our approach is rather robust some uncertainty remains since HYSPLIT does not yield potential vorticity.

The analysis covers all cases either forecasted by the four-day trajectories, with the exception of intrusions originating roughly 1000 km and less away from Garmisch-

Partenkirchen because they do not have a chance to subside to 3000 m, and all cases additionally identified from the observations and verified by HYSPLIT calculations. Therefore, stratospheric air masses from either outside the ETH model domain or from inside the domain, but with a travel time of more than four days, do not enter our count if they overpass the station at altitudes high enough that there is no indication in the Zugspitze record. As shown in our long-range transport studies there may be considerable advection of thick aged air layers, rapidly travelling in the middle and upper troposphere, fed by STT events above the Pacific Ocean and Asia (Trickl et al., 2003, 2009a).

2.4 Models used for the analysis

Details on individual intrusions cannot fully be analysed by the comparison of the forecast trajectory plots with the measurements due to the limited altitude and time information of these plots. Therefore, in part additional calculations were made with the two models used within ATMOfAST.

Some details have been evaluated with the FLEXPART tracer model (Stohl et al., 1998, 2005; Stohl and Thomson, 1999), operated with ECMWF re-analysis data. This model has been extensively used for transport studies, in particular for several STT case studies of our team (Stohl and Trickl, 1999; Stohl et al., 2000; Trickl et al., 2003, 2009a; Zanis et al., 2003).

With the exception of the example discussed by Zanis et al. (2003) the FLEXPART analyses made for the cases described in Sect. 3 were made in the retroplume mode described by Stohl et al. (2002) and also in our parallel paper (Trickl et al., 2009a). The backward plumes, calculated over twenty days, yield a more complete picture of the principal pathways of the air masses arriving at an observational site. Numerical particles were released in 250-m bins above Garmisch-Partenkirchen, up to 10 km a.s.l. Important for the investigations in this paper are the options showing the average horizontal and vertical positions and the fraction of numerical particles travelling in the stratosphere on a given day. A threshold of 2 pvu was selected for identifying strato-

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



spheric air. Occasionally, we also inspected the fraction of particles with boundary-layer contact in order to get some idea of the mixing with tropospheric air suggested by the model runs.

The EURAD model, in contrast to the Lagrangian FLEXPART model, uses an Eulerian approach to study the origin and pathways of stratospheric air intrusions and intercontinental transport events. EURAD consists of the meso-scale meteorological model MM5 (Grell et al., 1994), the EURAD CTM (Hass et al., 1993) and the EURAD emission model (Memmesheimer et al., 1991). The model has been adapted to the tropopause and free tropospheric conditions (Kowol-Santen et al., 2000).

Within the ATMOfAST project, the EURAD model has been extended from a regional to a hemispheric model using a polar-stereographic projection centred at the north-pole. The first hemispheric simulation with EURAD was performed for a well-documented STT event coinciding with import from North America to Europe that took place by the end of May 1997 (Eisele et al., 1999; Stohl and Trickl, 1999). The ozone profiles calculated for Garmisch-Partenkirchen agree well with the lidar observations at IMK-IFU (ATMOfAST, 2005).

Single episodes and longer periods have been studied with this modified version of EURAD. In particular, a full-year run was completed for 2001. The horizontal grid distance for the long-term simulations is 125 km, in the vertical 27 layers from the surface (937 mbar for the lidar location) up to a pressure level of 30 mbar are considered. The vertical resolution chosen for the long-term runs varies between 10 mbar next to the ground and 40 mbar in the uppermost troposphere, which is a compromise in view of the considerable computation time. GFS (Global Forecast System) analysis data are used as initial and boundary values for 24-hourly consecutive forecasts of MM5. The emissions were derived from the EDGAR 3.2 inventory.

Two intrusion examples extracted from the run for 2001 are shown in this paper, one more also discussed in the companion paper (Trickl et al., 2009a) is just briefly described. By comparison with the lidar measurements further information on the quality of the simulations was obtained.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3 Case studies

The forecasts made possible a very dense lidar sounding period in 2001 (11 February to 23 August, ended by a major laser damage). As mentioned in the introduction, measurements were made during 20 of the 33 predicted intrusion periods (see Table 1).

5 Some of the periods were missed, mostly due to limiting the STACCATO warnings in autumn 2000 and 2001 to direct intrusions covering more than a single station, but also in two cases due to unfavourable meteorological conditions. During 26 of the predicted intrusion periods at least one stratospheric air tongue was verified at the summit station (column 3 of Table 1). In one case with very pronounced O_3 , 7Be and RH signature
10 (13 May) the corresponding intrusion was predicted to pass more to the east (outside Germany), but a HYSPLIT run indicates that it also reached our site. Table 1 also lists the intrusion type according to a classification described in Sect. 4.1 (column 2).

In this section we briefly present the lidar ozone series for the eight most interesting intrusion periods in 2001, which document a large variety of rather different strato-
15 spheric air intrusions. Among these examples are the first cases for which the early phase of a tropopause folding event was caught by our lidar. Airborne lidar measurements have yielded complete mappings of tropopause folds (e.g., Browell et al., 1987; Grant et al., 2000; Flentje et al., 2005). This is much more difficult for ground-based measurements due to the fact that the stratospheric air tongues eventually reaching
20 Europe start to penetrate the troposphere usually far outside the continent and that residual low-lying clouds after the preceding frontal passage mostly prohibit an early beginning of the lidar sounding.

The lidar results for these cases are displayed in Fig. 2. A selection of model fore-
25 casts corresponding to these cases is shown in Fig. 3. For the final example in this section, a particularly spectacular intrusion in November 2001, no lidar measurements were available. Instead, we include the Zugspitze series for the respective period.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The intrusion period 31 March to 2 April 2001, is rather complex and shows three in part simultaneously subsiding tongues of stratospheric air (S1, S2, S3). One lidar measurement was also made up to 3.2 km (below a cloud layer) on 2 April at 07:30 CET and shows the extension of intrusion S2 with a peak mixing ratio of 69.6 ppb (Zugspitze: 65.3 ppb, up to 70 ppb in the preceding hours, see Fig. 4). The lower two intrusions (S1 and S2) were both observed at the Zugspitze summit (relative humidities between 10 and 20%, peak ozone around 75 ppb in both cases), whereas the third one did not lead to a pronounced signature on the following days (Fig. 4). The Hohenpeißenberg ozone sonde launched on 2 April at 05:35 CET yielded a rather constant ozone mixing ratio around 50 ppb above 4 km, without any sign of S3, but up to 60 ppb below 4 km indicating the presence of S2. The absence of S3 in the sonde profile suggests that this intrusion had already moved out of the observational area, as confirmed by the forecast trajectories.

The trajectory forecasts also look complex and cannot be easily assigned to the observations. Most of the forecasts show a large and rather thick intrusion system starting to the south of Greenland and straightly approaching Central Europe. This system can clearly be seen also in the two examples in Fig. 3. During the early phase of the forecasts (not shown) there were some low-lying contributions of the intrusion at its eastern edge overlapping with Garmisch-Partenkirchen. These trajectories can be assigned to S1. S2 and S3 could, in principle, be assigned to higher parts of this intrusion system entering the observational area at later times, but there is some indication that this view is too simple (see below).

We do not have any proof of having observed tropopause folding in the case of layer S3. There is the possibility that a second intrusion contributed that is visible in the first two forecasts in Fig. 3. This intrusion originated over Labrador and stayed at high altitudes during the entire passage across the Atlantic, before it turned southward and started to descend. This north-south section rapidly moved eastward across Central

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Europe. For the overpass period of the lidar the colour code in the forecast trajectories indicates upper-tropospheric altitudes, in agreement with the position of layer S3. The thermal tropopause above Munich stayed above 12 km during the entire observational period. It is, thus, likely that this air mass travelled entirely inside the troposphere.

To deepen the understanding a twenty-day FLEXPART retroplume analysis was carried out for four times on 1 April (not shown). This analysis clearly assigns S1 to the direct intrusion system. A part of layers S2 and S3 can be assigned to the Canadian air stream. However, with growing altitude above our station and particularly relevant for S3 is an air stream from lower latitudes over the Atlantic (where it formed a circle and reached more than 10 km) and the Pacific, with stratospheric inflow from -3 d to -20 d in all the four simulations made. This component, arriving via the United Kingdom, is not captured by the forecast trajectories since the retroplume reaches latitudes below 40° N which are outside the ETH modelling domain.

Figure 5 shows the maximum values of the daily stratospheric fractions within the twenty days of the FLEXPART backward calculation for the four times selected. It is interesting to note that the FLEXPART stratospheric fractions for S1 and S2 on individual days are less than 14%, in some contrast to the rather strong signature in both the ETH forecast and the Zugspitze data (RH 18%, rise in ozone by more than 25 ppb and ^7Be values exceeding 6 mBq m^{-3} on 1 April, although the intrusion covers just one third of the day). The stratospheric fraction for layer S3 is substantially higher at the beginning. The low fraction for S1 could be due to the limited vertical resolution of the model and, as a consequence, to overestimating the influence of mixing the thin (0.8 km) air stream with tropospheric air which is indicated by the non-negligible fraction of particles with boundary-layer contact found in the model results. Significant mixing does not look that likely because of the very sharp edges of S1 in both the lidar and the station results (see Fig. 4).

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2 23 April 2001

This is the first example of a tropopause fold that we can consider to be fully captured with the lidar (S). Again, the outflow from the stratosphere started rather close to our site, over the south-west corner of the North Sea and the United Kingdom. The stratospheric streamer quickly moved outside the observational area.

3.3 20–23 May 2001

Two intrusions (S1 and S2) occurred during this period. Unfortunately, no lidar sounding was possible during the early part of S2 due to low-lying clouds. The Zugspitze data also indicate a pronounced stratospheric influence during the first halves of both 20 May and 21 May with a rise in ozone from 47 to 62 ppb and 59 to 67 ppb, respectively, a relative humidity falling below 20%, and ^7Be values of about 6 and 9 mBq m^{-3} , respectively. S2 is seen in the Zugspitze record for just less than two hours on 23 May, with a sharp ozone rise to almost 68 ppb, but just about 5 mBq m^{-3} of ^7Be due to the very short duration of that event. The Munich radiosonde (launched twice a day 100 km roughly to the north of IMK-IFU) registered a significant drop in humidity above 1.5 km a.s.l., reaching RH values below 10% in the middle troposphere.

The first intrusion (S1) first proceeded southwards from Greenland and then split into different portions travelling at different altitudes, two of them reaching Garmisch-Partenkirchen on 19 and 20 May (see 15 May-panel in Fig. 3). The second intrusion (S2) was a “classical” direct intrusion with anti-cyclonic advection path originating over Greenland (see 18 May-panel in Fig. 3).

Another increase in Zugspitze ozone is seen in the evening of 23 May, but corresponds to air of high relative humidity. Thus, the corresponding layer at the end of the lidar measurements cannot be associated with an extension of the second intrusion.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.4 4–6 June 2001

The forecast shows an intrusion cyclonically descending to Central and South-East Europe from the North Atlantic between Greenland and Norway. There are both high-altitude and lower-altitude components which are nicely reproduced by the lidar measurements. There is even an indication of the outflow from the stratosphere at the beginning of the series when the ozone values stayed constantly high all the way from 6 km to the final stratospheric ozone rise starting at 8.8 km. At the Zugspitze summit the stratospheric air was briefly seen around midnight between 4 and 5 June (relative humidity 20%, ozone increase to 64 ppb, ^7Be on 5 June at 7.5 mBq m^{-3}) in agreement with the midnight ozone maximum around 3 km in the lidar measurements. For the rest of 5 June the Zugspitze relative humidity oscillated between 55 and 75% and rose to 100% on 6 June indicating no further input from the stratosphere at 3 km on that day.

The layer around noon on 5 June with 70 to 80 ppb of ozone between 4 and 10 km was attributed by HYSPLIT trajectories between 5 and 8 km to air masses with rather different advection pathways. However, most trajectories descend from high altitudes (in part more than 10 km) over Canada or East Siberia. This layer is not included in the analysis in Sect. 4 because it was neither forecasted nor did it reach 3000 m.

3.5 11–13 June 2001

This case is a rather thin direct intrusion originating over Greenland. The stratospheric air did not proceed beyond the Alps on its way to lower latitudes and bent eastward instead. The Zugspitze ozone data show sharp spikes on 12 June at about 11:00 and 23:00 CET (69 and 70 ppb, respectively), coinciding with two somewhat longer low-humidity periods. The ^7Be value for 12 June is about 10 mBq m^{-3} . The trajectory forecast for 10 June, 12:00 UTC (not shown), reveals that there was a short period during which the intrusion stayed slightly outside the Alps. This explains the gap in the layer in the lidar series at 3 km between 15:00 and 21:00 CET (39–45 h in the figure) that is also visible in the Zugspitze data (elevated humidity and ozone below 60 ppb).

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The upper-tropospheric ozone maxima were traced back to altitudes of frequently more than 10 km in rather different higher-latitude source regions and are likely to contain pronounced stratospheric contributions. These layers are not included in the analysis in Sect. 4 because there is no overlap with 3000 m.

5 3.6 20–25 June 2001

The lidar measurements on the first two days of this period (Zanis et al., 2003) served as a reference for an extensive model intercomparison and validation exercise within the STACCATO project (Roelofs et al., 2003). Recently, the EURAD model was also tested for this complex stratospheric intrusion case (Fig. 6). Due to the finer spatial resolution the agreement with the lidar sounding at Garmisch-Partenkirchen is (in part substantially) better than that reported for the different models used in the study by Roelofs et al. (2003). However, in the evening of 24 June a descending moderate-ozone layer (about 50 ppb) is suggested by the model results at about 450 mbar that is not confirmed by the lidar data before the end of the observational period. Also the low-ozone subtropical layer (SA) starts earlier in the EURAD results than in the measurements. However, the layer with less than 50 ppb of O₃ in the lidar results between 8 and 10 km in the afternoon of 20 June corresponds to air ascending from an altitude of about 5 km next to 30° N. This layer, therefore, corresponds to the beginning of the SA layer in the EURAD results.

20 Three layers very likely of stratospheric origin (S1–S3) are identified by the lidar and Zugspitze measurements on 20 and 21 June, another one (S4) on 23 June, following a cloudy period over the Alps on 22 June which interrupted the measurements. Intrusions S2 and S3, the latter observed all the way down from the stratosphere, were discussed at length by Zanis et al. (2003). The EURAD result shows elevated ozone in all four layers, in particular also S1 which was reproduced (downward shifted) by just one of the model simulations described in the two preceding publications. S4 is not fully resolved, but indicated by the step from 800 to 700 mbar on 23 June shortly before 10:00 UTC.

25 We spent some effort to identify the origin of layer S1. The Zugspitze data strongly

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

suggest the presence of stratospheric air: almost 8 mBq m^{-3} of ^7Be on 20 June (intrusion S2 contributed on the following day!), a strong ozone rise to 69 ppb and $\text{RH} \approx 22\%$. However, this layer cannot be assigned to any of the ETH forecasts. A 20-d FLEXPART retroplume calculation now indicates air masses slowly descending from altitudes above 8 km over the north coast of Canada, i.e., outside the model domains of both the ETH forecast and the forward FLEXPART simulation published by Zanis et al. (2003). The stratospheric fraction rises with backward time, but stays below 10%, the layer width clearly exceeding that in the measurements. This, again, indicates problems in the analysis of narrow stratospheric layers travelling over that many days.

Intrusion S3 is difficult to distinguish in the forecast plots (and not visible in the two examples shown in Fig. 3) since it, in part, horizontally coincides with intrusion S2. As confirmed by trajectory calculations S3 first leaves Greenland on a cyclonically turning pathway followed by an anti-cyclonic turn and some horizontal spreading during the final approach. It is interesting that, in spite of the long travel, the lidar measurements trace the ozone tongue all the way down from the stratosphere.

As suggested by the lidar soundings and the ETH forecast (for 20 June, 00:00 UTC) the intrusion on 23 June overpassed Garmisch-Partenkirchen almost horizontally. This intrusion is already seen in the forecast for 19 June, 00:00 UTC (Fig. 3), originating between Greenland and Iceland.

The high-ozone layers above 8 km on 23 and 24 June were examined by trajectory calculations. The results indicate import of stratospheric air from the Pacific as discussed in the companion paper (Trickl et al., 2009a).

3.7 21–24 July 2001

The four-day sounding series from 21 to 24 July 2001 is another, particularly nice example of the typical ozone time series observed at Garmisch-Partenkirchen after the beginning of high-pressure periods during the warm season. In Fig. 2 we just show the results for the first two days of observation. The observations above 2 km

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

after 21 July are explained in detail in our parallel papers on air-mass advection from beyond North America (Trickl et al., 2009a) and from the subtropical Atlantic (Trickl et al., 2009b). Here, we briefly discuss the stratospheric air intrusion on 21 and 22 July. This intrusion is (as the 20–21 June case) special since also a sudden downward step is observed (in the early afternoon of 21 July). This is ascribed to a splitting of the intrusion into two components, a higher-lying component more to the east and a strongly descending component more to the west. Very interestingly, as the source region shifts towards Europe, first the trajectories by-pass South Germany (panel for 18 July 12:00 UTC), later on (panel for 20 July 00:00 UTC) they, again, overpass our site at high altitude and could result in another contribution in the upper troposphere during early 21 July. This cannot be clearly distinguished since the lidar measurements did not start before 13:00 CET (12:00 UTC).

The result extracted from the EURAD run for this case (not shown) nicely reproduces the characteristic features of this intrusion, in particular the downward step in the afternoon of 21 July. However, the calculation yields a slightly lower final altitude of the stratospheric air tongue than the lidar measurements.

3.8 12–15 August 2001

This case is also in more general discussed by Trickl et al. (2009a). Although this it is not fully clear from the last panel of Fig. 2 the “classical” three-layer structure is seen that is reproducibly observed during high-pressure situations (in this paper in particular in the example shown in Sect. 3.7: S, SA, USA). The reason for this deception is that the subtropical layer above the intrusion is very narrow and short in duration in this case. Also, the intrusion (S) is limited to the beginning of the measurement that was started too late since the forecast, despite the favourable situation with a beginning high-pressure period, did not suggest a spectacular event for our site.

Indeed, this is an example for which the ETH forecast did not fully agree with the observations although a considerable intrusion event was predicted. In the last panel of Fig. 3 a rather large trajectory bundle covers a major part of Central, East and South-

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

East Europe. Garmisch-Partenkirchen is slightly outside this area, the high-pressure zone obviously having already proceeded to the east at the time of the intrusion. However, the Zugspitze data reveal that the edge of the intrusion was even fully caught. A considerable drop in relative humidity was observed on 12 August between 0:00 CET and 18:00 CET (minimum: 3%), the ozone mixing ratio rose at the same time by 17 ppb (to 66 ppb), and the ^7Be value on 12 August was as high as 11.5 mBq m^{-3} . This observation corresponds to the descending ozone feature (S) below 3 km just after the beginning of the lidar measurements. During the early part of 13 August the relative humidity, again, dropped, but not much below 30%. ^7Be was still above 8 mBq m^{-3} , but there was no pronounced increase in ozone. The most prominent feature in the lidar measurements (LA) during that period is located at about 4 km and, indeed, does not fully overlap with the summit station. Trajectory calculation revealed long-range advection from at least the upper troposphere above the northern Pacific. The high ^7Be value is indicative of stratospheric air.

Also the EURAD simulation (Fig. 7) clearly shows the presence of the intrusion. As in the measurements it was already located below 3 km (700 mbar) by the beginning of 12 August. Also the elevated-ozone features in the middle troposphere on 12 August are nicely reproduced. Layer LA on 13 August is missing. Instead, the subtropical layer (SA) is wider. This deviation might be caused by a slight geographical shift in the model results: the high-ozone layer at about 450 mbar starting on 13 August at 19:00 UTC and persisting also on 14 August confirms the corresponding layer in the lidar measurements, but with mixing ratios lower by roughly 50 ppb. The stratospheric nature of this layer (featuring very low EURAD CO) is further discussed in our companion paper (Trickl et al., 2009a).

3.9 14–22 November 2001

Figure 8 shows one trajectory forecast during one of the most spectacular intrusion cases observed since the beginning of these forecasts. The outflow from the stratosphere over Greenland continued over many days and covered most of the European

continent. The intrusion was observed at the Zugspitze station from 14 to 22 November 2001 (Fig. 9). The ^7Be values were above 8 mBq m^{-3} from 15 to 19 November, but the ozone mixing ratio stayed between 45 and 65 ppb. Due to ongoing problems with the laser system no lidar measurements were made. This classical direct intrusion was immediately followed by the advection of stratospheric air from Canada, which resulted in a total of nine subsequent intrusion days in the Zugspitze record. An extensive coverage of a month by intrusions is typical of the cold part of the year.

During the period 2001 to 2005 a total of more than 10 large intrusions covering major parts of Europe were found. Four of them, including the one in this section, covered about two thirds of the continent and more.

4 Comparison of ETH model predictions and Zugspitze measurements

The daily forecasts by ETH Zürich have offered the unique opportunity to examine which observations at the Zugspitze summit are most likely related to stratospheric intrusions and to exclude pure advection from the upper troposphere. In this way, also several filtering criteria used for the discrimination of stratospheric air tongues could be tested for the first time. We compared the forecasts and the Zugspitze data for the period 2001 to 2005.

4.1 Classification of the source regions and definition of the data-filtering criteria

For the detailed analysis presented in Sect. 4.2 a classification of the intrusions observed at the Zugspitze summit into eight categories was made:

- Type 1: The intrusion originates in the region covering Greenland, Iceland and the adjacent part of the North Atlantic and anti-cyclonically or almost straightly approaches the Bavarian Alps (e.g., Fig. 8 in Sect. 3.9). Type 1 also comprises a few exotic cases of anti-cyclonic long-range descent via Russia.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Type 2: The intrusion originates over Greenland or slightly to its east and proceeds to western Europe or the Atlantic off the European or African coast; the final advection to Garmisch-Partenkirchen is cyclonic (e.g., 15 May and 18 July panels in Fig. 3). In some cases, the distinction between Type 1 and Type 2 is rather difficult.
- Type 3: Direct advection from source regions over Scandinavia or the North Sea.
- Type 4: Advection from the Atlantic to the south of Greenland and Iceland, but typically above 45° N.
- Type 5: Advection from Canada and the adjacent part of the Polar Sea to the east of 80° W (i.e., within the range of the ETH forecast).
- Type 6: Westerly advection from Canada to the west of 80° W, Alaska, Siberia and the adjacent parts of the Pacific and Polar Sea (i.e., outside the range of the ETH forecast); in this case, HYSPLIT backward trajectories were used for the analysis. A few of these trajectories are quite exotic and, starting over Canada or Alaska, cross the Atlantic and penetrate deeply into East Europe or Asia before they return to Garmisch-Partenkirchen.
- Type 7: Advection from the Atlantic with initial latitudes below 45° north; this separation from Type 4 was made by intuition since this sector belongs to latitudes below that of the observational site.
- Type 8: Easterly to north-easterly advection from Russia, Siberia and the adjacent part of the Polar Sea; here, HYSPLIT trajectories were exclusively used for estimating the presence of stratospheric air.

Types 1 and 2 comprise mostly direct intrusions, but may also involve long advection paths or travel times. As mentioned, in a few cases rather exotic pathways were found, an example is shown in Fig. 10 (12 February 2003). The trajectory in Fig. 10 reaching the highest altitudes ended at almost 8 km above the ground over north-western

Canada 315 h backward in time. This case is difficult to assign (Types 3, 5, 6 or 8), since the local tropopause height along the path is not known. We selected Type 5 since the highest altitudes were reached over Labrador, but similar altitudes were also reached over northern Finland (Type 3) where the final descent starts.

Three filtering criteria were applied in order to determine stratospheric intrusions in the observations, all based on the ^7Be and RH measurements. No ozone criterion has been used in recent years since the increases in O_3 during intrusion periods are rather different in their appearance, requiring a complex algorithm for the discrimination. The behaviour of the ozone and CO mixing ratios was, however, in most cases inspected on the monthly plots used for the work presented here. The criteria are:

Criterion 1: The ^7Be value corresponds to more than the 85th percentile with respect to all data in the respective year and $\text{RH} < 60\%$. The ^7Be threshold was chosen to match the “classical” threshold for the 24-h specific activity of 8 mBq m^{-3} (Sladkovic and Munzert, 1990) during the late 1990s. The RH threshold corresponds roughly to the full width at half maximum of many of the humidity troughs caused by the passage of a stratospheric layer. Lower RH thresholds yielded too low fractions of stratospheric ozone. This criterion has been routinely used during the past decade (e.g., Scheel, 2003).

Criterion 2: $\text{RH} < 60\%$ and $\text{RH} < 30\%$ for at least one of the half-hour averages within $\pm 6 \text{ h}$. The second threshold is added to guarantee really dry conditions as expected for stratospheric air. This threshold is higher than those used by Beekmann et al. (1997), but this is adequate due to the rather low tropospheric altitude of the station.

Criterion 3: Same as Criterion 1, but with 5.5 mBq m^{-3} as the threshold for ^7Be . The application of this lower threshold was examined because Criterion 2 yielded so many more STT cases. The value was chosen from an inspection of many of the monthly plots for the Zugspitze data.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

4.2 Results of the comparison

The forecasts by ETH Zürich yielded substantially more intrusion cases than we would have registered with our formerly used forecasting or data filtering approaches which yielded only very deep intrusions that rarely miss the Zugspitze summit. An important consequence is that there are also few more stratospheric air tongues not fully reaching 3000 m during all seasons. These “medium intrusions” (Sprenger et al., 2003) may even overpass the Alps when there is ongoing rain or snowfall in the lower elevations prohibiting optical remote sensing.

Table 2 lists the number of cases for these types and of the tropospheric cases, as identified by the model calculations, identified during the period 2001–2005. The number of intrusion cases per year (which may last for several days) is 82.2, which substantially exceeds any estimate in the past. Also the average number of intrusion days per year, 154, is very high and demonstrates the significance of STT for the annual ozone budget. The fraction of the predicted (or verified) intrusions reaching the Zugspitze summit per year, identified by at least one of the three filtering criteria, is 77.9%. It was also found that about 95% of the forecasted intrusion cases led to a clear signature in either ^7Be or RH, or both, though not always fulfilling the criteria. The difference with respect to the fully accepted cases is mostly caused by overpasses missing the summit. In a few predicted cases measurements were missing, in others significant mixing with tropospheric air due to a travel time exceeding four days is hypothesized. Just in 16 cases during the five years of analysis no stratospheric assignment of observations fulfilling one of the criteria could be made (bottom line of Table 2). In all but one of these cases a rise in ozone by at least a few ppb simultaneous to a drop of RH to below 50% was seen. These episodes were mostly rather short and, therefore, perhaps not captured by the backward trajectory analysis. The backward trajectories did not reach sufficient height or, in a few cases, even stayed in the middle and lower troposphere within the preceding 315 h. Finally, it is important to note that the true number of intrusion cases may be, still, higher since our model analysis does not include overpasses

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

that originate outside the domain of the ETH model. However, this does not make any difference in the validation of the filtering criteria since here only the cases leading to an observation matter.

Two examples for the Zugspitze series were already given in Sect. 3 (Figs. 4 and 9). In the example in Fig. 4 the most typical problem with ^7Be as a tracer may be seen on 31 March 2001. Due to just a partial coverage of that day by intrusion 1 the 24-h average of ^7Be is below the threshold for Criterion 1. It is advisable to improve the time resolution of the ^7Be measurements to half a day or even less.

In order to highlight how many different intrusions can occur in a single month we show in Fig. 11 another example typical of the cold season (January 2003). The types of the different intrusions are marked in the figure next to the RH curve. The first two intrusions passed above the summit and, therefore, yield less pronounced signatures. In general, the Types 5 and 6 make a significant contribution during the dark period of the year.

This example (among many others, see also Sect. 3.9) demonstrates that the stratospheric influence in the lower free troposphere during this part of the year is considerable. The time periods for which the “classical” $^7\text{Be}/\text{RH}$ criterion (Criterion 1) and the RH criterion (Criterion 2) are fulfilled are marked in the upper two traces. Criterion 2 (see also violet triangles on the RH curve) is substantially more efficient in identifying intrusion periods than Criterion 1. Finally it is interesting to note that the drop in carbon-monoxide during intrusions is mostly not very pronounced, one exception being 15 January 2003.

Figures 12 to 16 show the seasonal cycles (2001–2005) derived for the intrusions predicted or verified by trajectories for the four most important intrusion types. As expected, the Types 1, 2 and 5 yield the strongest contributions. For most intrusion types (also Types 3 and 4) the winter maximum and summer minimum from previous studies is confirmed. However, quite interestingly, for Type 2 we obtain a late-spring maximum. This maximum obviously causes the asymmetry in the seasonal distribution for (Fig. 12). This asymmetry is also visible on the FLEXPART result shown in Fig. 1.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Criterion 1 consistently yields just a moderate number of intrusion cases. This is, to a major part, due to the accumulating ^7Be over 24 h which may result in an omission of short intrusions or of cases that occur around midnight. However, this criterion is robust and suitable for trend studies. The seasonal cycle for Criterion 1 in Fig. 12 is not very pronounced. The corresponding seasonal cycle for ozone should exhibit a stronger modulation.

Criterion 3, based on a lower ^7Be threshold, yields many more cases, close to the number of cases for Criterion 2. With the exception of intrusion Type 2 it works particular well around the middle of the year. The cases for Criteria 2 and 3 overlap to a major extent, but there are also mismatches. To our great surprise the very simple Criterion 2 almost perfectly corresponds to stratospheric intrusions. During the warm season some cases are missed that are identified by Criterion 3. This could be due to a slightly higher passage with less complete overlap with the summit station or enhanced mixing with tropospheric air resulting in an exceedance of the 30-% threshold.

Criterion 3 was thoroughly tested for the period 2001–2005. Outside the verified intrusion periods there were just 39 days per year on which 5.5 mBq m^{-3} were reached or exceeded. The seasonal distribution is shown in Fig. 17. Most of these exceptions, which may sometimes cover an extended period of time with even rather high ^7Be values, occur during the warm season. If $\text{RH} < 60\%$ is added the number of erroneously counted days is reduced to just 5.4 per year. However, in all but two cases the period with $\text{RH} < 60\%$ lasted for not more than just a few hours, which implies a very low significance of these events. The bad cases are governed by the results for the exceptionally hot year 2003.

Very-low-humidity cases are relatively rare. They are, in principle, expected to be more frequent for short advection paths due to the reduced time for mixing, but also air from rather remote intrusion events may be considerably dry if the layer is sufficiently thick to avoid a significant admixture of tropospheric air. Normally, a tropospheric travel time of about three days seems to be short enough for avoiding a significant loss of the stratospheric nature of the air.

**Forecasted deep
stratospheric
intrusions over
Central Europe**T. Trickl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Figure 18 gives a summary of the fractions of the cases for the statistically relevant intrusion types fulfilling different RH conditions (Type 4 being at the edge of being statistically acceptable here). The highest fractions with $RH < 10\%$ (43.1% and 36.5%, respectively) were obtained for Types 1 and 6. The explanation for Type 1 is the mostly rather rapid travel of these “classical” intrusions from Greenland to the Alps within two to three days. In the case of Type 6 the travel time is mostly rather long. Thus, these layers must be sufficiently thick in order to avoid significant mixing with tropospheric air. This idea is also hardened by the fact that intrusion Type 6 also comprises the lowest fraction of cases with $RH \geq 40\%$. The mixing with tropospheric air is further impeded in an aged intrusion layer since it is likely that the layer adopts the velocity of the surrounding air masses during the long travel, resulting in reduced turbulent mixing.

5 Discussion and conclusions

The intrusion forecasts have been invaluable for identifying periods during which lidar measurements of STT events were made. The period between February and August 2001 has been particularly successful in this regard and yielded, together with our earlier work (e.g., Eisele et al., 1999; Stohl and Trickl, 1999; Zanis et al., 2003; Roelofs et al., 2003; Trickl et al., 2003) a large variety of different cases with strongly differing complexity. Still, as pointed out by Eisele et al. (1999), it is very likely in complex situations that at least one of several observed intrusions descends down to 3000 m and can be observed at the Zugspitze summit. This probability is lower in summer. The individual intrusions may strongly descend during the observational period (Sects. 3.6 and 3.7) or they may intersect our area without much change in vertical position, all depending on the distance of the source region to the Alps. Quite frequently, layers of stratospheric air may overpass our area at about 3 km a.s.l. for several days (Trickl et al., 2003). In these cases they are observed at the Alpine summit stations during an extended period of time. The descending intrusions may occasionally exhibit downward steps (e.g., examples in Sects. 3.6 and 3.7). In Case 7 (Sect. 3.7) this was

explained by a splitting of an intrusion into two components. Most importantly, first observations of the initial phase of tropopause folding were made with our ozone lidar.

Intrusion Case 4 (Sect. 3.4) has been used for a variety of model validation exercises within the STACCATO (Zanis et al., 2003; Roelofs et al., 2003) and ATMOfAST (this work; Trickl et al., 2009a) projects. From the EURAD simulation discussed in this paper we derive a strong indication that a high spatial resolution, here 125 km horizontally and 10 to 40 mbar vertically within the troposphere, is needed to obtain results resembling the measurements. An even higher resolution is desirable for catching also the details in the lower troposphere. Episodic studies with the EURAD model (ATMOfAST, 2005; Ebel et al., 2008) with increased horizontal and vertical resolution confirmed that especially a higher vertical resolution in the tropopause region enhances the ability of the model to reproduce of the dynamics and composition of the upper troposphere/lower stratosphere in more detail.

The forecasts are also an important tool for testing the identification criteria for stratospheric air in the Zugspitze data. It was found that all three data-selection criteria examined work very reliably although there are differences in the individual cases identified. These differences are due to differences in mixing with tropospheric air, resulting in a wide range of humidity values, or due to the bad temporal resolution of the ^7Be measurements. A few discrepancies might be due to using forecast instead of re-analysis data. However, to repeat this exercise with improved model calculations is out of the scope of this work. We finally note that overpasses of stratospheric air from outside the domain of the forecast model at high altitude (e.g., Bithell et al., 2000; Trickl et al., 2009a) are excluded from the analysis.

The significant coverage of the dark part of the year by stratospheric events, with low humidity frequently extending over many days, suggests that the approach by Elbern et al. (1997) based on eleven-day statistics is not always adequate. The use of ozone as an additional tracer or STT events is also rather restrictive. Their analysis yielded a low number of significant intrusions of 19.5 per year, in considerable contrast to the numbers in Table 2. Our first criterion already yields 29.4 cases per year, but the cov-

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

erage of the events is, still, moderate in comparison with the predictions. Nevertheless, the percentiles chosen are related to full-year statistics and, thus, the results are less influenced by short-term variations than those by Elbern et al. (1997). A one-year period is sufficiently short in comparison with known periodicities in surface ^7Be (Reiter et al., 1983; Koch and Mann, 1996; Gerasopoulos et al., 2003) in order to account for their influence.

Despite the high reliability of the data-filtering criteria examined they do not allow one to distinguish between direct (those with closer-by source regions, in particular Types 1 to 3) and indirect intrusions. Even high beryllium values are not fully indicative of short advection paths. The specific activity reaches or exceeds 8 mBq m^{-3} in about ten Type-5 and Type-6 intrusions per year.

The considerable number of intrusions observed at the Zugspitze summit, with a strong monthly coverage in late autumn and winter (Fig. 8) contradict the frequently heard statement that STT has a minor influence on the tropospheric ozone budget. By contrast, Roelofs and Lelieveld (1997) concluded from model calculations that as much as 40% of the tropospheric ozone originates in the stratosphere. This remarkable fraction was recently approximately confirmed for 3000 m from the detailed filtering of the Zugspitze data between 1978 and 2004 (Scheel, in preparation; ATMOFAST, 2009). This filtering, based on experimental material alone, also confirms a pronounced positive trend in the stratospheric influence already found for the 1990s (Scheel, 2002, 2003) for the entire period. The Zugspitze ^7Be series, started in 1970, shows a positive trend in the specific activity of that isotope since the mid-seventies (Scheel, 2002). In the 1990s the positive ozone trend caused by the increase in the stratospheric contribution masked a decrease expected from a reduction in precursor emissions in Europe. In fact, for the nearby lower Wank station (1780 m a.s.l.), which is substantially less exposed to stratospheric intrusions (Elbern et al., 1997), no significant trend exists during that period. The positive trend in the Zugspitze ozone exceeds that at other, low-lying stations with long data records reviewed by Oltmans et al. (2006), with one strange exception: Mace Head (Simmonds et al., 2004).

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The seasonal cycles shown in Figs. 12 to 16 mostly show a winter maximum and a summer minimum. The overall seasonal cycle (Fig. 12) is less pronounced for Criterion 1 than for forecasts and the other two filtering criteria, in agreement with the findings by Reiter et al. (1983) for intrusions with high beryllium values. The asymmetry found for the predicted cases and Criteria 2 and 3 agrees well with the revised FLEXPART model calculation for the Zugspitze summit (Fig. 1). As discussed in Sect. 4.2 this asymmetry is caused by the spring maximum obtained for the Type-2 intrusions (Fig. 14). This asymmetry also exists in the simulation for Jungfrauoch (southwest of Zugspitze), but not in that for Sonnblick (east of Zugspitze). The cyclonically advected Type-2 intrusions are more relevant for western Europe. In fact, a maximum of STT between May and July was found for southern France (Van Haver et al., 1996). However, this view is too simple since Van Haver et al. derived almost the same seasonal cycle also for the troposphere above Uccle (Belgium) that is quite frequently overpassed by the Type-1 intrusions reaching Garmisch-Partenkirchen.

There are several possibilities to improve the performance of the data filtering. The best choice is expected to be a “or” combination of Criteria 2 and 3. From this procedure a coverage of most cases is expected. The analysis for 2001 to 2005 shows an enhancement of the coverage from 64.4% (Criterion 2) or 70.2% (Criterion 3) to 77.9% when using all criteria simultaneously. Of course, the 5.5-mBq m^{-3} threshold, chosen here as a constant value for simplicity, must be converted to the corresponding percentile as done earlier for the 8-mBq m^{-3} threshold, yielding the 85th percentile of a given year. Another possibility could be weighting the ^7Be integrated specific activity according to length of an intrusion. The length could be obtained from applying Criterion 2. However, this approach can be rather difficult due to temporary (short) exceedances of $\text{RH}=60\%$, data gaps and in the case of intrusions taking place on two subsequent days. It is obvious that better-resolved ^7Be data are desirable, e.g., 6-h instead of 24-h averages.

For a complete quantification of STT also the mixing with tropospheric air must be considered. Shapiro et al. (1980) showed that the admixture of tropospheric air already

starts in the tropopause region. There is clear observational evidence of an admixture of polluted air to intrusions in the free troposphere (e.g., Parrish et al., 2000; Brioude al., 2007). In many cases the water-vapour content of a stratospheric intrusion should be a good indicator of the amount of mixing. In the Zugspitze data the cases with RH<5% are rare. It is reasonable to assume that thick layers of stratospheric origin are drier than thin ones because mixing starts at the edges of the layer. One example of a layer several kilometres wide and exhibiting relative humidities of 1.0 to 2.4% (mixing ratios 79 to 116 ppm) was recently given by Vogelmann and Trickl (2009a), obtained from a measurement with the new water-vapour lidar operated by IMK-IFU about 300 m below the Zugspitze summit. On the other hand, in the same publication an intrusion just a few hundred metres wide and featuring just 1% minimum RH is shown in the lower troposphere (at 3200 m a.s.l., 31 October 2007 14:39 CET, exactly confirmed by the 13:00 CET Munich radiosonde launched 100 km to the north). This suggests a substantial variability in mixing efficiency.

A reasonable guess of the tropospheric contribution to the air in a stratospheric intrusion might be derived from the humidity, but this may be rather complex. Low-humidity upper tropospheric air may be already admixed in the region below the tropopause, in particular in the cases of thin layers. On the way downward the stratospheric layer could be embedded in simultaneously descending upper-tropospheric air and the interaction could continue. On the other hand the observations of intrusions in the middle and lower troposphere frequently show rather sharp layer boundaries, which seems to exclude major exchange with the surrounding air. We suspect that the altitude region where the mixing occurs may can be rather different.

Despite the variability of the minimum in-situ RH values in intrusions the results shown in the preceding section suggest that the travel time of intrusions in the troposphere, indeed, influences the RH values in deep intrusions. The interesting observation that Type-6 intrusions also exhibit rather low humidities could indicate either a rather pronounced layer thickness, allowing this layer to survive over the long distance, or travelling at about the same speed as the surrounding air, thus reducing turbulent

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

mixing.

Due to this complexity the admixture of tropospheric air to stratospheric intrusions has not yet been included in the data filtering exercises for the Zugspitze data. Given the moderate variability of tropospheric ozone, apart from a few really spectacular high-
5 O₃ events (that are unambiguous anyway), and the low humidity in intrusion layers, the uncertainty caused by the choice of the threshold conditions has been regarded to be more important than the error caused by neglecting the tropospheric admixtures. On the way to a more and more refined approach that is stimulated by the results presented here also the complexity in the analysis of the in-situ measurements will grow
10 and approaches using the water-vapour mixing ratio to estimate should be derived. However, it must be mentioned that meanwhile several very recent measurements with the water-vapour lidar have shown 0 to 2% RH inside rather thin intrusions (in perfect agreement with radiosonde measurements on the same days) and indicate that the Zugspitze in-situ measurements could have a (varying) wet bias of up to roughly 10%
15 RH under the driest conditions. This aspect will be investigated further.

In any case efforts to quantify free-tropospheric mixing are highly desirable. Our own approach will include combined vertical sounding of water vapour and ozone, also hoping for better-resolved model calculations.

It is also important to note that in some cases the Zugspitze station may be located
20 just at the edge of an intrusion layer, where the influence of mixing should be the highest. This may, e.g., be the case on sufficiently warm days when the alpine boundary layer may reach considerable height (Carnuth and Trickl, 2000; Carnuth et al., 2002; Kreipl, 2006). Very recently, we observed an intrusion case with a persistent 120-ppb ozone layer centred slightly above 3 km that was pushed upward by the PBL expansion during daytime resulting in the absence of a clear signature in the Zugspitze data.
25 During the following night the layer, again, subsided and was partly observed (83 ppb) at the summit station.

Acknowledgements. The authors thank A. Ebel, H. Davies, M. Kerschgens, P. Fabian and W. Seiler for their continual interest and support. T. Trickl appreciates the patience of his wife

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heidi during that challenging (www.trickl.de) spring and summer of 2001. H. Claude made available the Hohenpeißenberg data for the relevant period in 2001 for comparison. M. Sprenger and H. Wernli thank MeteoSwiss for providing access to the ECMWF forecast data in real time. This work has been funded by the European Union within the STACCATO project as well as by the German Bundesministerium für Bildung und Forschung within the programme Atmosphärenforschung 2000 (ATMOFAST project).

References

Ancellet, G., Pelon, J., Beekmann, M., Papayannis, A., and Mégie, G.: Ground based lidar studies of ozone exchanges between the stratosphere and the troposphere, *J. Geophys. Res.*, 96, 22401–22421, 1991.

ATMOFAST: Atmosphärischer Ferntransport und seine Auswirkungen auf die Spurengaskonzentrationen in der freien Troposphäre über Mitteleuropa (Atmospheric Long-range Transport and its Impact on the Trace-gas Composition of the Free Troposphere over Central Europe), Project Final Report, Trickl, T., Co-ordinator, Kerschgens, M., Stohl, A., and Trickl, T., subproject co-ordinators, funded by the German Ministry of Education and Research within the programme “Atmosphärenforschung 2000”, 130 pp., 2005 (in German).

Baray, J.-L., Daniel, V., Ancellet, G., and Legras, B.: Planetary-scale tropopause folds in the southern subtropics, *Geophys. Res. Lett.*, 27, 353–356, 2000.

Beekmann, M., Ancellet, G., Blonsky, S., De Muer, D., Ebel, A., Elbern, H., Hendricks, J., Kowol, J., Mancier, C., Sladkovic, R., Smit, H. G. J., Speth, P., Trickl, T., and Van Haver, P.: Regional and Global Tropopause Fold Occurrence and Related Ozone Flux across the Tropopause, *J. Atmos. Chem.*, 28, 29–44, 1997.

Bithell, M., Vaughan, G., and Gray, L. J.: Persistence of stratospheric ozone layers in the troposphere, *Atmos. Environ.*, 34, 2563–2570, 2000.

Brioude, J., Cooper, O. R., Trainer, M., Ryerson, T. B., Holloway, J. S., Baynard, T., Peischl, J., Warneke, C., Neuman, J. A., De Gow, J., Stohl, A., Eckhardt, S., Frost, G. J., McKeen, S. A., Hsie, E.-Y., Fehsenfeld, F. C., and Nédélec, P.: Mixing between a stratospheric intrusion and a biomass burning plume, *Atmos. Chem. Phys.*, 7, 4229–4235, 2007, <http://www.atmos-chem-phys.net/7/4229/2007/>.

Browell, E. V., Danielsen, E. F., Ismail, S., Gregory, G. L., and Beck, S. M.: Tropopause Fold

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Structure Determined from Airborne Lidar and in Situ Measurements, *J. Geophys. Res.*, 92, 2112–2120, 1987.

Carnuth, W. and Trickl, T.: Transport studies with the IFU three-wavelength aerosol lidar during the VOTALP Mesolcina experiment, *Atmos. Environ.*, 34, 1425–1434, 2000.

5 Carnuth, W., Kempfer, U., and Trickl, T.: Highlights of the Tropospheric Lidar Studies at IFU within the TOR Project, *Tellus B.*, 54, 163–185, 2002.

Cho, J. Y. N., Newell, R. E., Browell, E. V., Grant, W. B., Butler, C. F., and Fenn, M. A.: Observation of pollution plume capping by a tropopause fold, *Geophys. Res. Lett.*, 28, 3243–3246, 2001.

10 Cristofanelli, P., Bonasoni, P., Collins, W., Feichter, J., Forster, C., James, P., Kentarchos, A., Kubik, P. W., Land, C., Meloen, J., Roelofs, G. J., Siegmund, P., Sprenger, M., Schnabel, C., Stohl, A., Tobler, L., Tositti, L., Trickl, T., and Zanis, P.: Stratosphere-to-troposphere transport: A model and method evaluation, *J. Geophys. Res.*, 108, 8525, doi:10.1029/2002JD002600, 2003.

15 Cristofanelli, P., Bonasoni, P., Tositti, L., Bonafè, U., Calzolari, F., Evangelsti, F., Sandrini, S., and Stohl, A.: A 6-year analysis of stratospheric intrusions and their influence on the ozone at Mt. Cimone (2165 m above sea level), *J. Geophys. Res.*, 111, D03306, doi:10.1029/2005JD006553, 2006.

20 Danielsen, E. F. and Mohnen, V. A.: Project Dustorm Report: Ozone Transport, in Situ Measurements, and Meteorological Analyses of Tropopause Folding, *J. Geophys. Res.*, 82, 5867–5877, 1977.

Draxler, R. and Hess, G.: An overview of the HYSPLIT_4 modelling system for trajectories, dispersion, and deposition, *Aust. Meteorol. Mag.*, 47, 295–308, 1998.

25 Ebel, A., Feldmann, H., Jakobs, H. J., Memmesheimer, M., Offermann, D., Kuell, V., and Schaefer, B.: Simulation of transport and composition changes during a blocking episode over the East Atlantic and North Europe, *Ecol. Model.*, 217, 240–254, 2008.

Eisele, H. and Trickl, T.: Second Generation of the IFU Stationary Tropospheric Ozone Lidar, in: *Advances in Atmospheric Remote Sensing with Lidar, Selected Papers of the 18th International Laser Radar Conference, Berlin (Germany), July 22 to 26, 1996*, edited by: Ansmann, A., Neuber, R., Rairoux, P., and Wandinger, U., Springer, Berlin, Heidelberg, Germany, 379–382, 1997.

30 Eisele, H. and Trickl, T.: Improvements of the aerosol algorithm in ozone-lidar data processing by use of evolutionary strategies, *Appl. Optics*, 44, 2638–2651, 2005.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Eisele, H., Scheel, H. E., Sladkovic, R., and Trickl, T.: High-resolution Lidar Measurements of Stratosphere-troposphere Exchange, *J. Atmos. Sci.*, 56, 319–330, 1999.
- Elbern, H., Kowol, J., Sladkovic, R., and Ebel, A.: Deep stratospheric intrusions: A statistical assessment with model guided analysis, *Atmos. Environ.*, 31, 3207–3226, 1997.
- 5 Flentje, H., Dörnbrack, A., Ehret, G., Fix, A., Kiemle, C., Poberaj, G., and Wirth, M.: Water vapor heterogeneity related to tropopause folds over the North Atlantic revealed by airborne water vapor differential absorption lidar, *J. Geophys. Res.*, 110, D03115, doi:10.1029/2004JD004957, 2005.
- Galani, E., Balis, D., Zanis, P., Zerefos, C., Papayannis, A., Wernli, H., and Gerasopoulos, E.: Observations of stratosphere-to-troposphere transport events over the eastern Mediterranean using a ground-based lidar system, *J. Geophys. Res.*, 108, 8527, doi:10.1029/2002JD002596, 2003.
- 10 Gerasopoulos, E., Zerefos, C. S., Papastefanou, C., Zanis, P., and O'Brien, K.: Low-frequency variability of beryllium-7 surface concentrations over the Eastern Mediterranean, *Atmos. Environ.*, 37, 1745–1756, 2003.
- Grant, W. B., Browell, E. V., Butler, C. F., Fenn, M. A., Clayton, M. B., Hannan, J. R., Fuelberg, H. E., Blake, D. R., Blake, N. J., Gregory, G. L., Heikes, B. G., Sachse, G. W., Singh, H. B., Snow, J., and Talbot, R. W.: A case study of transport of tropical marine boundary layer and lower tropospheric air masses to the northern midlatitude upper troposphere, *J. Geophys. Res.*, 105, 3757–3769, 2000.
- 20 Grell, G., Duddhia, J., and Staufer, D.: A description of the fifth-generation Penn State/Ncar meso-scale model (MM5), Technical note NCAR/TN-398+STR, National Center of Atmospheric Research (Boulder, USA), 122 pp., 1994.
- Hass, H., Ebel, A., Feldmann, H., Jakobs, H. J., and Memmesheimer, M.: Evaluation studies with a regional chemical transport model (EURAD) using air quality data from the EMEP monitoring network, *Atmos. Environ.*, 27A, 867–887, 1993.
- 25 Holton, J. R., Douglass, A. R., Haynes, P. H., McIntyre, M. E., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, *Rev. Geophys.*, 33, 403–439, 1996.
- Koch, D. M. and Mann, M. E.: Spatial and temporal variability of ^7Be surface concentrations, *Tellus B*, 48, 387–396, 1996.
- 30 Kowol-Santen, J., Elbern, H., and Ebel, A.: Estimation of cross-tropopause air mass fluxes at midlatitudes: Comparison of different numerical methods and meteorological situations, *Mon. Weather Rev.*, 128, 4045–4057, 2000.

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Kreipl, S.: Messung des Aerosoltransports am Alpennordrand mittels Laserradar (Lidar), Dissertation zur Erlangung des Doktorgrads, Universität Erlangen (Germany), 195 pp., 2005 (in German).

Lal, D. and Peters, B.: Cosmic ray produced radioactivity on the earth, *Handb. Phys.*, 46, 551–612, 1967 (as cited by Zanis et al., 2003a).

Langford, A. O., Masters, C. D., Proffitt, M. H., Hsie, E. Y., and Tuck, A. F.: Ozone measurements in a tropopause fold associated with a cut-off low system, *Geophys. Res. Lett.*, 23, 2501–2504, 1996.

Meloan, J., Siegmund, P., van Velthoven, P., Kelder, H., Sprenger, M., Wernli, H., Kentarchos, A., Roelofs, G., Feichter, J., Land, C., Forster, C., James, P., Stohl, A., Collins, W., and Crstofanelli, P.: Stratosphere-troposphere exchange: A model and method intercomparison, *J. Geophys. Res.*, 108, 8526, doi:10.1029/2002JD002274, 2003.

Memmesheimer, M., Tippke, J., Ebel, A., Hass, H., Jakobs, H. J., and Laube, M.: On the use of EMEP emission inventories for European scale air pollution modeling with the EURAD model, in: *Proceedings of the EMEP workshop on Photooxidant Modelling for Long-Range Transport in Relation to Abatement Strategies*, Berlin, Germany, April 16–19, 307–324, 1991.

Oltmans, S. J., Lefohn, A. S., Harris, J. M., Galbally, I., Scheel, H. E., Bodeker, G., Brunke, E., Claude, H., Tarasick, D., Johnson, B. J., Simmonds, P., Shadwick, D., Anlauf, K., Hayden, K., Schmidlin, F., Fujimoto, F., Akagi, K., Meyer, C., Nichol, S., Davies, J., Redondas, A., and Cuevas, E.: Long-term changes in tropospheric ozone, *Atmos. Environ.*, 40, 3156–3173, 2006.

Papayannis, A., Balis, D., Zanis, P., Galani, E., Wernli, H., Zerefos, C., Stohl, A., Eckhardt, S., and Amiridis, V.: Sampling of an STT event over the Eastern Mediterranean region by lidar and electrochemical sonde, *Ann. Geophys.*, 23, 2039–2050, 2005, <http://www.ann-geophys.net/23/2039/2005/>.

Parrish, D. D., Holloway, J. S., Jakoubek, R., Trainer, M., Ryerson, T. B., Hübler, G., Fehsenfeld, F. C., Moody, J. L., and Cooper, O. C.: Mixing of anthropogenic pollution with stratospheric ozone: A case study from the North Atlantic wintertime troposphere, *J. Geophys. Res.*, 105, 24363–24374, 2000.

Pötzl, K. and Carnuth, W.: Isolierung und Identifizierung des atmosphärischen ^7Be , *Radiochim. Acta*, 6, 133–135, 1966 (in German).

Ravetta, F., Ancellet, G., Kowol-Santen, J., Wilson, R., and Nedeljkovic, D.: Ozone, Temper-

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- ature and Wind Field Measurements in a Tropopause Fold: Comparison with a Mesoscale Model Simulation, *Mon. Weather Rev.*, 127, 2641–2653, 1999.
- Reiter, R., Sladkovic, R., Pötl, K., Carnuth, W., and Kanter, H.-J.: Studies on the Influx of Stratospheric Air into the Lower Troposphere Using Cosmic-Ray Radionuclides and Fallout, *Arch. Meteor. Geophys. A.*, 20, 211–246, 1971.
- Reiter, R., Munzert, K., Kanter, H.-J., and Pötl, K.: Cosmogenic Radionuclides and Ozone at a Mountain Station at 3.0 km a.s.l., *Arch. Meteor. Geophys. B.*, 32, 131–160, 1983.
- Reiter, R., Sladkovic, R., and Kanter, H.-J.: Concentration of Trace Gases in the Lower Troposphere, Simultaneously Recorded at Neighboring Mountain Stations, Part II: Ozone, *Meteorol. Atmos. Phys.*, 37, 27–47, 1987.
- Roelofs, G.-J. and Lelieveld, J.: Model study of the influence of cross-tropopause O₃ transports on tropospheric O₃ levels, *Tellus B.*, 49, 38–55, 1997.
- Roelofs, G. J., Kentarchos, A. S., Trickl, T., Stohl, A., Collins, W. J., Crowther, R. A., Hauglustaine, D., Klonecki, A., Law, K. S., Lawrence, M. G., von Kuhlmann, R., and van Weele, M.: Intercomparison of tropospheric ozone models: Ozone transport in a complex tropopause folding event, *J. Geophys. Res.*, 108, 8529, doi:10.1029/2003JD003462, 2003.
- Scheel, H. E.: Measurements of O₃, ⁷Be, RH and CO at the Zugspitze summit, in: EUROTRAC – Influence of Stratosphere-Troposphere Exchange in a Changing Climate on Atmospheric Transport and Oxidation Capacity, Final Report, Contract EVK2-CT-1999-00050, A. Stohl, Co-ordinator, European Union, 58–62, available at: <http://www.forst.tu-muenchen.de/EXT/LST/METEO/staccato/>, 2002.
- Scheel, H. E.: Ozone Climatology Studies for the Zugspitze and Neighbouring Sites in the German Alps, in: Tropospheric Ozone Research 2, EUROTRAC-2 Subproject Final Report, A. Lindskog, Co-ordinator, EUROTRAC International Scientific Secretariat, München, Germany, 134–139, 2003.
- Scheel, H. E., Areskoug, H., Geiß, H., Gomiscek, B., Granby, K., Haszpra, L., Klasinc, L., Kley, D., Laurila, T., Lindskog, A., Roemer, M., Schmitt, R., Simmonds, P., Solberg, S., and Toupance, G.: On the Spatial Distribution and Seasonal Variation of Lower-Tropospheric Ozone over Europe, *J. Atmos. Chem.*, 28, 11–28, 1997.
- Seibert, P., Feldmann, H., Neininger, B., Bäumle, M., and Trickl, T.: South foehn and ozone in the Eastern Alps – case study and climatological aspect, *Atmos. Environ.*, 34, 1379–1394, 2000.
- Shapiro, M. A.: Turbulent Mixing within Tropopause Folds as a Mechanism for the Exchange

of Chemical Constituents between the Stratosphere and Troposphere, *J. Atmos. Sci.*, 37, 994–1004, 1980.

Simmonds, P. G., Derwent, R. G., Manning, A. L., and Spain, G.: Significant growth in surface ozone at Mace Head, Ireland, 1987–2003, *Atmos. Environ.*, 38, 4769–4778, 2004.

5 Sladkovic, R. and Munzert, K.: Lufthygienisch-klimatologische Überwachung im bayrischen Alpenraum, Abschnitt VI.4, Ozonspitzen auf der Zugspitze durch Zustrom aus der Stratosphäre, Final Report, Fraunhofer-Institut für Atmosphärische Umweltforschung, Report 908080, 49–50, 1990.

10 Sprenger, M., Croci Maspoli, M., and Wernli, H.: Tropopause folds and cross-tropopause exchange: A global investigation based upon ECMWF analyses for the time period March 2000 to February 2001, *J. Geophys. Res.*, 108, 8518, doi:10.1029/2002JD002587, 2003.

STACCATO: Special section with 15 publications, *J. Geophys. Res.*, 108(D12), papers STA 1–15, 27 June, 2003.

15 Stohl, A. and Seibert, P.: Accuracy of trajectories as determined from the conservation of meteorological tracers, *Q. J. Roy. Meteor. Soc.*, 124, 1465–1484, 1998.

Stohl, A. and Thomson, D. J.: A density correction for Lagrangian particle dispersion models, *Bound.-Lay. Meteorol.*, 90, 155–167, 1999.

20 Stohl, A. and Trickl, T.: A textbook example of long-range transport: Simultaneous observation of ozone maxima of stratospheric and North American origin in the free troposphere over Europe, *J. Geophys. Res.*, 104, 30445–30462, 1999.

Stohl, A., Spichtinger-Rakowsky, N., Bonasoni, P., Feldmann, H., Memmesheimer, M., Scheel, H. E., Trickl, T., Hübener, S., Ringer, W., and Mandl, M.: The influence of stratospheric intrusions on alpine ozone concentrations, *Atmos. Environ.*, 34, 1323–1354, 2000.

25 Stohl, A., Eckhardt, S., Forster, C., James, P., Spichtinger, N., and Seibert, P.: A replacement for simple back trajectory calculations in the interpretation of atmospheric trace substance measurements, *Atmos. Environ.*, 36, 4635–4648, 2002.

30 Stohl, A., Bonasoni, P., Cristofanelli, P., Collins, W., Feichter, J., Frank, A., Forster, C., Gerasopoulos, E., Gäggeler, H., James, P., Kentarchos, T., Kromp-Kolb, H., Krüger, B., Land, C., Meloen, J., Papayannis, A., Priller, A., Seibert, P., Sprenger, M., Roelofs, G. J., Scheel, H. E., Schnabel, C., Siegmund, P., Tobler, L., Trickl, T., Wernli, H., Wirth, V., Zanis, P., and Zerefos, C.: Stratosphere-troposphere exchange – a review, and what we have learned from STACCATO, *J. Geophys. Res.*, 108, 8516, doi:10.1029/2002JD002490, 2003.

Tositti, L., Hübener, S., Kanter, H.-J., Ringer, W., Sandrini, S., and Tobler, L.: Intercomparison

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of sampling and measurements of ^7Be in air at four high altitude locations in Europe, Appl. Radiat. Isotopes., 61, 1497–1502, 2004.

Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, Atmos. Chem. Phys., 5, 2461–2474, 2005, <http://www.atmos-chem-phys.net/5/2461/2005/>.

Trickl, T., Cooper, O. C., Eisele, H., James, P., Mücke, R., and Stohl, A.: Intercontinental transport and its influence on the ozone concentrations over central Europe: Three case studies, J. Geophys. Res., 108, 8530, doi:10.1029/2002JD002735, 2003.

Trickl, T., Eisele, H., Feldmann, H., Bärtsch-Ritter, N., Furger, M., Mücke, R., Scheel, H. E., and Stohl, A.: High-ozone Layers in the Middle and Upper Troposphere above Central Europe: Simultaneous Import from the Stratosphere over the Pacific Ocean, East Asia and North America, Atmos. Chem. Phys., accepted, 2009a.

Trickl, T., Forster, C., Scheel, H. E., and Stohl, A.: Low Ozone and Carbon Monoxide in Air Masses from the Subtropical North Atlantic, in preparation (2009b).

Van Haver, P., De Muer, D., Beekmann, M., and Mancier, C.: Climatology of tropopause folds at midlatitudes, Geophys. Res Lett., 23, 1033–1036, 1996.

Viezee, W., Johnson, W. B., and Singh, H. B.: Stratospheric Ozone in the Lower Troposphere – II. Assessment of Downward Flux and Ground-Level Impact, Atmos. Environ., 17, 1979–1993, 1983.

Vogelmann, H. and Trickl, T.: Wide-Range Sounding of Free-Tropospheric Water Vapor with a Differential-Absorption Lidar (DIAL) at a High-Altitude Station, Appl. Optics, 47, 2116–2132, 2008.

VOTALP: Vertical Ozone Transport in the Alps, special section with nine publications, Atmos. Environ., 34, 1319–1453, 2000a.

VOTALP II: Vertical Ozone Transport in the Alps II, Final Report for the European Union, Contract Nr.: ENV4 CT970413, Reporting Period 1/3/1998-29/2/2000, H. Kromp-Kolb, Coordinator, Universität für Bodenkultur Wien, Austria, Institut für Meteorologie und Physik, 96 pp., available at: <http://www.boku.ac.at/imp/votalp/>, 2000b.

Wernli, H. and Davies, H. C.: A Lagrangian-based analysis of extratropical cyclones. I. The method and some applications, 123, 467–489, 1997.

Wernli, H.: A Lagrangian-based analysis of extratropical cyclones. II: A detailed case study, Q. J. Roy. Meteor. Soc., 123, 1677–1706, 1997.

Zanis, P., Trickl, T., Stohl, A., Wernli, H., Cooper, O., Zerefos, C., Gaeggeler, H., Schnabel,

C., Tobler, L., Kubik, P. W., Priller, A., Scheel, H. E., Kanter, H. J., Cristofanelli, P., Forster, C., James, P., Gerasopoulos, E., Delcloo, A., Papayannis, A., and Claude, H.: Forecast, observation and modelling of a deep stratospheric intrusion event over Europe, *Atmos. Chem. Phys.*, 3, 763–777, 2003,

<http://www.atmos-chem-phys.net/3/763/2003/>.

Zanis, P., Gerasopoulos, E., Priller, A., Schnabel, C., Stohl, A., Zerefos, C., Gäggeler, H. W., Tobler, L., Kubik, P. W., Kanter, H. J., Scheel, H. E., Luterbacher, J., and Berger, M.: An estimate of the impact of stratosphere-to-troposphere transport (STT) on the lower free tropospheric ozone over the Alps using ^{10}Be and ^7Be measurements, *J. Geophys. Res.*, 108, 8520, doi:10.1029/2002JD002604, 2003a.

ACPD

9, 2223–2288, 2009

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. List of intrusion periods during the period of intensive lidar measurements between 10 February and 23 August 2001: The intrusion type is specified as defined in Sect. 4.1. For the Zugspitze data the days are listed on which one of the stratospheric criteria specified in Sect. 4.1 was met. For the lidar the number of observations on days with positive predictions is given. * denotes cases for which the lidar revealed a high overpass. Abbreviations: n.d. (not detected), c.n.f.m. (criteria not fully met), n.m. (no measurements).

Predicted	Type	Zugspitze	Lidar Meas.	Comment
11–12 Feb	4	11–12 Feb	24, 17	
14–16 Feb	1	14 Feb	0, 13, 0	14 Feb: fog in valley
18 Feb	5	18 Feb	9	
19 Feb	1	19 Feb	8	
21 Feb	1	21 Feb (short)	n.m.	fog in valley
25–28* Feb	2	27 Feb (short)	4,5,22,5	mostly above Zugspitze
5–7 Mar	2	6 Mar	n.m.	no full hit pred., long path
13 Mar	4	n.d.	4	lidar: elevated $O_3 > 5$ km
16 Mar	4	c.n.f.m.	5	lidar: elevated $O_3 > 5$ km
24 Mar	5	c.n.f.m.	n.m.	$[^7\text{Be}] \approx 7 \text{ mBq m}^{-3}$
28–29 Mar	5	c.n.f.m.	n.m.	
31 Mar–2 Apr*	4, 5, 7	2 detected	10, 22, 1	3 simultaneous intrusions
5 Apr	5	c.n.f.m.	n.m.	$[^7\text{Be}] \approx 5 \text{ mBq m}^{-3}$
14–15 Apr	1	n.d., RH100 %	n.m.	clearing too short
23 Apr	2, 3	first intr. seen	13	
24 Apr	4	24 Apr	n.m.	
25 Apr	4	c.n.f.m.	n.m.	
27 Apr	5	27 Apr	n.m.	
2 May	2	2 May	n.m.	highly indirect (Morocco)
missed	1	13 May	n.m.	
19–21 May	2,	19–22 May	6, 8, 9	
22 May	1	22 May	5,3	lidar: 22–23 May
4–5 Jun	2, 2	4–5 Jun	7, 12, 3	2 parallel intrusions
12–13 Jun	2	12–13 Jun	2, 13, 6	lidar: 11–13 Jun
15 Jun	4	15 Jun (short)	n.m.	$[^7\text{Be}] \approx 8.5 \text{ mBq m}^{-3}$
20–21 Jun (2 of 3)*	6, 1, 1	first two seen	8, 9	
23–24 Jun	2	24 Jun	9, 12	
25 Jun	3	26–27 Jun	4	
2–3 Jul	1	3 Jul	4, 7	lidar: 3–5 Jul
21–22 Jul	2	22 Jul	10, 20, 12, 4	lidar: 21–24 Jul
31 Jul	3	31 Jul	n.m.	
2 Aug	4	2 Aug	n.m.	
12–13 Aug (1 of 2)	1	2 seen	9, 12, 12, 8	lidar: 12–15 Aug
22–23 Aug	4	4 RH dips on 24–27 Aug	13, 3	$[^7\text{Be}] > 5.5 \text{ mBq m}^{-3}$ on 24–27 Aug

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Table 2. Overview of the number of cases suggested by the LAGRANTO and HYSPLIT model calculations for the different intrusion types and for advection within the troposphere within 315 h.

Type	2001	2002	2003	2004	2005	Sum
1	17	22	23	25	18	103
2	22	16	22	16	22	98
3	8	8	6	10	7	39
4	11	1	6	1	2	21
5	21	24	12	18	12	87
6	7	4	7	12	21	51
7	1	2	1	1	0	5
8	0	0	2	1	5	8
Sum	87	75	79	84	87	412
“Tropospheric”	3	1	3	4	5	16

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Results of a five-year run with the FLEXPART model (1995–1999) for stratospheric tracers arriving at the four VOTALP summit stations Monte Cimone, Zugspitze, Sonnblick and Jungfrauoch; the tropospheric age of the tracer is colour coded. The summer minimum at the higher Jungfrauoch station is less pronounced than at the three other, lower-lying sites.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 2. Examples of lidar sounding series selected from the measurements in 2001; the labels S, S1, S2 and S3 denote stratospheric intrusions, LA means long-range advection from high altitudes over the North Pacific, SA “subtropical Atlantic”, P “Pacific”.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 2. Continued.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 2. Continued. The upper panel is a short version of a series published by Trickl et al. (2008a); USA denotes air from the boundary layer in the United States.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.





Fig. 3. Four-day trajectory forecasts by ETH Zürich for the cases displayed in Fig. 2; the crosses mark the position every 24 h after the start times shown at the top of each panel. The pressure level is colour coded in mbar. The Zugspitze summit corresponds to about 700 mbar.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

**Fig. 3.** Continued.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.

**Fig. 3.** Continued.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. Zugspitze ozone, carbon monoxide, ^7Be , and relative humidity (RH) on 30 March to 9 April 2001; the upper two traces indicate the validity on filtering criteria 1 (elevated values in dark red curve) and 2 (elevated values in black curve) described in Sect. 4. The violet triangles on the RH curve mark the validity range of Criterion 2. The two intrusions (S1 and S2) are defined as in the first panel of Fig. 2. Due to the short coverage of 31 March by the first intrusion (S1) the ^7Be values do not reach the “classical” threshold condition for stratospheric air of about 8 mBq m^{-3} (85th) percentile.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 5. Maximum values of the daily stratospheric fractions obtained along the FLEXPART retroplumes calculated for four times on 1 April 2001, releasing the numerical particles in vertical intervals of 250 m.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 6. EURAD simulation for the period 20 June to 25 June 2001; the labels are the same as in the corresponding panels of Fig. 2. Time: UTC=CET–1 h.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 7. EURAD simulation for the period 11–12 August 2001; the labels are the same as in the corresponding panel of Fig. 2. Time: UTC=CET–1 h.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 8. Four-day trajectory forecast initialized on 12 November 2001, at 12:00 UTC: the intrusion covers most of the European continent. It lasted for six days (Zugspitze summit: 14–19 November) and was followed by the advection of stratospheric air from Canada. The colour indicates potential vorticity here.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 9. Zugspitze ozone, carbon monoxide, ⁷Be, and relative humidity (RH) for November 2001; the upper two traces indicate the validity on filtering criteria 1 (elevated values in dark red curve, top trace) and 2 (elevated values in black curve, second trace) described in Sect. 4. The violet triangles on the RH curve mark the validity range of Criterion 2 (Sect. 4). The spectacular CO spike by the end of 13 November corresponds to polluted air advected in the frontal system preceding the intrusion.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 10. 315-h backward trajectories for the three altitude levels 3000 m (red triangles), 3300 m (blue squares) and 3600 m a.s.l. (green circles) calculated with the HYSPLIT model: The heights in the figure are given with respect to the ground (AGL) and are obviously wrong for the Zugspitze position (asterisk) due to a insufficient model orography. Thus, the exact height representing the Zugspitze summit is not certain. The air mass for the blue trajectory (3300 m a.s.l.) started at almost 8 km above north-western Canada at -315 h and proceeded as far east as West Siberia before it descended back to the observational site.

Forecasted deep
 stratospheric
 intrusions over
 Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 11. Zugspitze data for January 2003 (for details see Fig. 4); the numbers next to the RH curve specify the intrusion type obtained from the trajectories.

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 12. Total number of intrusion cases predicted or verified by trajectories and total number of cases verified by the data filtering (Criteria 1–3) obtained for 2001 to 2005; please, note that a single intrusion case may extend over several days and may not fulfil the individual filtering criteria during the entire period.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 13. Same as Fig. 12, but for intrusion Type 1 only.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 14. Same as Fig. 12, but for intrusion Type 2 only.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.




Fig. 15. Same as Fig. 12, but for intrusion Type 5 only.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Forecasted deep
stratospheric
intrusions over
Central Europe**

T. Trickl et al.




Fig. 16. Same as Fig. 12, but for intrusion Type 6 only.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 17. Seasonal cycle of the cases not related to stratospheric air when using 5.5 mBq m^{-3} as the ${}^7\text{Be}$ threshold (number of days on which a wrong identification was obtained); the additional use of $\text{RH} < 60\%$ (Criterion 3) is important to reduce the wrong counts.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Forecasted deep stratospheric intrusions over Central Europe

T. Trickl et al.




Fig. 18. Fractions of intrusions for six of the eight intrusion types identified in this paper fulfilling different RH conditions (minimum RH values).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)