

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

# New cloud chamber experiments on the heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner, O. Möhler, H. Saathoff, M. Schnaiter, and T. Leisner

Karlsruhe Institute of Technology, Institute for Meteorology and Climate Research, Karlsruhe, Germany

Received: 17 November 2010 – Accepted: 23 November 2010 – Published: 2 December 2010

Correspondence to: R. Wagner (robert.wagner2@kit.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

The heterogeneous ice nucleation ability of oxalic acid in the immersion mode has been investigated by controlled expansion cooling runs with airborne, ternary solution droplets composed of, (i), sodium chloride, oxalic acid, and water (NaCl/OA/H<sub>2</sub>O) and, (ii), sulphuric acid, oxalic acid, and water (H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O). Polydisperse aerosol populations with median diameters ranging from 0.5–0.7 μm and varying solute concentrations were prepared. The expansion experiments were conducted in the AIDA aerosol and cloud chamber of the Karlsruhe Institute of Technology at initial temperatures of 244 and 235 K. In the ternary NaCl/OA/H<sub>2</sub>O system, solid inclusions of oxalic acid, presumably nucleated as oxalic acid dihydrate, were formed by temporarily exposing the ternary solution droplets to a relative humidity below the efflorescence point of NaCl. The matrix of the crystallised NaCl particulates triggered the precipitation of the organic crystals which later on remained as solid inclusions in the solution droplets when the relative humidity was again raised above the deliquescence point of NaCl. The embedded oxalic acid crystals reduced the critical ice saturation ratio required for the homogeneous freezing of pure NaCl/H<sub>2</sub>O solution droplets at a temperature of around 231 K from 1.38 to about 1.32. Aqueous solution droplets with OA inclusions larger than about 0.27 μm in diameter efficiently nucleated ice by condensation freezing when they were activated to micron-sized cloud droplets at 241 K, i.e., they froze well above the homogeneous freezing temperature of pure water droplets of about 237 K. Our results on the immersion freezing potential of oxalic acid corroborate the findings from a recent study with emulsified aqueous solutions containing crystalline oxalic acid. In those experiments, the crystallisation of oxalic acid dihydrate was triggered by a preceding homogeneous freezing cycle with the emulsion samples. The expansion cooling cycles with ternary H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O solution droplets were aimed to analyse whether those findings can be transferred to ice nucleation experiments with airborne oxalic acid containing aerosol particles. Under our experimental conditions, the efficiency by which the surface of homogeneously nucleated ice crystals triggered the precipitation

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of oxalic acid dihydrate was very low, i.e., less than one out of a hundred ice crystals that were formed by homogeneous freezing in a first expansion cooling cycle left behind an ice-active organic crystal that acted as immersion freezing nucleus in a second expansion cooling cycle.

## 1 Introduction

Dicarboxylic acids are ubiquitous components of the water-soluble organic fraction of the tropospheric aerosol. Their contribution to the total particulate organic carbon typically ranges from 1–3 wt% in urban and semi-urban areas to up to 10 wt% in remote marine atmospheres (Kerminen et al., 2000, 1999). Oxalic acid (HOOC-COOH), malonic acid (HOOC-CH<sub>2</sub>-COOH), and succinic acid (HOOC-(CH<sub>2</sub>)<sub>2</sub>-COOH) are usually the most abundant species, partly comprising more than 80% of the total diacid mass concentrations (Kawamura et al., 1996). Apart from investigating their hygroscopic behaviour and ability to act as atmospheric cloud condensation nuclei, several recent studies have addressed the ice nucleation potential of solid low-molecular weight dicarboxylic acids. On the one hand, the dicarboxylic acids can nucleate ice directly from the vapour phase (deposition mode ice nucleation) provided that they are directly exposed to an ice supersaturated environment, e.g. in form of a crystalline coating layer that was formed by physical and chemical processing on prevalent particulates such as mineral dust and soot. At typical tropospheric relative humidities, diacids of low water solubility like oxalic and succinic acid may, on the other hand, also be present as solids that are immersed in aqueous inorganic and/or organic solution droplets (Marcolli et al., 2004; Zobrist et al., 2006). These embedded crystals can act as ice nuclei in the immersion mode.

For both modes of ice nucleation, oxalic acid in form of its dihydrate has revealed the by far highest ice nucleation potential amongst a variety of other dicarboxylic acids investigated. Oxalic acid dihydrate particles deposited onto a hydrophobic cold stage at 233 K were found to be efficient deposition mode ice nuclei at critical ice saturation

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nucleation from a gaseous oxalic acid/air mixture or by rapid crystallisation of highly supersaturated aqueous solution droplets at temperatures  $\leq 244$  K proved to be even more ice active in the deposition mode than the particles probed by Kanji et al. (2008), showing ice formation at threshold ice saturation ratios lower than 1.10. In contrast, oxalic acid dihydrate particles that had crystallised from less supersaturated solution droplets at a higher temperature and were then cooled to 244 K partly revealed no ice activity at all. We have suggested that the high variability in the ice nucleation potential is related to differences in the surface morphology and particle microstructure of the variously generated oxalic acid dihydrate crystals, a dependence that has been evidenced in previous ice nucleation experiments with e.g. crystalline ammonium sulphate particles (Zuberi et al., 2001).

Concerning the immersion mode ice nucleation potential of dicarboxylic acids, Zobrist et al. (2006) have detected a pathway that leads to the nucleation of crystalline organics in aqueous solution droplets, namely homogeneous ice nucleation. After having measured the homogeneous freezing temperature of emulsified binary solutions of oxalic, succinic, and adipic acid in water with a differential scanning calorimeter (DSC), the samples were heated above the eutectic melting point of ice and then revealed a higher freezing temperature in a second cooling cycle. Only in the case of oxalic acid, however, the freezing temperature in the second cooling cycle was higher than the homogeneous freezing temperature of pure water, indicating that this compound, presumably precipitated as oxalic acid dihydrate, is able to act as a heterogeneous immersion mode ice nucleus. Oxalic acid dihydrate also proved to be ice-active in the immersion mode when precipitated from ternary aqueous solutions with sodium chloride, malonic acid, and sulphuric acid as additional solute species. In their summary section, Zobrist et al. (2006) state that the crystallisation of oxalic acid dihydrate occurs concurrently with or induced by ice freezing. As a plausible mechanism, oxalic acid dihydrate crystallises heterogeneously on the homogeneously nucleated ice crystals during the first cooling cycle. The trajectory for ice freezing experiments with emulsified solutions in the temperature-composition plane deviates from that of cooling air parcels in the

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmosphere because in the latter case the aerosol particles are directly susceptible to the ambient relative humidity and will correspondingly change their composition. In this manuscript, we present new ice nucleation experiments on the immersion freezing potential of oxalic acid which closely mimic the trajectory of a cooling air parcel in the atmosphere. We want to analyse whether the findings from Zobrist et al. (2006) can be transferred to ice nucleation experiments with airborne oxalic acid containing aerosol particles. For this purpose, we have conducted a series of expansion cooling cycles in the AIDA aerosol and cloud chamber of the Karlsruhe Institute of Technology.

In Fig. 1, the experimental trajectories of two feasible AIDA expansion cooling experiments are shown as green and blue lines. In the preceding ice nucleation experiments with binary oxalic acid/water solution droplets, we have always observed the crystallisation of aqueous oxalic solution droplets to oxalic acid dihydrate after adding the aerosol to the AIDA chamber at  $T \leq 244$  K and were therefore not able to simulate the envisaged experimental approach which requires that supercooled solution droplets are initially present. In this manuscript, however, we describe a set of experiments with ternary sulphuric acid/oxalic acid/water solution droplets that remained in a supercooled liquid state in the AIDA chamber. Thereby, repetitive expansion cooling cycles similar to the Zobrist et al. (2006) freezing experiments were feasible. Figure 1 shows the phase diagram for a ternary solution with a sulphuric acid to oxalic acid mass ratio of 5:1.14. The abscissa denotes the total solute concentration (wt% sulphuric acid + wt% oxalic acid). The initial composition of the injected aerosol particles in experiments A and B adjusts to the prevailing relative humidity. In the preparation of the expansion cooling cycles, the inner walls of the AIDA chamber are coated by a thin ice layer (see detailed description in Sect. 2) and, therefore, the initial aerosol compositions are given by points A1 and B1 on the ice melting-point curve in the aqueous solutions, yielding a total solute concentration of about 31 and 35 wt%, respectively. To mimic the expansion cooling of rising air parcels in the atmosphere, the AIDA pressure is reduced by controlled pumping, leading to a decrease in the gas temperature of up to 10 K, depending on the pumping speed. As a result of the increasing relative humidity

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

during expansion cooling, the initially concentrated solution droplets will strongly dilute. The initial temperature of experiment A (244 K) is chosen in a way that saturation with respect to supercooled water is reached during the expansion before the gas temperature has dropped below the homogeneous freezing temperature of pure water droplets of about 237 K. When reaching 100% RH after about 3 to 4° of cooling (240–241 K, A2), the ternary solution droplets are activated to a cloud of almost pure supercooled water droplets. After the gas temperature drops below 237 K (A3) during the continued expansion, the droplet cloud starts to glaciate. In experiment B that is started at 235 K below the homogeneous freezing temperature of pure water droplets, the ternary solution droplets will not dilute to almost pure water droplets but will homogeneously freeze once the solute concentration has fallen below a critical threshold value (B2).

After the ice cloud formation in experiments A and B, the pumping is stopped and the ice crystals slowly evaporate on a timescale of typically 10 to 15 min. During this time period, the available ice particle surface may trigger the precipitation of oxalic acid dihydrate from the ternary sulphuric acid/oxalic acid/water solutions that have remained unfrozen and that will be included within the nucleated ice crystals or form a liquid film on their surface. After the ice crystals have completely evaporated, second expansion cooling cycles with the same aerosol loading are again started at 244 and 235 K to probe whether the homogeneous freezing event has indeed triggered the crystallisation of oxalic acid dihydrate. The nucleated organic particulates could act as immersion freezing nuclei in experiment B, thereby inducing earlier ice formation compared to the preceding homogeneous freezing run. In experiment A, the aerosol particles will again be activated to a droplet cloud at point A2. Our previous experiments have shown that potentially immersed oxalic acid dihydrate crystals might not get instantly and completely dissolved on the typical timescale of an expansion cooling experiment (Wagner et al., 2010). The embedded crystals might therefore act as immersion or condensation mode ice nuclei in this experiment and could provoke the glaciation of the droplet cloud before the homogeneous freezing temperature of pure water droplets has been reached.

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Already in the introductory part of our manuscript, we want to stress an important difference between the AIDA and the Zobrist et al. (2006) experiments, namely the ability to detect the potentially precipitated oxalic acid dihydrate particles in the second cooling cycles. For the experiments with emulsified aqueous solutions, it is reasonable to assume that all droplets of the emulsion will homogeneously freeze in the first cooling cycle. The samples are then slowly heated with  $1 \text{ K min}^{-1}$  to the conditioning temperature of 279 K. For an overall time period of about one hour, the ice crystals and the remaining unfrozen supersaturated solutions containing oxalic acid are in contact. The exemplarily shown exothermic ice freezing peak in the DSC heat flow record (see Fig. 1 in their publication) that was monitored during the second freezing cycle almost shows the same shape (peak height and peak area) compared to the first freezing cycle, i.e., the exothermic peak is shifted as a whole towards a clearly higher freezing temperature. This suggests that the outlined experimental conditions must have led to the precipitation of an ice-active oxalic acid dihydrate crystal in the vast majority of the droplets of the emulsion.

In contrast, heterogeneous ice nucleation induced by potentially precipitated oxalic acid dihydrate particles in the second expansion cooling cycles in the AIDA chamber experiments will only be detectable for a low number fraction of the initially added supercooled solution droplets. In the course of trajectory A, all seed aerosol particles will be activated to cloud droplets when reaching water saturation at point A2. Upon glaciation of the droplet cloud at about 237 K, however, ice crystals that have started to nucleate first in the chamber volume will grow at the expense of still unfrozen cloud droplets via the Bergeron-Findeisen process. Even for the highest employable pumping speed or cooling rate, the absolute number concentration of nucleated ice crystals is limited to a value of at most  $100 \text{ cm}^{-3}$ , irrespective of the seed aerosol number concentration that is typically between  $10^3$  and  $10^4 \text{ cm}^{-3}$ . Concerning experiment B, only supercooled solution droplets in a narrow range of the overall size distribution at around  $1 \mu\text{m}$  will homogeneously freeze at point B2; see Fig. 8 in Haag et al. (2003) and Fig. 3 in Wagner et al. (2008). Larger droplets remain liquid because their water activity lags



behind the equilibrium value due to kinetic growth limitations. Smaller droplets retain a higher solute concentration due to the Kelvin effect. The frozen droplets will rapidly deplete the excess of water vapour in the gas phase and prevent further solution droplets from freezing. Likewise to experiment A, the maximum ice particle number concentration, independent of the reservoir of seed aerosol particles, is thereby also limited to about  $100\text{ cm}^{-3}$ . Let us assume that in 10% of these nucleated ice crystals the precipitation of oxalic acid dihydrate might be triggered during their residence time of about 10–15 min in the chamber. Then, only up to 10 particles per  $\text{cm}^3$  AIDA volume would contain a solid nucleus before starting the second expansion cooling cycles. Not necessarily all of these embedded crystals will act as a heterogeneous immersion mode ice nucleus. Again assuming an ice-active fraction of 10%, one would obtain an ice particle number concentration of  $1\text{ cm}^{-3}$  that is formed by heterogeneous nucleation in the second freezing cycles. Although this is above the threshold number concentration of about  $0.1\text{ cm}^{-3}$  for our ice diagnostic instruments, the differences between the first and second cooling cycles in AIDA are expected to be much less pronounced in comparison with the repetitive emulsion freezing cycles. In case that the yield of ice-active oxalic acid nuclei from the first nucleation event would be even lower than estimated above, no change in the ice nucleation behaviour could be monitored.

Before addressing the experiments with ternary sulphuric acid/oxalic acid/water solution droplets in Sect. 3.2, we therefore describe in Sect. 3.1 a set of benchmark experiments with an aerosol population where indeed each individual particle contained a solid oxalic acid (dihydrate) nucleus. These experiments address another pathway that could lead to the precipitation of low-soluble organics within aqueous solution droplets in the atmosphere. An air parcel containing aqueous solution droplets of inorganic solutes like sodium chloride and/or ammonium sulphate in a mixture with small amounts of oxalic acid might at least temporarily be exposed to such a low relative humidity (RH) that efflorescence can occur. The matrix of the crystallized inorganic particulates could then also trigger the precipitation of oxalic acid dihydrate whereas pure oxalic acid aerosol particles have frequently been observed to be difficult to effloresce

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



even at relative humidities  $<5\%$  (Braban et al., 2003; Mikhailov et al., 2009; Prenni et al., 2001). When entering a regime with a higher relative humidity, the inorganic components might again get dissolved at the deliquescence point which, for a multi-component mixture, can be well below the deliquescence relative humidities of the pure compounds (Marcolli et al., 2004; Wexler and Seinfeld, 1991). In contrast to dicarboxylic acids with a high solubility in water like e.g. malonic acid and maleic acid, less soluble compounds like oxalic acid and succinic acid do not notably affect the deliquescence behaviour of the other solutes and remain as solid inclusions within the droplets even at elevated humidity levels (Brooks et al., 2002; Choi and Chan, 2002; Lightstone et al., 2000; Marcolli et al., 2004; Treuel et al., 2008). In the AIDA, we have investigated the ternary system sodium chloride, oxalic acid, and water. Reference expansion experiments that were conducted with pure effloresced sodium chloride particles at 244 and 235 K revealed that the inorganic crystals did not act as deposition mode ice nuclei but underwent a deliquescence transition to form aqueous solution droplets at about  $72 \pm 3\%$  RH which then, later on during the expansion, homogeneously froze as almost pure water droplets or dilute aqueous solution droplets at respective points A3 and B2 in the sodium chloride-water phase diagram (see Fig. 1). Performing these expansion experiments with mixed effloresced particles of sodium chloride and small amounts of oxalic acid, we will obtain an aerosol population of oxalic acid crystallites immersed in aqueous solution droplets after exceeding the deliquescence relative humidity of the inorganic salt. Thereby, we can investigate the ice nucleability of oxalic acid in the immersion mode with an ensemble of aerosol particles that all contain a solid nucleus. Based on these results, we will then analyse the efficiency by which oxalic acid crystals can precipitate from supercooled aqueous solution droplets via homogeneous ice nucleation.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2 Experimental

### 2.1 The AIDA facility as a moderate expansion cloud chamber for ice nucleation studies

The centre of the AIDA facility at the Karlsruhe Institute of Technology is its 84 m<sup>3</sup> sized aluminium chamber that is located inside a large, thermally insulated housing whose interior can be cooled to any temperature between ambient and 183 K (Fig. 2). A mixing fan, located at the bottom of the chamber, is continuously operating throughout the experiments for providing homogeneous conditions in terms of temperature, relative humidity, as well as aerosol and cloud particle number concentration inside the aerosol vessel. Under stirred conditions, the spatial temperature variability inside the vessel, as monitored by lines of horizontally and vertically arranged temperature sensors, is less than  $\pm 0.3$  K. In the preparation of an expansion run to be started at a given temperature  $T_{\text{start}}$ , the chamber, initially kept at a temperature higher than  $T_{\text{start}}$ , is evacuated with two vacuum pumps to a final pressure of about 0.01 hPa, followed by a series of flushing cycles with particle-free synthetic air to ensure aerosol free experimental conditions. The background particle number concentration is usually less than 0.1 cm<sup>-3</sup>. Then, a certain amount of Milli-Q quality water is evaporated into the evacuated chamber. Upon cooling to  $T_{\text{start}}$  and refilling the chamber with synthetic air to ambient pressure, the excess water vapour forms an ice layer at the inner chamber walls. Thereby, the relative humidity is close to an ice saturation ratio  $S_{\text{ice}}$  of one before the expansion run is started.  $S_{\text{ice}}$  is defined as the quotient of the prevailing water vapour pressure  $p_w(T)$  and the saturation pressure over ice  $p_{w,\text{ice}}(T)$ ,  $S_{\text{ice}} = p_w(T)/p_{w,\text{ice}}(T)$ . At a closer look,  $S_{\text{ice}}$  is typically in the range from 0.90 to 0.95 at static pressure and temperature conditions prior to the expansion cooling cycle because the gas temperature is always a few tenths of a Kelvin higher than the temperature of the ice-covered walls due to some internal heat sources (heating foils, heated sampling tubes, mixing fan). Finally, the desired seed aerosol particles whose ice nucleability will be investigated are added to the chamber (see Sect. 2.2).

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



i.e., the sum of interstitial and evaporated particle water, is measured ex situ with a fast high precision chilled mirror hygrometer (MBW, model 373) that samples through a heated sampling tube. Cloud particles (either supercooled droplets and/or ice crystals) are counted and sized by two optical particle counters (OPC1 and OPC2, type WE-LAS2000, Palas) in the size range from 0.7–40  $\mu\text{m}$  (OPC1) and 2.3–104  $\mu\text{m}$  (OPC2). In situ laser light scattering and depolarisation measurements as well as FTIR extinction measurements are further powerful tools to detect the nucleation and growth of the cloud particles and to distinguish between spherical supercooled cloud droplets and aspherical ice crystals. The records from the light scattering measurements with a 488 nm semiconductor laser (Coherent) that is directed horizontally through the AIDA chamber are the scattered light intensities in the forward ( $I_{\text{for}}$ , scattering angle  $2^\circ$ ) and backward ( $I_{\text{back}}$ , scattering angle  $178^\circ$ ) direction. The backscattered light is detected polarisation-resolved with  $I_{\text{back,par}}$  and  $I_{\text{back,per}}$  denoting the backscattered intensities parallel and perpendicular to the incident polarisation state of the light beam. The backscattering linear depolarisation ratio  $\delta$  is then obtained by  $\delta = I_{\text{back,per}}/I_{\text{back,par}}$ .  $\delta$  is zero for light scattering by spherical cloud droplets and, apart from very small or highly aspherical particles, different from zero for light scattering by aspherical ice crystals, thereby featuring a strong dependency on the particle size and shape (Mishchenko et al., 1996; Zakharova and Mishchenko, 2000). Infrared extinction of the cloud particles is measured between 6000 and 800  $\text{cm}^{-1}$  at 4  $\text{cm}^{-1}$  resolution with a FTIR spectrometer that is coupled to an internal White-type multiple reflection cell. Fitting algorithms are used to retrieve the number size distributions of the cloud particles from their measured infrared extinction spectra as outlined in Wagner et al. (2006a), thereby exploiting the different spectral habitus of the absorption bands of liquid water droplets and ice crystals to detect the glaciation of a droplet cloud (Zasetsky et al., 2005). The light scattering and infrared measurements are also valuable tools for the characterisation of the seed aerosol particles with respect to e.g. their phase (solid or liquid) and chemical composition, as will be repeatedly shown in the current manuscript.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 2.2 Aerosol generation and characterisation

In the present study, we have investigated the ternary systems sodium chloride/oxalic acid/water ( $\text{NaCl/OA/H}_2\text{O}$ ) and sulphuric acid/oxalic acid/water ( $\text{H}_2\text{SO}_4/\text{OA/H}_2\text{O}$ ) as seed aerosols in expansion cooling experiments.  $\text{NaCl/OA/H}_2\text{O}$  aerosol particles of three different NaCl: OA mass ratios were generated from aqueous solutions of 5 wt% NaCl and 0.07, 0.36, and 0.51 wt% OA, prepared by dissolving sodium chloride (Merck, >99.5%,  $58.44 \text{ g mol}^{-1}$ ) and oxalic acid dihydrate (Merck, >99.5%,  $126.07 \text{ g mol}^{-1}$ ) in Milli-Q quality water. In two additional experiments, a pure salt solution of 5 wt% NaCl and a quaternary solution of 5 wt% NaCl, 0.36 wt% OA, and 0.1 wt% succinic acid (SA, Merck, >99.5%,  $118.09 \text{ g mol}^{-1}$ ) were employed. The solutions were filtrated by a submicron syringe filter (PTFE) to remove any undissolved particulates and were subsequently injected into the AIDA chamber at  $T = 244 \text{ K}$  and  $\text{RH} = 70\%$  with an ultrasonic nebuliser (GA2400, Sinaptec). As already known from previous crystallisation experiments with supersaturated salt solution droplets, the ensemble of injected aqueous solution droplets does not remain in a supercooled liquid state in AIDA under such conditions but gradually crystallises on a time scale of several hours (see Sect. 3.1). The size distribution of the so-generated mixed effloresced particles of sodium chloride and oxalic acid was measured with a scanning mobility particle sizer (SMPS, TSI) and an aerodynamic particle spectrometer (APS, TSI). In all experiments the median particle diameter was about  $0.7 \mu\text{m}$ ; an exemplary size distribution measurement of pure effloresced NaCl particles is shown in Fig. 3 (left panel). Typical number concentrations, obtained after an overall injection period of about 30 min and measured with a condensation particle counter (CPC3010, TSI), were in the range from  $400$  to  $600 \text{ cm}^{-3}$ .

In the  $\text{H}_2\text{SO}_4/\text{OA/H}_2\text{O}$  system, three different aqueous solutions with 5 wt%  $\text{H}_2\text{SO}_4$  as well as 0.51, 1.14, and 3.43 wt% OA were prepared, supplemented by a binary reference solution of 5 wt%  $\text{H}_2\text{SO}_4$ . The aqueous solutions were directly sprayed into the cloud chamber via a two-stream dispersion nozzle. An injection period of 5 s typically yielded aerosol number concentrations in the order of  $10^4$  particles  $\text{cm}^{-3}$ . The AIDA

### Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are, at most, only slightly above the threshold value where the onset of degradation has been observed at room temperature.

Concluding this section, Table 1 presents a summary of the ice nucleation experiments that were performed in the present study, serving as a look-up table for the detailed discussion in Sect. 3. The first column denotes the experiment number, the second and third columns list the employed solute species (NaCl, OA, SA, H<sub>2</sub>SO<sub>4</sub>) as well as their relative mass ratio, and the final column briefly describes the experimental procedure with the number of expansion cooling runs that were performed at various AIDA temperatures.

### 3 Results and discussion

#### 3.1 Ice nucleation studies in the ternary NaCl/OA/H<sub>2</sub>O system

This section comprises the analysis of the ice nucleation Exps. 1–5 from Table 1. We will first describe the hygroscopic behaviour and ice nucleation potential of pure effloresced NaCl particles (Exp. 1) in expansion cooling cycles that were started at 244 and 235 K before addressing the experiments in the ternary system. As indicated in the introduction, the solid NaCl particles were formed by crystallisation of aqueous solution droplets that were added to the AIDA chamber at 244 K and 70% RH with respect to supercooled water. The bottom left panel in Fig. 4 shows the trace of the depolarisation ratio  $\delta$  that was monitored upon spraying the aqueous solution of 5 wt% NaCl into the AIDA vessel with the ultrasonic nebuliser. With the start of aerosol addition,  $\delta$  immediately adopts a small background value of about 0.02 that is typically observed when spherical droplets are present in the chamber. The slight deviation from zero is due to Rayleigh scattering by air molecules, a less than 100% linear polarisation of the laser light, and minor misalignments of the polarisers in the optical set-up. Already during the injection period, as well as in the course of the following observation period of about 4 h,  $\delta$  continuously increases due to the formation of aspherical NaCl crystals. Efflorescence is also documented by the loss of the condensed-water

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





features in the simultaneously recorded infrared extinction spectra, shown in the top left panel of Fig. 4. The liquid water extinction bands are located at about  $3300\text{ cm}^{-1}$  (O-H stretching mode) and  $1640\text{ cm}^{-1}$  (H-O-H bending mode). Similar to the IR spectroscopic study on the efflorescence of NaCl by Cziczo and Abbatt (2000), anhydrous crystals of sodium chloride have formed instead of  $\text{NaCl}\cdot 2\text{H}_2\text{O}$  that is the stable crystalline solid below the  $\text{NaCl}/\text{NaCl}\cdot 2\text{H}_2\text{O}$  peritectic temperature of  $273.3\text{ K}$  (Koop et al., 2000). The latter species can be unambiguously identified by its prominent, highly structured infrared spectrum in the regime of the O-H stretching mode due to the hydration water (Schiffer and Hornig, 1961). We will discuss this issue in more detail later in this manuscript.

The observation of NaCl efflorescence at a comparatively high relative humidity of 70% is in contrast to a set of literature studies that have reported an efflorescence relative humidity of about 45%, see Table 1 in Gao et al. (2007). We have recently observed a similar discrepancy for the efflorescence behaviour of oxalic acid. Whereas several previous studies have reported on the difficulty to effloresce oxalic acid particles even at very low relative humidities of less than 5%, supercooled aqueous oxalic acid solution droplets readily crystallised in AIDA at  $244\text{ K}$  and 70% RH on a time scale of about one hour (Wagner et al., 2010). As outlined in our recent work, we will attempt to quantitatively analyse the crystallisation process in future chamber studies by exploiting the potential of a novel light scattering instrument (Small Ice Detector probe SID3, University of Hertfordshire) to determine the crystallisation rate. At present, we cannot completely exclude that the observed high efflorescence relative humidity might be due to heterogeneous crystallisation on suited sites in the chamber interior.

After an observation time of about 4 h, the major part of the volume fraction of the injected  $\text{NaCl}/\text{H}_2\text{O}$  solution droplets had crystallised and the generated solid NaCl particles were probed on their hygroscopic behaviour and ice nucleability in an expansion cooling experiment at  $244\text{ K}$  (Fig. 5, left panel). The AIDA records from the cooling cycle are arranged in the following order. In the top row panel, the time evolution of the AIDA pressure (black line) and the mean AIDA gas temperature (red line) is shown.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The second row depicts time series of the saturation ratios with respect to ice ( $S_{\text{ice}}$ , black line) and supercooled water ( $S_{\text{liq}}$ , red line) from the TDL water vapour absorption measurements. As a comparison, the trace for  $S_{\text{ice}}$  that was calculated for the ex situ measured total water concentration is shown as blue line. In the absence of any clouds of supercooled water droplets and/or ice crystals and provided that any aerosol water content, if present, can be neglected in comparison with the interstitial water vapour concentration, both traces for  $S_{\text{ice}}$  coincide. In the third row panel, the scatter plot from the OPC2 is shown where each dot represents a single particle count event that was classified into one of the OPC2 size channels. Whereas the size of spherical particles of known refractive index is exactly determined, the size information for aspherical particles is ambiguous due to the dependence of the phase function on the particle shape and orientation in the detection volume. Both employed OPCs detect the pulses of white light that was scattered by single particles at side-scattering angles from  $78\text{--}102^\circ$  and correlate the measured scattering intensity with the particle diameter assuming Mie theory. In the side-scattering regime, the values of the normalised phase function for aspherical particle habits are usually higher than those for equivalent spheres whose phase function features a prominent minimum in this regime (Liu et al., 2006). For this reason, the size of aspherical particles is typically overestimated in the OPC scatter plots by up to a factor of two. A population of deliquesced NaCl solution droplets will therefore be classified by the OPC2 at smaller diameters than an ensemble of crystalline NaCl particles. Large aspherical ice particles that have nucleated in AIDA during an expansion cooling cycle can be distinguished from small interstitial aerosol particles and/or cloud droplets by introducing an optical threshold size. Adding all particle count events above that threshold size yields the ice particle number concentration  $N_{\text{ice}}$ . The ice-active number fraction  $f_{\text{ice}}$  of the aerosol population, obtained by further dividing  $N_{\text{ice}}$  through the prevailing aerosol number concentration, is shown in fourth row panel of Fig. 5. The bottom panel contains time series of the two records  $I_{\text{back,par}}$  (black line) and  $\delta$  (red line) from the laser light scattering and depolarisation measurements.

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Shortly after the onset of pumping at time zero, when surpassing a relative humidity ( $S_{liq}$ ) of  $(72 \pm 3)\%$ , the depolarisation ratio  $\delta$  drops to its background value of 0.02, indicating the deliquescence transition of the NaCl crystals to form saturated aqueous solution droplets. The sudden water uptake leads to an increase in the size of the particles which manifests itself in a stepwise increase in the backscattered intensity  $I_{back,par}$ . In contrast, for the reason outlined above, the deliquesced solution droplets are counted by the OPC2 at smaller diameters than the effloresced NaCl crystals. Note that the OPC2 only records the tail of the aerosol size distribution towards large particle diameters whereas micron-sized cloud droplets and ice crystals are quantitatively counted. The observed deliquescence relative humidity agrees with low-temperature deliquescence measurements for NaCl in a flow cell apparatus and falls onto the extrapolated line for anhydrous NaCl to low temperatures, indicating that indeed NaCl has formed upon efflorescence instead of  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  (Koop et al., 2000). Upon further expansion cooling, the deliquesced NaCl solution droplets further grow by the uptake of water vapour from the gas phase, as can be seen by the continuous increase in  $I_{back,par}$  after the deliquescence step. When reaching saturation with respect to super-cooled water at about  $t = 70$  s (first vertical green line), the seed aerosol particles are activated to a droplet cloud that rapidly grows into the size range of the OPC2 and is evident as a dense point cloud with diameters of up to  $10 \mu\text{m}$ . The droplet cloud is maintained for a very long time period in AIDA until at  $t = 240$  (second green line) the gas temperature has further dropped below the homogeneous freezing temperature of pure water droplets during the continued expansion. The subsequent glaciation of the droplet cloud provokes an increase of the depolarisation ratio and a decrease of the relative humidity to  $S_{ice} = 1$ . The scatter signals of the nucleated ice crystals appear as a point cloud at very large diameters in the OPC2 scatter plot. Before  $t = 240$  s, only very few large ice crystals with a number concentration of less than  $0.1 \text{ cm}^{-3}$  are detected by the OPC2, probably formed by nucleation on the small number of background aerosol particles that were present in the chamber. After the stop of expansion cooling at  $t = 370$  s, the ice crystals evaporate on a time scale of about 400 s. The

largest particles may also be lost by sedimentation.

After the evaporation of the ice cloud, the AIDA chamber was slowly cooled to 235 K over a time period of 3 h (Fig. 6) and another expansion cooling cycle with the same aerosol loading was started whose records are shown in the left panel of Fig. 7. At the end of the first expansion cooling cycle ( $t = 1000$  s, Fig. 5), the depolarisation ratio  $\delta$  had again temporarily returned to the background value that was monitored at the beginning of aerosol injection as well as immediately after the deliquescence transition in the early stage of the expansion experiment. While cooling the AIDA chamber to 235 K, however, the deliquesced NaCl solution droplets again crystallised, leading to an increase of  $\delta$  to 0.26 before the second expansion cooling cycle was started, as shown in Fig. 6. In the following discussion, we will refer to this time period as the second crystallisation period whereas the time period of crystallisation immediately after aerosol injection will be called the first crystallisation period. During the second expansion run, the NaCl crystals again deliquesce at  $RH = (72 \pm 3)\%$  (for the sake of clarity, as explained in the figure caption, the trace for  $S_{liq}$  is not explicitly shown). The so-generated supercooled solution droplets further take up water until at  $t = 130$  s their critical dilution has been surpassed and homogeneous freezing occurs at  $S_{ice} = 1.38$  and  $T = 231$  K. The nucleated ice crystals rapidly deplete the excess of water vapour from the gas phase, thereby limiting the maximum of the ice saturation ratio,  $S_{ice,max}$ , to a value of 1.40. Due to the lower AIDA temperature in the second expansion cooling cycle, the chamber air was already slightly supersaturated with respect to ice before the deliquescence transition of NaCl had occurred. The solid NaCl crystals, however, are not ice-active in the deposition mode at low ice supersaturations because early ice formation was not detected. This is the prerequisite for the experiments in the ternary NaCl/OA/H<sub>2</sub>O system on the immersion freezing potential of oxalic acid that will be discussed in the following. When performing expansion cooling cycles with mixed effloresced NaCl/OA particles, the surrounding salt matrix will not be active as a heterogeneous ice nucleus but will undergo deliquescence to form aqueous NaCl solution droplets with embedded OA crystallites whose immersion freezing potential can

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



then be investigated. The right panels of Figs. 4, 5, and 7 are duplicates of their left-hand counterparts for an experiment where, instead of a pure 5 wt% NaCl solution, a ternary solution with 5 wt% NaCl and 0.51 wt% OA was sprayed into the AIDA chamber at 244 K and 70% RH (Exp. 2a in Table 1). As the solubility of OA is significantly decreased in NaCl solutions compared to pure water, this solution was almost saturated with OA at room temperature.

As shown in the right panels of Fig. 4, the ensemble of injected ternary solution droplets continuously crystallised on a time scale of several hours, similarly to the binary NaCl/H<sub>2</sub>O system from Exp. 1. The enlarged frame in the panel with the FTIR spectra recordings shows the subset in the regime of the C-O stretching mode of oxalic acid. The vertical line indicates the peak position of this transition in pure oxalic acid dihydrate crystals at around 1270 cm<sup>-1</sup>. Interestingly, the observed band position does not shift from its original location of 1240 cm<sup>-1</sup> (characteristic of aqueous oxalic acid) to 1270 cm<sup>-1</sup> during crystallisation. The results from the expansion cooling cycles will clearly evidence that mixed effloresced particles of sodium chloride and oxalic acid crystals have formed. So either the surrounding NaCl matrix strongly effects the peak position of the C-O vibrational mode in oxalic acid dihydrate or a different species has formed, e.g. anhydrous  $\beta$  oxalic acid crystals whose C-O stretching mode is slightly shifted to lower wavenumbers compared to oxalic acid dihydrate and thus approximately overlaps with the band position in aqueous oxalic acid (Bellamy and Pace, 1963; Braban et al., 2003). Upon starting the first expansion cooling cycle at 244 K (Fig. 5, right panel), the internally mixed NaCl/OA particles undergo a deliquescence transition at  $(72 \pm 3)$  % RH, i.e., the deliquescence behaviour of NaCl remains unchanged in the mixture with oxalic acid. Such behaviour has been typically observed for mixtures of inorganics and slightly water soluble dicarboxylic acids, such as mixtures of ammonium sulphate and oxalic, succinic, or adipic acid (Brooks et al., 2002), mixtures of succinic acid and sodium chloride (Choi and Chan, 2002) or ammonium nitrate (Lightstone et al., 2000), as well as for mixtures of oxalic or succinic acid and a five-component combination of high water-soluble dicarboxylic acids (Marcolli et al.,

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2004). At the deliquescence point, the inorganic fraction (or the mixture of high water-soluble dicarboxylic acids) forms a droplet, containing only a minor weight fraction of the less water soluble organic acid that gets dissolved according to the eutonic composition. The best part of the slightly water soluble organic compound remains as a solid inclusion within the droplet. Evidence for such solid, non-spherical inclusion has been provided by light scattering measurements on mixed particles of ammonium nitrate/succinic acid (25% succinic acid by mass) that were levitated in an electrodynamic balance (Lightstone et al., 2000). Also at elevated humidity levels well above the deliquescence point of ammonium nitrate, the recorded light scattering signal still contained the random high-frequency signal characteristic of solid, nonspherical particles; see Fig. 5 in Lightstone et al. (2000).

Concerning the internally mixed NaCl/OA particles with only 10% OA by mass from Exp. 2a, the sizes of the embedded oxalic acid crystals that remain undissolved after the deliquescence of NaCl are too small to provoke any change in the depolarisation ratio in relation to the background value of 0.02 for completely liquid droplets that was observed after the deliquescence transition in Exp. 1. The solid OA inclusions, however, significantly affect the ice nucleability of the solution droplets. Almost immediately after the seed aerosol particles have become activated into cloud droplets at water saturation ( $t = 75$  s, vertical green line), the droplet cloud glaciates at a temperature of 241 K, i.e., well above the homogeneous freezing temperature of pure water droplets at which ice nucleation has started in Exp. 1. Note that the cloud droplets in Exp. 2a have barely grown into the measurement range of the OPC2 (as highlighted by the green circle in the scatter plot) before the ice cloud is formed whereas the droplet cloud in Exp. 1 persists for a much longer time period so that the water droplets can grow to much larger diameters during expansion cooling. This behaviour is further illustrated by the two series of FTIR spectra recordings that are shown in Fig. 8. The spectra are arranged from bottom to top and were recorded at a rate of 6 spectra/min during the time periods that are indicated as grey horizontal bars on the time scales of the OPC2 scatter plots in Fig. 5. The first six spectra recordings from Exp. 2a reflect the water

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

uptake and droplet activation of the NaCl/OA particles after the deliquescence transition, as evidenced by the growth of the liquid water extinction band at  $3300\text{ cm}^{-1}$ . The signature of the sixth spectrum can be accurately reproduced with a Mie fit using the optical constants of supercooled water (blue line), yielding a median droplet diameter of  $2.0\text{ }\mu\text{m}$  (Zasetsky et al., 2005). Twenty seconds later, the recorded infrared spectrum already bares the signature of a mixed phase cloud. Superimposed on the liquid water extinction bands are two pronounced extinction minima at  $3500$  and  $950\text{ cm}^{-1}$ , the so-called Christiansen bands of large ice crystals (Arnott et al., 1995). The spectrum can only be accurately mimicked by a bimodal fit (red line), with one mode of  $2.7\text{ }\mu\text{m}$  median diameter for water droplets and a second mode of  $10\text{ }\mu\text{m}$  median (equal-volume sphere) diameter for ice crystals. Another ten seconds later, the extinction features due to liquid water have almost disappeared and a pure ice cloud has remained. In contrast, the spectral signatures of cloud droplets can be monitored for a much longer time period during Exp. 1. The spectra series displays the growth of the cloud droplets during expansion cooling, starting with a median diameter of  $1.9\text{ }\mu\text{m}$  (blue line, Mie fit of the lowermost spectrum) and ending with a diameter of  $6.5\text{ }\mu\text{m}$  (red line, Mie fit of the uppermost spectrum) prior to glaciation at the homogeneous freezing temperature.

The comparison of the AIDA records from Exp. 1 and Exp. 2a underlines that the embedded OA crystals act as very efficient heterogeneous ice nuclei in the condensation mode. A significant number fraction,  $f_{\text{ice}}$ , of about 10% of the overall aerosol population has formed ice in the expansion cooling cycle. Note that ice crystals which have formed early on by the most ice-active immersed OA crystals rapidly consume the unfrozen supercooled cloud droplets via the Bergeron-Findeisen process. In that way, immersed OA particles with a slightly lower ice activity will be prevented from contributing to  $f_{\text{ice}}$ . Obviously, the OA inclusions do not get instantly and completely dissolved during CCN activation and subsequent droplet growth but can trigger the glaciation of the droplet cloud at a temperature much higher than monitored in Exp. 1. Such incomplete dissolution of oxalic acid has also been observed in our previous study; see e.g. Fig. 14 in Wagner et al. (2010) and the related discussion. Note that during Exp. 2a pumping

was stopped shortly after the droplet cloud had glaciated and  $S_{\text{ice}}$  had levelled off at a value of one. Thereby, the gas temperature had always been well above the freezing temperature from Exp. 1.

In the same way as in Exp. 1, the AIDA chamber was slowly cooled to 235 K after the ice cloud from the first expansion cooling cycle of Exp. 2a had evaporated. The depolarisation ratio  $\delta$  had not completely levelled off at the background value of 0.02 after the evaporation of the ice cloud, indicating that a fraction of the NaCl/OA/H<sub>2</sub>O aerosol population had already started to effloresce. As discussed below, the crystallisation of the particle ensemble in the second crystallisation period might be facilitated by the immersed OA crystals that have remained as solid particulates within the solution droplets. During cooling,  $\delta$  increased to a value of 0.28 (Fig. 6). The AIDA records from the second expansion cooling cycle (Fig. 7, right panel) reveal two pronounced differences compared to Exp. 1. Firstly, the ensemble of NaCl/OA particles undergoes a dual deliquescence transition at 72 and 82% RH, as nicely documented by the two-step decrease in the depolarisation ratio. Secondly, the OA crystals that remain immersed in the aqueous NaCl solution droplets upon deliquescence induce earlier ice formation in comparison with the homogeneous freezing run from Exp. 1. The critical ice saturation ratio,  $S_{\text{ice,crit}}$ , is reduced from 1.38 to 1.32, thereby provoking that during Exp. 2a the peak relative humidity stays below the homogeneous freezing threshold from Exp. 1, as indicated by the horizontal green line. Similarly to the first expansion run at 244 K,  $f_{\text{ice}}$  amounts to about 10%. Our results thereby confirm the findings from Zobrist et al. (2006) that oxalic acid acts a heterogeneous ice nucleus. A detailed comparison with the Zobrist et al. (2006) results is given later on after the discussion of Exps. 2b, 3, 4, and 5. We now focus on the deliquescence behaviour of the mixed NaCl/OA particles.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The additionally observed deliquescence transition at 82% RH approximately falls onto the expected deliquescence curve of  $\text{NaCl} \cdot 2\text{H}_2\text{O}$ , as constructed in the temperature – RH plane of the  $\text{NaCl}$ - $\text{H}_2\text{O}$  phase diagram by connecting the deliquescence relative humidity line of  $\text{NaCl}$  at the  $\text{NaCl}/\text{NaCl} \cdot 2\text{H}_2\text{O}$  peritectic temperature of 273.3 K to the ice melting point curve at the ice/ $\text{NaCl} \cdot 2\text{H}_2\text{O}$  eutectic temperature of 251.9 K; see Fig. 6 in Koop et al. (2000). Therefore, the aerosol population before the second cooling cycle in Exp. 2a has consisted of an external mixture containing both  $\text{NaCl}/\text{OA}$  as well as  $\text{NaCl} \cdot 2\text{H}_2\text{O}/\text{OA}$  crystals. Note that prior to the second crystallisation period of Exp. 2a after performing the expansion cooling cycle at 244 K, the salt solution droplets already contained a solid nucleus of oxalic acid because, as discussed above, the oxalic acid crystals did not get completely dissolved on the timescale of the expansion cooling experiment. In contrast, both the first and second crystallisation periods of Exp. 1 and the first crystallisation period of Exp. 2a have involved entirely liquid solution droplets. As evident from Fig. 6, the rate of increase of the depolarisation ratio  $\delta$  during the second crystallisation period of Exp. 2a was significantly higher than during the respective crystallisation period of Exp. 1, indicating that a different, more efficient crystallisation mechanism was involved. We therefore speculate that the partial formation of  $\text{NaCl} \cdot 2\text{H}_2\text{O}/\text{OA}$  particles in the second crystallisation period of Exp. 2a was induced by heterogeneous nucleation on the surface of the embedded oxalic acid crystals. On the contrary, homogeneous crystallisation obviously leads to the formation of anhydrous sodium chloride. That heterogeneous crystallisation favours the formation of  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  is in agreement with observations from Koop et al. (2000). Whereas  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  did not form in their flow cell efflorescence experiments (homogeneous crystallisation), the precipitation of ice and  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  was detected in additional freezing and melting experiments with bulk and emulsion samples of aqueous  $\text{NaCl}$ . Individual freezing peaks for ice and  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  were detected for low  $\text{NaCl}$  concentrations whereas above threshold concentrations of 10 wt%  $\text{NaCl}$  (emulsion samples) or 16 wt%  $\text{NaCl}$  (bulk samples), single freezing peaks were detected in the cooling experiments, indicating that  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  instantly formed upon ice

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

nucleation, probably induced by heterogeneous nucleation on the ice crystal surface. In our experiments, it is not the ice particle surface but the surface of the immersed OA crystals that most likely triggers the precipitation of  $\text{NaCl}\cdot 2\text{H}_2\text{O}$  within a certain number fraction of the  $\text{NaCl}/\text{OA}/\text{H}_2\text{O}$  aerosol particles. Instantaneous crystallisation of  $\text{NaCl}\cdot 2\text{H}_2\text{O}$  triggered by ice formation, as discovered by Koop et al. (2000) in the experiments with bulk and emulsion samples, is difficult to detect in our expansion experiments and should have also been observed during ice cloud formation in Exp. 1 with binary  $\text{NaCl}/\text{H}_2\text{O}$  solution droplets. If immediate, ice-mediated crystallisation of  $\text{NaCl}\cdot 2\text{H}_2\text{O}$  had occurred in all aerosol particles that have nucleated ice in the first expansion run of Exp. 1, the overall aerosol population would already have contained a 10% number fraction of crystalline particles directly after the ice cloud had evaporated. If so, the depolarisation ratio would not have returned to its background value of 0.02 after ice cloud formation and evaporation in this expansion cooling cycle (Fig. 5, left panel). Obviously, the crystallisation efficiency of freeze-concentrated  $\text{NaCl}$  solutions in contact with ice particles to form  $\text{NaCl}\cdot 2\text{H}_2\text{O}$  particles is much lower in our experiments, meaning that the number fraction of particles that have actually crystallised via this process is too low to provoke a significant increase in the depolarisation ratio. We have discussed this issue in such detail because it closely relates to the experiments that are presented in Sect. 3.2 where we want to study the efficiency of the ice surface mediated precipitation of oxalic acid dihydrate from ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solutions. The different observations for the efficiency of  $\text{NaCl}\cdot 2\text{H}_2\text{O}$  formation by heterogeneous crystallisation on ice in either freezing experiments with emulsion samples (Koop et al., 2000) or ice nucleation experiments with airborne particles (AIDA) already support our introductory comments that it will be difficult to replicate the findings from the Zobrist et al. (2006) emulsion freezing experiments in AIDA expansion cooling cycles.

With the procedure of Exp. 2a, we could only indirectly prove the precipitation of  $\text{NaCl}\cdot 2\text{H}_2\text{O}/\text{OA}$  in the second crystallisation period by the observation of the second deliquescence step at 82% RH. Independent evidence from the FTIR spectra recordings was not available because the cooling of the AIDA chamber to 235 K induced a



cannot conclude which species of oxalic acid has formed during crystallisation of the ternary NaCl/OA/H<sub>2</sub>O solution droplets.

Having probed the heterogeneous ice nucleability of oxalic acid in a 0.51:5 mass ratio mixture with NaCl, the experiments were repeated with two ternary solutions of lower OA content (Exps. 3 and 4). Figure 10 compares the respective AIDA records from the first expansion cooling cycles at 244 K. The records for Exp. 3 (0.36:5 mass ratio of OA:NaCl) closely reproduce the data from Exp. 2a (Fig. 5, right panel), i.e., the ice cloud is formed almost instantaneously as the seed aerosol particles are activated to micron-sized cloud droplets at water saturation. In contrast, heterogeneous ice formation is almost absent in Exp. 4 (0.07:5 mass ratio of OA:NaCl) where only very few early ice crystals are detected by the OPC2. Instead, the droplet cloud glaciates at the homogeneous freezing temperature. In order to observe heterogeneous condensation mode ice nucleation by immersed OA particles at temperatures higher than 237 K, the size of the embedded particulates must evidently exceed a certain threshold size that will be defined by the dissolution kinetics of OA under the given hydrodynamic conditions of the experiment. Treating the effloresced NaCl/OA as spheres of diameter 0.7  $\mu\text{m}$  (see Fig. 3, left panel), the diameter of the immersed OA particles from Exp. 4 can be roughly estimated to about 0.17  $\mu\text{m}$  ( $\rho_{\text{NaCl}} = 2.17 \text{ g cm}^{-3}$ ,  $\rho_{\text{OA}} = 1.9 \text{ g cm}^{-3}$ ). These small embedded OA crystallites apparently get completely dissolved during droplet activation and, therefore, cannot trigger the early glaciation of the droplet cloud. A similar calculation for Exp. 3 where heterogeneous ice formation was observed yields a diameter for the OA inclusion of about 0.30  $\mu\text{m}$ . During expansion cooling cycles started at temperatures below 237 K, however, the seed aerosol particles are not activated to almost pure water droplets and so even small OA inclusions will not dissolve and remain present as heterogeneous ice nuclei. This is evidenced by Fig. 11, showing the second expansion cooling cycles from Exp. 3 and Exp. 4 at 235 K. In both experiments, heterogeneous ice nucleation by the embedded OA crystals can be detected, reducing the critical ice saturation  $S_{\text{ice,crit}}$  from 1.38 (homogeneous freezing, see Exp. 1, denoted as horizontal green line in Fig. 11) to 1.31 (Exp. 3) and 1.33 (Exp. 4). As in Exp. 2a, the maximum

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ice saturation ratios encountered during the expansion runs remain below the homogeneous freezing threshold from the second expansion cooling cycle of Exp. 1. The deliquescence transition during Exp. 3 features the well-known dual deliquescence behaviour due to the presence of both NaCl/OA and NaCl·2H<sub>2</sub>O/OA crystals. In contrast, the deliquescence step due to NaCl·2H<sub>2</sub>O is almost absent in Exp. 4 since the major part of the aerosol population has already deliquesced at 72% RH. This behaviour can be well understood when considering that most NaCl/OA crystals have completely deliquesced in the first expansion cooling cycle of Exp. 4. Homogeneous crystallisation will then again lead to the formation of anhydrous sodium chloride. In Exp. 3 the embedded OA crystals do not get completely dissolved during the first expansion cooling cycle and can initiate the heterogeneous crystallisation of sodium chloride dihydrate in the second crystallisation period. In the final experiment (Exp. 5), we have investigated whether a small addition of succinic acid to the ternary NaCl/OA/H<sub>2</sub>O solution from Exp. 3 will influence the heterogeneous ice nucleability of oxalic acid, recalling that Zobrist et al. (2006) have found that succinic acid was not ice-active in the immersion mode. During Exp. 5, only a single expansion cooling cycle at 244 K was performed whose records are depicted in Fig. 12. The AIDA records from Exp. 5 are almost duplicates of the respective data sets from Exp. 3 (left panel of Fig. 10), underlining that small admixtures of succinic acid do not suppress the high heterogeneous ice nucleability of oxalic acid.

Table 2 summarises our findings on the hygroscopic and ice nucleation behaviour of the investigated NaCl/H<sub>2</sub>O, NaCl/OA/H<sub>2</sub>O, and NaCl/OA/SA/H<sub>2</sub>O aerosol particles. The data for the heterogeneous ice nucleation ability of immersed oxalic acid crystals in the condensation mode from the first expansion cooling cycles have their counterpart in emulsion freezing experiments within the binary oxalic acid/water system from Zobrist et al. (2006). In the second cooling cycles with emulsified aqueous solutions of 4.95, 9.76, and 19.87 wt% oxalic acid, the emulsified droplets contained a solid oxalic acid dihydrate nucleus embedded in saturated oxalic acid solutions. Due to the small solubility of oxalic acid at low temperatures, the authors have chosen pure water as a



**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

transition because the outcome of the chamber experiments might be affected by heterogeneous crystallisation. Considering, however, our interpretation that the precipitation of  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  indicates heterogeneous crystallisation, as also emphasised by Koop et al. (2000), the absence of any  $\text{NaCl} \cdot 2\text{H}_2\text{O}$  formation in the two crystallisation periods of pure  $\text{NaCl}$  solution droplets from Exp. 1 contradicts this hypothesis so that further measurements will be needed to clarify this issue. If solution droplets with immersed oxalic acid crystals are formed, our measurements indicate that at least a 10% number fraction of the aerosol population is ice-active, both in the expansion cooling cycles at 244 and 235 K. In view of the experiments presented in Sect. 3.2, this means that at least  $10 \text{ cm}^{-3}$  sulphuric acid solution droplets with an immersed OA crystal must be formed in the first homogeneous freezing expansion run with the supercooled  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets to yield an ice particle number concentration of about  $1 \text{ cm}^{-3}$  in the second expansion run due to heterogeneous ice nucleation which was identified as the threshold number concentration to clearly evidence this process. As outlined in the introduction, the number concentration of ice crystals formed by homogeneous freezing will be at most  $100 \text{ cm}^{-3}$ . Therefore, the precipitation of oxalic acid dihydrate from the  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solutions must occur in every tenth ice crystal. In the following chapter, we describe whether this alternative pathway for the nucleation of oxalic acid dihydrate crystals, as previously evidenced in the emulsion freezing experiments by Zobrist et al. (2006), can also be observed in the AIDA expansion cooling experiments.

### 3.2 Ice nucleation studies in the ternary $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$ system

As in the previous section, we first briefly describe the results from reference expansion cooling experiments with binary  $\text{H}_2\text{SO}_4/\text{H}_2\text{O}$  solution droplets at 244 and 235 K (Exp. 6, Fig. 13). The AIDA data shown in Fig. 13 in principle mimic those from the respective expansion cooling cycles in the binary  $\text{NaCl}/\text{H}_2\text{O}$  system (see left panels of Fig. 5 and Fig. 7), with the only exception that the  $\text{H}_2\text{SO}_4/\text{H}_2\text{O}$  expansion runs were directly started with supercooled solution droplets so that the initial deliquescence step

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

from crystalline NaCl particles to aqueous sodium chloride solution droplets is missing. By contrast, the supercooled sulphuric acid solution droplets immediately adjust to the increasing relative humidity after the start of pumping at time zero by the uptake of water from the gas phase. The concomitant increase in particle size is evident by the increase in the intensity  $I_{\text{back,par}}$  of the back-scattered laser light. When starting the expansion run at 244 K (left panel), the  $\text{H}_2\text{SO}_4/\text{H}_2\text{O}$  solution droplets are first activated to a droplet cloud when approaching water saturation at  $t = 70$  s (first vertical green line). The supercooled water droplets then homogeneously freeze at  $t = 230$  s (second vertical green line) when the median gas temperature has dropped below 237 K. In the expansion cooling experiment started at 235 K (right panel), the dilute sulphuric acid solution droplets homogeneously freeze at  $S_{\text{ice}} = 1.38$  ( $t = 150$  s, vertical green line).

The left and right panels of Fig. 14 show two repetitive expansion cooling runs with ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets having a sulphuric acid to oxalic acid mass ratio of 5:0.51 (Exp. 7). As outlined in Sect. 2, the aerosol particles were injected at 265 K and then cooled to 244 K. In the first expansion run (left panel), about  $60 \text{ cm}^{-3}$  ice particles were generated by homogeneous freezing of the droplet cloud ( $t = 220$  s, vertical green line) that had been formed in the early stage of the experiment upon approaching water saturation. Only very few ice crystals ( $N_{\text{ice}} < 0.1 \text{ cm}^{-3}$ ) have nucleated before the homogeneous freezing threshold, indicating that the aerosol population has remained in a supercooled liquid state upon cooling, i.e., immersed oxalic acid (dihydrate) crystallites that could have acted as heterogeneous ice nuclei were not present. After the ice cloud from the first expansion run had evaporated, a second expansion run was started at 244 K to investigate whether the surface of the afore generated ice crystals had triggered the crystallisation of oxalic acid dihydrate from the unfrozen  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solutions. Also in the second cooling cycle, however, ice formation was only observed after the droplet cloud had been cooled below the homogeneous freezing temperature of about 237 K. Early heterogeneous freezing of at least a small fraction of the cloud droplets (those containing a solid oxalic acid dihydrate inclusion), as observed during Exps. 2a and 3 in the NaCl/OA/ $\text{H}_2\text{O}$  system (see right panel of



**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fig. 5 and left panel of Fig. 10), could not be detected. Subsequently, the AIDA chamber was cooled to 235 K to probe whether a change in the ice nucleation behaviour could be detected in two repetitive expansion cooling cycles with the same aerosol loading at a lower temperature. The left panel of Fig. 15 shows the AIDA records from the first expansion cooling experiment at 235 K. The ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets homogeneously froze at  $S_{\text{ice}} = 1.38$  (vertical green line), yielding a peak ice particle number concentration of about  $80 \text{ cm}^{-3}$ . The AIDA data from the second expansion cooling cycle that was started after the evaporation of the ice cloud from the preceding expansion is shown on an expanded time scale in the right panel. Whereas in the first expansion cooling cycle a fast pumping speed ( $200 \text{ m}^3 \text{ h}^{-1}$ ) was used to produce a large number of ice crystals that could trigger the precipitation of oxalic acid dihydrate, the pumping speed and thus the humidification rate  $dS_{\text{ice}}/dt$  was reduced in the second cooling cycle in order to be more sensitive for the detection of a small number concentration of ice crystals that potentially would be formed before the homogeneous freezing threshold. Given the results from the benchmark expansion cooling experiments with immersed OA crystals in the  $\text{NaCl}/\text{OA}/\text{H}_2\text{O}$  system at 235 K, the onset relative humidity due to immersion freezing would be at around  $S_{\text{ice}} = 1.32$  (see Table 2). This threshold is marked by the first vertical green line in the right panel of Fig. 15. As already outlined in the introduction, only a very small subset of the overall aerosol population is expected to be able to act as immersion freezing nucleus in the  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  system. Therefore, the number concentration of generated ice crystals will be too low to instantly deplete the water vapour supersaturation during the expansion so that soon after surpassing the heterogeneous freezing limit the homogeneous freezing threshold at  $S_{\text{ice}} = 1.38$  will be reached (second vertical green line in the right panel of Fig. 15). Only during a short time period of about 20 s, whose duration can be adjusted by controlling the humidification rate, early heterogeneous ice nucleation by potentially precipitated oxalic acid dihydrate crystals can be expected. The AIDA records from Fig. 15 indicate that prior to the formation of the mode of homogeneously frozen solution droplets no early heterogeneous ice formation took place

( $N_{\text{ice}} \leq 0.1 \text{ cm}^{-3}$ ). Given the difficulty to detect a small heterogeneous mode of ice crystals just prior to the homogeneous freezing mode at the lower temperature, the chamber was warmed again to 244 K. At this temperature, it is easier to detect heterogeneous ice formation because the supercooled droplet cloud that is formed at water saturation persists for an elongated time period of up to 4 min before the gas temperatures has further dropped below the homogeneous freezing limit. But also in this expansion run (results not explicitly shown), early heterogeneous ice formation was not detected.

Besides the obvious conclusion that the efficiency of the ice surface mediated precipitation of oxalic acid (dihydrate) under the given experimental conditions was too low to become detectable, different explanations for the absence of heterogeneous ice formation should be considered. One cannot definitely exclude that a less ice-active species of solid oxalic acid has precipitated from the ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets compared to the experiments in the  $\text{NaCl}/\text{OA}/\text{H}_2\text{O}$  system, given that oxalic acid is known to exist in various crystalline forms (Bellamy and Pace, 1963; Villepin and Novak, 1978a, b; Wenger and Bernstein, 2007) and that also the formation of solid amorphous particles has been recently discussed (Mikhailov et al., 2009). As outlined in Sect. 3.1, the chemical nature of (potentially) precipitated OA inclusions could not be identified. Given the observations from Exp. 2a with particles of the same OA solute concentration (Fig. 5, right panel and the related discussion), these embedded particulates would also be too small to change the value of the depolarisation ratio in relation to the background level of 0.02 and, therefore, their presence cannot be verified by this technique. One might also suggest that the OA inclusions were so small that they had instantly dissolved upon droplet activation during the expansion runs at 244 K (similar to Exp. 4, see right panel of Fig. 10) and that the heterogeneous and homogeneous freezing thresholds for the experiments started at 235 K were just too close to each other to observe early heterogeneous nucleation in a small subset of the overall aerosol population. As indicated in Sect. 2, the injection of the  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets typically yielded very broad number size distributions centred at around

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

0.5  $\mu\text{m}$ . For the first two expansion runs conducted at 244 K (Fig. 14), it is not feasible to assess what size fraction of the activated aerosol particles has nucleated ice in order to estimate the diameter of potentially precipitated OA inclusions. Such estimate, however, can be done for the final expansion run at 244 K that was performed after the two homogeneous freezing runs at 235 K. As shown by modelling results on the homogeneous freezing of solution droplets in AIDA chamber experiments, only solution droplets within a narrow range of the overall size distribution will freeze, i.e., the largest droplets that can still instantaneously react to the changing relative humidity during the expansion; see e.g. Fig. 8 in Haag et al. (2003). This regime of droplet sizes is typically centred at about 1  $\mu\text{m}$ . Based on the AIM model, 1  $\mu\text{m}$ -sized ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets with a  $\text{H}_2\text{SO}_4:\text{OA}$  mass ratio of 5:0.51 (Exp. 7) have an oxalic acid content of 1.5 wt% at the threshold relative humidity for homogeneous freezing. With an estimated density  $\rho$  of 1.15  $\text{g cm}^{-3}$  for the dilute  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets (approximated by the densities of pure  $\text{H}_2\text{SO}_4/\text{H}_2\text{O}$  solutions from Luo et al., 1996), this would translate into a spherical OA inclusion with a diameter of about 0.2  $\mu\text{m}$  that might have precipitated from the ternary solution droplets. This size is only slightly larger than that estimated for the OA inclusions in Exp. 4 where early condensation mode ice nucleation has not been observed.

In a succeeding experiment within the  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  system, we have therefore more than doubled the OA solute concentration (Exp. 8, see Table 1). A solid OA inclusion, precipitated from a 1  $\mu\text{m}$ -sized ternary solution droplet during a homogeneous freezing expansion cooling run at 235 K, would now have a diameter of about 0.27  $\mu\text{m}$ . This is close to the inclusion size estimated for Exp. 3 where early condensation mode ice nucleation by immersed OA particulates was clearly detected (left panel of Fig. 10). At first, we have again performed two repetitive expansion cooling runs at 244 K. As in Exp. 7, the number concentration of ice crystals that were formed by early heterogeneous nucleation in the second freezing run did not exceed the background level of 0.1–0.2 particles  $\text{cm}^{-3}$ , i.e., was comparable to the ice crystal number concentration that has already been observed in the preceding expansion cooling run. Afterwards,



experiments (air-borne particles, limited residence time of the generated ice clouds), the efficiency by which ice crystals with a surrounding or encapsulated  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution trigger the precipitation of an ice-active oxalic acid nucleus is therefore at most in the order of 1%.

5 In the final experiment within the  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  system (Exp. 9), we have further increased the oxalic acid content in the ternary solution droplets, yielding a  $\text{H}_2\text{SO}_4:\text{OA}$  mass ratio of 5:3.43. This is similar to the 4:4 mass ratio that was employed in the corresponding emulsion freezing experiment by Zobrist et al. (2006); see Table 2 therein. The injected  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets, however, did not remain in a super-cooled liquid state upon cooling the AIDA chamber to 244 K, as shown by the increase  
10 in the depolarisation ratio that was monitored after the gas temperature had dropped below about 252 K (black trace in the top panel of Fig. 17). According to the AIM model, the ternary solution droplets are composed of about 19 wt%  $\text{H}_2\text{SO}_4$  and 13 wt% OA at 252 K under ice-saturated conditions (ice-coated chamber walls). Obviously, homogeneous crystallisation of oxalic acid (dihydrate) occurs for such elevated OA concentrations. The sizes of the OA inclusions are evidently large enough to provoke an increase in the depolarisation ratio and, furthermore, are large enough in order to not get instantly dissolved upon droplet activation during the expansion cooling run that was later on started at 244 K (lower part of Fig. 17). Upon formation of the droplet  
15 cloud at water saturation (as apparent by a denser point cloud in the OPC2 scatter plot for particles  $\leq 10 \mu\text{m}$  and the temporary decrease in the depolarisation ratio, both marked by green circles in Fig. 17), the immersed OA particulates immediately trigger the glaciation of the droplet cloud at  $T = 241 \text{ K}$  (vertical green line in Fig. 17), i.e., well above the homogeneous freezing temperature. This finding confirms the pronounced heterogeneous ice nucleability of immersed OA particulates that has already been observed in the  $\text{NaCl}/\text{OA}/\text{H}_2\text{O}$  system.  
20  
25

In the introductory section, we have stated that the most plausible explanation for the precipitation of oxalic acid dihydrate in the first cooling cycles of the Zobrist et al. (2006) emulsion freezing experiments is heterogeneous crystallisation triggered by

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the surface of the homogeneously nucleated ice crystals. In view of the low efficiency of this mechanism in the AIDA ice nucleation experiments, we suggest that crystallisation in the emulsion experiments might also be the result of the increased oxalic acid concentration in the freeze concentrated solution that coexists with the ice crystals after homogeneous freezing. Once a dilute, emulsified aqueous oxalic acid solution is cooled below the homogeneous freezing temperature in the first cooling cycle (e.g. below point B2 of trajectory C in Fig. 1), the solute concentration will follow the ice-melting point curve. The so-induced strong increase of the concentration of oxalic acid and of the supersaturation of oxalic acid dihydrate might increase the homogeneous crystallisation rate to such a degree to become detectable on the experimental time scale. This explanation would also account for the observations in the binary OA/H<sub>2</sub>O system from our previous study (Wagner et al., 2010). As outlined in Sect. 1, we have always observed the crystallisation of oxalic acid dihydrate when adding dilute oxalic acid solution droplets into the AIDA chamber at temperatures  $\leq 244$  K under ice-saturated conditions (ice-coated chamber walls). Under such conditions, the injected solution droplets instantly attain a solute concentration of  $>60$  wt% oxalic acid; see Fig. 1 in Wagner et al. (2010). In the emulsion freezing experiments with binary oxalic acid/water solutions, such a high solute concentration is also obtained after ice has formed in the first homogeneous freezing runs. The impact of ice formation on the crystallisation of oxalic acid dihydrate might therefore solely be related to the increase in the solute concentration instead of providing a heterogeneous surface for the crystallisation. Concerning the experiments in the ternary H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O system, we have detected the homogeneous crystallisation of oxalic acid dihydrate in aqueous solution droplets prepared from a dilute 5 wt% H<sub>2</sub>SO<sub>4</sub> and 3.43 wt% OA solution (Exp. 9) when injected at 265 K and cooled to 244 K at ice-saturated conditions (Fig. 17, top part). As addressed in Sect. 2.2, the solution droplets will concentrate to about 24.0 wt% H<sub>2</sub>SO<sub>4</sub> and 16.5 wt% OA after cooling to 244 K. Zobrist et al. (2006) have prepared a 4 wt% H<sub>2</sub>SO<sub>4</sub>/4 wt% OA ternary solution for their freezing measurements in this system. After ice formation in the first cooling cycle, the unfrozen solution in the emulsion experiment

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



will attain a similar composition as the ternary solution droplets in AIDA Exp. 9 before the expansion run at 244 K was started. Therefore, homogeneous crystallisation of oxalic acid dihydrate instead of ice-surface mediated heterogeneous crystallisation might have also happened in the emulsion freezing experiments. In AIDA Exps. 7 and 8 with a lower OA solute concentration, the oxalic acid dihydrate supersaturation and corresponding homogeneous crystallisation rate is obviously too low so that homogeneous crystallisation of oxalic acid dihydrate is not detectable at the time scale of the experiments. In contrast to the emulsion freezing experiments, the solute concentrations in AIDA expansion experiments are very similar at the start of the expansion runs (almost ice-saturated conditions) and shortly after ice formation when the ice supersaturation has again levelled off to an ice saturation ratio of one. Crystallisation of oxalic acid dihydrate can therefore only be induced heterogeneously by the surface of the nucleated ice crystals. The observed low efficiency of this process supports the assumption that the crystallisation of oxalic acid dihydrate in the emulsion freezing cycles might be due to homogeneous crystallisation of the freeze concentrated solutions. Zobrist et al. (2006) do not report additional freezing experiments in the ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  system with a lower OA concentration to validate this hypothesis.

## 4 Summary

We have presented measurements on the heterogeneous ice nucleation ability of oxalic acid in the immersion mode by performing controlled expansion cooling cycles with airborne particles in the ternary  $\text{NaCl}/\text{OA}/\text{H}_2\text{O}$  and  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  systems. The expansion runs were conducted in the AIDA aerosol and cloud chamber of the Karlsruhe Institute of Technology at initial temperatures of 244 and 235 K. Immersed oxalic acid particulates were observed to reduce the critical ice saturation ratio  $S_{\text{ice,crit}}$  required for the homogenous freezing of aqueous solution droplets at a temperature of around 231 K from 1.38 to about 1.32. Aqueous solution droplets with OA inclusions larger than about 0.27  $\mu\text{m}$  in diameter efficiently froze by condensation mode ice nucleation when

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



activated to cloud droplets at 241 K, i.e., well above the homogeneous freezing temperature of pure water droplets of about 237 K. The solid inclusions were presumably composed of oxalic acid dihydrate whose formation has been evidenced by infrared extinction spectroscopy in our preceding experiments on the crystallisation and ice nucleation behaviour of oxalic acid in the binary OA/H<sub>2</sub>O system (Wagner et al., 2010).

The present findings corroborate the assumption from a previous study that oxalic acid might play a considerable role in the Earth's climate system (Zobrist et al., 2006). Our experiments even disclose a slightly higher heterogeneous ice nucleation ability of immersed oxalic particles compared to the emulsion freezing runs conducted by Zobrist et al. (2006). For a closer inter-comparison of the ice nucleation results, however, it is indispensable to cover a wider range of temperatures in future AIDA chamber studies. The experiments in the ternary H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O system shed some light on the mechanism by which ice-active oxalic acid dihydrate crystals might nucleate in aqueous solution droplets. In supercooled sulphuric acid solution droplets with a high oxalic acid content of 13 wt%, the homogeneous crystallisation of oxalic acid dihydrate was observed. Ternary solution droplets with a lower OA mass concentration that did not crystallise homogeneously on the experimental time scale were processed in a series of low-temperature homogeneous freezing runs to investigate whether the surface of the nucleated ice crystals would trigger the precipitation of oxalic acid dihydrate, a potential mechanism to explain the results from the emulsion freezing experiments by Zobrist et al. (2006). We have shown that the efficiency of this crystallisation pathway was very low under our experimental conditions. For ice cloud residence times of typically 10–15 min, only at most one out of a hundred ice crystals that had been generated in a homogeneous freezing expansion run has triggered the precipitation of an ice-active oxalic acid nucleus from the surrounding H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O solutions. This was evidenced by the appearance of a small, early heterogeneous mode of ice crystals prior to the homogeneous freezing threshold in a succeeding expansion cooling run. We therefore suggest that the crystallisation of oxalic acid dihydrate in the emulsion freezing experiments might just be related to the strong increase of the oxalic acid



**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

dihydrate supersaturation upon ice formation, thereby enabling homogeneous crystallisation to become detectable on the time scale of the experiment. As an alternative pathway, mimicked by the experiments in the NaCl/OA/H<sub>2</sub>O system, oxalic acid dihydrate might also nucleate in a mixture of crystallisable inorganic salts when the relative humidity has dropped below the efflorescence point and the matrix of the inorganic particulates triggers its precipitation. As previously shown, oxalic acid dihydrate also readily crystallised from highly supersaturated binary oxalic water/water solution droplets at temperatures  $\leq 244$  K (Wagner et al., 2010). Our new results underline that, in addition to the repeatedly observed high ice nucleability of bare oxalic acid dihydrate crystals in the deposition mode (Kanji et al., 2008; Wagner et al., 2010), also imbedded organic crystals notably enhance the ice nucleation ability of aqueous solution droplets.

*Acknowledgements.* The authors are grateful to all members of the AIDA staff for their continuous support during the numerous successful measurement campaigns on ice nucleation which have been conducted at the chamber facility in the past decade. We thank Julian Skrotzki and Volker Ebert for the data analysis of the TDL system. The work has been funded by the Helmholtz-Gemeinschaft deutscher Forschungszentren as part of the program “Atmosphere and Climate”.

## References

- Arnott, W. P., Dong, Y. Y., and Hallett, J.: Extinction Efficiency in the Infrared (2–18  $\mu$ m) of Laboratory Ice Clouds – Observations of Scattering Minima in the Christiansen Bands of Ice, *Appl. Optics*, 34, 541–551, 1995.
- Baustian, K. J., Wise, M. E., and Tolbert, M. A.: Depositional ice nucleation on solid ammonium sulfate and glutaric acid particles, *Atmos. Chem. Phys.*, 10, 2307–2317, doi:10.5194/acp-10-2307-2010, 2010.
- Bellamy, L. J. and Pace, R. J.: Hydrogen Bonding in Carboxylic Acids.1. Oxalic Acids, *Spectrochim. Acta*, 19, 435–442, 1963.
- Braban, C. F., Carroll, M. F., Styler, S. A., and Abbatt, J. P. D.: Phase transitions of malonic and oxalic acid aerosols, *J. Phys. Chem. A*, 107, 6594–6602, 2003.

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Brooks, S. D., Wise, M. E., Cushing, M., and Tolbert, M. A.: Deliquescence behavior of organic/ammonium sulfate aerosol, *Geophys. Res. Lett.*, 29, 1917–1920, 2002.
- Choi, M. Y. and Chan, C. K.: The effects of organic species on the hygroscopic behaviors of inorganic aerosols, *Environ. Sci. Technol.*, 36, 2422–2428, 2002.
- 5 Clegg, S. L., Brimblecombe, P., and Wexler, A. S.: Thermodynamic model of the system  $\text{H}^+ - \text{NH}_4^+ - \text{SO}_4^{2-} - \text{NO}_3^- - \text{H}_2\text{O}$  at tropospheric temperatures, available at: <http://www.aim.env.uea.ac.uk/aim/aim.php>, *J. Phys. Chem. A*, 102, 2137–2154, 1998.
- Clegg, S. L. and Seinfeld, J. H.: Thermodynamic models of aqueous solutions containing inorganic electrolytes and dicarboxylic acids at 298.15 K. 1. The acids as nondissociating components, *J. Phys. Chem. A*, 110, 5692–5717, 2006a.
- 10 Clegg, S. L. and Seinfeld, J. H.: Thermodynamic models of aqueous solutions containing inorganic electrolytes and dicarboxylic acids at 298.15 K. 2. Systems including dissociation equilibria, *J. Phys. Chem. A*, 110, 5718–5734, 2006b.
- Cziczo, D. J. and Abbatt, J. P. D.: Infrared observations of the response of  $\text{NaCl}$ ,  $\text{MgCl}_2$ ,  $\text{NH}_4\text{HSO}_4$ , and  $\text{NH}_4\text{NO}_3$  aerosols to changes in relative humidity from 298 to 238 K, *J. Phys. Chem. A*, 104, 2038–2047, 2000.
- 15 Gao, Y. G., Chen, S. B., and Yu, L. E.: Efflorescence relative humidity of airborne sodium chloride particles: A theoretical investigation, *Atmos. Environ.*, 41, 2019–2023, 2007.
- Haag, W., Kärcher, B., Schaefers, S., Stetzer, O., Möhler, O., Schurath, U., Krämer, M., and Schiller, C.: Numerical simulations of homogeneous freezing processes in the aerosol chamber AIDA, *Atmos. Chem. Phys.*, 3, 195–210, doi:10.5194/acp-3-195-2003, 2003
- 20 Herz, W. and Neukirch, E.: Die Beeinflussung der Oxalsäurelöslichkeit durch starke Säuren, *Z. Anorg. Allg. Chem.*, 131, 303–309, 1923.
- Kanji, Z. A., Florea, O., and Abbatt, J. P. D.: Ice formation via deposition nucleation on mineral dust and organics: dependence of onset relative humidity on total particulate surface area, *Environ. Res. Lett.*, 3, 025004, doi:10.1088/1748-9326/3/2/025004, 2008.
- 25 Kawamura, K., Kasukabe, H., and Barrie, L. A.: Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in arctic aerosols: One year of observations, *Atmos. Environ.*, 30, 1709–1722, 1996.
- 30 Kerminen, V. M., Teinila, K., Hillamo, R., and Makela, T.: Size-segregated chemistry of particulate dicarboxylic acids in the Arctic atmosphere, *Atmos. Environ.*, 33, 2089–2100, 1999.
- Kerminen, V. M., Ojanen, C., Pakkanen, T., Hillamo, R., Aurela, M., and Merilainen, J.: Low-molecular-weight dicarboxylic acids in an urban and rural atmosphere, *J. Aerosol Sci.*, 31,

349–362, 2000.

Koop, T., Kapilashrami, A., Molina, L. T., and Molina, M. J.: Phase transitions of sea-salt/water mixtures at low temperatures: Implications for ozone chemistry in the polar marine boundary layer, *J. Geophys. Res. Atmos.*, 105, 26393–26402, 2000.

5 Koop, T., Ng, H. P., Molina, L. T., and Molina, M. J.: A new optical technique to study aerosol phase transitions: The nucleation of ice from  $\text{H}_2\text{SO}_4$  aerosols, *J. Phys. Chem. A*, 102, 8924–8931, 1998.

Lightstone, J. M., Onasch, T. B., Imre, D., and Oatis, S.: Deliquescence, efflorescence, and water activity in ammonium nitrate and mixed ammonium nitrate/succinic acid microparticles, *J. Phys. Chem. A*, 104, 9337–9346, 2000.

10 Liu, L., Mishchenko, M. I., Cairns, B., Carlson, B. E., and Travis, L. D.: Modeling single-scattering properties of small cirrus particles by use of a size-shape distribution of ice spheroids and cylinders, *J. Quant. Spectrosc. Rad.*, 101, 488–497, 2006.

Luo, B., Krieger, U. K., and Peter, T.: Densities and refractive indices of  $\text{H}_2\text{SO}_4/\text{HNO}_3/\text{H}_2\text{O}$  solutions to stratospheric temperatures, *Geophys. Res. Lett.*, 23, 3707–3710, 1996.

15 Marcolli, C., Luo, B. P., and Peter, T.: Mixing of the organic aerosol fractions: Liquids as the thermodynamically stable phases, *J. Phys. Chem. A*, 108, 2216–2224, 2004.

Mikhailov, E., Vlasenko, S., Martin, S. T., Koop, T., and Pöschl, U.: Amorphous and crystalline aerosol particles interacting with water vapor: conceptual framework and experimental evidence for restructuring, phase transitions and kinetic limitations, *Atmos. Chem. Phys.*, 9, 9491–9522, doi:10.5194/acp-9-9491-2009, 2009.

20 Mishchenko, M. I., Travis, L. D., and Mackowski, D. W.: T-matrix computations of light scattering by nonspherical particles: A review, *J. Quant. Spectrosc. Ra.*, 55, 535–575, 1996.

Murphy, D. M. and Koop, T.: Review of the vapour pressures of ice and supercooled water for atmospheric applications, *Q. J. Roy. Meteor. Soc.*, 131, 1539–1565, 2005.

25 Parsons, M. T., Mak, J., Lipetz, S. R., and Bertram, A. K.: Deliquescence of malonic, succinic, glutaric, and adipic acid particles, *J. Geophys. Res. Atmos.*, 109, D06212, doi:10.1029/2003JD004075, 2004.

Prenni, A. J., DeMott, P. J., Kreidenweis, S. M., Sherman, D. E., Russell, L. M., and Ming, Y.: The effects of low molecular weight dicarboxylic acids on cloud formation, *J. Phys. Chem. A*, 105, 11240–11248, 2001.

30 Pruppacher, H. R. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, Kluwer Academic Publishers, Dordrecht, 1997.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Schiffer, J. and Hornig, D. F.: On a Reported New Form of Ice, *J. Chem. Phys.*, 35, 1136–1137, 1961.
- Treuel, L., Schulze, S., Leisner, T., and Zellner, R.: Deliquescence behaviour of single levitated ternary salt/carboxylic acid/water microdroplets, *Faraday Discuss.*, 137, 265–278, 2008.
- 5 Villepin, J. D. and Novak, A.: Vibration-Spectra of Oxalic Acids.1. Infrared and Raman-Spectra of Beta-Phase of Some Acids, *Spectrochim. Acta A*, 34, 1009–1017, 1978a.
- Villepin, J. D. and Novak, A.: Vibration-Spectra of Oxalic Acids.2. Infrared and Raman-Spectra of Alpha-Phase of Some Acids, *Spectrochim. Acta A*, 34, 1019–1024, 1978b.
- 10 Wagner, R., Benz, S., Möhler, O., Saathoff, H., and Schurath, U., Probing ice clouds by broadband mid-infrared extinction spectroscopy: case studies from ice nucleation experiments in the AIDA aerosol and cloud chamber, *Atmos. Chem. Phys.*, 6, 4775–4800, 2006a, <http://www.atmos-chem-phys.net/6/4775/2006/>.
- Wagner, R., Bunz, H., Linke, C., Möhler, O., Naumann, K. H., Saathoff, H., Schnaiter, M., and Schurath, U.: Chamber Simulations of Cloud Chemistry: The AIDA Chamber, in: *Proceedings of the NATO Advances Research Workshop on Environmental Simulation Chambers: Application to Atmospheric Chemical Processes*, held in Zakopane, Poland, from 1 to 4 October 2004, edited by: Barnes, I. and Rudzinski, K. J., Springer, 2006b.
- 15 Wagner, R., Benz, S., Bunz, H., Möhler, O., Saathoff, H., Schnaiter, M., Leisner, T., and Ebert, V.: Infrared Optical Constants of Highly Diluted Sulfuric Acid Solution Droplets at Cirrus Temperatures, *J. Phys. Chem. A*, 112, 11661–11676, 2008.
- 20 Wagner, R., Linke, C., Naumann, K. H., Schnaiter, M., Vragel, M., Gangl, M., and Horvath, H., A review of optical measurements at the aerosol and cloud chamber AIDA, *J. Quant. Spectrosc. Ra.*, 110, 930–949, 2009.
- Wagner, R., Möhler, O., Saathoff, H., Schnaiter, M., and Leisner, T.: High variability of the heterogeneous ice nucleation potential of oxalic acid dihydrate and sodium oxalate, *Atmos. Chem. Phys.*, 10, 7617–7641, doi:10.5194/acp-10-7617-2010, 2010.
- 25 Wenger, M. and Bernstein, J.: Cocrystal design gone awry?, A new dimorphic hydrate of oxalic acid, *Mol. Pharm.*, 4, 355–359, 2007.
- Wexler, A. S. and Seinfeld, J. H.: Second-Generation Inorganic Aerosol Model, *Atmos. Environ.*, 25A, 2731–2748, 1991.
- 30 Zakharova, N. T. and Mishchenko, M. I.: Scattering properties of needlelike and platelike ice spheroids with moderate size parameters, *Appl. Optics*, 39, 5052–5057, 2000.
- Zasetsky, A. Y., Khalizov, A. F., Earle, M. E., and Sloan, J. J.: Frequency Dependent Complex

**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Refractive Indices of Supercooled Liquid Water and Ice Determined from Aerosol Extinction Spectra, *J. Phys. Chem. A*, 109, 2760–2764, 2005.

Zobrist, B., Marcolli, C., Koop, T., Luo, B. P., Murphy, D. M., Lohmann, U., Zardini, A. A., Krieger, U. K., Corti, T., Cziczo, D. J., Fueglistaler, S., Hudson, P. K., Thomson, D. S., and Peter, T.:

5 Oxalic acid as a heterogeneous ice nucleus in the upper troposphere and its indirect aerosol effect, *Atmos. Chem. Phys.*, 6, 3115–3129, doi:10.5194/acp-6-3115-2006, 2006.

Zuberi, B., Bertram, A. K., Koop, T., Molina, L. T., and Molina, M. J., Heterogeneous freezing of aqueous particles induced by crystallized  $(\text{NH}_4)_2\text{SO}_4$ , ice, and letovicite, *J. Phys. Chem. A*, 105, 6458–6464, 2001.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Compilation of the various ice nucleation experiments in the five investigated systems NaCl/H<sub>2</sub>O, NaCl/OA/H<sub>2</sub>O, NaCl/OA/SA/H<sub>2</sub>O, H<sub>2</sub>SO<sub>4</sub>/H<sub>2</sub>O, and H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O. The procedure for each experiment is discussed in detail in Sect. 3.

Exp.	Solute(s)	Mass ratio of solutes	Experimental procedure/ Number of expansion cooling cycles
1	NaCl		Aerosol addition at 244 K. Observation of crystallisation for about 5 h, followed by expansion cooling cycle at 244 K. Cooling of the chamber to 235 K and expansion cooling cycle at 235 K.
2a	NaCl, OA	5:0.51	Same procedure as in Exp. 1.
2b	NaCl, OA	5:0.51	Aerosol addition at 244 K. Observation of crystallisation for about 5 h, followed by expansion cooling cycle at 244 K. Again observation of crystallisation and another expansion cooling cycle at 244 K.
3	NaCl, OA	5:0.36	Same procedure as in Exp. 1.
4	NaCl, OA	5:0.07	Same procedure as in Exp. 1.
5	NaCl, OA, SA	5:0.36:0.1	Aerosol addition at 244 K. Observation of crystallisation for about 5 h, followed by expansion cooling cycle at 244 K. No further cooling as in Exp. 1.
6	H <sub>2</sub> SO <sub>4</sub>		Aerosol addition at 265 K, cooling of the chamber to 244 K and two repetitive expansion cooling cycles at 244 K. Cooling of the chamber to 235 K and expansion cooling cycle at 235 K.
7	H <sub>2</sub> SO <sub>4</sub> , OA	5:0.51	Aerosol addition at 265 K, cooling of the chamber to 244 K and two repetitive expansion cooling cycles at 244 K. Cooling of the chamber to 235 K and two repetitive expansion cooling cycles at 235 K. Warming of the chamber to 244 K and another expansion cooling cycle at 244 K.
8	H <sub>2</sub> SO <sub>4</sub> , OA	5:1.14	Aerosol addition at 265 K, cooling of the chamber to 244 K and two repetitive expansion cooling cycles at 244 K. Cooling of the chamber to 235 K and three repetitive expansion cooling cycles at 235 K. Warming of the chamber to 244 K and another expansion cooling cycle at 244 K.
9	H <sub>2</sub> SO <sub>4</sub> , OA	5:3.43	Aerosol addition at 265 K, cooling of the chamber to 244 K and two repetitive expansion cooling cycles at 244 K.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.

**Table 2.** Overview about the hygroscopic and ice nucleation behaviour of the investigated aerosol particles in the systems NaCl/H<sub>2</sub>O, NaCl/OA/H<sub>2</sub>O, and NaCl/OA/SA/H<sub>2</sub>O. For the first expansion runs that were started at 244 K, the deliquescence relative humidity (DRH) as well as the glaciation temperature  $T_{ice}$  of the generated droplet cloud are given. For the second expansion runs that were started at 235 K, the deliquescence relative humidity and the critical ice saturation ratio  $S_{ice,crit}$  at the onset of ice nucleation are given. Note that the second expansion run of Exp. 2b was also performed at 244 K, yielding a glaciation temperature of 241 K as in run 1.

Exp.	Solute(s)	Mass ratio of solutes	1st run DRH/%	1st run $T_{ice}$ /K	2nd run DRH/%	2nd run $S_{ice,crit}$
1	NaCl		72	237	72	1.38
2a	NaCl, OA	5:0.51	72	241	72 + 82	1.32
2b	NaCl, OA	5:0.51	72	241	72 + 82	
3	NaCl, OA	5:0.36	72	241	72 + 82	1.31
4	NaCl, OA	5:0.07	72	237	72*	1.33
5	NaCl, OA, SA	5:0.36:0.1	72	241		

\* As outlined in the discussion, only a very small second deliquescence step was observed at 82% RH.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.




**Fig. 1.** Schematic experimental trajectories of various ice nucleation experiments with supercooled aqueous sulphuric/oxalic acid solution droplets in the temperature – composition plane, adopted from Wagner et al. (2010). The abscissa denotes the total solute concentration (wt% sulphuric acid + wt% oxalic acid). The ice melting-point curve, shown as solid black line, was calculated for a sulphuric acid to oxalic acid mass ratio of 5:1.14 with the Extended *A/M* Aerosol Thermodynamics Model II (Clegg et al., 1998; Clegg and Seinfeld, 2006a, b). The right y-axis denotes the prevailing relative humidity with respect to supercooled water,  $RH_{liq}$ , for an environment that is saturated with respect to ice, i.e., when moving on the ice-melting point curve in the phase diagram. The red trajectory corresponds to the freezing experiment with emulsified solution droplets of 4 wt% sulphuric acid and 4 wt% oxalic acid by Zobrist et al. (2006) which homogeneously froze at 229.4 K in the first cooling cycle, as indicated by the grey circle. In lack of further freezing temperature measurements for different compositions in the ternary system, the dashed black line denotes the critical ice nucleation temperatures as a function of solute concentration for pure sulphuric acid solution droplets from the study by Koop et al. (1998). The blue and green trajectories represent expansion cooling experiments with airborne particles in the AIDA cloud chamber, see text for details.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.



**Fig. 2.** Schematic cross section of the AIDA facility with the main instrumentation for the ice nucleation experiments in the ternary NaCl/OA/H<sub>2</sub>O and H<sub>2</sub>SO<sub>4</sub>/OA/H<sub>2</sub>O systems.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.




**Fig. 3.** Left panel: Normalised number size distribution of effloresced NaCl particles from Exp. 1 (Table 1) obtained by combination of SMPS and APS measurements. Middle Panel: Infrared extinction spectra of binary aqueous sulphuric acid solution droplets (Exp. 6, trace a) and three ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets with  $\text{H}_2\text{SO}_4$  to OA mass ratios of 5:0.51 (trace b, Exp. 7), 5:1.14 (trace c, Exp. 8), and 5:3.43 (trace d, Exp. 9), recorded after spraying the respective aqueous solutions into the AIDA chamber at about 265K and 90% RH. The spectra are offset for clarity. Right panel: Difference spectra of the traces b, c, and d shown in the middle panel in relation to spectrum a in the regime of the C-O stretching mode of aqueous oxalic acid (enlarged y-axis compared to the middle panel).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode


R. Wagner et al.



**Fig. 4.** Bottom panels: Time series of the backscattering linear depolarisation ratio  $\delta$  after spraying aqueous solutions of 5 wt% NaCl (Exp. 1, Table 1, left panel) and 5 wt% NaCl + 0.51 wt% OA (Exp. 2a, right panel) into the AIDA chamber at 244 K with an ultrasonic nebuliser. The horizontal arrows denote the overall time period of aerosol injection ( $t_{inj}$ ). Time zero denotes the start of the first expansion cooling cycles that were performed later on to study the ice nucleability of the crystallised particles. The horizontal grey bars on the time scales denote the time periods of the FTIR spectra recordings that are shown in the top panels. The spectra are to be read from top to bottom and were recorded at a time interval of 20 min. The inset in the top right panel shows an enlarged view in the regime of the C-O stretching mode of oxalic acid.

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.



**Fig. 5.** Time series of various AIDA records during the first expansion cooling cycles from Exp. 1 and Exp. 2a that were started at 244 K. In each graph, time zero denotes the start of pumping (see Fig. 4). The five panels contain the following data. Panel 1: AIDA pressure (black line) and mean gas temperature (red line). Panel 2: Saturation ratios with respect to ice ( $S_{ice}$ , black and blue line) and supercooled water ( $S_{liq}$ , red line).  $S_{liq}$  is only shown for the gas-phase water concentration measured by TDL absorption whereas  $S_{ice}$  is plotted for both the gas-phase and total water concentration as measured ex situ with the MBW chilled mirror hygrometer. Panel 3: Size of individual particles detected by the OPC2. The horizontal grey bars denote the time periods of the FTIR spectra recordings that are shown in Fig. 8. Panel 4: Number fraction of ice-active aerosol particles. Panel 5: Backscattering intensity parallel to the incident polarisation state of the laser light (black line) and backscattering linear depolarisation ratio  $\delta$  (red line). All y-axes in adjacent sub-panels are identically scaled to facilitate the comparison of the records from Exp. 1 and Exp. 2a. See text for details.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion


**Heterogeneous ice nucleation ability of oxalic acid in the immersion mode**

R. Wagner et al.



**Fig. 6.** Time series of the backscattering linear depolarisation ratio  $\delta$  from Exp. 1 (black line) and Exp. 2a (blue line), recorded during the second crystallisation periods after the end of the first expansion cooling cycles (Fig. 5) while cooling the AIDA chamber to 235 K. The records end with the start of the second expansion cooling cycles which are shown in Fig. 7.


[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 7.** Time series of various AIDA records during the second expansion cooling cycles from Exp. 1 and Exp. 2a that were started at 235 K. Time zero again denotes the start of pumping and corresponds to the time stamp of the last records in the respective time series of the depolarisation ratio that are shown in Fig. 6. The individual panels contain the same measurements as in Fig. 5. In order to elucidate the subtle differences in the critical ice saturation ratio between the two experiments, the traces for  $S_{liq}$  are omitted to show the recorded ice saturation ratios on an enlarged y-axis scale in the second row panels. Except for the top-row panels, all y-axes of adjacent panels are identically scaled.


## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.



**Fig. 8.** Series of infrared extinction spectra recorded during the first expansion cooling cycles of Exp. 1 and Exp. 2a (black lines). The spectra are to be read from bottom to top and were recorded at a 10 s time interval during  $t = 70 - 200$  s (Exp. 1) and  $t = 30 - 110$  s (Exp. 2a), as indicated by horizontal grey bars on the time scales of the OPC2 scatter plots in Fig. 5. Various computed extinction spectra that best-fitted the measurements are shown as red and blue lines. Both coloured lines in the left panel as well as the blue trace in the right panel represent fits by unimodal log-normal number size distributions of supercooled water, yielding the indicated count median diameters  $D$ . The red line in the right panel represents a bimodal fit by two log-normal modes, with one mode for water droplets and the other for ice crystals. The latter were modelled as circular cylinders of aspect ratio 0.7 (Wagner et al., 2006a).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




**Fig. 9.** Compilation of infrared spectra for various chemical species in the NaCl/OA/H<sub>2</sub>O system. Traces a1 and a2: Reference spectra of NaCl·2H<sub>2</sub>O, obtained by digitising Fig. 1 from Schiffer and Hornig (1961). Trace b: Infrared extinction spectrum of effloresced NaCl·2H<sub>2</sub>O/OA recorded after the second crystallisation period of Exp. 2b. Trace c: Infrared extinction spectrum of effloresced NaCl/OA particles in an external mixture with a small fraction of unfrozen NaCl/OA/H<sub>2</sub>O solution droplets, recorded at the end of the first crystallisation period of Exp. 2a (lowermost spectrum from the series shown in the right panel in Fig. 4). Trace d: Infrared extinction spectrum of oxalic acid dihydrate crystals at 244 K from the study by Wagner et al. (2010).



## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.



**Fig. 10.** Time series of various AIDA records during the first expansion cooling cycles from Exp. 3 (OA:NaCl mass ratio: 0.36:5) and Exp. 4 (OA:NaCl mass ratio: 0.07:5) that were started at 244 K. The individual panels contain the same measurements as in Fig. 5.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.




**Fig. 11.** Time series of various AIDA records during the second expansion cooling cycles from Exp. 3 and Exp. 4 that were started at 235 K. The individual panels contain the same measurements as in Fig. 7. The horizontal green line connecting the second row panels denotes the critical ice saturation ratio for homogeneous freezing of aqueous NaCl solution droplets, as detected in the second expansion cooling cycle of Exp. 1 (Fig. 7, left panel).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode


R. Wagner et al.



**Fig. 12.** Time series of various AIDA records during the expansion cooling cycle from Exp. 5 that was started at 244 K. The individual panels contain the same measurements as in Fig. 5. The ice nucleation behaviour of the mixture of NaCl, oxalic acid, and succinic acid is very much the same as that of the NaCl + oxalic acid mixture in Exp. 3 (left panel of Fig. 10).

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.




**Fig. 13.** Time series of various AIDA records during two expansion cooling cycles from Exp. 6 that were started at 244 and 235 K. The individual panels show the same measurements as in Fig. 5, except from panel 4 where the ice particle number concentration  $N_{\text{ice}}$  instead of the ice-active fraction  $f_{\text{ice}}$  is shown.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.




**Fig. 14.** Time series of various AIDA records during two consecutive expansion cooling cycles from Exp. 7 that were started at 244 K. The individual panels show the same measurements as in Fig. 13.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.




**Fig. 15.** Time series of various AIDA records during two consecutive expansion cooling cycles from Exp. 7 that were started at 235 K. The individual panels show the same measurements as in Fig. 13. The y-axis scale in the fourth panel ( $N_{ice}$ ) of the second run was changed to logarithmic style in order to be more sensitive for small number concentrations of ice crystals that might have been formed by heterogeneous nucleation.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

## Heterogeneous ice nucleation ability of oxalic acid in the immersion mode

R. Wagner et al.



**Fig. 16.** Time series of various AIDA records during two expansion cooling cycles from Exp. 8 that were started at 244 K. The individual panels show the same measurements as in Fig. 13.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 17.** Top part: Time series of the depolarisation ratio  $\delta$  (black line) and the mean AIDA gas temperature (red line) after spraying ternary  $\text{H}_2\text{SO}_4/\text{OA}/\text{H}_2\text{O}$  solution droplets with a  $\text{H}_2\text{SO}_4$ :OA mass ratio of 5:3.43 (Exp. 9) into the aerosol vessel at  $t = -55000$  s ( $t_{\text{inj}}$ ). After aerosol addition, the chamber was cooled from 265 to 244 K, see text for details. Bottom part: Time series of various AIDA records during the first expansion cooling cycle from Exp. 9 that was started at 244 K. The individual panels show the same measurements as in Fig. 13.