

WRF-HMS: a fully-coupled regional atmospheric-hydrological modeling system for long-term scale applications

Sven Wagner^{1,2} (sven.wagner@kit.edu), Benjamin Fersch¹, Fei Yuan³, Zhongbo Yu³, Harald Kunstmann^{1,2}

¹Institute of Meteorology and Climate Research (IMK-IFU), KIT, Garmisch-Partenkirchen, Germany; ²Institute of Geography, Chair for Regional Climate and Hydrology, University, Augsburg, Augsburg, Germany; ³State Key Laboratory of Hydrology- Water Resources and Hydraulic Engineering, Hohai University, Nanjing, China

Motivation



- Studying the water and energy fluxes among the atmosphere, land surface and subsurface is important to understand the role of non-linear feedback within the hydrological cycle.
- Investigations of climate and land-use change impact on the regional water balance require **fully-coupled atmospheric-hydrological modeling systems**, which describe such feedback mechanisms and allow long-term simulations at climate-relevant scales.

Research Area

Poyang Lake catchment, China

- size: ~160,000 km²
- Tributary of Yangtze River
- Humid subtropical climate:
 - mean temperature: 17.5°C
 - annual precipitation: 1600 mm
- 5 main tributaries:

A: Waizhou
B: Lijadu
C: Meigang
D: Huashan
E: Wangjabu



Coupling Approach

- The regional atmospheric model **WRF-ARW** is coupled with the hydrological model **HMS**. Both models use the **Noah-LSM** and therefore share compatible water and energy flux formulations and communicate at the same spatial scale.
- 2-way interaction between groundwater and the Noah-LSM** is implemented by a Fixed-head or Darcy-flux method.
- The integration of the hydrological model HMS into WRF-ARW required primarily
 - a **hydrology driver routine** for the implementation of HMS into the WRF code structure,
 - MPI-parallelization of HMS** to run the coupled modelling system on HPC clusters,
 - changes to the pre-processing system, e.g. additional static surface and sub-surface hydrological parameters.
- In comparison to standard WRF simulations, the **fully coupled WRF-HMS model system** enables
 - lateral water flows at the surface and subsurface,
 - 2-way interaction between groundwater, the unsaturated zone, the land surface, and the atmosphere.
- The reasonable computational demand allows regional, long-term simulations for climate-relevant scales of multiyears, and longer time periods.


Step1&2: Stand-alone WRF & HMS simulations

WRF:

- Double nesting approach: D01 (30km), D02(10km)
- Reanalysis simulations using ERA interim:


HMS:

- Met. Forcing: station data



Step3: Fully coupled WRF-HMS simulations

- Results of fully coupled simulations and impact analysis of groundwater coupling on recharge (flux out of bottom layer of LSM) and further hydrometeorological and streamflow variables:



	Runoff (mm)	Recharge (mm)	SM top [-]	SM bottom [-]	ET (mm)	T [°C]	P (mm)
R-WRF-HMS	442	371	0.315	0.364	1141	18.0	1942
R-WRF-HMS-GW	381	467	0.304	0.330	1112	18.1	1941
Absolute Differences	-61	95	-0.01	-0.03	-29	0.1	-1
Relative Differences	-14%	26%	-3.5%	-9.5%	-3%	0.5%	-0.03%

Conclusions

- WRF-HMS enables a closed description of the water cycle at regional and long-term scale.**
- A **significant impact** on the regional water balance was found if **groundwater-unsaturated zone interaction is considered**. But the differences between the two groundwater coupling approaches are minor.
- For the fully coupled model system, **streamflow results strongly depend on the simulation quality for precipitation**.
- Two-way interaction results in net upward water fluxes in up to 25% of the basin area after the rainy season.
- In total, two-way interaction increases basin averaged recharge amounts.
- The evaluation with CPC and GLEAM indicates a **better performance of the fully coupled simulation**.
- The impact of groundwater coupling on LSM and atmospheric variables differs. Largest differences occur for the variable recharge (26%), whereas for atmospheric variables, the basin-averaged impact is minor (<1%).
- But locally, a **spatial redistribution up to 5% occurs for precipitation**.
- The **WRF-HMS model system is suitable** for many further applications, including **joint climate and land use change impact studies**, to investigate hydro-meteorological flux responses at basin scale for different climate regions worldwide.

Reference: Wagner, S., B. Fersch, F. Yuan, Z. Yu, and H. Kunstmann (2016), Fully coupled atmospheric-hydrological modeling at regional and long-term scales: Development, application, and analysis of WRF-HMS, *Water Resour. Res.*, 52, doi:10.1002/2015WR018185.

Acknowledgments: This research was funded by the German Research Foundation (DFG) and the National Natural Science Foundation of China (NSFC), and Helmholtz REKLIM.