

RECEIVED: September 26, 2017 Accepted: September 26, 2017 Published: October 10, 2017

Erratum: Closing the window for compressed dark sectors with disappearing charged tracks

Rakhi Mahbubani, a,b Pedro Schwaller and José Zurita d,e

^a Theoretical Physics Department, CERN, Geneva, Switzerland

E-mail: rakhi@cern.ch, pedro.schwaller@uni-mainz.de, jose.zurita@kit.edu

ERRATUM TO: JHEP06(2017)119

ARXIV EPRINT: 1703.05327

^b Theoretical Particle Physics Laboratory, Institute of Physics, EPFL, Lausanne, Switzerland

^cPRISMA Cluster of Excellence & Mainz Institute for Theoretical Physics, Johannes Gutenberg University, 55099 Mainz, Germany

^dInstitute for Nuclear Physics (IKP), Karlsruhe Institute of Technology, Hermann-von-Helmholtz-Platz 1, D-76344 Eggenstein-Leopoldshafen, Germany

^eInstitute for Theoretical Particle Physics (TTP), Karlsruhe Institute of Technology, Engesserstraße 7, D-76128 Karlsruhe, Germany




Figure 1. Corrected pair-production cross sections for a weak doublet with hypercharge $\frac{1}{2}$ at NLO, obtained using PROSPINO in the pure-Higgsino limit, for LHC14 (a) and FCC-hh (b).

The NLO pair-production cross sections for pure higgsinos at LHC14 and FCC-hh shown in figure 2 of our paper were underestimated by $\sim 30\%$. The corrected cross sections are shown in figure 1 above.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.