

LETTER • OPEN ACCESS

# Potential strong contribution of future anthropogenic land-use and land-cover change to the terrestrial carbon cycle

To cite this article: Benjamin Quesada *et al* 2018 *Environ. Res. Lett.* **13** 064023

View the [article online](#) for updates and enhancements.

## Related content

- [Nitrogen and phosphorous limitation reduces the effects of land use change on land carbon uptake or emission](#)  
Ying-Ping Wang, Qian Zhang, Andrew J Pitman *et al.*
- [Hydrological and biogeochemical constraints on terrestrial carbon cycle feedbacks](#)  
Stefanos Mystakidis, Sonia I Seneviratne, Nicolas Gruber *et al.*
- [Drivers and patterns of land biosphere carbon balance reversal](#)  
Christoph Müller, Elke Stehfest, Jelle G van Minnen *et al.*



Are you our new  
project leader  
marine technology  
development?



MORE INFO? VISIT [WORKINGATNIOZ.NL](http://WORKINGATNIOZ.NL)

# Environmental Research Letters



## LETTER

# Potential strong contribution of future anthropogenic land-use and land-cover change to the terrestrial carbon cycle

### OPEN ACCESS

RECEIVED  
13 April 2018

ACCEPTED FOR PUBLICATION  
15 May 2018

PUBLISHED  
6 June 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Benjamin Quesada<sup>1,4</sup> , Almut Arneith<sup>1</sup>, Eddy Robertson<sup>2</sup> and Nathalie de Noblet-Ducoudré<sup>3</sup>

<sup>1</sup> Institute of Meteorology and Climate Research, Atmospheric Environmental Research, Karlsruhe Institute of Technology, 82467 Garmisch-Partenkirchen, Germany

<sup>2</sup> Met Office Hadley Centre, Exeter, United Kingdom

<sup>3</sup> Laboratoire des Sciences du Climat et de l'Environnement LSCE/IPSL, Unité mixte CEA-CNRS-UVSQ, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France

<sup>4</sup> Author to whom any correspondence should be addressed.

E-mail: [benjamin.quesada@kit.edu](mailto:benjamin.quesada@kit.edu)

**Keywords:** carbon cycle, attribution, land cover changes, deforestation, greening, land carbon storage, Earth system model

Supplementary material for this article is available [online](#)

## Abstract

Anthropogenic land-use and land cover changes (LULCC) affect global climate and global terrestrial carbon (C) cycle. However, relatively few studies have quantified the impacts of future LULCC on terrestrial carbon cycle. Here, using Earth system model simulations performed with and without future LULCC, under the RCP8.5 scenario, we find that in response to future LULCC, the carbon cycle is substantially weakened: browning, lower ecosystem C stocks, higher C loss by disturbances and higher C turnover rates are simulated. Projected global greening and land C storage are dampened, in all models, by 22% and 24% on average and projected C loss by disturbances enhanced by ~49% when LULCC are taken into account. By contrast, global net primary productivity is found to be only slightly affected by LULCC (robust +4% relative enhancement compared to all forcings, on average). LULCC is projected to be a predominant driver of future C changes in regions like South America and the southern part of Africa. LULCC even cause some regional reversals of projected increased C sinks and greening, particularly at the edges of the Amazon and African rainforests. Finally, in most carbon cycle responses, direct removal of C dominates over the indirect CO<sub>2</sub> fertilization due to LULCC. In consequence, projections of land C sequestration potential and Earth's greening could be substantially overestimated just because of not fully accounting for LULCC.

## 1. Introduction

Terrestrial soils and vegetation contribute to the global carbon cycle and climate mainly through biogeochemical emissions and uptake of greenhouse gases (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, etc) and exchange of energy, water and momentum (i.e. biophysical effects) [1–3]. The productivity and carbon stocks of terrestrial ecosystems can be in turn affected by climate and human use. All other things equal, changes in terrestrial carbon storage are found to be positively correlated with changes in atmospheric CO<sub>2</sub> concentration and negatively to temperature changes (~1 PgC.ppm<sup>-1</sup> and ~-80 PgC.K<sup>-1</sup> as approximated sensitivities among the studies [4–7]).

Land-use and land cover changes (LULCC), like conversion of forests into crops or pastures, also affect ecosystem-climate-carbon cycle processes through changes in biophysical properties of the land-cover, changes in phenology and changes in biogeochemical emissions and uptake. Using bookkeeping, censuses, remote sensing or carbon/vegetation models, several studies quantified the contribution of past and present-day LULCC to carbon fluxes and global warming [e.g. 8–13]. Over the last 150 years, estimated cumulative LULCC emissions represent approximately one-third of total cumulative anthropogenic CO<sub>2</sub> emissions but only one-eighth over the recent period 1990–2010 [8, 10].

By contrast, relatively few studies have focused on future LULCC impacts on the carbon cycle, and in most cases (historical or future perspective) the emphasis was on net carbon uptake rather than on the underlying terrestrial carbon processes: photosynthesis, heterotrophic and autotrophic respiration, carbon turnover time, land cover productivity, phenology or even disturbances. Under present-day climate conditions, [14, 15] estimate that land-use reduced terrestrial NPP of potential natural vegetation at global scale by ~5%–10%. [16] found that under A2/B1/B2 future scenarios, based on LPJmL vegetation model simulations with climate forcings from four global climate models, land-use (from 1970–2100) contributes to modulate NPP by ~−5%/−1%/−3%, vegetation carbon by −47%/−19%/−27% and soil carbon by −7%/+1%/−2% compared to a baseline scenario, respectively. Moreover, while historical LULCC decreased soil carbon sequestration, meta-analysis reviews [17–19] found that deforestation would not necessarily lead to decreased soil carbon stocks: conversion of native forest to plantation or crops can imply reductions varying from −13% to −42% (~−2.1 kgC m<sup>−2</sup>) while transformation of native forest to pasture or grassland tends to increase them by 8% up to 19% (+1.2 kgC m<sup>−2</sup>).

In consequence, LULCC also alter the ecosystem carbon turnover time at global scale, in general reducing it by several years [16, 20–24]. By comparing the vegetation carbon turnover time of the actual vegetation and with that of a hypothetical vegetation state without land-use under current climate conditions, [24] find that land-use halved the biomass turnover time.

Robust estimates of the interplay between LULCC and terrestrial ecosystems state are of paramount importance to constrain the future projections in terrestrial carbon cycle and reduce their uncertainties. However, few studies have attributed and quantified the net impacts of LULCC on the terrestrial carbon cycle and all the above-mentioned underlying physical processes, *a fortiori* in a multi-Earth system model (ESM) framework or based on a common realistic and global LULCC scenario. Finally, future land-use and land-cover changes are not often explicitly taken into account in global coupled models and sometimes only included in terms of CO<sub>2</sub> emissions only [9, 25, 26]. Besides, biophysical effects can substantially modify future hydrological cycle and energy balance at the surface particularly in tropical deforested areas [27, 28], which can in turn modulate the future terrestrial carbon cycle.

Our study fills those gaps and explores new findings on terrestrial carbon cycle, making use of simulations with and without future LULCC (based on the RCP8.5 scenario) from five ESMs (i.e. General circulation models—GCMs—including interactive carbon cycle). We aim to analyze (i) the likely effects of LULCC on the global terrestrial carbon cycle (section 3.1) and (ii) the relative contribution of LULCC forcing

in the projected changes at global (section 3.2) and regional scale (section 3.3). Furthermore, we not only attribute the *net* changes in carbon cycle in response to future LULCC, but we also disentangle the direct effect of LULCC without altering CO<sub>2</sub> emissions ('LULCC only') and the biogeochemical effect of the emissions induced by LULCC ('LULCC-emissions only').

## 2. Methods

### 2.1. Models and experiments

The Land-Use and Climate, Identification of Robust Impacts (LUCID) is a major international intercomparison exercise that aims to investigate the robust impacts of LULCC using as many climate models as possible forced with a common LULCC scenario ([www.lucidproject.org.au/](http://www.lucidproject.org.au/)).

Analyzing the future impacts of LULCC, several modeling groups from the 5th Phase of the Coupled Model Intercomparison Project (CMIP5) performed ESM simulations without anthropogenic land-use changes from 2006–2100. We use outputs from all the five state-of-the-art CMIP5 models used in LUCID-CMIP5 (CanESM2, HadGEM2-ES, IPSL-CM5A-LR, MPI-ESM-LR and MIROC-ESM) on the last 30 years period of each experiment (2071–2100). Several different ESM experiments are explored here as detailed in table 1.

RCP8.5 simulations are the CMIP5 runs with all forcings including the future anthropogenic land-use and land-cover change forcing based on the business-as-usual RCP8.5 scenario (see table 1). This scenario includes spatially explicit future LULCC characterized by an expansion of croplands and pastures driven by the food demands of an increasing population and corresponds to a radiative forcing of more than 8.5 W.m<sup>−2</sup> in 2100 [29] (CO<sub>2</sub> atmospheric concentration ~936 ppm in 2100). L2A85 simulations are the same runs as RCP8.5 but without the anthropogenic land-use and land-cover change forcing (after year 2005), with atmospheric CO<sub>2</sub> concentration prescribed from the RCP8.5 scenario (table 1).

The difference between RCP8.5 and L2A85 simulations (i.e. RCP8.5–L2A85) corresponds thus to the sole *biophysical* effects of future anthropogenic land-use and land cover changes ('LULCC only', i.e. without changes in CO<sub>2</sub> atmospheric concentration). L1A85 (respectively, L1B85) is a similar simulation to RCP8.5 but without future anthropogenic land-use and land cover changes and prescribed (resp., interactively simulated) atmospheric CO<sub>2</sub> concentration. Thus, L1A85–L1B85 corresponds to the sole *biogeochemical* effects corresponding to the changes in atmospheric CO<sub>2</sub> in response to future anthropogenic land-use and land cover changes ('LULCC-emissions only', i.e. without LULCC). The 'net' effects, including feedbacks between biophysical

**Table 1.** Description of simulations used from CMIP5 and LUCID-CMIP5 following the RCP8.5 scenario.

CMIP5 simulations	Atmospheric CO <sub>2</sub> concentration	LULCC
<b>Historical</b>	Prescribed from historical scenario	Transient changes from 1850–2005
<b>rcp85</b>	Prescribed from RCP8.5 scenario	Transient changes from 2006–2100
<b>esmrcp85</b>	Interactive (emissions-driven with atmospheric CO <sub>2</sub> concentration determined by model) <sup>a</sup>	As in rcp85
<b>L1A85</b>	Prescribed (concentration-driven) from esmrcp85	Fixed to year 2005
<b>L1B85</b>	Interactive (emissions-driven atmospheric CO <sub>2</sub> concentration from RCP8.5 scenario)	Fixed to year 2005
<b>L2A85</b>	As in rcp85	Fixed to year 2005
<b>rcp85–L2A85</b>	(1) Biophysical effects: ‘LULCC only’ (same CO <sub>2</sub> but different LULCC)	
<b>L1A85–L1B85<sup>b,c</sup></b>	(2) Biogeochemical effects: ‘LULCC-emissions only’ (same LULCC but different CO <sub>2</sub> )	
<b>esmrcp85–L1B85</b>	(3) Net effects of LULCC	
<b>(3) minus [(1)+(2)]</b>	(4) Synergistic effects: non-linear feedbacks between biophysical and biogeochemical effects of LULCC	

<sup>a</sup> The CO<sub>2</sub> concentration results among other factors from prescribed anthropogenic CO<sub>2</sub> emissions and land-use and land-cover changes. As the predicted CO<sub>2</sub> concentration also depends on the climate simulated by the models, the esmrcp85 simulations allow for carbon-climate feedbacks.

<sup>b</sup> CanESM2 model does not provide L1A85 simulations, so for this model, the effects of ‘LULCC-emissions only’ are reasonably approximated by (esmrcp85–L1B85)–(rcp85–L2A85).

<sup>c</sup> HadGEM2-ES model does not provide L1A85 nor L1B85 simulations, so for this model, the effects of ‘LULCC-emissions only’ cannot be calculated and the net effects correspond to the effects of ‘LULCC only’.

and biogeochemical effects, correspond to the difference between esmrcp85 and L1B85, while the difference between net effects and the sum of ‘LULCC only’ and ‘LULCC-emissions only’ yields the ‘synergistic effects’ (table 1). Supplementary table S1 available at [stacks.iop.org/ERL/13/064023/mmedia](https://stacks.iop.org/ERL/13/064023/mmedia) details the main characteristics of the five LUCID-CMIP5 models including number of PFTs representation of dynamical vegetation, fire modules and horizontal resolution.

Historical simulations (HIST, *historical* according to CMIP5 abbreviation) are also used and averaged over the 1976–2005 period. For HadGEM2-ES model, it is not possible to calculate biogeochemical effects due to LULCC (absence of L1A85 and L1B85 simulations, see table 1). We can however infer from the smallest cumulative net land-use emissions simulated by this model (~25 PgC from 2006–2100 under RCP8.5 [30]) that those effects are overall negligible vs. ‘LULCC only’ effects.

In general, all figures present ensemble-mean results i.e. averaged results among the 5 LUCID-CMIP5 models interpolated on a common medium resolution grid of MIROC-ESM (2.8° × 2.8°, see supplementary table S1). When all 5 ESM simulate the same anomaly sign for spatial averages in response to LULCC, the signal is called ‘robust’. Supplementary figure S1 shows the tree fraction changes between RCP8.5 and historical simulation (from 2006–2100) averaged over the 5 LUCID-CMIP5 models. Future changes in tree cover are about 4 million km<sup>2</sup> by 2100 (i.e. ~1/10th > of global tree cover [31]). A strong tropical deforestation signal (up to –20% change in some regions) robustly dominates while at mid- and high-latitudes, given the representation or not of dynamical vegetation (see supplementary table S1), some forest expansion signal takes place (HadGEM2-ES, MPI-ESM-LR and MIROC-ESM) or not (CanESM2 and IPSL-CM5A-LR). Note that the LULCC scenario implemented

here do not include simulation of specific land-use as the irrigation, fertilizers, urbanization or other land management. Harmonization and implementation of future LULCC scenario into the five CMIP5 models are fully detailed in [29] and [30].


## 2.2. Carbon cycle variables

For the analysis of the changes in carbon cycle, we investigate seven key variables: net primary productivity (*npp*, CMIP5 abbreviation), leaf area index (*lai*), total ecosystem carbon stocks ( $C_{TOT}$ , sum of soil *cSoil* and vegetation carbon *cVeg*), ecosystem carbon residence time ( $\tau$ , defined as the ratio between  $C_{TOT}$  and gross primary productivity *gpp*, similar to the definition of [22]), heterotrophic respiration ( $r_H$ ), autotrophic respiration ( $r_A$ ) and carbon loss due to disturbances (e.g. fires) ( $L_d$ , defined as the difference between net ecosystem production NEP and net biome production NBP).

## 3. Results and Discussion

### 3.1. Weakened terrestrial carbon cycle in response to future LULCC

The biophysical (left column), biogeochemical (middle column) and net effects (right column) of LULCC on each carbon cycle variable (row) at global scale are presented in figure 1. Global changes in NPP in response to future LULCC are simulated: –1.1, +6.6 and +9.8 gC m<sup>–2</sup> yr<sup>–1</sup> on average, respectively i.e. –0.2, +1.0 and +1.5 PgC yr<sup>–1</sup> for ‘LULCC only’, ‘LULCC emissions only’ and net effects of LULCC. However, models simulate a large spatial variability of net NPP responses: strong NPP decreases are simulated in South African region (up to –100 gC m<sup>–2</sup> yr<sup>–1</sup>) enhanced by biogeochemical effects of LULCC as well as in Eastern South America and Sahelian regions.



**Figure 1.** Spatial patterns of changes in carbon cycle variables in response to (a) LULCC only, (b) LULCC-emissions only and (c) sum of both (net effects of LULCC) averaged on 2071–2100 period. Units are for NPP,  $r_H$ ,  $r_A$  and  $L_d$ :  $\text{gC yr}^{-1} \text{m}^{-2}$ , LAI: non-dimensional,  $C_{\text{TOT}}$ :  $\text{kgC m}^{-2}$  and  $\tau$ : years. Global continental averaged values are indicated in the bottom left corner of each panel (for  $\tau$ , the ratio between global continental averages of  $C_{\text{TOT}}$  and of GPP is calculated). Only gridpoints where four out of five LUCID-CMIP5 models simulate the anomaly sign are shown (white blank otherwise).

By contrast, Eurasia, large parts of South America, tropical Africa and Canada display increases in NPP. A disagreement in the sign of model NPP response occurs above several tropical deforested areas (blank areas in figure 1), particularly due to ‘LULCC only’ effects, while biogeochemical effects via  $\text{CO}_2$  fertilization tend to slightly boost vegetation productivity. LAI is decreased in response to future LULCC by 0.07 on average, predominantly affected by direct

LULCC effects of tropical deforestation in South America, Africa, Eastern Australia and Indonesia up to  $-0.6$ . Biogeochemical effects of LULCC cause no significant change on LAI at global scale. Changes in  $C_{\text{TOT}}$  maps mirror the changes in LAI: although some slight increases in  $C_{\text{TOT}}$  due to  $\text{CO}_2$  fertilization, particularly in boreal latitudes ( $+0.05 \text{ kgC m}^{-2}$  on global average or  $\sim 7.5 \text{ GtC}$  globally), direct land-cover changes strongly decrease the content in carbon of

the vegetation and soils ( $-0.45 \text{ kgC m}^{-2}$  on global average or  $\sim 67 \text{ GtC}$  globally). In response to future LULCC, we also find that soil carbon reductions are on average five times less important than vegetation carbon (not shown). Overall, the net effects of LULCC result in reduced ecosystem carbon content, particularly pronounced in Eastern South America, Africa and Indonesia. Global negative impacts of climate change on land carbon storage and foliage density, particularly in the Tropics [7, 32], are thus aggravated by LULCC.

Ecosystem carbon residence time  $\tau$  at global scale is found to decrease both in response to ‘LULCC only’ ( $-0.17 \text{ yrs}$ ) and to ‘LULCC-emissions only’ ( $-0.05 \text{ yrs}$ ) causing a global net decrease of  $\sim -0.27 \text{ yrs}$  (see figure 1 for  $\tau$ ). Large changes are simulated at regional level, particularly in deserts and semi-arid areas (e.g. Central Australia, Sahara) because of very low GPP on average which makes  $\tau$  being very sensitive to any small modulation of  $C_{\text{TOT}}$  or GPP in response to global LULCC. Plant respirations ( $r_A$ ) are substantially weakened globally in response to LULCC ( $-10.8 \text{ gC m}^{-2} \text{ yr}^{-1}$  i.e.  $-1.6 \text{ PgC yr}^{-1}$ ), particularly due to LULCC-only effects ( $-15.7 \text{ gC m}^{-2} \text{ yr}^{-2}$  i.e.  $-2.3 \text{ PgC yr}^{-1}$ ), and to the conversion of landscape in the Tropics: Eastern South America and Tropical Africa being the most affected regions. Dead organic matter by decomposition ( $r_H$ ) results in small positive net changes ( $+4.6 \text{ gC m}^{-2} \text{ yr}^{-1}$ ) due to LULCC-only ( $-2.1 \text{ gC m}^{-2} \text{ yr}^{-1}$  on average) and LULCC-biogeochemical effects ( $+3.2 \text{ gC m}^{-2} \text{ yr}^{-1}$ ).


The net carbon loss due to terrestrial disturbances ( $L_d$ ) is enhanced under LULCC ( $+7.2 \text{ gC m}^{-2} \text{ yr}^{-1}$ ) where both LULCC-only effects ( $+3.2 \text{ gC m}^{-2} \text{ yr}^{-1}$ ) and biogeochemical effects of LULCC ( $+5.0 \text{ gC m}^{-2} \text{ yr}^{-1}$ ) play a significant positive role, particularly marked above deforested areas of Tropical Africa, South America and to some lesser extent in Eurasia. Aside from direct biomass removal induced by LULCC, land carbon sequestration potential is expected to decrease due to increased decomposition rates ( $r_H$ ) and disturbances ( $L_d$ ) but only very weakly influenced by changes in gross primary productivity ( $-0.97 \text{ gC m}^{-2} \text{ yr}^{-1}$ , not shown). Besides, depleted total carbon stocks ( $C_{\text{TOT}}$ ) explain the increases in carbon turnover rates, while enhancement in global NPP (i.e.  $\text{GPP} - r_A$ ) is fully due to decrease in plant respiration ( $r_A$ ).

### 3.2. Global relative contribution of future LULCC

Under the RCP8.5 scenario, particularly due to the effect of increased  $\text{CO}_2$  fertilization and global warming induced by greenhouse gases emissions, terrestrial carbon cycle is found to be enhanced on average as reported by previous studies, despite a large inter-model uncertainty [9, 16, 26, 33]. When looking at the trends over the 21st century (supplementary figure S2), the multimodel-mean projects more productive terrestrial ecosystems (NPP is boosted by  $236 \text{ gC m}^{-2} \text{ yr}^{-1}$  i.e.  $35.2 \text{ Pg yr}^{-1}$ ) with denser foliage density (LAI is

increased by 0.22), more carbon content in soils and vegetation ( $C_{\text{TOT}}$  is increased by  $1.33 \text{ kg m}^{-2}$  on average i.e.  $198 \text{ GtC}$  global total) in parallel with stronger decomposition rates and plant respiration (projected changes in  $r_H$  and  $r_A$  are  $+192$  and  $+210 \text{ gC m}^{-2} \text{ yr}^{-1}$ , respectively:  $+28.7$  and  $31.3 \text{ PgC yr}^{-1}$ ). The five ESM projections used in our study are consistent with previous literature estimates: Wieder *et al* [34] find similar projected patterns of NPP and total carbon stocks with 11 CMIP5 GCM including carbon cycle, while, Friend *et al* [35] using seven DGVMs simulate on average increases in NPP, carbon stocks and vegetation carbon residence time comparable in magnitude to the changes presented in our analysis (supplementary figure S3). Ecosystem carbon residence time is projected to decrease by  $\sim 3.7$  years by the end of the century under RCP8.5 scenario (supplementary figure S3), which can be explained by ESMs simulating a greater relative increase in gross primary productivity (here GPP) with respect to the increase in ecosystem carbon stocks ( $C_{\text{TOT}}$ ). Moreover, LUCID-CMIP5 models project enhanced disturbance-related carbon losses ( $L_d$  is enhanced by  $16 \text{ gC m}^{-2} \text{ yr}^{-1}$ ), which is coherent with projected increases in fire carbon emissions by ESM in CMIP5 [36].

When future LULCC impacts on the terrestrial carbon cycle changes are compared *versus* projected impacts (i.e. with all forcings, supplementary figure S2), the relative carbon cycle response becomes more prominent (see global relative contribution of LULCC and relative changes in figure 2 and supplementary figure S3, respectively). Figure 2 displays for each carbon cycle variable ( $x$ -axis), each model (symbols) and for the ensemble-mean (bars), the relative contribution of LULCC, disentangled in ‘LULCC only’ effects (e.g. blue bars), ‘LULCC-emissions only’ (red bars) and net effects (black bars; synergistic effects being indicated by grey hatching, see table 1). While global LAI and  $C_{\text{TOT}}$  are simulated to increase at the end of the 21st century (supplementary figure S2), net effects of future LULCC contribute to reduce those projected changes by 22% and 24% respectively (black bars for those variables in figure 2). This negative contribution is found robust as every LUCID-CMIP5 model simulates it with relative contributions for LAI varying from  $-10\%$  (HadGEM2-ES) to  $-45\%$  (MIROC-ESM) and for  $C_{\text{TOT}}$  varying from  $-9\%$  (HadGEM2-ES) to  $-65\%$  (MIROC-ESM). The future contribution of LULCC to the projected changes of greening and carbon storage are thus much more important than the current contribution estimates: for instance, in a multi-dynamic global vegetation model (DGVM) framework, Zhu *et al* [37] find a positive relative contribution of present-day LULCC to the Earth’s greening (increased LAI) compared to all forcings lesser than 5% *vs.*  $-22\%$  in our study for future LULCC under RCP8.5. Besides, we found smaller contributions of LULCC to projected changes in global  $C_{\text{TOT}}$  ( $-24\%$ ) compared to [16] and [38] who simulate



very large LULCC relative contribution of  $-183\%$  and  $-97\%$ , respectively (when changes in total carbon stocks due to future LULCC are compared to projected changes under A2 scenario, most similar to RCP8.5 scenario used here).


This effect is even more pronounced for  $L_d$  where net effects of LULCC are found to enhance the projected changes by  $+49\%$  on average. The other carbon cycle variables are relatively less affected by LULCC: projected increases in NPP are robustly enhanced by  $4\%$ , projected increases in  $r_A$  are dampened by  $4\%$  while projected decrease in carbon residence times  $\tau$  are enhanced by  $7\%$  (figure 2 for  $\tau$ ). The global relative contributions calculated on 2071–2100 period tend to remain stable on time as depicted by the transitional values calculated on 2011–2040 and 2041–2070 periods (supplementary figure S4).

As suggested by [39], we find that synergy between biophysical and biogeochemical effects on carbon cycle variables are in general low (in most cases,  $< 2\%$  additional relative contribution) but relatively non-negligible, particularly for NPP (see synergistic LULCC effects in figure 2). When such synergistic effects are accounted for, spatial variability of net impacts of LULCC on NPP (figure 1) is reduced by  $25\%$  (in terms of standard deviation), attenuating negative impacts in Tropics and enhancing positive ones in mid- and high-latitudes.

We also find a strong model disagreement and an overall small relative contribution of LULCC-only induced changes in NPP and  $r_H$ . HadGEM2-ES, IPSL-CM5A-LR and CanESM2 simulate global increase in NPP and  $r_H$  due to future LULCC while MIROC-ESM and MPI-ESM-LR simulate global decrease. However,

even if the LULCC-only effects on NPP are largely uncertain (symbols around blue bar for NPP, figure 2) in agreement with a recent multi-DGVM study [40], positive synergistic effects (with biogeochemical effects of LULCC, grey hatching and red bars for NPP) make all 5 ESM agree on a net NPP enhancement in response to future LULCC ( $[1\%–13\%]$  relative contribution range among models, see symbols around black bar for NPP in figure 2), which corresponds to an opposite sensitivity compared with previous modelling studies [14–16]. In other means, ESM simulate that in a slightly warmer climate with more  $\text{CO}_2$ , NPP is slightly more enhanced by LULCC.

The effects of LULCC on NPP and respirations can be attributed to several complex competing mechanisms differently simulated by ESM and across regions: plant functional trait (PFT) classification and parametrization of key plant properties [41], temperature and precipitation changes induced by PFT changes [28], climate and stomatal conductance changes induced by  $\text{CO}_2$  increase [42], as well as non-linear interactions between them highlighted in our study. For instance, global NPP tends to be negatively related to temperature but positively with precipitation [4, 43–46] and  $\text{CO}_2$  [46–48] while LULCC impact on climate and stomatal conductance is regional-, PFT transitions- and model-dependent. That being said, the increases in NPP,  $r_H$  and  $r_A$  caused by the combination of climate change and  $\text{CO}_2$  fertilization in response to fossil fuel emissions outweigh the changes caused by LULCC, which are even smaller here than the ones simulated by [16]. Except for  $L_d$ , we also find that biogeochemical effects of LULCC (red bars) have a very limited contribution ( $< 3\%$ , in general



positive) on terrestrial carbon cycle. This result confirms that overall the biogeochemical effects of LULCC on future atmospheric  $\text{CO}_2$  concentrations are relatively small in comparison to fossil fuel emissions ( $\sim 4\%$  relative contribution on average among models here) [38, 49]. Finally, biogeochemical effects on terrestrial carbon cycle simulated here by ESM could even be overestimated as those models do not fully represent ecophysiological mechanisms in response to temperature increase, such as changes in nutrient availability, permafrost dynamics, soil moisture, phenology, microbial decomposition and species distribution [47, 48, 50]. Key biogeochemical processes as for instance Carbon-Nitrogen interactions (not implemented in the ESMS here) would also affect the carbon cycle response to climate variations induced by  $\text{CO}_2$  increase [51].

**3.3. Regional relative contribution of future LULCC**  
Figure 3 shows the relative contribution of LULCC on the projected changes in the seven carbon cycle

variables (Y-Axis) in each of the 26 IPCC regions (x-Axis, see domains in supplementary figure S5). In the great majority of the regions and variables (except  $L_d$ ), the direct LULCC effects (upper panel) dominate over the slight  $\text{CO}_2$  fertilization effect provoked by LULCC-biogeochemical impacts (middle panel). Black dots in figure 3 highlight inter-model robustness, when every LUCID-CMIP5 model simulates the same response sign for a given region and variable. Regions whose carbon cycle is most adversely affected by net LULCC effects (lower panel) are South America (in particular, Amazon AMZ, North East Brasil NEB, West Coast and South-Eastern parts of South America WSA and SSA, see supplementary figure S5 for geographical domains and names), Africa (WAF, SAF, EAF) and Southeast Asia (SEA), regions with most intense deforestation under RCP8.5 (see supplementary figure S1). Most negatively impacted variables are LAI,  $C_{\text{TOT}}$  and  $L_d$  (figure 3 lower panel). In NEB, SAH, WAF, EAF, SAF and SEA regions, LULCC contribute to dampen by more than 40% the

corresponding regional projected changes of the three variables LAI,  $C_{TOT}$  and  $L_d$ . In those regions, ‘LULCC-only’ effects explain the net contribution but, in South America, small warming and further drying due to biogeochemical effects (‘LULCC-emissions only’) tend to aggravate the browning and the decreasing capacity of land carbon sequestration (figure 3 middle panel for above-mentioned regions). To a lesser extent,  $\tau$  is also robustly reduced by 7% (ENA), 27% (CAM), 19% (AMZ), 10% (WSA), 20% (SSA) and 3% (Western Asia, WAS) compared to regional projected changes.

Enhanced productivity and heterotrophic respirations are also simulated in regions like SSA, NEU (Northern Europe), TIB (Tibetan Plateau), NAS (Northern Asia) and CAS (Central Asia). For instance, in CAS and TIB region, robust positive contribution of NPP (17% and 7%, respectively), LAI (27% and 5%),  $r_A$  (10% and 4%) and  $r_H$  (16% and 6%) are simulated. Those are regions where conversions of landscape are very low (supplementary figure S1), where projected LAI changes and LAI changes due to LULCC are both slightly positive (supplementary figure S2 and figure 3) and where the global biogeochemical effects mostly due to tropical deforestation tend to fertilize remote areas with  $CO_2$ .

There is a generalized model disagreement on the sign of change in NPP,  $r_H$  and  $r_A$  in response to ‘LULCC-only’ (upper panel) in almost every region. However, in response to LULCC, a majority of models simulates increase in mid- and high-latitudes and an overall decrease in some tropical areas (figure 3), that hides a large spatial variability (see also figure 1). At the gridcell level, we find that models simulate a net decrease in NPP and  $r_H$  with strong tropical deforestation greater than 10% change in tree fraction ( $\sim -4gC\ m^{-2}\ yr^{-1}/\%$ ; see supplementary figure S6, red curves) but in mid- and high-latitudes decrease in tree cover are associated with higher NPP (supplementary figure S6, orange and blue curves). Furthermore,  $r_H$  is negatively affected by tropical deforestation ( $-3.5gC\ m^{-2}\ yr^{-1}/\%$ ) by a factor 3 compared to  $r_A$  ( $-1.2gC\ m^{-2}\ yr^{-1}/\%$ ).

By contrast, LAI and components of  $C_{TOT}$ ,  $cSoil$  and  $cVeg$ , are gradually decreased by deforestation: on average, carbon in soils is decreased by  $-0.03kgC\ m^{-2}/\%$  while carbon in vegetation is decreased by  $-0.15kgC\ m^{-2}/\%$  due to direct tropical deforestation effect (five times more than  $cSoil$ ).

At regional level, figure 4 shows stronger effects due to LULCC than a weakened terrestrial carbon cycle. If under RCP8.5 scenario, models do not simulate a global ‘projected terrestrial carbon reversal’ (i.e. global changes in total carbon stocks in response to LULCC fully dampen their projected changes until 2071–2100 period), as discussed by [52] and [53] with DGVMs, we find however strong evidence of regional terrestrial carbon reversal mainly located in tropical regions (see figure 4(a) below, ratio of changes in carbon stocks

due to LULCC vs. changes in carbon stocks due to all forcings except LULCC).

About 19% of land gridpoints between 25°S and 10°N are subject to a projected terrestrial carbon reversal, particularly located around deforested edges of Amazon and African rainforests. Similar results are found for a ‘projected terrestrial greening reversal’ (i.e. changes in LAI in response to LULCC fully dampen projected greenings): at global scale, there is not such evidence while at regional level, a portion of deforested areas ( $\sim 18\%$  of land gridpoints between 25°S and 10°N) show a reversal in greening towards browning when LULCC is accounted for (figure 4(b)). Those reversals evidence the regional overwhelming impact of LULCC that dominate projected carbon cycle changes over all other forcings (greenhouse gases, aerosols and others).


In consequence, our regionalized results prove that LULCC play an extremely important role in the South American and African terrestrial carbon cycle, which makes stopping deforestation in those regions a paramount mitigation measure that could lead to even more benefits than previously thought.

#### 4. Conclusion

The contributions of future LULCC to the projections of global and regional terrestrial carbon cycle (2071–2100) are now assessed in a multi-model framework of five different ESMs and under a common realistic LULCC scenario (RCP8.5), distinguishing the direct impacts of carbon removal and the indirect  $CO_2$  emissions induced by those LULCC.

The terrestrial biosphere currently absorbs large amounts of carbon dioxide ( $CO_2$ ) from the atmosphere, partially compensating  $CO_2$  emissions from fossil fuel combustion and LULCC and tempering anthropogenic climate change. If land carbon uptake is projected to increase under future greenhouse gases scenarios, mainly driven by the positive effects of  $CO_2$  fertilization of photosynthesis [5, 42], although large uncertainties [9, 33, 54, 55], our results show that the ability of the terrestrial biosphere to sequester carbon from the atmosphere is substantially dampened by future LULCC.

We find that in response to future LULCC, the terrestrial carbon cycle is robustly weakened: browning, lower ecosystem carbon stocks, higher carbon loss by disturbances and higher turnover rates are simulated. At the end of the 21st century, projected global greening and land carbon storage are dampened, in all models, by  $\sim 20\%$ – $25\%$  on average and projected carbon loss by disturbances enhanced by  $\sim 50\%$  when LULCC are taken into account. By contrast, global NPP is found to be robustly but very slightly enhanced by LULCC ( $\sim +4\%$  relative contribution on average) compared to effects of greenhouse gases. LULCC are found to be a predominant driver of future C changes



in regions like South America and Southern part of Africa. Accounting for future LULCC leads the models to simulate regional reversals of projected increased carbon sinks and greening, particularly located at the edges of the Amazon and African rainforests. The multi-ESM framework under RCP8.5 LULCC scenario forges a lower road path compared to previous studies [16, 21, 38, 52] who found LULCC to greatly affect and sometimes reverse the projected terrestrial carbon stocks of the 21st century under business-as-usual warming scenarios.

Nonetheless, on top of the negative impact of LULCC on future land carbon sequestration and greening, the latest research reveals other adverse effects: the present-day reduction of the Amazon Basin carbon sink efficiency [56], the consideration of the Nitrogen-Phosphorus cycles [34, 51], the biogenic volatile organic compound feedback [57] and the land management [11, 23, 58] neglected in current inter-comparison model exercises. All these factors could further reduce the land carbon sink and, under high

emission scenarios and severe climate change [5, 59] or severe LULCC scenario [52], they could even reverse the terrestrial sink in to a source.

### Acknowledgments

This work is performed in the framework of the EC FP7 LUC4C project (<http://luc4c.eu>, grant n°603542). The authors thank the data producers from LUCID-CMIP5 project ([www.mpimet.mpg.de/en/science/the-land-in-the-earth-system/working-groups/climate-biogeosphere-interaction/lucid-cmip5/](http://www.mpimet.mpg.de/en/science/the-land-in-the-earth-system/working-groups/climate-biogeosphere-interaction/lucid-cmip5/)): Vivek Arora, Victor Brovkin, Patricia Cadule, Veronika Gayler, Etsushi Kato, Andy Pitman and Julia Pongratz, for making available their model outputs and for their contribution. We also acknowledge the World Climate Research Programmes Working Group on Coupled Modeling, which is responsible for CMIP5 (<http://cmip-pcmdi.llnl.gov/cmip5>), and the climate modeling groups.

## ORCID iDs

Benjamin Quesada  <https://orcid.org/0000-0002-8827-4801>

## References

- [1] Pielke R A *et al* 2011 Land use/land cover changes and climate: modeling analysis and observational evidence *Wiley Interdiscip. Rev. Clim. Change* **2** 828–50
- [2] Mahmood R *et al* 2014 Land cover changes and their biogeophysical effects on climate *Int. J. Climatol.* **34** 929–53
- [3] Perugini L, Caporaso L, Marconi S, Cescatti A, Quesada B, de Noblet N, House J and Arneth A 2017 Biophysical effects on temperature and precipitation due to land cover change *Environ. Res. Lett.* **12** 053002
- [4] Friedlingstein P *et al* 2006 Climate-carbon cycle feedback analysis: results from the C<sup>4</sup> MIP model intercomparison *J. Clim.* **19** 3337–53
- [5] Sitch S *et al* 2008 Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five dynamic global vegetation models (DGVMs) *Glob. Change Biol.* **14** 2015–39
- [6] Zickfeld K, Eby M, Matthews H D, Schmittner A and Weaver A J 2011 Nonlinearity of carbon cycle feedbacks *J. Clim.* **24** 4255–75
- [7] Peng J, Dan L and Huang M 2014 Sensitivity of global and regional terrestrial carbon storage to the direct CO<sub>2</sub> effect and climate change based on the CMIP5 model intercomparison *PLoS ONE* **9** e95282
- [8] Houghton R A, House J I, Pongratz J, van der Werf G R, DeFries R S, Hansen M C, Le Quéré C and Ramankutty N 2012 Carbon emissions from land use and land-cover change *Biogeosciences* **9** 5125–42
- [9] Ciais P *et al* 2014 Carbon and other biogeochemical cycles climate change 2013: the physical science basis *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge: Cambridge University Press) pp 465–570
- [10] Le Quéré C *et al* 2018 Global carbon budget 2017 *Earth Syst. Sci. Data* **10** 405–48
- [11] Arneth A *et al* 2017 Historical carbon dioxide emissions caused by land-use changes are possibly larger than assumed *Nat. Geosci.* **10** 79–84
- [12] Houghton R A and Nassikas A A 2017 Global and regional fluxes of carbon from land use and land cover change 1850–2015: carbon emissions from land use *Glob. Biogeochem. Cycles* **31** 456–72
- [13] Li W *et al* 2017 Land-use and land-cover change carbon emissions between 1901 and 2012 constrained by biomass observations *Biogeosciences* **14** 5053–67
- [14] DeFries R 2002 Past and future sensitivity of primary production to human modification of the landscape *Geophys. Res. Lett.* **29** 36–1
- [15] Haberl H, Erb K H, Krausmann F, Gaube V, Bondeau A, Plutzer C, Gingrich S, Lucht W and Fischer-Kowalski M 2007 Quantifying and mapping the human appropriation of net primary production in earth's terrestrial ecosystems *Proc. Natl Acad. Sci. USA* **104** 12942–7
- [16] Müller C, Eickhout B, Zaehle S, Bondeau A, Cramer W and Lucht W 2007 Effects of changes in CO<sub>2</sub>, climate, and land use on the carbon balance of the land biosphere during the 21st century *J. Geophys. Res. Biogeosci.* **112** G02032
- [17] Guo L B and Gifford R M 2002 Soil carbon stocks and land use change: a meta analysis *Glob. Change Biol.* **8** 345–60
- [18] Houghton R A and Goodale C L 2004 Effects of land-use change on the carbon balance of terrestrial ecosystems *Geophysical Monograph Series* vol 153 ed R S DeFries, G P Asner and R A Houghton (Washington, DC: American Geophysical Union) pp 85–98
- [19] Deng L, Zhu G, Tang Z and Shangguan Z 2016 Global patterns of the effects of land-use changes on soil carbon stocks *Glob. Ecol. Conserv.* **5** 127–38
- [20] Post W M and Kwon K C 2000 Soil carbon sequestration and land-use change: processes and potential *Glob. Change Biol.* **6** 317–27
- [21] Gitz V and Ciais P 2003 Amplifying effects of land-use change on future atmospheric CO<sub>2</sub> levels *Glob. Biogeochem. Cycles* **17** 1024
- [22] Carvalhais N *et al* 2014 Global covariation of carbon turnover times with climate in terrestrial ecosystems *Nature* **514** 213–7
- [23] Pugh T A M, Arneth A, Olin S, Ahlström A, Bayer A D, Klein Goldewijk K, Lindeskog M and Schurgers G 2015 Simulated carbon emissions from land-use change are substantially enhanced by accounting for agricultural management *Environ. Res. Lett.* **10** 124008
- [24] Erb K-H, Fetzel T, Plutzer C, Kastner T, Lauk C, Mayer A, Niedertscheider M, Körner C and Haberl H 2016 Biomass turnover time in terrestrial ecosystems halved by land use *Nat. Geosci.* **9** 674–8
- [25] Pongratz J, Reick C H, Houghton R A and House J I 2014 Terminology as a key uncertainty in net land use and land cover change carbon flux estimates *Earth Syst. Dyn.* **5** 177–95
- [26] Friedlingstein P 2015 Carbon cycle feedbacks and future climate change *Phil. Trans. R. Soc. Math. Phys. Eng. Sci.* **373** 20140421
- [27] Quesada B, Devaraju N, de Noblet-Ducoudré N and Arneth A 2016 Reduction of monsoon rainfall in response to past and future land use and land cover changes *Geophys. Res. Lett.* **44** 1041–50
- [28] Quesada B, Arneth A and de Noblet-Ducoudré N 2017 Atmospheric, radiative, and hydrologic effects of future land use and land cover changes: a global and multimodel climate picture *J. Geophys. Res. Atmos.* **122** 2016JD025448
- [29] Hurtt G C *et al* 2011 Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands *Clim. Change* **109** 117–61
- [30] Brovkin V *et al* 2013 Effect of anthropogenic land-use and land-cover changes on climate and land carbon storage in CMIP5 projections for the twenty-first century *J. Clim.* **26** 6859–81
- [31] Ward D S, Mahowald N M and Kloster S 2014 Potential climate forcing of land use and land cover change *Atmos. Chem. Phys.* **14** 12701–24
- [32] Mahowald N, Lo F, Zheng Y, Harrison L, Funk C, Lombardozzi D and Goodale C 2016 Projections of leaf area index in earth system models *Earth Syst. Dyn.* **7** 211–29
- [33] Ahlström A, Schurgers G, Arneth A and Smith B 2012 Robustness and uncertainty in terrestrial ecosystem carbon response to CMIP5 climate change projections *Environ. Res. Lett.* **7** 044008
- [34] Wieder W R, Cleveland C C, Smith W K and Todd-Brown K 2015 Future productivity and carbon storage limited by terrestrial nutrient availability *Nat. Geosci.* **8** 441–4
- [35] Friend A D *et al* 2014 Carbon residence time dominates uncertainty in terrestrial vegetation responses to future climate and atmospheric CO<sub>2</sub> *Proc. Natl Acad. Sci.* **111** 3280–5
- [36] Kloster S and Lasslop G 2017 Historical and future fire occurrence (1850–2100) simulated in CMIP5 Earth system models *Glob. Planet. Change* **150** 58–69
- [37] Zhu Z *et al* 2016 Greening of the Earth and its drivers *Nat. Clim. Change* **6** 791–5
- [38] Sitch S, Brovkin V, von Bloh W, van Vuuren D, Eickhout B and Ganopolski A 2005 Impacts of future land cover changes on atmospheric CO<sub>2</sub> and climate *Glob. Biogeochem. Cycles* **19** GB2013
- [39] Pongratz J, Reick C H, Raddatz T and Claussen M 2010 Biogeophysical versus biogeochemical climate response to historical anthropogenic land cover change *Geophys. Res. Lett.* **37** L08702

- [40] Krause A *et al* 2018 Large uncertainty in carbon uptake potential of land-based climate-change mitigation efforts *Glob. Change Biol.* **1**–14
- [41] Alton P B 2011 How useful are plant functional types in global simulations of the carbon, water, and energy cycles? *J. Geophys. Res.* **116** G01030
- [42] Pugh T A M, Muller C, Arneth A, Haverd V and Smith B 2016 Key knowledge and data gaps in modelling the influence of CO<sub>2</sub> concentration on the terrestrial carbon sink *J. Plant Physiol.* **203** 3–15
- [43] Wu Z, Dijkstra P, Koch G W, Peñuelas J and Hungate B A 2011 Responses of terrestrial ecosystems to temperature and precipitation change: a meta-analysis of experimental manipulation: meta-analysis of experimental manipulation *Glob. Change Biol.* **17** 927–42
- [44] Liu Y, Wang T, Huang M, Yao Y, Ciais P and Piao S 2016 Changes in interannual climate sensitivities of terrestrial carbon fluxes during the 21st century predicted by CMIP5 Earth system models: change in climate sensitivity of C flux *J. Geophys. Res. Biogeosci.* **121** 903–18
- [45] Chang J *et al* 2017 Benchmarking carbon fluxes of the ISIMIP2a biome models *Environ. Res. Lett.* **12** 045002
- [46] Boisier J P, Halladay K, Kay G, Ciais P and Good P 2014 *Report on quantifying sensitivity of regional climate of Amazonia to feedbacks from CO<sub>2</sub> physiological forcing* (Delivery Report 3.2—Work Package 3: AMAZALERT European FP7 Project)
- [47] Piao S *et al* 2013 Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO<sub>2</sub> trends *Glob. Change Biol.* **19** 2117–32
- [48] Kolby Smith Y Y, Reed S C, Cleveland C C, Ballantyne A P, Anderegg W R L, Wieder W R, Liu Y Y and Running S W 2016 Large divergence of satellite and Earth system model estimates of global terrestrial CO<sub>2</sub> fertilization *Nat. Clim. Change* **6** 306–10
- [49] House J I, Colin Prentice I and Le Quere C 2002 Maximum impacts of future reforestation or deforestation on atmospheric CO<sub>2</sub> *Glob. Change Biol.* **8** 1047–52
- [50] Luo Y *et al* 2016 Toward more realistic projections of soil carbon dynamics by Earth system models: soil carbon modeling *Glob. Biogeochem. Cycles* **30** 40–56
- [51] Wang Y-P, Zhang Q, Pitman A J and Dai Y 2015 Nitrogen and phosphorous limitation reduces the effects of land use change on land carbon uptake or emission *Environ. Res. Lett.* **10** 014001
- [52] Müller C, Stehfest E, van Minnen J G, Strengers B, von Bloh W, Beusen A H W, Schaphoff S, Kram T and Lucht W 2016 Drivers and patterns of land biosphere carbon balance reversal *Environ. Res. Lett.* **11** 044002
- [53] Bayer A D, Pugh T A M, Krause A and Arneth A 2015 Historical and future quantification of terrestrial carbon sequestration from a Greenhouse-Gas-Value perspective *Glob. Environ. Change* **32** 153–64
- [54] Schimel D, Stephens B B and Fisher J B 2015 Effect of increasing CO<sub>2</sub> on the terrestrial carbon cycle *Proc. Natl Acad. Sci.* **112** 436–41
- [55] Wu Z, Ahlström A, Smith B, Ardö J, Eklundh L, Fensholt R and Lehsten V 2017 Climate data induced uncertainty in model-based estimations of terrestrial primary productivity *Environ. Res. Lett.* **12** 064013
- [56] Brienen R J W *et al* 2015 Long-term decline of the Amazon carbon sink *Nature* **519** 344–8
- [57] Scott C E *et al* 2018 Impact on short-lived climate forcers increases projected warming due to deforestation *Nat. Commun.* **9** 157
- [58] Pongratz J *et al* 2018 Models meet data: challenges and opportunities in implementing land management in Earth system models *Glob. Change Biol.* **24** 1470–87
- [59] Scheffer M, Brovkin V and Cox P M 2006 Positive feedback between global warming and atmospheric CO<sub>2</sub> concentration inferred from past climate change *Geophys. Res. Lett.* **33** L10702