


# GPS-based gravity field recovery from reprocessed GOCE precise science orbits


Thomas Grombein<sup>1</sup>, Daniel Arnold<sup>2</sup>, Adrian Jäggi<sup>2</sup>

<sup>1</sup> Geodetic Institute, Karlsruhe Institute of Technology <sup>2</sup> Astronomical Institute, University of Bern




## ■ GOCE Precise Science Orbits (PSO)

- GPS-based orbit determination via Satellite-to-Satellite Tracking (SST-hl)
- **Kinematic** and reduced-dynamic orbits




## ■ Gravity field recovery

- Kinematic orbit positions are used to recover the long-wavelength gravity field
- Input for combined GOCE gravity field models (GPS + gradiometry)
- Degradation of kinematic positions propagates into gravity field solutions



- Gravity field recovery based on operational GOCE PSO (Jäggi et. al., 2015)



Ionosphere-induced  
artifacts along the  
geomagnetic equator

Differences w.r.t.  
ITSG-GRACE2016  
(300 km Gauss-filtered)

**Workaround:** Exclusion of affected GPS observations from orbit determination


**Problem:** Degradation of orbit quality (→ not applied for official PSO product)


**GOCE Reprocessing campaign:** Reprocessing of GOCE PSO and gravity field recovery

## ■ Complete reprocessing of GOCE PSO


- Bernese GNSS software using reprocessed GPS products (IGb08)
- **Down-weighting strategy** to mitigate ionosphere-induced effects
  - Affected GPS observations get higher covariance values
  - **Criterion 1:** Large changes in geometry-free linear combination (2<sup>nd</sup> derivative)
  - **Criterion 2:** Large ROTI values (Rate Of TEC Index)
  - Performance is validated by orbit overlaps and SLR residuals

- Gravity field recovery (Celestial Mechanics Approach)
  - 1-sec reprocessed kinematic GOCE positions (epoch-wise covariance information)
  - **Arc-specific orbit** and **gravity field parameters** are estimated simultaneously
    - Six initial **Keplerian elements** for each 24-hour arc
    - Constant and once-per-revolution **empirical accelerations** over 24 hours
    - **Pseudo-stochastic pulses** each 6 minutes (constrained)
    - **SH coefficients** up to degree and order 120 (without regularization)
  - Use of **GOCE accelerometer (ACC) data** to improve lowest SH coefficients
- Gravity field solutions







2009 Nov/Dec




2010 Nov/Dec




2011 Nov/Dec




2012 Nov/Dec




2009 Nov/Dec




2010 Nov/Dec



2011 Nov/Dec




2012 Nov/Dec






# Reprocessed PSO + ACC data


2009 Nov/Dec




2010 Nov/Dec




2011 Nov/Dec




2012 Nov/Dec



# Accumulated solutions








# Long-term solution

Nov 2009 – Jul 2012



# Long term solution: Nov 2009 – Jul 2012



### Operational PSO





### Reprocessed PSO + ACC





## ■ Statistics of differences in the region $|\varphi| \leq 50^\circ$

Version	Min [mm]	Max [mm]	WRMS [mm]
Operational PSO	-32.5	50.1	10.9
Reprocessed PSO + ACC	-11.7	16.7	3.3

Operational PSO



Reprocessed PSO + ACC





■ **Current activities:** Additional screening of kinematic positions



- Screening based on covariance values of positions in the region  $|\varphi| \leq 50^\circ$
- Threshold of 3 cm  $\rightarrow$  2.97 % reduction of observations



**Reprocessed PSO + ACC**





**Reprocessed PSO (screened) + ACC**





■ Statistics of differences in the region  $|\varphi| \leq 50^\circ$

Version	Min [mm]	Max [mm]	WRMS [mm]
Reprocessed PSO + ACC	-11.7	16.7	3.3
Reprocessed PSO (screened) + ACC	-11.9	9.4	2.7

**Reprocessed PSO + ACC**



**Reprocessed PSO (screened) + ACC**



- Statistics of differences in the region  $|\varphi| \leq 50^\circ$

Version	Min [mm]	Max [mm]	WRMS [mm]
Reprocessed PSO + ACC	-11.7	16.7	3.3
Reprocessed PSO (screened) + ACC	-11.9	9.4	2.7

## ■ GOCE Reprocessing Campaign


- Complete reprocessing of the GOCE Precise Science Orbits (PSO)
- Improved GPS-based gravity field recovery based on reprocessed PSO

## ■ Main findings

- Observation weighting positively impacts lower degrees (esp.  $10 < n < 50$ )
- Strong reduction of artifacts along the geomagnetic equator
- Use of GOCE accelerometer data improves lowest degrees ( $n < 10$ )
- Further significant reduction of artifacts by screening based on covariance values

- New long-term GOCE GPS-only gravity field solution based on reprocessed PSO is available for the whole mission period (2009 – 2013)

Thank you for your attention



Contact: [grombein@kit.edu](mailto:grombein@kit.edu)