

Search for pair production of first-generation scalar leptoquarks at $\sqrt{s} = 13$ TeV

A. M. Sirunyan *et al.**
(CMS Collaboration)

 (Received 3 November 2018; published 14 March 2019)

A search for the pair production of first-generation scalar leptoquarks is performed using proton-proton collision data recorded at 13 TeV center-of-mass energy with the CMS detector at the LHC. The data correspond to an integrated luminosity of 35.9 fb^{-1} . The leptoquarks are assumed to decay promptly to a quark and either an electron or a neutrino, with branching fractions β and $1 - \beta$, respectively. The search targets the decay final states comprising two electrons, or one electron and large missing transverse momentum, along with two quarks that are detected as hadronic jets. First-generation scalar leptoquarks with masses below 1435 (1270) GeV are excluded for $\beta = 1.0(0.5)$. These are the most stringent limits on the mass of first-generation scalar leptoquarks to date. The data are also interpreted to set exclusion limits in the context of an R -parity violating supersymmetric model, predicting promptly decaying top squarks with a similar dielectron final state.

DOI: [10.1103/PhysRevD.99.052002](https://doi.org/10.1103/PhysRevD.99.052002)

I. INTRODUCTION

The quark and lepton sectors of the standard model (SM) [1–3] are similar: both have the same number of generations composed of electroweak doublets. This could indicate the existence of an additional fundamental symmetry linking the two sectors, as proposed in many scenarios of physics beyond the SM. These include grand unified theories with symmetry groups $SU(4)$ of the Pati–Salam model [4,5], $SU(5)$, $SO(10)$, and $SU(15)$ [6–11]; technicolor [12–14]; superstring-inspired models [15]; and models exhibiting quark and lepton substructures [16]. A common feature of these models is the presence of a new class of bosons, called leptoquarks (LQs), that carry both lepton (L) and baryon numbers (B). In general, LQs have fractional electric charge and are color triplets under $SU(3)_C$. Their other properties, such as spin, weak isospin, and fermion number ($3B + L$), are model dependent.

Direct searches for LQs at colliders are usually interpreted in the context of effective theories that impose constraints on their interactions. In order to ensure renormalizability, these interactions are required to respect SM group symmetries, restricting the couplings of the LQs to SM leptons and quarks only. A detailed account of LQs and their interactions can be found in Ref. [17]. Results from

experiments sensitive to lepton number violation, flavor changing neutral currents, and proton decay allow the existence of three distinct generations of LQs with negligible intergenerational mixing for mass scales accessible at the CERN LHC [18,19]. Indirect searches for new physics in rare B meson decays [20–24] by LHCb and Belle suggest a possible breakdown of lepton universality. These anomalies, if confirmed, could provide additional support for LQ-based models [25]. A comprehensive review of LQ phenomenology and experimental constraints on their properties is given in Ref. [26].

We search for the pair production of first-generation scalar LQs that decay promptly. The final state arising from each LQ decay comprises a quark that is detected as a hadronic jet and either an electron or a large missing transverse momentum attributed to the presence of an undetected neutrino. For light-quark final states, the quark flavors cannot be determined from the observed jets. We assume the LQs decay only to $e(\nu_e)$ and up or down quarks. The branching fractions for the LQ decay are expressed in terms of a free parameter β , where β denotes the branching fraction to an electron and a quark, and $1 - \beta$ the branching fraction to a neutrino and a quark. For pair production of LQs, we consider two decay modes. The first arises when each LQ decays to an electron and a quark, having an overall branching fraction of β^2 . In the second mode one LQ decays to an electron and a quark, and the other to a neutrino and a quark. This mode has a branching fraction of $2\beta(1 - \beta)$. We, therefore, utilize final states with either two high transverse momentum (p_T) electrons and two high- p_T jets, denoted as $eejj$, or one high- p_T electron, large missing transverse momentum, and two high- p_T jets, denoted as $e\nu jj$.

*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

Previous experiments at the LEP [27], HERA [28,29], and Tevatron [30,31] colliders have searched for LQ production and placed lower limits of several hundreds of GeV on allowed LQ masses (m_{LQ}) at 95% confidence level (C.L.). The CMS experiment at the LHC has extended the limits on pair production of first-generation scalar LQs using proton-proton (pp) collision data recorded during 2012 at a center-of-mass energy of $\sqrt{s} = 8$ TeV. Based on a sample corresponding to an integrated luminosity of 19.7 fb^{-1} , the lower limit obtained on m_{LQ} was 1010 (850) GeV for $\beta = 1.0(0.5)$ [32]. The CMS Collaboration has also published results on a search for singly produced LQs with the final states of either two electrons and one jet, or two muons and one jet [33]. Recently, using a data set of 3.2 fb^{-1} collected at $\sqrt{s} = 13$ TeV, the ATLAS experiment has placed a lower limit on m_{LQ} of 1100 GeV [34] for $\beta = 1.0$.

This analysis is based on data recorded in pp collisions at $\sqrt{s} = 13$ TeV with the CMS detector, corresponding to an integrated luminosity of 35.9 fb^{-1} . At LHC energies, the pair production of LQs would mainly proceed via gluon-gluon fusion with a smaller contribution from quark-antiquark annihilation. The corresponding Feynman diagrams are shown in Fig. 1. The production cross section as a function of m_{LQ} has been calculated at next-to-leading order (NLO) in perturbation theory [35]. At the LHC, the LQ-lepton-quark Yukawa coupling has negligible effect on the production rate for promptly decaying LQs, which are the focus of our search.

The paper is organized as follows. Section II introduces the CMS detector, and Sec. III describes the data and simulated samples used in the search. The core of the analysis in terms of event reconstruction and selection is discussed in Sec. IV, while the background estimation is presented in Sec. V. Section VI deals with the systematic uncertainties affecting this analysis. Sections VII and VIII describe the results of the LQ search and its interpretation

in an exotic scenario of supersymmetry, respectively. We conclude with a summary of the main results in Sec. IX.

II. THE CMS DETECTOR

The key feature of the CMS apparatus is a superconducting solenoid of 6 m diameter, providing a magnetic field of 3.8 T. Within the solenoid volume lie a silicon pixel and microstrip tracker, a lead-tungstate crystal electromagnetic calorimeter (ECAL), and a brass-scintillator hadron calorimeter (HCAL), each composed of a barrel and two end-cap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and end-cap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. The first level of the trigger system [36], composed of custom electronics, uses information from the calorimeters and muon detectors to select the most interesting events in an interval of less than $4 \mu\text{s}$. The high-level trigger processor farm further reduces the event rate from around 100 kHz to 1 kHz, before data storage. A detailed description of the CMS detector, along with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [37].

III. DATA AND SIMULATED SAMPLES

Events are selected using a combination of triggers requiring either a single electron or a single photon. Electron candidates are required to have a minimum p_T of 27 (115) GeV for the low (high) threshold trigger. Each of these triggers examines clusters of energy deposited in the ECAL that are matched to tracks reconstructed within a range $|\eta| < 2.5$. Cluster shape requirements as well as calorimetric and track-based isolation (only for the low threshold trigger) are also applied. By comparison, the photon trigger requires $p_T > 175$ GeV without any requirements on track-cluster matching, cluster shape, or




FIG. 1. Leading order Feynman diagrams for the scalar LQ pair production channels at the LHC.

isolation. The latter three criteria are applied to electron triggers to reduce background rates and are not necessary at high p_T . Therefore, the single photon and electron triggers are combined to improve efficiency at high electron p_T . Events selected using other single-photon triggers with lower thresholds are used for determining the multijet background.

Monte Carlo (MC) simulation samples of scalar LQ signals are generated using PYTHIA version 8.212 [38] at leading order (LO) with the NNPDF2.3LO parton distribution function (PDF) set [39]. Samples are generated for m_{LQ} ranging from 200 to 2000 GeV in 50 GeV steps. The LQ is assumed to have quantum numbers corresponding to the combination of an electron ($L = 1$) and an up quark ($B = 1/3$), implying it has an electric charge of $-1/3$. Possible formation of hadrons containing LQs is not included in the simulation. The cross sections are normalized to the values calculated at NLO [35,40] using the CTEQ6L1 PDF set [41].

The main backgrounds for searches in the $eejj$ and $e\nu jj$ channels include Drell–Yan (Z/γ^*) production with jets, top quark pair production ($t\bar{t}$), single top quark and diboson ($VV = WW, WZ, \text{ or } ZZ$) production. Additional background contributions arise from $W + \text{jets}$, $\gamma + \text{jets}$, and multijet production, where jets are misidentified as electrons. The $t\bar{t}$ background in the $eejj$ channel as well as the multijet background in both channels are estimated from data, while MC simulated events are used to calculate all other backgrounds. The $Z/\gamma^* + \text{jets}$, $W + \text{jets}$, and VV samples are generated at next to leading order (NLO) with MADGRAPH5_aMC@NLO version 2.3.3 using the FxFx merging method [42,43]. Both $t\bar{t}$ and single top quark events are generated at NLO using MADGRAPH5_aMC@NLO, and POWHEG v2 complemented with MADSPIN [44], except for single top quark production in association with a W boson, where events are generated with POWHEG v1 at NLO [45–50], and s -channel single top quark production, where MADGRAPH5_aMC@NLO at NLO is used. The $\gamma + \text{jets}$ events are generated with MADGRAPH5_aMC@NLO at LO with MLM merging [51]. The NNPDF3.0 at NLO [52] PDF set is used, except for $\gamma + \text{jets}$ events that are generated using the LO PDF set.

The $W + \text{jets}$ and $Z/\gamma^* + \text{jets}$ samples are normalized to next-to-NLO (NNLO) inclusive cross sections calculated with FEWZ versions 3.1 and 3.1.b2, respectively [53]. Single top quark samples are normalized to NLO inclusive cross sections [54,55], except for the tW production, where the NNLO calculations of Refs. [56] are used. The calculations from Refs. [57–63] with TOP++2.0 are used to normalize the $t\bar{t}$ sample at NNLO in quantum chromodynamics (QCD) including resummation of the next-to-next-to-leading-logarithmic soft gluon terms.

PYTHIA 8.212 with the CUETP8M1 underlying event tune [64] is used for hadronization and fragmentation in all simulated samples, with the exception of a dedicated tune

used for the $t\bar{t}$ sample [65]. All samples include an overlay of minimum bias events (pileup), generated with an approximate distribution for the number of additional pp interactions expected within the same or nearby bunch crossings, and reweighted to match the distribution observed in data. In all cases, the GEANT4 software v.10.00.p02 [66,67] is used to simulate the response of the CMS detector.

IV. EVENT RECONSTRUCTION AND SELECTION

A particle-flow (PF) algorithm [68] aims to reconstruct and identify each individual particle in a given event, by optimally combining information from the various elements of the CMS detector. The energy of photons is directly obtained from the ECAL measurement. On the other hand, the energy of electrons is determined from a combination of their momentum at the primary interaction vertex as determined by the tracker, the energy of the corresponding ECAL clusters, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the associated track. The momentum of muons is obtained from the curvature of the corresponding track. The energy of charged hadrons is determined from a combination of their momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for zero suppression effects as well as for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energy.

Electrons are identified by spatially matching a reconstructed charged-particle track to a cluster of energy deposits in the ECAL. The ECAL cluster is required to have longitudinal and transverse profiles compatible with those expected from an electromagnetic shower. Electrons used in this analysis are required to have $p_T > 50$ GeV and $|\eta| < 2.5$, excluding the transition regions between barrel and end-cap detectors $1.4442 < |\eta| < 1.5660$. Additional selection criteria are applied to electron candidates in order to reduce backgrounds while maintaining high efficiency for identification of electrons with large p_T [69]. The absolute difference in η between the ECAL cluster seed and the matched track is required to be less than 0.004 (0.006) in the barrel (end cap), and the corresponding quantity in the azimuthal angle, ϕ , must be less than 0.06 rad. Leptons resulting from the decay of LQs are expected to be isolated from hadronic activity in the event. Requirements are, therefore, applied based on calorimeter energy deposits and tracks in the vicinity of electron candidates. The scalar sum of p_T associated with calorimeter clusters in a cone of radius $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.3$ centered on the electron candidate, excluding clusters associated to the candidate itself, must be less than 3% of the electron p_T . A correction to the isolation sum accounts for contributions from pileup interactions. The track-based isolation, calculated as the scalar p_T sum of all tracks in the cone defined

above, must be less than 5 GeV to reduce misidentification of jets as electrons. At most one layer of the pixel detector may have missing hits along the trajectory of the matched track. The track must also be compatible with originating from the primary pp interaction vertex, which is taken to be the reconstructed vertex with the largest value of summed physics-object p_T^2 . Here the physics objects are the jets, reconstructed using the algorithm [70,71] with the tracks assigned to the vertex as inputs, and the negative vector sum of the p_T of those jets. To correct for the possible difference of electron reconstruction and identification efficiencies between collision and simulated data, appropriate corrections or scale factors are applied to the simulated samples.

Muons are used in defining a control region to estimate the $t\bar{t}$ background contribution. They are identified as tracks in the central tracker consistent with either a track or several hits in the muon system [72]. These muon candidates must have $p_T > 35$ GeV and $|\eta| < 2.4$, and are required to pass a series of identification criteria designed for high- p_T muons as follows. Segments in at least two muon stations must be geometrically matched to a track in the central tracker, with at least one hit from a muon chamber included in the muon track fit. In order to reject muons from decays in flight and increase momentum measurement precision, at least five tracker layers must have hits associated with the muon, and there must be at least one hit in the pixel detector. Isolation is imposed by requiring the p_T sum of tracks in a cone of $\Delta R = 0.3$ (excluding the muon itself) divided by the muon p_T to be less than 0.1. For rejection of cosmic ray muons, the transverse impact parameter of the muon track with respect to the primary vertex must be less than 2 mm and the longitudinal distance of the track formed from tracker system only to the primary vertex must be less than 5 mm. Finally, the relative uncertainty on the p_T measurement from the muon track must be less than 30%.

Jets are reconstructed using the anti- k_T algorithm [70,71] with a distance parameter of 0.4. Their momentum is determined as the vectorial sum of all particle momenta in the jet, and is found in simulation to be within 5%–10% of the true momentum [73] over the entire p_T spectrum and detector acceptance. Pileup interactions can contribute spurious tracks and calorimeter energy deposits to the jet momentum. To mitigate this effect, tracks identified to be originating from pileup vertices are discarded, while a correction [74] is applied to compensate for the remaining contributions. Jet energy corrections are extracted from simulation to compensate for differences between the true and reconstructed momenta of jets. *In situ* measurements of the momentum balance in dijet, γ + jets, Z/γ^* + jets, and multijet events are used to estimate and correct for any residual differences in jet energy scale between data and simulation [74]. Additional selection criteria are applied to all jets to remove those potentially affected by spurious energy deposits originating from instrumental effects or

reconstruction failures [75]. Jets must have $p_T > 50$ GeV and $|\eta| < 2.4$, and only jets separated from electrons or muons by $\Delta R > 0.3$ are retained.

The missing transverse momentum (\vec{p}_T^{miss}) is given by the negative vector sum of p_T of all PF candidates in the event. The magnitude of \vec{p}_T^{miss} is referred to as p_T^{miss} .

To identify b jets arising from top quark decays in the determination of the $e\nu jj$ background control regions, the combined secondary vertex algorithm is used with the loose working point of Ref. [76]. Based on simulation, the corresponding b -jet identification efficiency is above 80% with a probability of 10% of misidentifying a light-flavor jet.

A. The $eejj$ channel

For the $eejj$ analysis, we select events with at least two electrons and at least two jets passing the criteria described above. No charge requirements are imposed on the electrons. When additional objects satisfy these requirements, the two highest p_T electrons and jets are considered. Further, there should not be any muon fulfilling the requirements mentioned earlier in this section. The dielectron invariant mass m_{ee} is required to be greater than 50 GeV. The p_T of the dielectron system must be greater than 70 GeV. The scalar p_T sum over the electrons and two jets, $S_T = p_T(e_1) + p_T(e_2) + p_T(j_1) + p_T(j_2)$, must be at least 300 GeV. This initial selection is used for the determination of backgrounds in control regions, as explained in Section V.

Final selections are then optimized by maximizing the Punzi criterion for observation of a signal at a significance of five standard deviations [77]. These selections are determined by examining three variables: m_{ee} , S_T , and m_{ej}^{min} . The electron-jet pairing is chosen to minimize the difference in the invariant mass of the LQ candidates, and the quantity m_{ej}^{min} is defined as the smaller of the two masses. Thresholds for the three observables are varied independently, and the Punzi criterion is then calculated at each set of thresholds as well as for each m_{LQ} hypothesis. The optimized thresholds as a function of m_{LQ} are shown in Fig. 2 (left). For the m_{LQ} hypotheses above 1050 GeV, the statistical uncertainty in the background prediction becomes large, making an optimization for these masses impossible, and thus the thresholds for the 1050 GeV hypothesis are applied.

B. The $e\nu jj$ channel

In the $e\nu jj$ channel, we select events containing exactly one electron, at least two jets, and $p_T^{\text{miss}} > 100$ GeV. The electron and jets must pass the aforementioned identification criteria. Events with isolated muons are rejected, applying the same criteria as for the $eejj$ channel. The absolute difference in the angle between the \vec{p}_T^{miss} and the leading p_T jet, $\Delta\phi(\vec{p}_T^{\text{miss}}, j_1)$, is required to be larger




FIG. 2. Optimized threshold values applied for the selection variables in the $eejj$ (left) and $evjj$ (right) channels as a function of m_{LQ} .

than 0.5 rad. This helps reject events with p_T^{miss} arising primarily from instrumental effects. The $\Delta\phi(\vec{p}_T^{\text{miss}}, e)$ must be greater than 0.8 rad for similar reasons. The p_T and transverse mass of the \vec{p}_T^{miss} -electron system must be greater than 70 and 50 GeV, respectively. Here and later, the transverse mass of a two-object system is given by $m_T = \sqrt{2p_{T,1}p_{T,2}(1 - \cos \Delta\phi)}$, with $\Delta\phi$ being the angle between the p_T vectors of two objects, namely \vec{p}_T^{miss} , electron and jet. The m_T criterion helps suppress the W + jets contribution. Finally, selected events must have $S_T > 300 \text{ GeV}$, where $S_T = p_T(e) + p_T^{\text{miss}} + p_T(j_1) + p_T(j_2)$. This initial selection is used for the determination of backgrounds in control regions, similarly to the $eejj$ channel.

The selection criteria are then optimized in a similar fashion as for the $eejj$ channel, except that four observables are considered for final selections at each m_{LQ} hypothesis: S_T , m_T of the \vec{p}_T^{miss} -electron system, p_T^{miss} , and the electron-jet invariant mass m_{ej} . The \vec{p}_T^{miss} -jet and electron-jet pairing is chosen to minimize the difference in m_T between the two LQ candidates. The optimized thresholds as a function of




FIG. 3. Data and background comparison for events passing the initial selection requirements for the $eejj$ channel, shown for the variables used for final selection optimization: m_{ee} (upper), m_{ej}^{\min} (lower left), and S_T (lower right). “Other background” includes diboson, single top quark, and W + jets. Signal predictions for $m_{LQ} = 650$ and 1200 GeV hypotheses are overlaid on the plots. The last bin includes all events beyond the upper x -axis boundary.

m_{LQ} are shown in Fig. 2 (right). As with the $eejj$ channel, for the m_{LQ} hypotheses above 1200 GeV , the thresholds for the 1200 GeV hypothesis are used.

V. BACKGROUND ESTIMATION

The SM processes that produce electrons and jets can have final states similar to those of an LQ signal and are, therefore, considered as backgrounds for this search. These include dilepton events from $Z/\gamma^* + \text{jets}$, $t\bar{t}$, and VV ; single top quark production; and $W + \text{jets}$. Another background arises from multijet production in which at least one jet is misidentified as an electron.

The major backgrounds in the $eejj$ channel are $Z/\gamma^* + \text{jets}$ and $t\bar{t}$ production. The $Z/\gamma^* + \text{jets}$ background is estimated from simulation and normalized to the data in a control region that comprises the initial selection plus a window of $80 < m_{ee} < 100$ GeV around the nominal Z boson mass; the latter criterion is applied to enrich the sample with $Z/\gamma^* + \text{jets}$ events. The m_{ee} distribution is corrected for the presence of non- $Z/\gamma^* + \text{jets}$ events in the data control region using simulation. The resulting normalization factor applied to the $Z/\gamma^* + \text{jets}$ simulated events is $R_Z = 0.97 \pm 0.01(\text{stat})$.

The contribution from $t\bar{t}$ events containing two electrons is estimated using a control region in data, which consists of events containing one electron and one muon, to which

all applicable $eejj$ selection criteria are applied. Residual backgrounds from other processes are subtracted using simulated event samples. Corrections for the branching fractions between the two states as well as for the differences in electron/muon identification and isolation efficiencies and acceptances are determined using simulation. The difference in the trigger efficiency between the one- and two-electron final states is corrected by reweighting each event in the $e\mu$ sample with the calculated efficiencies for the single electron final state.

After application of event selection requirements, the background contribution to the $eejj$ channel arising from single top quark production, $W + \text{jets}$, and VV is found to be small and is estimated from simulations.

The multijet background in the $eejj$ channel is estimated using control samples in data. The electron identification requirements for the calorimeter shower profile and track-cluster matching are relaxed to define a loose selection. We measure the probability that an electron candidate that passes the loose selection requirements also satisfies the electron identification and isolation criteria used in the analysis. This probability is obtained as a function of the candidate p_T and η . The events are required to have




FIG. 4. Data and background for events passing the initial selection requirements in the $eejj$ channel, shown for the variables used for final selection optimization: m_T (upper left), m_{ej} (upper right), S_T (lower left), and p_T^{miss} (lower right). “Other background” includes diboson, single top quark, and $Z/\gamma^* + \text{jets}$. Signal predictions for $m_{LQ} = 650$ and 1200 GeV hypotheses are overlaid on the plots. The last bin includes all events beyond the upper x -axis boundary.

exactly one loose electron, at least two jets, and low p_T^{miss} (<100 GeV). Contributions from electrons satisfying the full identification requirements are removed. The number of such electrons is calculated by comparing the number of candidates that pass the tight selection criteria minus the track-isolation requirement, with those that satisfy the track-isolation requirement but fail one of the other selection criteria. This sample is dominated by QCD multijet events. The distribution of multijet events in the $eejj$ channel following final selections is obtained by applying the measured probability twice to an event sample with two electrons passing loose electron requirements, and two or more jets that satisfy all the requirements of the signal selection. The normalization is obtained by scaling the weighted multijet sample to an orthogonal control region defined by inverting track-isolation requirement for electrons.

Distributions of kinematic variables for the $eejj$ channel in data, including those used in the final selections, have been studied at the initial selection level, and are found to agree with the background models within background estimation uncertainties. The distributions of S_T , m_{ej}^{min} , and m_{ee} are shown in Fig. 3.

The largest background in the $e\nu jj$ channel comes from W + jets and $t\bar{t}$ production. Single top quark, VV , and

$Z/\gamma^* + \text{jets}$ backgrounds have small contributions and are estimated from simulations. The QCD multijet background is estimated from control samples in data using the same probability for jets to be misidentified as electrons as is used in the background estimation for the $eejj$ channel. The number of multijet events at the final selection is obtained by selecting events having exactly one loose electron, large p_T^{miss} , and at least two jets satisfying the signal selection criteria, and weighting these with the probability of a jet being misidentified as an electron.

The background contributions from W + jets and $t\bar{t}$ are estimated from simulation and normalized to the data in a control region defined by requiring $50 < m_T < 110$ GeV after the initial selection. Then b -tagging information is used to distinguish W + jets from $t\bar{t}$ in the control region. The W + jets contribution is enhanced by requiring zero b -tagged jets in the event, while the $t\bar{t}$ control region is defined by requiring at least one b -tagged jet in the event. These regions each have a purity of about 75%. The normalization factors for the two backgrounds are determined from these control regions using

$$\begin{aligned} N_1 &= R_{t\bar{t}}N_{1,t\bar{t}} + R_W N_{1,W} + N_{1,0} \\ N_2 &= R_{t\bar{t}}N_{2,t\bar{t}} + R_W N_{2,W} + N_{2,0}, \end{aligned} \quad (1)$$

TABLE I. Systematic uncertainties for the $eejj$ and $e\nu jj$ channels. The values shown are calculated for the selections used in the $m_{LQ} = 1000$ GeV search hypothesis and reflect the variations in the event yields due to each source. Major backgrounds, namely $Z/\gamma^* + \text{jets}$ ($eejj$), W + jets and $t\bar{t}$ ($e\nu jj$), are normalized at the initial selection level when calculating the effect of shifts for various systematics.

Source of the uncertainty	$eejj$		$e\nu jj$	
	Signal (%)	Background (%)	Signal (%)	Background (%)
Electron energy scale	1.5	2.5	1.9	6.9
Electron energy resolution	0.2	5.3	0.1	4.9
Electron reconstruction efficiency	3.0	3.0	0.6	0.8
Electron identification efficiency	1.3	0.3	0.6	0.1
Trigger	1.1	1.4	9.5	7.6
Jet energy scale	0.5	0.9	0.5	2.3
Jet energy resolution	0.1	1.7	0.1	2.4
p_T^{miss}	0.8	13.1
$Z/\gamma^* + \text{jets}$ shape	...	5.6
$Z/\gamma^* + \text{jets}$ normalization	...	1.0
W + jets shape	7.1
W + jets normalization	1.1
W + jets sideband selection	10.0
W + jets b tagging	3.0
$t\bar{t}$ shape	10.4
$t\bar{t}$ normalization	...	1.0	...	1.0
$t\bar{t}$ b tagging	3.0
Diboson shape	...	3.4	...	3.2
QCD multijet	...	<0.1	...	2.6
Integrated luminosity	2.5	0.6	2.5	0.5
Pileup	0.2	1.0	0.4	1.4
PDF	2.8	3.0	2.9	4.7

where $N_{1(2)}$ is the number of events in the $t\bar{t}$ ($W + \text{jets}$) control region in data. The terms $N_{i,t\bar{t}}$ and $N_{i,W}$ are the numbers of $t\bar{t}$ and $W + \text{jets}$ events in the simulated samples, while $N_{i,O}$ is the number of events arising from other background sources, namely diboson, single top quark, $Z/\gamma^* + \text{jets}$ and multijet. The subscript $i = 1, 2$ refers to the two control regions described above. The background normalization factors $R_{t\bar{t}} = 0.92 \pm 0.01(\text{stat})$ and $R_W = 0.87 \pm 0.01(\text{stat})$ are then determined by solving Eq. (1).

The observed distributions of kinematic variables for the $eejj$ channel following the initial selection are found to agree with the background prediction within estimation uncertainties. The distributions of m_T , m_{ej} , S_T , and p_T^{miss} are shown in Fig. 4.

VI. SYSTEMATIC UNCERTAINTIES

The sources of systematic uncertainties considered in this analysis are listed in Table I. Uncertainties in the reconstruction of electrons, jets and p_T^{miss} affect the selected sample of events used in the analysis. The uncertainty due to the electron energy scale is obtained by shifting the

electron energy up and down by 2%. The uncertainty in the electron energy resolution is measured by smearing the electron energy by $\pm 10\%$ [78]. The uncertainties due to electron reconstruction and identification efficiencies are obtained by varying the corresponding scale factors applied to simulated events by ± 1 standard deviation with respect to their nominal values. The trigger efficiency for electrons is measured by utilizing the tag-and-probe method [79] in data, and parametrized as a function of electron p_T and η . The corresponding uncertainty depends on the number of data events and is almost entirely statistical in origin for the kinematic range studied in this analysis.

The uncertainty due to the jet energy scale is obtained by varying the nominal scale correction by ± 1 standard deviation and taking the maximum difference with respect to the nominal event yield. The jet energy resolution models the variation between the reconstructed and generated jets. The corresponding uncertainty is obtained by modifying the parametrization of this difference [74].

To determine uncertainties in p_T^{miss} , we consider up and down shifts in the jet energy scale and resolution, electron energy correction, and the scale corrections applied to the




FIG. 5. m_{ej}^{min} (left) and S_T (right) distributions for events passing the $eejj$ final selection for LQs of mass 650 (upper) and 1200 (lower) GeV. The predicted signal model distributions are shown, along with major backgrounds and “other background” which consists of the sum of the $W + \text{jets}$, diboson, single top quark, and $\gamma + \text{jets}$ contributions. The background contributions are stacked, while the signal distributions are plotted unstacked. The dark shaded region indicates the statistical and systematic uncertainty in the total background. The last bin includes all events beyond the upper x -axis boundary.

energy not associated with any PF candidates. For each variation, a new p_T^{miss} vector is computed for each event. The uncertainties corresponding to different variations in the quantities are then added in quadrature to determine the variation in p_T^{miss} , and the maximum difference of the event yield with respect to nominal is taken as the uncertainty.

Variations in the shape of the $Z/\gamma^* + \text{jets}$ ($eejj$ channel only), $W + \text{jets}$ and $t\bar{t}$ ($evjj$ channel only), and diboson (both channels) backgrounds are determined using simulated samples with renormalization and factorization scales independently varied up and down in the matrix element by a factor of two, yielding eight different combinations. The event yields are then calculated for each of these variations and the maximum variation with respect to nominal is taken as the systematic uncertainty. The corresponding normalization uncertainties are evaluated from the statistical uncertainties in the scale factors obtained while normalizing these backgrounds to data in the control regions. In the $evjj$ channel, an additional uncertainty of 10% is included to account for the observed differences associated with the

choice of the m_T range, defining the control region used to calculate the normalization scale factors. As described above, b -tagging is used to define the control region for $W + \text{jets}$ and $t\bar{t}$ normalization in the $evjj$ channel; therefore, the uncertainty in the b -tagging efficiency (3%) is taken into account.

The uncertainty in the QCD multijet background is assessed by using an independent data sample. This sample is required to have exactly two electron candidates satisfying loosened criteria applied to the track-cluster matching, the isolation (both track-based and calorimetric), and the shower profile. We compare the number of events in this sample, where one candidate satisfies the electron selection requirements, to that predicted by the multijet background method. This test is repeated on a subsample of the data after applying an S_T threshold of 320 GeV, which corresponds to the optimized final selection for an LQ mass of 200 GeV. The relative difference of 25% observed between the results of the two tests is taken as the systematic uncertainty in the probability for a jet to be misidentified as




FIG. 6. m_{ej} (left) and S_T (right) distributions for events passing the $evjj$ final selection for LQs of mass 650 (upper) and 1200 (lower) GeV. The predicted signal model distributions are shown, along with major backgrounds and “other background” which consists of the sum of $Z/\gamma^* + \text{jets}$, diboson, single top quark, and $\gamma + \text{jets}$ contributions. The background contributions are stacked, while the signal distributions are plotted unstacked. The dark shaded region indicates the statistical and systematic uncertainty in the total background. The last bin includes all events beyond the upper x-axis boundary.

an electron and applied in the $e\nu jj$ channel. For the $eejj$ case, we assume full correlation between the two electrons and take 50% as the uncertainty.

The uncertainty in the integrated luminosity is 2.5% [80]. An uncertainty in the modeling of pileup is evaluated by reweighting the simulated events after varying the inelastic pp cross section by $\pm 4.6\%$ [81]. The acceptance for both signal and backgrounds, and the expected background cross sections are affected by PDF uncertainties. We estimate this effect by evaluating the complete set of NNPDF 3.0 PDF eigenvectors, following the PDF4LHC prescription [52,82–85].

VII. RESULTS OF THE LEPTOQUARK SEARCH

After applying the final selection criteria shown in Fig. 2, the data are compared to SM background expectations for both channels and each m_{LQ} hypothesis. Distributions of $m_{e_j}^{\min}$ and S_T are shown in Fig. 5 for the $eejj$ channel with the selections applied for the 650 and 1200 GeV m_{LQ} hypotheses. Figure 6 shows the corresponding distributions




FIG. 7. Data, background, and expected signal yields after applying the final selection criteria for the $eejj$ (left) and $e\nu jj$ (right) channels. “Other background” includes diboson, single top quark, and $W + jets$ (for the $eejj$ channel) or $Z/\gamma^* + jets$ (for the $e\nu jj$ channel). The bin contents are correlated, because events selected for higher-mass LQ searches are a subset of those selected for lower mass searches.

of m_{e_j} and S_T for the $e\nu jj$ channel for the same mass hypotheses.

Figure 7 shows background, data, and expected signal for each LQ mass point after applying the final selection criteria. Signal efficiency times acceptance, along with tables listing event yields for signal, background, and data are provided in the Appendix. The data are found to be in agreement with SM background expectations in both channels. We set upper limits on the product of the cross section and branching fraction for scalar LQs as a function of m_{LQ} and β . The limits are calculated using the asymptotic approximation [86] of the CL_s modified frequentist approach [87–89]. Systematic uncertainties described in Sec. VI are modeled with log-normal probability density functions, while statistical uncertainties are modeled with gamma functions whose widths are calculated from the number of events in the control regions or simulated samples.

We set upper limits on the production cross section multiplied by the branching fraction β^2 or $2\beta(1-\beta)$ at 95% C.L. as a function of m_{LQ} . The expected and observed limits are shown with NLO predictions for the scalar LQ pair production cross section in Fig. 8 for both $eejj$ and $e\nu jj$




FIG. 8. Observed upper limits for scalar LQ pair-production cross section times β^2 (left) and $\beta(1-\beta)$ (right) at 95% C.L. obtained with the $eejj$ (left) and $e\nu jj$ (right) analysis. The median (dashed line), 68% (inner green band) and 95% (outer yellow band) confidence-interval expected limits are also shown.




FIG. 9. Expected and observed exclusion limits at 95% C.L. for pair production of first-generation scalar LQ shown in the β versus m_{LQ} plane for the individual $eejj$ and $evjj$ channels and their combination. The inner green and outer yellow bands represent the 68% and 95% confidence intervals on the expected limits.

channels. The observed limits are within two standard deviations of expectations from the background-only hypothesis. The uncertainty in the theoretical prediction for the LQ pair production cross section is calculated as the quadrature sum of the PDF uncertainty in the signal cross section and the uncertainty due to the choice of renormalization and factorization scales. The latter is estimated by independently varying the scales up and down by a factor of two.

Under the assumption $\beta = 1.0$, where only the $eejj$ channel contributes, first-generation scalar LQs with masses less than 1435 GeV are excluded at 95% C.L. compared to a median expected limit of 1465 GeV. For $\beta = 0.5$, using the $evjj$ channel alone, LQ masses are excluded below 1195 GeV with the corresponding expected limit being 1210 GeV. As both $eejj$ and $evjj$ decays contribute at β values smaller than 1, the LQ mass limit is improved using the combination of the two channels. In this combination, systematic uncertainties are considered to be fully correlated between the channels, while statistical uncertainties are treated as fully uncorrelated. Limits for a range of β values from 0 to 1 are set at 95% C.L. for both $eejj$ and $evjj$ channels, as well as for their combination, as shown in Fig. 9. In the $\beta = 0.5$ case, the combination excludes first-generation scalar LQs with masses less than 1270 GeV, compared to a median expected value of 1285 GeV.

VIII. R-PARITY VIOLATING SUPERSYMMETRY INTERPRETATION

Many new physics models predict the existence of particles with couplings of the type expected for LQs. One such model is R -parity violating supersymmetry (RPV SUSY) [90,91], where the superpartners of quarks or ‘squarks’ can decay into LQ-like final states. For example,




FIG. 10. Expected and observed upper limits at 95% C.L. on the RPV SUSY \tilde{t} squark pair production cross section as a function of $M_{\tilde{t}}$ for $c\tau = 0$ cm. The expected limits represent the median values, while the inner green and outer yellow bands are the 68% and 95% confidence intervals, respectively.

the top squark (\tilde{t}) can decay to a bottom quark and an electron. The topology of the resulting events is similar to an LQ decay and hence these events will pass our nominal selection for the $eejj$ channel.

The analysis is recast in terms of the possible production of prompt top-squark pairs ($c\tau = 0$ cm), with each \tilde{t} subsequently decaying to a bottom quark and an electron. Limits on the production cross section for \tilde{t} pairs are calculated from the $eejj$ data, accounting for the difference in branching fractions of LQ and \tilde{t} decays to electrons.

Figure 10 shows the expected and observed 95% C.L. upper limits on the RPV SUSY \tilde{t} pair production cross section as a function of the \tilde{t} squark mass ($m_{\tilde{t}}$). The observed exclusion limit is 1100 GeV for $c\tau = 0$ cm.

IX. SUMMARY

A search has been performed for the pair production of first-generation scalar leptoquarks in final states consisting of two high-momentum electrons and two jets, or one electron, large missing transverse momentum and two jets. The data sample used in the study corresponds to an integrated luminosity of 35.9 fb^{-1} recorded by the CMS experiment at $\sqrt{s} = 13$ TeV. The data are found to be in agreement with standard model background expectations and a lower limit at 95% confidence level is set on the scalar leptoquark mass at 1435 (1270) GeV for $\beta = 1.0$ (0.5), where β is the branching fraction of the leptoquark decay to an electron and a quark. These results constitute the most stringent limits on the mass of first-generation scalar leptoquarks to date. The data are also interpreted in the context of an R -parity violating supersymmetric model with promptly decaying top squarks, which can decay into leptoquark-like final states. Top squarks are excluded for masses below 1100 GeV.

ACKNOWLEDGMENTS

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC and the CMS detector provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES and CSF (Croatia); RPF (Cyprus); SENESCYT (Ecuador); MoER, ERC IUT, and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); NKFI (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Dubna); MON, RosAtom, RAS, RFBR, and NRC KI (Russia); MESTD (Serbia); SEIDI, CPAN, PCTI, and FEDER (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); ThEPCenter, IPST, STAR, and NSTDA (Thailand); TUBITAK and TAEK (Turkey); NASU and SFFR (Ukraine); STFC (United Kingdom); DOE and NSF (USA). Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, Contract No. 675440 (European Union); the Leventis Foundation; the A. P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la

Formation *à la* Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F. R. S.-FNRS and FWO (Belgium) under the “Excellence of Science—EOS”—be.h Project No. 30820817; the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Lendület (“Momentum”) Program and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, the New National Excellence Program ÚNKP, the NKFI Research Grants No. 123842, No. 123959, No. 124845, No. 124850, and No. 125105 (Hungary); the Council of Science and Industrial Research, India; the HOMING PLUS program of the Foundation for Polish Science, cofinanced from European Union, Regional Development Fund, the Mobility Plus program of the Ministry of Science and Higher Education, the National Science Center (Poland), Contracts Harmonia No. 2014/14/M/ST2/00428, Opus No. 2014/13/B/ST2/02543, No. 2014/15/B/ST2/03998, and No. 2015/19/B/ST2/02861, Sonata-bis No. 2012/07/E/ST2/01406; the National Priorities Research Program by Qatar National Research Fund; the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, Grant No. MDM-2015-0509 and the Programa Severo Ochoa del Principado de Asturias; the Thalís and Aristeia programs cofinanced by EU-ESF and the Greek NSRF; the Rachadapisek Sompot Fund for Postdoctoral Fellowship, Chulalongkorn University and the Chulalongkorn Academic into Its 2nd Century Project Advancement Project (Thailand); the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

APPENDIX: EFFICIENCIES AND EVENT YIELDS

In Fig. 11 the product of signal acceptance and efficiency is shown after final optimized selections as a function of m_{LQ} for the $eejj$ (left) and $e\nu jj$ (right) channels. Tables II and III list the number of events passing the final selection criteria in data and the various background components as a function of m_{LQ} for the $eejj$ and $e\nu jj$ channels, respectively.




FIG. 11. The product of signal acceptance and efficiency after final optimized selections, as a function of m_{LQ} for the $eejj$ (left) and $e\nu jj$ (right) channels.

TABLE II. Event yields after the optimized $eejj$ selections. Uncertainties are statistical except for the total background, where both statistical and systematic uncertainties are shown. An entry of 0.0 quoted for the uncertainty indicates that its value is negligibly small. LQ masses are given in units of GeV and init. sel. refers to initial selection.

LQ mass	Signal	$Z/\gamma^* + \text{jets}$	$t\bar{t}$	Multijet	VV, W, single t, $\gamma + \text{jets}$	Total background	Data
init. sel.	...	41600 ± 49	7100 ± 68	26 ± 0.1	2400 ± 36	$51100 \pm 91 \pm 2700$	50585
200	311500 ± 3300	1900 ± 16	2300 ± 39	15 ± 0.1	630 ± 18	$4800 \pm 46 \pm 120$	4709
250	137400 ± 1200	910 ± 11	1200 ± 29	9.1 ± 0.1	380 ± 14	$2500 \pm 34 \pm 69$	2426
300	63160 ± 510	470 ± 4.2	630 ± 22	4.8 ± 0.0	$220^{+10}_{-9.5}$	$1300^{+24}_{-24} \pm 24$	1278
350	30150 ± 230	250 ± 2.7	310 ± 15	2.5 ± 0.0	$140^{+9.5}_{-8.6}$	$700^{+18}_{-18} \pm 27$	652
400	15440 ± 110	140 ± 1.8	150 ± 11	1.0 ± 0.0	$89^{+7.2}_{-6.2}$	$380^{+13}_{-13} \pm 11$	376
450	8260 ± 60	85 ± 1.5	79 ± 7.7	0.6 ± 0.0	$49^{+2.3}_{-2.3}$	$210^{+8.2}_{-8.1} \pm 5.3$	209
500	4700 ± 33	54 ± 1.1	36 ± 5.5	0.3 ± 0.0	$30^{+2.0}_{-1.9}$	$120^{+6.0}_{-5.9} \pm 4.4$	128
550	2830 ± 19	33 ± 0.8	15 ± 4.0	0.2 ± 0.0	$22^{+1.8}_{-1.8}$	$70^{+4.5}_{-4.5} \pm 2.6$	84
600	1750 ± 12	21 ± 0.6	9.6 ± 3.3	0.1 ± 0.0	$16^{+1.6}_{-1.6}$	$47^{+3.7}_{-3.7} \pm 1.9$	58
650	1110 ± 7.2	15 ± 0.6	7.7 ± 2.9	0.1 ± 0.0	$11^{+1.4}_{-1.3}$	$34^{+3.2}_{-3.2} \pm 1.3$	37
700	718 ± 4.5	12 ± 0.5	3.7 ± 2.2	0.1 ± 0.0	$7.3^{+1.2}_{-1.2}$	$23^{+2.6}_{-2.6} \pm 1.0$	28
750	470 ± 2.9	7.8 ± 0.3	2.0 ± 1.9	0.0 ± 0.0	$5.5^{+1.1}_{-1.1}$	$15^{+2.2}_{-2.2} \pm 0.6$	17
800	320 ± 1.9	6.4 ± 0.4	$1.1^{+0.5}_{-0.4}$	0.0 ± 0.0	$3.5^{+1.1}_{-0.9}$	$11^{+1.2}_{-1.1} \pm 0.6$	13
850	220 ± 1.3	4.9 ± 0.3	$1.5^{+0.7}_{-0.5}$	0.0 ± 0.0	$2.8^{+1.0}_{-0.6}$	$9.2^{+1.3}_{-0.8} \pm 0.5$	10
900	150 ± 0.9	4.0 ± 0.3	$0.0^{+1.2}_{-0.0}$	0.0 ± 0.0	$2.6^{+0.8}_{-0.5}$	$6.6^{+1.4}_{-0.6} \pm 0.4$	8
950	110 ± 0.6	3.6 ± 0.5	$0.0^{+0.9}_{-0.0}$	0.0 ± 0.0	$2.1^{+0.7}_{-0.5}$	$5.7^{+1.3}_{-0.7} \pm 0.3$	5
1000	77 ± 0.4	2.2 ± 0.1	$0.0^{+0.7}_{-0.0}$	0.0 ± 0.0	$1.9^{+0.7}_{-0.4}$	$4.1^{+1.0}_{-0.5} \pm 0.2$	5
1050	55 ± 0.3	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1100	41 ± 0.2	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1150	31 ± 0.2	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1200	23 ± 0.1	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1250	17 ± 0.1	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1300	13 ± 0.1	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1350	9.8 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1400	7.4 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1450	5.6 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1500	4.2 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1550	3.2 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1600	2.4 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1650	1.8 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1700	1.4 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1750	1.1 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1800	0.8 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1850	0.6 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1900	0.5 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
1950	0.4 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4
2000	0.3 ± 0.0	1.8 ± 0.1	$0.0^{+0.3}_{-0.0}$	0.0 ± 0.0	$1.4^{+0.6}_{-0.4}$	$3.2^{+0.7}_{-0.4} \pm 0.2$	4

TABLE III. Event yields after the optimized $e\nu jj$ selections. Uncertainties are statistical except for the total background, where both statistical and systematic uncertainties are shown. An entry of 0.0 quoted for the uncertainty indicates that its value is negligibly small. LQ masses are given in units of GeV and init. sel. refers to initial selection.

LQ mass	Signal	$W + \text{jets}$	$t\bar{t}$	Multijet	VV, Z, single t , $\gamma + \text{jets}$	Total background	Data
init. sel.	...	47900 ± 160	66900 ± 110	2800 ± 15	11300 ± 72	$128900 \pm 210 \pm 8800$	125076
200	130800 ± 1600	40100 ± 150	52800 ± 94	2100 ± 11	9600 ± 57	$104500 \pm 190 \pm 7300$	101618
250	44200 ± 520	1800 ± 25	3800 ± 25	300 ± 2.3	1300 ± 38	$7100 \pm 52 \pm 430$	7151
300	19800 ± 220	800 ± 15	1400 ± 16	120 ± 1.4	660 ± 37	$3000 \pm 43 \pm 170$	3164
350	9800 ± 100	410 ± 13	610 ± 10	62 ± 1.0	330 ± 11	$1400 \pm 20 \pm 88$	1539
400	5100 ± 51	230 ± 8.9	300 ± 7.2	37 ± 0.8	200 ± 10	$760 \pm 15 \pm 74$	847
450	2900 ± 27	150 ± 6.0	160 ± 5.2	28 ± 0.8	120 ± 9.6	$460 \pm 12 \pm 31$	496
500	1700 ± 15	90 ± 4.1	88 ± 3.9	21 ± 0.8	$75_{-3.3}^{+3.9}$	$270_{-6.6}^{+6.9} \pm 21$	298
550	990 ± 8.8	59 ± 5.2	49 ± 2.9	9.1 ± 0.4	$53_{-2.9}^{+3.5}$	$170_{-6.6}^{+6.9} \pm 13$	195
600	620 ± 5.3	45 ± 5.1	32 ± 2.3	6.1 ± 0.4	$36_{-2.2}^{+2.8}$	$120_{-6.0}^{+6.3} \pm 12$	132
650	400 ± 3.3	34 ± 5.0	20 ± 1.8	5.0 ± 0.4	$26_{-1.9}^{+2.5}$	$84_{-5.7}^{+5.9} \pm 8.1$	94
700	270 ± 2.1	22 ± 1.2	12 ± 1.5	4.2 ± 0.5	$18_{-1.5}^{+2.1}$	$56_{-2.5}^{+2.9} \pm 6.1$	71
750	180 ± 1.4	15 ± 0.9	10 ± 1.3	3.7 ± 0.5	$13_{-1.3}^{+2.1}$	$42_{-2.1}^{+2.7} \pm 4.9$	49
800	130 ± 0.9	13 ± 1.0	6.3 ± 1.0	3.4 ± 0.6	$9.8_{-1.1}^{+2.0}$	$32_{-1.9}^{+2.5} \pm 4.6$	38
850	86 ± 0.6	13 ± 1.1	5.2 ± 0.9	3.2 ± 0.7	$7.0_{-1.2}^{+2.0}$	$28_{-2.0}^{+2.6} \pm 4.8$	28
900	61 ± 0.4	11 ± 1.2	3.8 ± 0.8	3.0 ± 0.7	$6.3_{-1.1}^{+2.0}$	$24_{-2.0}^{+2.6} \pm 4.1$	21
950	44 ± 0.3	8.4 ± 1.0	3.0 ± 0.7	0.7 ± 0.1	$5.7_{-1.1}^{+2.0}$	$18_{-1.6}^{+2.3} \pm 3.3$	20
1000	31 ± 0.2	7.9 ± 0.9	2.2 ± 0.6	0.6 ± 0.1	$4.8_{-1.1}^{+2.0}$	$16_{-1.5}^{+2.3} \pm 2.8$	15
1050	23 ± 0.2	7.1 ± 0.9	$1.4_{-0.5}^{+0.7}$	0.5 ± 0.1	$4.4_{-1.1}^{+2.0}$	$13_{-1.4}^{+2.3} \pm 2.5$	14
1100	17 ± 0.1	5.9 ± 0.8	$1.2_{-0.4}^{+0.6}$	0.5 ± 0.1	$4.0_{-1.0}^{+2.0}$	$12_{-1.4}^{+2.3} \pm 2.1$	12
1150	12 ± 0.1	5.4 ± 0.9	$0.9_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.3_{-1.0}^{+2.0}$	$10_{-1.4}^{+2.3} \pm 1.7$	12
1200	9.1 ± 0.1	5.2 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.2_{-1.0}^{+2.0}$	$9.5_{-1.5}^{+2.3} \pm 1.6$	10
1250	7.1 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1300	5.4 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1350	4.1 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1400	3.1 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1450	2.4 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1500	1.9 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1550	1.4 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1600	1.1 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1650	0.8 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1700	0.6 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1750	0.5 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1800	0.4 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1850	0.3 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1900	0.2 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
1950	0.2 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9
2000	0.1 ± 0.0	5.0 ± 1.1	$0.7_{-0.4}^{+0.6}$	0.4 ± 0.1	$3.0_{-1.0}^{+2.0}$	$9.1_{-1.5}^{+2.3} \pm 1.5$	9

- [1] S.L. Glashow, Partial-symmetries of weak interactions, *Nucl. Phys.* **22**, 579 (1961).
- [2] S. Weinberg, A Model of Leptons, *Phys. Rev. Lett.* **19**, 1264 (1967).
- [3] A. Salam, Weak and electromagnetic interactions, in *Elementary Particle Physics: Relativistic Groups and Analyticity*, edited by N. Svartholm (Almqvist & Wiksell, Stockholm, 1968), p. 367, proceedings of the eighth Nobel symposium.
- [4] J.C. Pati and A. Salam, Unified lepton-hadron symmetry and a gauge theory of the basic interactions, *Phys. Rev. D* **8**, 1240 (1973).
- [5] J.C. Pati and A. Salam, Lepton number as the fourth "color", *Phys. Rev. D* **10**, 275 (1974).
- [6] H. Georgi and S. Glashow, Unity of all Elementary-Particle Forces, *Phys. Rev. Lett.* **32**, 438 (1974).
- [7] H. Murayama and T. Yanagida, A viable SU(5) GUT with light leptoquark bosons, *Mod. Phys. Lett. A* **07**, 147 (1992).
- [8] H. Fritzsch and P. Minkowski, Unified interactions of leptons and hadrons, *Ann. Phys. (N.Y.)* **93**, 193 (1975).
- [9] G. Senjanović and A. Sokorac, Light lepto-quarks in SO(10), *Z. Phys. C* **20**, 255 (1983).
- [10] P.H. Frampton and B.-H. Lee, SU(15) Grand Unification, *Phys. Rev. Lett.* **64**, 619 (1990).
- [11] P.H. Frampton and T.W. Kephart, Higgs sector and proton decay in SU(15) grand unification, *Phys. Rev. D* **42**, 3892 (1990).
- [12] S. Dimopoulos and L. Susskind, Mass without scalars, *Nucl. Phys.* **B155**, 237 (1979).
- [13] S. Dimopoulos, Technicoloured signatures, *Nucl. Phys.* **B168**, 69 (1980).
- [14] E. Eichten and K. Lane, Dynamical breaking of weak interaction symmetries, *Phys. Lett.* **90B**, 125 (1980).
- [15] J.L. Hewett and T.G. Rizzo, Low-energy phenomenology of superstring-inspired E_6 models, *Phys. Rep.* **183**, 193 (1989).
- [16] B. Schrempp and F. Schrempp, Light leptoquarks, *Phys. Lett.* **153B**, 101 (1985).
- [17] W. Buchmüller, R. Rückl, and D. Wyler, Leptoquarks in lepton-quark collisions, *Phys. Lett. B* **191**, 442 (1987).
- [18] W. Buchmüller and D. Wyler, Constraints on SU(5)-type leptoquarks, *Phys. Lett. B* **177**, 377 (1986).
- [19] O. Shanker, $\pi\ell 2$, $K\ell 3$, and $K^0\bar{K}^0$ constraints on leptoquarks and supersymmetric particles, *Nucl. Phys.* **B204**, 375 (1982).
- [20] LHCb Collaboration, Measurement of Form-Factor-Independent Observables in the Decay $B^0 \rightarrow K^{*0}\mu^+\mu^-$, *Phys. Rev. Lett.* **111**, 191801 (2013).
- [21] LHCb Collaboration, Test of Lepton Universality Using $B^+ \rightarrow K^+\ell^+\ell^-$ Decays, *Phys. Rev. Lett.* **113**, 151601 (2014).
- [22] LHCb Collaboration, Angular analysis of the $B^0 \rightarrow K^{*0}\mu^+\mu^-$ decay using 3 fb^{-1} of integrated luminosity, *J. High Energy Phys.* **02** (2016) 104.
- [23] LHCb Collaboration, Test of lepton universality with $B^0 \rightarrow K^{*0}\ell^+\ell^-$ decays, *J. High Energy Phys.* **08** (2017) 055.
- [24] S. Wehle *et al.* (Belle Collaboration), Lepton-flavor-dependent angular analysis of $B \rightarrow K^*\ell^+\ell^-$, *Phys. Rev. Lett.* **118**, 111801 (2017).
- [25] G. Hiller, Lepton nonuniversality anomalies and implications, [arXiv:1804.02011](https://arxiv.org/abs/1804.02011).
- [26] I. Doršner, S. Fajfer, A. Greljo, J.F. Kamenik, and N. Košnik, Physics of leptoquarks in precision experiments and at particle colliders, *Phys. Rep.* **641**, 1 (2016).
- [27] G. Abbiendi *et al.* (OPAL Collaboration), Search for pair-produced leptoquarks in e^+e^- interactions at $\sqrt{s} \simeq 189\text{--}209\text{ GeV}$, *Eur. Phys. J. C* **31**, 281 (2003).
- [28] S. Chekanov *et al.* (ZEUS Collaboration), Search for resonance decays to lepton + jet at DESY HERA and limits on leptoquarks, *Phys. Rev. D* **68**, 052004 (2003).
- [29] F.D. Aaron *et al.* (H1 Collaboration), Search for first generation leptoquarks in ep collisions at HERA, *Phys. Lett. B* **704**, 388 (2011).
- [30] D. Acosta *et al.* (CDF Collaboration), Search for first-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$, *Phys. Rev. D* **72**, 051107 (2005).
- [31] V.M. Abazov *et al.* (D0 Collaboration), Search for pair production of first-generation leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$, *Phys. Lett. B* **681**, 224 (2009).
- [32] CMS Collaboration, Search for pair production of first and second generation leptoquarks in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$, *Phys. Rev. D* **93**, 032004 (2016).
- [33] CMS Collaboration, Search for single production of scalar leptoquarks in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$, *Phys. Rev. D* **93**, 032005 (2016); *Phys. Rev. D* Erratum **95**, 039906(E) (2017).
- [34] ATLAS Collaboration, Search for scalar leptoquarks in pp collisions at $\sqrt{s} = 13\text{ TeV}$ with the ATLAS experiment, *New J. Phys.* **18**, 093016 (2016).
- [35] M. Krämer, T. Plehn, M. Spira, and P.M. Zerwas, Pair production of scalar leptoquarks at the CERN LHC, *Phys. Rev. D* **71**, 057503 (2005).
- [36] CMS Collaboration, The CMS trigger system, *J. Instrum.* **12**, P01020 (2017).
- [37] CMS Collaboration, The CMS experiment at the CERN LHC, *J. Instrum.* **3**, S08004 (2008).
- [38] T. Sjöstrand, S. Ask, J.R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C.O. Rasmussen, and P.Z. Skands, An introduction to PYTHIA 8.2, *Comput. Phys. Commun.* **191**, 159 (2015).
- [39] R.D. Ball, V. Bertone, S. Carrazza, C.S. Deans, L. Del Debbio, S. Forte, A. Guffanti, N.P. Hartland, J.I. Latorre, J. Rojo, and M. Ubiali, Parton distributions with LHC data, *Nucl. Phys.* **B867**, 244 (2013).
- [40] M. Krämer, T. Plehn, M. Spira, and P.M. Zerwas, Pair Production of Scalar Leptoquarks at the Fermilab Tevatron, *Phys. Rev. Lett.* **79**, 341 (1997).
- [41] J. Pumplin, D.R. Stump, J. Huston, H.L. Lai, P. Nadolsky, and W.K. Tung, New generation of parton distributions with uncertainties from global QCD analysis, *J. High Energy Phys.* **07** (2002) 012.
- [42] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
- [43] R. Frederix and S. Frixione, Merging meets matching in MC@NLO, *J. High Energy Phys.* **12** (2012) 061.

- [44] P. Artoisenet, R. Frederix, O. Mattelaer, and R. Rietkerk, Automatic spin-entangled decays of heavy resonances in Monte Carlo simulations, *J. High Energy Phys.* **03** (2013) 015.
- [45] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, *J. High Energy Phys.* **11** (2004) 040.
- [46] S. Frixione, P. Nason, and C. Oleari, Matching NLO QCD computations with parton shower simulations: The POWHEG method, *J. High Energy Phys.* **11** (2007) 070.
- [47] S. Alioli, P. Nason, C. Oleari, and E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: The POWHEG BOX, *J. High Energy Phys.* **06** (2010) 043.
- [48] S. Alioli, P. Nason, C. Oleari, and E. Re, NLO single-top production matched with shower in POWHEG: s - and t -channel contributions, *J. High Energy Phys.* **09** (2009) 111; *J. High Energy Phys. Erratum* **02** (2010) 11(E).
- [49] E. Re, Single-top Wt -channel production matched with parton showers using the POWHEG method, *Eur. Phys. J. C* **71**, 1547 (2011).
- [50] S. Frixione, P. Nason, and G. Ridolfi, A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction, *J. High Energy Phys.* **09** (2007) 126.
- [51] J. Alwall, S. Höche, F. Krauss, N. Lavesson, L. Lönnblad, F. Maltoni, M. L. Mangano, M. Moretti, C. G. Papadopoulos, F. Piccinini, S. Schumann, M. Treccani, J. Winter, and M. Worek, Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions, *Eur. Phys. J. C* **53**, 473 (2008).
- [52] R. D. Ball *et al.* (NNPDF Collaboration), Parton distributions for the LHC Run II, *J. High Energy Phys.* **04** (2015) 040.
- [53] Y. Li and F. Petriello, Combining QCD and electroweak corrections to dilepton production in the framework of the FEWZ simulation code, *Phys. Rev. D* **86**, 094034 (2012).
- [54] M. Aliev, H. Lacker, U. Langenfeld, S. Moch, P. Uwer, and M. Wiedermann, HATHOR—hadronic top and heavy quarks cross section calculator, *Comput. Phys. Commun.* **182**, 1034 (2011).
- [55] P. Kant, O. M. Kind, T. Kintscher, T. Lohse, T. Martini, S. Mölbitz, P. Rieck, and P. Uwer, HATHOR for single top-quark production: Updated predictions and uncertainty estimates for single top-quark production in hadronic collisions, *Comput. Phys. Commun.* **191**, 74 (2015).
- [56] N. Kidonakis, Top quark production, in *Helmholtz International Summer School on Physics of Heavy Quarks and Hadrons (HQ 2013): JINR, Dubna, Russia, 2013*, <http://www-library.desy.de/preparch/desy/proc/proc13-03/Kidonakis.pdf>.
- [57] M. Beneke, P. Falgari, S. Klein, and C. Schwinn, Hadronic top-quark pair production with NNLL threshold resummation, *Nucl. Phys.* **B855**, 695 (2012).
- [58] M. Cacciari, M. Czakon, M. Mangano, A. Mitov, and P. Nason, Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation, *Phys. Lett. B* **710**, 612 (2012).
- [59] P. Bärnreuther, M. Czakon, and A. Mitov, Percent-Level-Precision Physics at the Tevatron: Next-to-Next-to-Leading Order QCD Corrections to $q\bar{q} \rightarrow t\bar{t} + X$, *Phys. Rev. Lett.* **109**, 132001 (2012).
- [60] M. Czakon and A. Mitov, NNLO corrections to top-pair production at hadron colliders: The all-fermionic scattering channels, *J. High Energy Phys.* **12** (2012) 054.
- [61] M. Czakon and A. Mitov, NNLO corrections to top pair production at hadron colliders: The quark-gluon reaction, *J. High Energy Phys.* **01** (2013) 080.
- [62] M. Czakon, P. Fiedler, and A. Mitov, Total Top-Quark Pair-Production Cross Section at Hadron Colliders Through $\mathcal{O}(\alpha_s^4)$, *Phys. Rev. Lett.* **110**, 252004 (2013).
- [63] M. Czakon and A. Mitov, Top++: A program for the calculation of the top-pair cross-section at hadron colliders, *Comput. Phys. Commun.* **185**, 2930 (2014).
- [64] CMS Collaboration, Event generator tunes obtained from underlying event and multiparton scattering measurements, *Eur. Phys. J. C* **76**, 155 (2016).
- [65] CMS Collaboration, Investigations of the impact of the parton shower tuning in PYTHIA 8 in the modelling of $t\bar{t}$ at $\sqrt{s} = 8$ and 13 TeV, CMS Physics Analysis Summary Report No. CMS-PAS-TOP-16-021, 2016, <https://cds.cern.ch/record/2235192>.
- [66] S. Agostinelli *et al.* (GEANT4 Collaboration), GEANT4—a simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [67] J. Allison *et al.*, Geant4 developments and applications, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006).
- [68] CMS Collaboration, Particle-flow reconstruction and global event description with the CMS detector, *J. Instrum.* **12**, P10003 (2017).
- [69] CMS Collaboration, Performance of electron reconstruction and selection with the CMS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV, *J. Instrum.* **10**, P06005 (2015).
- [70] M. Cacciari, G. P. Salam, and G. Soyez, The anti- k_T jet clustering algorithm, *J. High Energy Phys.* **04** (2008) 063.
- [71] M. Cacciari, G. P. Salam, and G. Soyez, FastJet user manual, *Eur. Phys. J. C* **72**, 1896 (2012).
- [72] CMS Collaboration, Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. Instrum.* **13**, P06015 (2018).
- [73] CMS Collaboration, Jet algorithms performance in 13 TeV data, CMS Physics Analysis Summary Report No. CMS-PAS-JME-16-003, 2017, <http://cds.cern.ch/record/2256875>.
- [74] CMS Collaboration, Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV, *J. Instrum.* **12**, P02014 (2017).
- [75] CMS Collaboration, Performance of missing transverse momentum reconstruction algorithms in proton-proton collisions at $\sqrt{s} = 8$ TeV with the CMS detector, CMS Physics Analysis Summary Report No. CMS-PAS-JME-12-002, 2013, <http://cds.cern.ch/record/1543527>.
- [76] CMS Collaboration, Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV, *J. Instrum.* **13**, P05011 (2018).
- [77] G. Punzi, Sensitivity of searches for new signals and its optimization, in *PHYSTAT 2003: Statistical problems in particle physics, astrophysics and cosmology* (Stanford, USA, 2003), Conference C03-09-08, [arXiv:physics/0308063](https://arxiv.org/abs/physics/0308063).

- [78] CMS Collaboration, Performance of photon reconstruction and identification with the CMS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV, *J. Instrum.* **10**, P08010 (2015).
- [79] CMS Collaboration, Measurements of inclusive W and Z cross sections in pp collisions at $\sqrt{s} = 7$ TeV, *J. High Energy Phys.* **01** (2011) 080.
- [80] CMS Collaboration, CMS luminosity measurements for the 2016 data taking period, CMS physics analysis summary Report No. CMS-PAS-LUM-17-001, 2017, <http://cds.cern.ch/record/2257069>.
- [81] CMS Collaboration, Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **07** (2018) 161.
- [82] M. Botje, J. Butterworth, A. Cooper-Sarkar, A. de Roeck, J. Feltesse, S. Forte, A. Glazov, J. Huston, R. McNulty, T. Sjöstrand, and R.S. Thorne, The PDF4LHC Working Group Interim Recommendations, [arXiv:1101.0538](https://arxiv.org/abs/1101.0538).
- [83] S. Alekhin *et al.*, The PDF4LHC Working Group Interim Report, [arXiv:1101.0536](https://arxiv.org/abs/1101.0536).
- [84] M.R. Whalley, D. Bourilkov, and R.C. Group, The Les Houches accord PDFs (LHAPDF) and LHAGLUE, in *HERA and the LHC: A Workshop on the implications of HERA for LHC physics. Proceedings, Part B*, [arXiv:hep-ph/0508110](https://arxiv.org/abs/hep-ph/0508110).
- [85] D Bourilkov, R.C. Group, and M.R. Whalley, LHAPDF: PDF use from the Tevatron to the LHC, in *TeV4LHC Workshop—4th meeting Batavia, Illinois, 2005*, [arXiv:hep-ph/0605240](https://arxiv.org/abs/hep-ph/0605240).
- [86] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011); *Eur. Phys. J. C* Erratum **73**, 2501(E) (2013).
- [87] T. Junk, Confidence level computation for combining searches with small statistics, *Nucl. Instrum. Methods Phys. Res., Sect. A* **434**, 435 (1999).
- [88] A.L. Read, Presentation of search results: The CL_s technique, in *Advanced statistical techniques in particle physics proceedings, conference, Durham, UK, 2002*, DOI: 10.1088/0954-3899/28/10/313.
- [89] ATLAS and CMS Collaboration, The LHC Higgs Combination Group, Procedure for the LHC Higgs boson search combination in Summer 2011, Technical Reports No. CMS NOTE-2011/005 and No. ATL-PHYS-PUB-2011-11, 2011, <https://cds.cern.ch/record/1379837>.
- [90] H. K. Dreiner, An introduction to explicit *R*-parity violation, *Pramana* **51**, 123 (1998).
- [91] R. Barbier, C. Bérat, M. Besançon, M. Chemtob, A. Deandrea, E. Dudas, P. Fayet, S. Lavignac, G. Moreau, E. Perez, and Y. Sirois, *R*-parity-violating supersymmetry, *Phys. Rep.* **420**, 1 (2005).

A. M. Sirunyan,¹ A. Tumasyan,¹ W. Adam,² F. Ambroggi,² E. Asilar,² T. Bergauer,² J. Brandstetter,² M. Dragicevic,² J. Erö,² A. Escalante Del Valle,² M. Flechl,² R. Frühwirth,^{2,a} V. M. Ghete,² J. Hrubec,² M. Jeitler,^{2,a} N. Krammer,² I. Krätschmer,² D. Liko,² T. Madlener,² I. Mikulec,² N. Rad,² H. Rohringer,² J. Schieck,^{2,a} R. Schöfbeck,² M. Spanring,² D. Spitzbart,² A. Taurok,² W. Waltenberger,² J. Wittmann,² C.-E. Wulz,^{2,a} M. Zarucki,² V. Chekhovsky,³ V. Mossolov,³ J. Suarez Gonzalez,³ E. A. De Wolf,⁴ D. Di Croce,⁴ X. Janssen,⁴ J. Lauwers,⁴ M. Pieters,⁴ H. Van Haevermaet,⁴ P. Van Mechelen,⁴ N. Van Remortel,⁴ S. Abu Zeid,⁵ F. Blekman,⁵ J. D’Hondt,⁵ J. De Clercq,⁵ K. Deroover,⁵ G. Flouris,⁵ D. Lontkovskiy,⁵ S. Lowette,⁵ I. Marchesini,⁵ S. Moortgat,⁵ L. Moreels,⁵ Q. Python,⁵ K. Skovpen,⁵ S. Tavernier,⁵ W. Van Doninck,⁵ P. Van Mulders,⁵ I. Van Parijs,⁵ D. Beghin,⁶ B. Bilin,⁶ H. Brun,⁶ B. Clerbaux,⁶ G. De Lentdecker,⁶ H. Delannoy,⁶ B. Dorney,⁶ G. Fasanella,⁶ L. Favart,⁶ R. Goldouzian,⁶ A. Grebenyuk,⁶ A. K. Kalsi,⁶ T. Lenzi,⁶ J. Luetic,⁶ N. Postiau,⁶ E. Starling,⁶ L. Thomas,⁶ C. Vander Velde,⁶ P. Vanlaer,⁶ D. Vannerom,⁶ Q. Wang,⁶ T. Cornelis,⁷ D. Dobur,⁷ A. Fagot,⁷ M. Gul,⁷ I. Khvastunov,^{7,b} D. Poyraz,⁷ C. Roskas,⁷ D. Trocino,⁷ M. Tytgat,⁷ W. Verbeke,⁷ B. Vermassen,⁷ M. Vit,⁷ N. Zaganidis,⁷ H. Bakhshiansohi,⁸ O. Bondu,⁸ S. Brochet,⁸ G. Bruno,⁸ C. Caputo,⁸ P. David,⁸ C. Delaere,⁸ M. Delcourt,⁸ A. Giammanco,⁸ G. Krintiras,⁸ V. Lemaître,⁸ A. Magitteri,⁸ K. Piotrkowski,⁸ A. Saggio,⁸ M. Vidal Marono,⁸ S. Wertz,⁸ J. Zobec,⁸ F. L. Alves,⁹ G. A. Alves,⁹ M. Correa Martins Junior,⁹ G. Correia Silva,⁹ C. Hensel,⁹ A. Moraes,⁹ M. E. Pol,⁹ P. Rebello Teles,⁹ E. Belchior Batista Das Chagas,¹⁰ W. Carvalho,¹⁰ J. Chinellato,^{10,c} E. Coelho,¹⁰ E. M. Da Costa,¹⁰ G. G. Da Silveira,^{10,d} D. De Jesus Damiao,¹⁰ C. De Oliveira Martins,¹⁰ S. Fonseca De Souza,¹⁰ H. Malbouisson,¹⁰ D. Matos Figueiredo,¹⁰ M. Melo De Almeida,¹⁰ C. Mora Herrera,¹⁰ L. Mundim,¹⁰ H. Nogima,¹⁰ W. L. Prado Da Silva,¹⁰ L. J. Sanchez Rosas,¹⁰ A. Santoro,¹⁰ A. Sznajder,¹⁰ M. Thiel,¹⁰ E. J. Tonelli Manganote,^{10,c} F. Torres Da Silva De Araujo,¹⁰ A. Vilela Pereira,¹⁰ S. Ahuja,^{11a} C. A. Bernardes,^{11a} L. Calligaris,^{11a} T. R. Fernandez Perez Tomei,^{11a} E. M. Gregores,^{11a,11b} P. G. Mercadante,^{11a,11b} S. F. Novaes,^{11a} Sandra S. Padula,^{11a} A. Aleksandrov,¹² R. Hadjiiska,¹² P. Iaydjiev,¹² A. Marinov,¹² M. Misheva,¹² M. Rodozov,¹² M. Shopova,¹² G. Sultanov,¹² A. Dimitrov,¹³ L. Litov,¹³ B. Pavlov,¹³ P. Petkov,¹³ W. Fang,^{14,e} X. Gao,^{14,e} L. Yuan,¹⁴ M. Ahmad,¹⁵ J. G. Bian,¹⁵ G. M. Chen,¹⁵ H. S. Chen,¹⁵ M. Chen,¹⁵ Y. Chen,¹⁵ C. H. Jiang,¹⁵ D. Leggat,¹⁵ H. Liao,¹⁵ Z. Liu,¹⁵ F. Romeo,¹⁵ S. M. Shaheen,^{15,f} A. Spiezia,¹⁵ J. Tao,¹⁵ Z. Wang,¹⁵ E. Yazgan,¹⁵ H. Zhang,¹⁵ S. Zhang,^{15,f} J. Zhao,¹⁵ Y. Ban,¹⁶ G. Chen,¹⁶

A. Levin,¹⁶ J. Li,¹⁶ L. Li,¹⁶ Q. Li,¹⁶ Y. Mao,¹⁶ S. J. Qian,¹⁶ D. Wang,¹⁶ Y. Wang,¹⁷ C. Avila,¹⁸ A. Cabrera,¹⁸ C. A. Carrillo Montoya,¹⁸ L. F. Chaparro Sierra,¹⁸ C. Florez,¹⁸ C. F. González Hernández,¹⁸ M. A. Segura Delgado,¹⁸ B. Courbon,¹⁹ N. Godinovic,¹⁹ D. Lelas,¹⁹ I. Puljak,¹⁹ T. Sculac,¹⁹ Z. Antunovic,²⁰ M. Kovac,²⁰ V. Brigljevic,²¹ D. Ferencek,²¹ K. Kadija,²¹ B. Mesic,²¹ A. Starodumov,^{21,g} T. Susa,²¹ M. W. Ather,²² A. Attikis,²² M. Kolosova,²² G. Mavromanolakis,²² J. Mousa,²² C. Nicolaou,²² F. Ptochos,²² P. A. Razis,²² H. Rykaczewski,²² M. Finger,^{23,h} M. Finger Jr.,^{23,h} E. Ayala,²⁴ E. Carrera Jarrin,²⁵ Y. Assran,^{26,ij} S. Elgammal,^{26,i} S. Khalil,^{26,k} S. Bhowmik,²⁷ A. Carvalho Antunes De Oliveira,²⁷ R. K. Dewanjee,²⁷ K. Ehataht,²⁷ M. Kadastik,²⁷ M. Raidal,²⁷ C. Veelken,²⁷ P. Eerola,²⁸ H. Kirschenmann,²⁸ J. Pekkanen,²⁸ M. Voutilainen,²⁸ J. Havukainen,²⁹ J. K. Heikkilä,²⁹ T. Järvinen,²⁹ V. Karimäki,²⁹ R. Kinnunen,²⁹ T. Lampén,²⁹ K. Lassila-Perini,²⁹ S. Laurila,²⁹ S. Lehti,²⁹ T. Lindén,²⁹ P. Luukka,²⁹ T. Mäenpää,²⁹ H. Siikonen,²⁹ E. Tuominen,²⁹ J. Tuominiemi,²⁹ T. Tuuva,³⁰ M. Besancon,³¹ F. Couderc,³¹ M. DeJardin,³¹ D. Denegri,³¹ J. L. Faure,³¹ F. Ferri,³¹ S. Ganjour,³¹ A. Givernaud,³¹ P. Gras,³¹ G. Hamel de Monchenault,³¹ P. Jarry,³¹ C. Leloup,³¹ E. Locci,³¹ J. Malcles,³¹ G. Negro,³¹ J. Rander,³¹ A. Rosowsky,³¹ M. Ö. Sahin,³¹ M. Titov,³¹ A. Abdulsalam,^{32,l} C. Amendola,³² I. Antropov,³² F. Beaudette,³² P. Busson,³² C. Charlot,³² R. Granier de Cassagnac,³² I. Kucher,³² A. Lobanov,³² J. Martin Blanco,³² C. Martin Perez,³² M. Nguyen,³² C. Ochando,³² G. Ortona,³² P. Paganini,³² P. Pigard,³² J. Rembser,³² R. Salerno,³² J. B. Sauvan,³² Y. Sirois,³² A. G. Stahl Leiton,³² A. Zabi,³² A. Zghiche,³² J.-L. Agram,^{33,m} J. Andrea,³³ D. Bloch,³³ J.-M. Brom,³³ E. C. Chabert,³³ V. Cherepanov,³³ C. Collard,³³ E. Conte,^{33,m} J.-C. Fontaine,^{33,m} D. Gelé,³³ U. Goerlach,³³ M. Jansová,³³ A.-C. Le Bihan,³³ N. Taroni,³³ P. Van Hove,³³ S. Gadrat,³⁴ S. Beauceron,³⁵ C. Bernet,³⁵ G. Boudoul,³⁵ N. Chanon,³⁵ R. Chierici,³⁵ D. Contardo,³⁵ P. Depasse,³⁵ H. El Mamouni,³⁵ J. Fay,³⁵ L. Finco,³⁵ S. Gascon,³⁵ M. Gouzevitch,³⁵ G. Grenier,³⁵ B. Ille,³⁵ F. Lagarde,³⁵ I. B. Laktineh,³⁵ H. Lattaud,³⁵ M. Lethuillier,³⁵ L. Mirabito,³⁵ S. Perries,³⁵ A. Popov,^{35,n} V. Sordini,³⁵ G. Touquet,³⁵ M. Vander Donckt,³⁵ S. Viret,³⁵ T. Toriashvili,^{36,o} Z. Tsamalaidze,^{37,h} C. Autermann,³⁸ L. Feld,³⁸ M. K. Kiesel,³⁸ K. Klein,³⁸ M. Lipinski,³⁸ M. Preuten,³⁸ M. P. Rauch,³⁸ C. Schomakers,³⁸ J. Schulz,³⁸ M. Teroerde,³⁸ B. Wittmer,³⁸ A. Albert,³⁹ D. Duchardt,³⁹ M. Erdmann,³⁹ S. Erdweg,³⁹ T. Esch,³⁹ R. Fischer,³⁹ S. Ghosh,³⁹ A. Güth,³⁹ T. Hebbeker,³⁹ C. Heidemann,³⁹ K. Hoepfner,³⁹ H. Keller,³⁹ L. Mastrolorenzo,³⁹ M. Merschmeyer,³⁹ A. Meyer,³⁹ P. Millet,³⁹ S. Mukherjee,³⁹ T. Pook,³⁹ M. Radziej,³⁹ H. Reithler,³⁹ M. Rieger,³⁹ A. Schmidt,³⁹ D. Teyssier,³⁹ S. Thüer,³⁹ G. Flügge,⁴⁰ O. Hlushchenko,⁴⁰ T. Kress,⁴⁰ T. Müller,⁴⁰ A. Nehr Korn,⁴⁰ A. Nowack,⁴⁰ C. Pistone,⁴⁰ O. Pooth,⁴⁰ D. Roy,⁴⁰ H. Sert,⁴⁰ A. Stahl,^{40,p} M. Aldaya Martin,⁴¹ T. Arndt,⁴¹ C. Asawatrangkuldee,⁴¹ I. Babounikau,⁴¹ K. Beernaert,⁴¹ O. Behnke,⁴¹ U. Behrens,⁴¹ A. Bermúdez Martínez,⁴¹ D. Bertsche,⁴¹ A. A. Bin Anuar,⁴¹ K. Borras,^{41,q} V. Botta,⁴¹ A. Campbell,⁴¹ P. Connor,⁴¹ C. Contreras-Campana,⁴¹ V. Danilov,⁴¹ A. De Wit,⁴¹ M. M. Defranchis,⁴¹ C. Diez Pardos,⁴¹ D. Domínguez Damiani,⁴¹ G. Eckerlin,⁴¹ T. Eichhorn,⁴¹ A. Elwood,⁴¹ E. Eren,⁴¹ E. Gallo,^{41,r} A. Geiser,⁴¹ J. M. Grados Luyando,⁴¹ A. Grohsjean,⁴¹ M. Guthoff,⁴¹ M. Haranko,⁴¹ A. Harb,⁴¹ J. Hauk,⁴¹ H. Jung,⁴¹ M. Kasemann,⁴¹ J. Keaveney,⁴¹ C. Kleinwort,⁴¹ J. Knolle,⁴¹ D. Krücker,⁴¹ W. Lange,⁴¹ A. Lelek,⁴¹ T. Lenz,⁴¹ J. Leonard,⁴¹ K. Lipka,⁴¹ W. Lohmann,^{41,s} R. Mankel,⁴¹ I.-A. Melzer-Pellmann,⁴¹ A. B. Meyer,⁴¹ M. Meyer,⁴¹ M. Missiroli,⁴¹ G. Mittag,⁴¹ J. Mnich,⁴¹ V. Myronenko,⁴¹ S. K. Pflitsch,⁴¹ D. Pitzl,⁴¹ A. Raspereza,⁴¹ M. Savitskiy,⁴¹ P. Saxena,⁴¹ P. Schütze,⁴¹ C. Schwanenberger,⁴¹ R. Shevchenko,⁴¹ A. Singh,⁴¹ H. Tholen,⁴¹ O. Turkot,⁴¹ A. Vagnerini,⁴¹ G. P. Van Onsem,⁴¹ R. Walsh,⁴¹ Y. Wen,⁴¹ K. Wichmann,⁴¹ C. Wissing,⁴¹ O. Zenaiev,⁴¹ R. Aggleton,⁴² S. Bein,⁴² L. Benato,⁴² A. Benecke,⁴² V. Blobel,⁴² T. Dreyer,⁴² A. Ebrahimi,⁴² E. Garutti,⁴² D. Gonzalez,⁴² P. Gunnellini,⁴² J. Haller,⁴² A. Hinzmann,⁴² A. Karavdina,⁴² G. Kasieczka,⁴² R. Klanner,⁴² R. Kogler,⁴² N. Kovalchuk,⁴² S. Kurz,⁴² V. Kutzner,⁴² J. Lange,⁴² D. Marconi,⁴² J. Multhaup,⁴² M. Niedziela,⁴² C. E. N. Niemeyer,⁴² D. Nowatschin,⁴² A. Perieanu,⁴² A. Reimers,⁴² O. Rieger,⁴² C. Scharf,⁴² P. Schleper,⁴² S. Schumann,⁴² J. Schwandt,⁴² J. Sonneveld,⁴² H. Stadie,⁴² G. Steinbrück,⁴² F. M. Stober,⁴² M. Stöver,⁴² A. Vanhoefer,⁴² B. Vormwald,⁴² I. Zoi,⁴² M. Akbiyik,⁴³ C. Barth,⁴³ M. Baselga,⁴³ S. Baur,⁴³ E. Butz,⁴³ R. Caspart,⁴³ T. Chwalek,⁴³ F. Colombo,⁴³ W. De Boer,⁴³ A. Dierlamm,⁴³ K. El Morabit,⁴³ N. Faltermann,⁴³ B. Freund,⁴³ M. Giffels,⁴³ M. A. Harrendorf,⁴³ F. Hartmann,^{43,p} S. M. Heindl,⁴³ U. Husemann,⁴³ I. Katkov,^{43,n} S. Kudella,⁴³ S. Mitra,⁴³ M. U. Mozer,⁴³ Th. Müller,⁴³ M. Musich,⁴³ M. Plagge,⁴³ G. Quast,⁴³ K. Rabbertz,⁴³ M. Schröder,⁴³ I. Shvetsov,⁴³ H. J. Simonis,⁴³ R. Ulrich,⁴³ S. Wayand,⁴³ M. Weber,⁴³ T. Weiler,⁴³ C. Wöhrmann,⁴³ R. Wolf,⁴³ G. Anagnostou,⁴⁴ G. Daskalakis,⁴⁴ T. Geralis,⁴⁴ A. Kyriakis,⁴⁴ D. Loukas,⁴⁴ G. Paspalaki,⁴⁴ G. Karathanasis,⁴⁵ P. Kontaxakis,⁴⁵ A. Panagiotou,⁴⁵ I. Papavergou,⁴⁵ N. Saoulidou,⁴⁵ E. Tziaferi,⁴⁵ K. Vellidis,⁴⁵ K. Kousouris,⁴⁶ I. Papakrivopoulos,⁴⁶ G. Tsipolitis,⁴⁶ I. Evangelou,⁴⁷ C. Foudas,⁴⁷ P. Gianneios,⁴⁷ P. Katsoulis,⁴⁷ P. Kokkas,⁴⁷ S. Mallios,⁴⁷ N. Manthos,⁴⁷ I. Papadopoulos,⁴⁷ E. Paradas,⁴⁷ J. Strologas,⁴⁷ F. A. Triantis,⁴⁷ D. Tsitsonis,⁴⁷ M. Bartók,^{48,t} M. Csanad,⁴⁸ N. Filipovic,⁴⁸ P. Major,⁴⁸ M. I. Nagy,⁴⁸ G. Pasztor,⁴⁸ O. Surányi,⁴⁸ G. I. Veres,⁴⁸ G. Bencze,⁴⁹

C. Hajdu,⁴⁹ D. Horvath,^{49,u} Á. Hunyadi,⁴⁹ F. Sikler,⁴⁹ T. Á. Vámi,⁴⁹ V. Veszpremi,⁴⁹ G. Vesztergombi,^{49,†} v N. Beni,⁵⁰
 S. Czellar,⁵⁰ J. Karancsi,^{50,t} A. Makovec,⁵⁰ J. Molnar,⁵⁰ Z. Szillasi,⁵⁰ P. Raics,⁵¹ Z. L. Trocsanyi,⁵¹ B. Ujvari,⁵¹
 S. Choudhury,⁵² J. R. Komaragiri,⁵² P. C. Tiwari,⁵² S. Bahinipati,^{53,w} C. Kar,⁵³ P. Mal,⁵³ K. Mandal,⁵³ A. Nayak,^{53,x}
 D. K. Sahoo,^{53,w} S. K. Swain,⁵³ S. Bansal,⁵⁴ S. B. Beri,⁵⁴ V. Bhatnagar,⁵⁴ S. Chauhan,⁵⁴ R. Chawla,⁵⁴ N. Dhingra,⁵⁴
 R. Gupta,⁵⁴ A. Kaur,⁵⁴ M. Kaur,⁵⁴ S. Kaur,⁵⁴ P. Kumari,⁵⁴ M. Lohan,⁵⁴ A. Mehta,⁵⁴ K. Sandeep,⁵⁴ S. Sharma,⁵⁴ J. B. Singh,⁵⁴
 A. K. Virdi,⁵⁴ G. Walia,⁵⁴ A. Bhardwaj,⁵⁵ B. C. Choudhary,⁵⁵ R. B. Garg,⁵⁵ M. Gola,⁵⁵ S. Keshri,⁵⁵ Ashok Kumar,⁵⁵
 S. Malhotra,⁵⁵ M. Naimuddin,⁵⁵ P. Priyanka,⁵⁵ K. Ranjan,⁵⁵ Aashaq Shah,⁵⁵ R. Sharma,⁵⁵ R. Bhardwaj,^{56,y} M. Bharti,^{56,y}
 R. Bhattacharya,⁵⁶ S. Bhattacharya,⁵⁶ U. Bhawandeep,^{56,y} D. Bhowmik,⁵⁶ S. Dey,⁵⁶ S. Dutt,^{56,y} S. Dutta,⁵⁶ S. Ghosh,⁵⁶
 K. Mondal,⁵⁶ S. Nandan,⁵⁶ A. Purohit,⁵⁶ P. K. Rout,⁵⁶ A. Roy,⁵⁶ S. Roy Chowdhury,⁵⁶ G. Saha,⁵⁶ S. Sarkar,⁵⁶ M. Sharan,⁵⁶
 B. Singh,^{56,y} S. Thakur,^{56,y} P. K. Behera,⁵⁷ R. Chudasama,⁵⁸ D. Dutta,⁵⁸ V. Jha,⁵⁸ V. Kumar,⁵⁸ P. K. Netrakanti,⁵⁸
 L. M. Pant,⁵⁸ P. Shukla,⁵⁸ T. Aziz,⁵⁹ M. A. Bhat,⁵⁹ S. Dugad,⁵⁹ G. B. Mohanty,⁵⁹ N. Sur,⁵⁹ B. Sutar,⁵⁹
 Ravindra Kumar Verma,⁵⁹ S. Banerjee,⁶⁰ S. Bhattacharya,⁶⁰ S. Chatterjee,⁶⁰ P. Das,⁶⁰ M. Guchait,⁶⁰ Sa. Jain,⁶⁰
 S. Karmakar,⁶⁰ S. Kumar,⁶⁰ M. Maity,^{60,z} G. Majumder,⁶⁰ K. Mazumdar,⁶⁰ N. Sahoo,⁶⁰ T. Sarkar,^{60,z} S. Chauhan,⁶¹
 S. Dube,⁶¹ V. Hegde,⁶¹ A. Kapoor,⁶¹ K. Kotheekar,⁶¹ S. Pandey,⁶¹ A. Rane,⁶¹ A. Rastogi,⁶¹ S. Sharma,⁶¹ S. Chenarani,^{62,aa}
 E. Eskandari Tadavani,⁶² S. M. Etesami,^{62,aa} M. Khakzad,⁶² M. Mohammadi Najafabadi,⁶² M. Naseri,⁶²
 F. Rezaei Hosseinabadi,⁶² B. Safarzadeh,^{62,bb} M. Zeinali,⁶² M. Felcini,⁶³ M. Grunewald,⁶³ M. Abbrescia,^{64a,64b}
 C. Calabria,^{64a,64b} A. Colaleo,^{64a} D. Creanza,^{64a,64c} L. Cristella,^{64a,64b} N. De Filippis,^{64a,64c} M. De Palma,^{64a,64b}
 A. Di Florio,^{64a,64b} F. Errico,^{64a,64b} L. Fiore,^{64a} A. Gelmi,^{64a,64b} G. Iaselli,^{64a,64c} M. Ince,^{64a,64b} S. Lezki,^{64a,64b} G. Maggi,^{64a,64c}
 M. Maggi,^{64a} G. Miniello,^{64a,64b} S. My,^{64a,64b} S. Nuzzo,^{64a,64b} A. Pompili,^{64a,64b} G. Pugliese,^{64a,64c} R. Radogna,^{64a}
 A. Ranieri,^{64a} G. Selvaggi,^{64a,64b} A. Sharma,^{64a} L. Silvestris,^{64a} R. Venditti,^{64a} P. Verwilligen,^{64a} G. Zito,^{64a} G. Abbiendi,^{65a}
 C. Battilana,^{65a,65b} D. Bonacorsi,^{65a,65b} L. Borgonovi,^{65a,65b} S. Braibant-Giacomelli,^{65a,65b} R. Campanini,^{65a,65b}
 P. Capiluppi,^{65a,65b} A. Castro,^{65a,65b} F. R. Cavallo,^{65a} S. S. Chhibra,^{65a,65b} C. Ciocca,^{65a} G. Codispoti,^{65a,65b} M. Cuffiani,^{65a,65b}
 G. M. Dallavalle,^{65a} F. Fabbri,^{65a} A. Fanfani,^{65a,65b} E. Fontanesi,^{65a} P. Giacomelli,^{65a} C. Grandi,^{65a} L. Guiducci,^{65a,65b}
 S. Lo Meo,^{65a} S. Marcellini,^{65a} G. Masetti,^{65a} A. Montanari,^{65a} F. L. Navarria,^{65a,65b} A. Perrotta,^{65a} F. Primavera,^{65a,65b,p}
 A. M. Rossi,^{65a,65b} T. Rovelli,^{65a,65b} G. P. Siroli,^{65a,65b} N. Tosi,^{65a} S. Albergo,^{66a,66b} A. Di Mattia,^{66a} R. Potenza,^{66a,66b}
 A. Tricomi,^{66a,66b} C. Tuve,^{66a,66b} G. Barbagli,^{67a} K. Chatterjee,^{67a,67b} V. Ciulli,^{67a,67b} C. Civinini,^{67a} R. D'Alessandro,^{67a,67b}
 E. Focardi,^{67a,67b} G. Latino,^{67a} P. Lenzi,^{67a,67b} M. Meschini,^{67a} S. Paoletti,^{67a} L. Russo,^{67a,cc} G. Sguazzoni,^{67a} D. Strom,^{67a}
 L. Viliani,^{67a} L. Benussi,⁶⁸ S. Bianco,⁶⁸ F. Fabbri,⁶⁸ D. Piccolo,⁶⁸ F. Ferro,^{69a} R. Mulargia,^{69a,69b} F. Ravera,^{69a,69b}
 E. Robutti,^{69a} S. Tosi,^{69a,69b} A. Benaglia,^{70a} A. Beschi,^{70a,70b} F. Brivio,^{70a,70b} V. Ciriolo,^{70a,70b,p} S. Di Guida,^{70a,70b,p}
 M. E. Dinardo,^{70a,70b} S. Fiorendi,^{70a,70b} S. Gennai,^{70a} A. Ghezzi,^{70a,70b} P. Govoni,^{70a,70b} M. Malberti,^{70a,70b} S. Malvezzi,^{70a}
 A. Massironi,^{70a,70b} D. Menasce,^{70a} F. Monti,^{70a} L. Moroni,^{70a} M. Paganoni,^{70a,70b} D. Pedrini,^{70a} S. Ragazzi,^{70a,70b}
 T. Tabarelli de Fatis,^{70a,70b} D. Zuolo,^{70a,70b} S. Buontempo,^{71a} N. Cavallo,^{71a,71c} A. De Iorio,^{71a,71b} A. Di Crescenzo,^{71a,71b}
 F. Fabozzi,^{71a,71c} F. Fienga,^{71a} G. Galati,^{71a} A. O. M. Iorio,^{71a,71b} W. A. Khan,^{71a} L. Lista,^{71a} S. Meola,^{71a,71d,p} P. Paolucci,^{71a,p}
 C. Sciacca,^{71a,71b} E. Voevodina,^{71a,71b} P. Azzi,^{72a} N. Bacchetta,^{72a} D. Bisello,^{72a,72b} A. Boletti,^{72a,72b} A. Bragagnolo,^{72a}
 R. Carlin,^{72a,72b} P. Checchia,^{72a} M. Dall'Osso,^{72a,72b} P. De Castro Manzano,^{72a} T. Dorigo,^{72a} U. Dosselli,^{72a}
 F. Gasparini,^{72a,72b} U. Gasparini,^{72a,72b} A. Gozzelino,^{72a} S. Y. Hoh,^{72a} S. Lacaprara,^{72a} P. Lujan,^{72a} M. Margoni,^{72a,72b}
 A. T. Meneguzzo,^{72a,72b} J. Pazzini,^{72a,72b} P. Ronchese,^{72a,72b} R. Rossin,^{72a,72b} F. Simonetto,^{72a,72b} A. Tiko,^{72a} E. Torassa,^{72a}
 M. Tosi,^{72a,72b} M. Zanetti,^{72a,72b} P. Zotto,^{72a,72b} G. Zumerle,^{72a,72b} A. Braghieri,^{73a} A. Magnani,^{73a} P. Montagna,^{73a,73b}
 S. P. Ratti,^{73a,73b} V. Re,^{73a} M. Ressegotti,^{73a,73b} C. Riccardi,^{73a,73b} P. Salvini,^{73a} I. Vai,^{73a,73b} P. Vitulo,^{73a,73b} M. Biasini,^{74a,74b}
 G. M. Bilei,^{74a} C. Cecchi,^{74a,74b} D. Ciangottini,^{74a,74b} L. Fanò,^{74a,74b} P. Lariccia,^{74a,74b} R. Leonardi,^{74a,74b} E. Manoni,^{74a}
 G. Mantovani,^{74a,74b} V. Mariani,^{74a,74b} M. Menichelli,^{74a} A. Rossi,^{74a,74b} A. Santocchia,^{74a,74b} D. Spiga,^{74a} K. Androsov,^{75a}
 P. Azzurri,^{75a} G. Bagliesi,^{75a} L. Bianchini,^{75a} T. Boccali,^{75a} L. Borrello,^{75a} R. Castaldi,^{75a} M. A. Ciocci,^{75a,75b} R. Dell'Orso,^{75a}
 G. Fedi,^{75a} F. Fiori,^{75a,75c} L. Giannini,^{75a,75c} A. Giassi,^{75a} M. T. Grippo,^{75a} F. Ligabue,^{75a,75c} E. Manca,^{75a,75c}
 G. Mandorli,^{75a,75c} A. Messineo,^{75a,75b} F. Palla,^{75a} A. Rizzi,^{75a,75b} G. Rolandi,^{75a,dd} P. Spagnolo,^{75a} R. Tenchini,^{75a}
 G. Tonelli,^{75a,75b} A. Venturi,^{75a} P. G. Verdini,^{75a} L. Barone,^{76a,76b} F. Cavallari,^{76a} M. Cipriani,^{76a,76b} D. Del Re,^{76a,76b}
 E. Di Marco,^{76a,76b} M. Diemoz,^{76a} S. Gelli,^{76a,76b} E. Longo,^{76a,76b} B. Marzocchi,^{76a,76b} P. Meridiani,^{76a} G. Organtini,^{76a,76b}
 F. Pandolfi,^{76a} R. Paramatti,^{76a,76b} F. Preiato,^{76a,76b} S. Rahatlou,^{76a,76b} C. Rovelli,^{76a} F. Santanastasio,^{76a,76b} N. Amapane,^{77a,77b}
 R. Arcidiacono,^{77a,77c} S. Argiro,^{77a,77b} M. Arneodo,^{77a,77c} N. Bartosik,^{77a} R. Bellan,^{77a,77b} C. Biino,^{77a} A. Cappati,^{77a,77b}
 N. Cartiglia,^{77a} F. Cenna,^{77a,77b} S. Cometti,^{77a} M. Costa,^{77a,77b} R. Covarelli,^{77a,77b} N. Demaria,^{77a} B. Kiani,^{77a,77b}

C. Mariotti,^{77a} S. Maselli,^{77a} E. Migliore,^{77a,77b} V. Monaco,^{77a,77b} E. Monteil,^{77a,77b} M. Monteno,^{77a} M. M. Obertino,^{77a,77b} L. Pacher,^{77a,77b} N. Pastrone,^{77a} M. Pelliccioni,^{77a} G. L. Pinna Angioni,^{77a,77b} A. Romero,^{77a,77b} M. Ruspa,^{77a,77c} R. Sacchi,^{77a,77b} R. Salvatico,^{77a,77b} K. Shchelina,^{77a,77b} V. Sola,^{77a} A. Solano,^{77a,77b} D. Soldi,^{77a,77b} A. Staiano,^{77a} S. Belforte,^{78a} V. Candelise,^{78a,78b} M. Casarsa,^{78a} F. Cossutti,^{78a} A. Da Rold,^{78a,78b} G. Della Ricca,^{78a,78b} F. Vazzoler,^{78a,78b} A. Zanetti,^{78a} D. H. Kim,⁷⁹ G. N. Kim,⁷⁹ M. S. Kim,⁷⁹ J. Lee,⁷⁹ S. Lee,⁷⁹ S. W. Lee,⁷⁹ C. S. Moon,⁷⁹ Y. D. Oh,⁷⁹ S. I. Pak,⁷⁹ S. Sekmen,⁷⁹ D. C. Son,⁷⁹ Y. C. Yang,⁷⁹ H. Kim,⁸⁰ D. H. Moon,⁸⁰ G. Oh,⁸⁰ B. Francois,⁸¹ J. Goh,^{81,ee} T. J. Kim,⁸¹ S. Cho,⁸² S. Choi,⁸² Y. Go,⁸² D. Gyun,⁸² S. Ha,⁸² B. Hong,⁸² Y. Jo,⁸² K. Lee,⁸² K. S. Lee,⁸² S. Lee,⁸² J. Lim,⁸² S. K. Park,⁸² Y. Roh,⁸² H. S. Kim,⁸³ J. Almond,⁸⁴ J. Kim,⁸⁴ J. S. Kim,⁸⁴ H. Lee,⁸⁴ K. Lee,⁸⁴ K. Nam,⁸⁴ S. B. Oh,⁸⁴ B. C. Radburn-Smith,⁸⁴ S. h. Seo,⁸⁴ U. K. Yang,⁸⁴ H. D. Yoo,⁸⁴ G. B. Yu,⁸⁴ D. Jeon,⁸⁵ H. Kim,⁸⁵ J. H. Kim,⁸⁵ J. S. H. Lee,⁸⁵ I. C. Park,⁸⁵ Y. Choi,⁸⁶ C. Hwang,⁸⁶ J. Lee,⁸⁶ I. Yu,⁸⁶ V. Dudenias,⁸⁷ A. Juodagalvis,⁸⁷ J. Vaitkus,⁸⁷ I. Ahmed,⁸⁸ Z. A. Ibrahim,⁸⁸ M. A. B. Md Ali,^{88,ff} F. Mohamad Idris,^{88,gg} W. A. T. Wan Abdullah,⁸⁸ M. N. Yusli,⁸⁸ Z. Zolkapli,⁸⁸ J. F. Benitez,⁸⁹ A. Castaneda Hernandez,⁸⁹ J. A. Murillo Quijada,⁸⁹ H. Castilla-Valdez,⁹⁰ E. De La Cruz-Burelo,⁹⁰ M. C. Duran-Osuna,⁹⁰ I. Heredia-De La Cruz,^{90,hh} R. Lopez-Fernandez,⁹⁰ J. Mejia Guisao,⁹⁰ R. I. Rabadan-Trejo,⁹⁰ M. Ramirez-Garcia,⁹⁰ G. Ramirez-Sanchez,⁹⁰ R. Reyes-Almanza,⁹⁰ A. Sanchez-Hernandez,⁹⁰ S. Carrillo Moreno,⁹¹ C. Oropeza Barrera,⁹¹ F. Vazquez Valencia,⁹¹ J. Eysermans,⁹² I. Pedraza,⁹² H. A. Salazar Ibarquen,⁹² C. Uribe Estrada,⁹² A. Morelos Pineda,⁹³ D. Krofcheck,⁹⁴ S. Bheesette,⁹⁵ P. H. Butler,⁹⁵ A. Ahmad,⁹⁶ M. Ahmad,⁹⁶ M. I. Asghar,⁹⁶ Q. Hassan,⁹⁶ H. R. Hoorani,⁹⁶ A. Saddique,⁹⁶ M. A. Shah,⁹⁶ M. Shoaib,⁹⁶ M. Waqas,⁹⁶ H. Bialkowska,⁹⁷ M. Bluj,⁹⁷ B. Boimska,⁹⁷ T. Frueboes,⁹⁷ M. Górski,⁹⁷ M. Kazana,⁹⁷ M. Szleper,⁹⁷ P. Traczyk,⁹⁷ P. Zalewski,⁹⁷ K. Bunkowski,⁹⁸ A. Byszuk,^{98,ii} K. Doroba,⁹⁸ A. Kalinowski,⁹⁸ M. Konecki,⁹⁸ J. Krolikowski,⁹⁸ M. Misiura,⁹⁸ M. Olszewski,⁹⁸ A. Pyskir,⁹⁸ M. Walczak,⁹⁸ M. Araujo,⁹⁹ P. Bargassa,⁹⁹ C. Beirão Da Cruz E Silva,⁹⁹ A. Di Francesco,⁹⁹ P. Faccioli,⁹⁹ B. Galinhas,⁹⁹ M. Gallinaro,⁹⁹ J. Hollar,⁹⁹ N. Leonardo,⁹⁹ J. Seixas,⁹⁹ G. Strong,⁹⁹ O. Toldaiev,⁹⁹ J. Varela,⁹⁹ S. Afanasiev,¹⁰⁰ P. Bunin,¹⁰⁰ M. Gavrilenko,¹⁰⁰ I. Golutvin,¹⁰⁰ I. Gorbunov,¹⁰⁰ A. Kamenev,¹⁰⁰ V. Karjavine,¹⁰⁰ A. Lanev,¹⁰⁰ A. Malakhov,¹⁰⁰ V. Matveev,^{100,jj,kk} P. Moisenz,¹⁰⁰ V. Palichik,¹⁰⁰ V. Perelygin,¹⁰⁰ S. Shmatov,¹⁰⁰ S. Shulha,¹⁰⁰ N. Skatchkov,¹⁰⁰ V. Smirnov,¹⁰⁰ N. Voytishin,¹⁰⁰ A. Zarubin,¹⁰⁰ V. Golovtsov,¹⁰¹ Y. Ivanov,¹⁰¹ V. Kim,^{101,ll} E. Kuznetsova,^{101,mm} P. Levchenko,¹⁰¹ V. Murzin,¹⁰¹ V. Oreshkin,¹⁰¹ I. Smirnov,¹⁰¹ D. Sosnov,¹⁰¹ V. Sulimov,¹⁰¹ L. Uvarov,¹⁰¹ S. Vavilov,¹⁰¹ A. Vorobyev,¹⁰¹ Yu. Andreev,¹⁰² A. Dermenev,¹⁰² S. Gninenko,¹⁰² N. Golubev,¹⁰² A. Karneyeu,¹⁰² M. Kirsanov,¹⁰² N. Krasnikov,¹⁰² A. Pashenkov,¹⁰² D. Tlisov,¹⁰² A. Toropin,¹⁰² V. Epshteyn,¹⁰³ V. Gavrilov,¹⁰³ N. Lychkovskaya,¹⁰³ V. Popov,¹⁰³ I. Pozdnyakov,¹⁰³ G. Safronov,¹⁰³ A. Spiridonov,¹⁰³ A. Stepenov,¹⁰³ V. Stolin,¹⁰³ M. Toms,¹⁰³ E. Vlasov,¹⁰³ A. Zhokin,¹⁰³ T. Aushev,¹⁰⁴ R. Chistov,^{105,nn} M. Danilov,^{105,nn} P. Parygin,¹⁰⁵ D. Philippov,¹⁰⁵ S. Polikarpov,^{105,nn} E. Tarkovskii,¹⁰⁵ V. Andreev,¹⁰⁶ M. Azarkin,¹⁰⁶ I. Dremin,^{106,kk} M. Kirakosyan,¹⁰⁶ A. Terkulov,¹⁰⁶ A. Baskakov,¹⁰⁷ A. Belyaev,¹⁰⁷ E. Boos,¹⁰⁷ M. Dubinin,^{107,oo} L. Dudko,¹⁰⁷ A. Ershov,¹⁰⁷ A. Gribushin,¹⁰⁷ V. Klyukhin,¹⁰⁷ O. Kodolova,¹⁰⁷ I. Lokhtin,¹⁰⁷ I. Miagkov,¹⁰⁷ S. Obraztsov,¹⁰⁷ S. Petrushanko,¹⁰⁷ V. Savrin,¹⁰⁷ A. Snigirev,¹⁰⁷ A. Barnyakov,^{108,pp} V. Blinov,^{108,pp} T. Dimova,^{108,pp} L. Kardapoltsev,^{108,pp} Y. Skovpen,^{108,pp} I. Azhgirey,¹⁰⁹ I. Bayshev,¹⁰⁹ S. Bitiukov,¹⁰⁹ D. Elumakhov,¹⁰⁹ A. Godizov,¹⁰⁹ V. Kachanov,¹⁰⁹ A. Kalinin,¹⁰⁹ D. Konstantinov,¹⁰⁹ P. Mandrik,¹⁰⁹ V. Petrov,¹⁰⁹ R. Ryutin,¹⁰⁹ S. Slabospitskii,¹⁰⁹ A. Sobol,¹⁰⁹ S. Troshin,¹⁰⁹ N. Tyurin,¹⁰⁹ A. Uzunian,¹⁰⁹ A. Volkov,¹⁰⁹ A. Babaev,¹¹⁰ S. Baidali,¹¹⁰ V. Okhotnikov,¹¹⁰ P. Adzic,^{111,qq} P. Cirkovic,¹¹¹ D. Devetak,¹¹¹ M. Dordevic,¹¹¹ J. Milosevic,¹¹¹ J. Alcaraz Maestre,¹¹² A. Álvarez Fernández,¹¹² I. Bachiller,¹¹² M. Barrio Luna,¹¹² J. A. Brochero Cifuentes,¹¹² M. Cerrada,¹¹² N. Colino,¹¹² B. De La Cruz,¹¹² A. Delgado Peris,¹¹² C. Fernandez Bedoya,¹¹² J. P. Fernández Ramos,¹¹² J. Flix,¹¹² M. C. Fouz,¹¹² O. Gonzalez Lopez,¹¹² S. Goy Lopez,¹¹² J. M. Hernandez,¹¹² M. I. Josa,¹¹² D. Moran,¹¹² A. Pérez-Calero Yzquierdo,¹¹² J. Puerta Pelayo,¹¹² I. Redondo,¹¹² L. Romero,¹¹² M. S. Soares,¹¹² A. Triossi,¹¹² C. Albajar,¹¹³ J. F. de Trocóniz,¹¹³ J. Cuevas,¹¹⁴ C. Erice,¹¹⁴ J. Fernandez Menendez,¹¹⁴ S. Folgueras,¹¹⁴ I. Gonzalez Caballero,¹¹⁴ J. R. González Fernández,¹¹⁴ E. Palencia Cortezon,¹¹⁴ V. Rodríguez Bouza,¹¹⁴ S. Sanchez Cruz,¹¹⁴ P. Vischia,¹¹⁴ J. M. Vizán García,¹¹⁴ I. J. Cabrillo,¹¹⁵ A. Calderon,¹¹⁵ B. Chazin Quero,¹¹⁵ J. Duarte Campderros,¹¹⁵ M. Fernandez,¹¹⁵ P. J. Fernández Manteca,¹¹⁵ A. García Alonso,¹¹⁵ J. Garcia-Ferrero,¹¹⁵ G. Gomez,¹¹⁵ A. Lopez Virto,¹¹⁵ J. Marco,¹¹⁵ C. Martinez Rivero,¹¹⁵ P. Martinez Ruiz del Arbol,¹¹⁵ F. Matorras,¹¹⁵ J. Piedra Gomez,¹¹⁵ C. Prieels,¹¹⁵ T. Rodrigo,¹¹⁵ A. Ruiz-Jimeno,¹¹⁵ L. Scodellaro,¹¹⁵ N. Trevisani,¹¹⁵ I. Vila,¹¹⁵ R. Vilar Cortabitarte,¹¹⁵ N. Wickramage,¹¹⁶ D. Abbaneo,¹¹⁷ B. Akgun,¹¹⁷ E. Auffray,¹¹⁷ G. Auzinger,¹¹⁷ P. Baillon,¹¹⁷ A. H. Ball,¹¹⁷ D. Barney,¹¹⁷ J. Bendavid,¹¹⁷ M. Bianco,¹¹⁷ A. Bocci,¹¹⁷ C. Botta,¹¹⁷ E. Brondolin,¹¹⁷ T. Camporesi,¹¹⁷ M. Cepeda,¹¹⁷ G. Cerminara,¹¹⁷ E. Chapon,¹¹⁷ Y. Chen,¹¹⁷ G. Cucciati,¹¹⁷ D. d'Enterria,¹¹⁷ A. Dabrowski,¹¹⁷ N. Daci,¹¹⁷ V. Daponte,¹¹⁷ A. David,¹¹⁷ A. De Roeck,¹¹⁷ N. Deelen,¹¹⁷ M. Dobson,¹¹⁷ M. Dünser,¹¹⁷

N. Dupont,¹¹⁷ A. Elliott-Peisert,¹¹⁷ P. Everaerts,¹¹⁷ F. Fallavollita,^{117,ir} D. Fasanella,¹¹⁷ G. Franzoni,¹¹⁷ J. Fulcher,¹¹⁷ W. Funk,¹¹⁷ D. Gigi,¹¹⁷ A. Gilbert,¹¹⁷ K. Gill,¹¹⁷ F. Glege,¹¹⁷ M. Gruchala,¹¹⁷ M. Guilbaud,¹¹⁷ D. Gulhan,¹¹⁷ J. Hegeman,¹¹⁷ C. Heidegger,¹¹⁷ V. Innocente,¹¹⁷ A. Jafari,¹¹⁷ P. Janot,¹¹⁷ O. Karacheban,^{117,s} J. Kieseler,¹¹⁷ A. Kormmayer,¹¹⁷ M. Kramer,^{117,a} C. Lange,¹¹⁷ P. Lecocq,¹¹⁷ C. Lourenço,¹¹⁷ L. Malgeri,¹¹⁷ M. Mannelli,¹¹⁷ F. Meijers,¹¹⁷ J. A. Merlin,¹¹⁷ S. Mersi,¹¹⁷ E. Meschi,¹¹⁷ P. Milenovic,^{117,ss} F. Moortgat,¹¹⁷ M. Mulders,¹¹⁷ J. Ngadiuba,¹¹⁷ S. Nourbakhsh,¹¹⁷ S. Orfanelli,¹¹⁷ L. Orsini,¹¹⁷ F. Pantaleo,^{117,p} L. Pape,¹¹⁷ E. Perez,¹¹⁷ M. Peruzzi,¹¹⁷ A. Petrilli,¹¹⁷ G. Petruccianni,¹¹⁷ A. Pfeiffer,¹¹⁷ M. Pierini,¹¹⁷ F. M. Pitters,¹¹⁷ D. Rabady,¹¹⁷ A. Racz,¹¹⁷ T. Reis,¹¹⁷ M. Rovere,¹¹⁷ H. Sakulin,¹¹⁷ C. Schäfer,¹¹⁷ C. Schwick,¹¹⁷ M. Seidel,¹¹⁷ M. Selvaggi,¹¹⁷ A. Sharma,¹¹⁷ P. Silva,¹¹⁷ P. Sphicas,^{117,t} A. Stakia,¹¹⁷ J. Steggemann,¹¹⁷ D. Treille,¹¹⁷ A. Tsiros,¹¹⁷ V. Veckalns,^{117,uu} M. Verzetti,¹¹⁷ W. D. Zeuner,¹¹⁷ L. Caminada,^{118,uv} K. Deiters,¹¹⁸ W. Erdmann,¹¹⁸ R. Horisberger,¹¹⁸ Q. Ingram,¹¹⁸ H. C. Kaestli,¹¹⁸ D. Kotlinski,¹¹⁸ U. Langenegger,¹¹⁸ T. Rohe,¹¹⁸ S. A. Wiederkehr,¹¹⁸ M. Backhaus,¹¹⁹ L. Bäni,¹¹⁹ P. Berger,¹¹⁹ N. Chernyavskaya,¹¹⁹ G. Dissertori,¹¹⁹ M. Dittmar,¹¹⁹ M. Donegà,¹¹⁹ C. Dorfer,¹¹⁹ T. A. Gómez Espinosa,¹¹⁹ C. Grab,¹¹⁹ D. Hits,¹¹⁹ T. Klijnsma,¹¹⁹ W. Lustermann,¹¹⁹ R. A. Manzoni,¹¹⁹ M. Marionneau,¹¹⁹ M. T. Meinhard,¹¹⁹ F. Micheli,¹¹⁹ P. Musella,¹¹⁹ F. Nessi-Tedaldi,¹¹⁹ J. Pata,¹¹⁹ F. Pauss,¹¹⁹ G. Perrin,¹¹⁹ L. Perrozzini,¹¹⁹ S. Pigazzini,¹¹⁹ M. Quittnat,¹¹⁹ C. Reissel,¹¹⁹ D. Ruini,¹¹⁹ D. A. Sanz Becerra,¹¹⁹ M. Schönenberger,¹¹⁹ L. Shchutska,¹¹⁹ V. R. Tavolaro,¹¹⁹ K. Theofilatos,¹¹⁹ M. L. Vesterbacka Olsson,¹¹⁹ R. Wallny,¹¹⁹ D. H. Zhu,¹¹⁹ T. K. Aarrestad,¹²⁰ C. Amsler,^{120,ww} D. Brzhechko,¹²⁰ M. F. Canelli,¹²⁰ A. De Cosa,¹²⁰ R. Del Burgo,¹²⁰ S. Donato,¹²⁰ C. Galloni,¹²⁰ T. Hreus,¹²⁰ B. Kilminster,¹²⁰ S. Leontsinis,¹²⁰ I. Neutelings,¹²⁰ G. Rauco,¹²⁰ P. Robmann,¹²⁰ D. Salerno,¹²⁰ K. Schweiger,¹²⁰ C. Seitz,¹²⁰ Y. Takahashi,¹²⁰ A. Zucchetta,¹²⁰ T. H. Doan,¹²¹ R. Khurana,¹²¹ C. M. Kuo,¹²¹ W. Lin,¹²¹ A. Pozdnyakov,¹²¹ S. S. Yu,¹²¹ P. Chang,¹²² Y. Chao,¹²² K. F. Chen,¹²² P. H. Chen,¹²² W.-S. Hou,¹²² Arun Kumar,¹²² Y. F. Liu,¹²² R.-S. Lu,¹²² E. Paganis,¹²² A. Psallidas,¹²² A. Steen,¹²² B. Asavapibhop,¹²³ N. Srimanobhas,¹²³ N. Suwonjandee,¹²³ M. N. Bakirci,^{124,xx} A. Bat,¹²⁴ F. Boran,¹²⁴ S. Damarseckin,¹²⁴ Z. S. Demiroglu,¹²⁴ F. Dolek,¹²⁴ C. Dozen,¹²⁴ E. Eskut,¹²⁴ S. Girgis,¹²⁴ G. Gokbulut,¹²⁴ Y. Guler,¹²⁴ E. Gurpinar,¹²⁴ I. Hos,^{124,yy} C. Isik,¹²⁴ E. E. Kangal,^{124,zz} O. Kara,¹²⁴ U. Kiminsu,¹²⁴ M. Oglakci,¹²⁴ G. Onengut,¹²⁴ K. Ozdemir,^{124,aaa} A. Polatoz,¹²⁴ D. Sunar Cerci,^{124,bbb} B. Tali,^{124,bbb} U. G. Tok,¹²⁴ H. Topakli,^{124,xx} S. Turkapar,¹²⁴ I. S. Zorbakir,¹²⁴ C. Zorbilmez,¹²⁴ B. Isildak,^{125,ccc} G. Karapinar,^{125,ddd} M. Yalvac,¹²⁵ M. Zeyrek,¹²⁵ I. O. Atakisi,¹²⁶ E. Gülmez,¹²⁶ M. Kaya,^{126,eee} O. Kaya,^{126,fff} S. Ozkorucuklu,^{126,ggg} S. Tekten,¹²⁶ E. A. Yetkin,^{126,hhh} M. N. Agaras,¹²⁷ A. Cakir,¹²⁷ K. Cankocak,¹²⁷ Y. Komurcu,¹²⁷ S. Sen,^{127,iii} B. Grynyov,¹²⁸ L. Levchuk,¹²⁹ F. Ball,¹³⁰ J. J. Brooke,¹³⁰ D. Burns,¹³⁰ E. Clement,¹³⁰ D. Cussans,¹³⁰ O. Davignon,¹³⁰ H. Flacher,¹³⁰ J. Goldstein,¹³⁰ G. P. Heath,¹³⁰ H. F. Heath,¹³⁰ L. Kreczko,¹³⁰ D. M. Newbold,^{130,iii} S. Paramesvaran,¹³⁰ B. Penning,¹³⁰ T. Sakuma,¹³⁰ D. Smith,¹³⁰ V. J. Smith,¹³⁰ J. Taylor,¹³⁰ A. Titterton,¹³⁰ K. W. Bell,¹³¹ A. Belyaev,^{131,kkk} C. Brew,¹³¹ R. M. Brown,¹³¹ D. Cieri,¹³¹ D. J. A. Cockerill,¹³¹ J. A. Coughlan,¹³¹ K. Harder,¹³¹ S. Harper,¹³¹ J. Linacre,¹³¹ E. Olaiya,¹³¹ D. Petyt,¹³¹ C. H. Shepherd-Themistocleous,¹³¹ A. Thea,¹³¹ I. R. Tomalin,¹³¹ T. Williams,¹³¹ W. J. Womersley,¹³¹ R. Bainbridge,¹³² P. Bloch,¹³² J. Borg,¹³² S. Breeze,¹³² O. Buchmuller,¹³² A. Bundock,¹³² D. Colling,¹³² P. Dauncey,¹³² G. Davies,¹³² M. Della Negra,¹³² R. Di Maria,¹³² G. Hall,¹³² G. Iles,¹³² T. James,¹³² M. Komm,¹³² C. Laner,¹³² L. Lyons,¹³² A.-M. Magnan,¹³² S. Malik,¹³² A. Martelli,¹³² J. Nash,^{132,iii} A. Nikitenko,^{132,g} V. Palladino,¹³² M. Pesaresi,¹³² D. M. Raymond,¹³² A. Richards,¹³² A. Rose,¹³² E. Scott,¹³² C. Seez,¹³² A. Shtipliyski,¹³² G. Singh,¹³² M. Stoye,¹³² T. Strebler,¹³² S. Summers,¹³² A. Tapper,¹³² K. Uchida,¹³² T. Virdee,^{132,p} N. Wardle,¹³² D. Winterbottom,¹³² J. Wright,¹³² S. C. Zenz,¹³² J. E. Cole,¹³³ P. R. Hobson,¹³³ A. Khan,¹³³ P. Kyberd,¹³³ C. K. Mackay,¹³³ A. Morton,¹³³ I. D. Reid,¹³³ L. Teodorescu,¹³³ S. Zahid,¹³³ K. Call,¹³⁴ J. Dittmann,¹³⁴ K. Hatakeyama,¹³⁴ H. Liu,¹³⁴ C. Madrid,¹³⁴ B. McMaster,¹³⁴ N. Pastika,¹³⁴ C. Smith,¹³⁴ R. Bartek,¹³⁵ A. Dominguez,¹³⁵ A. Buccilli,¹³⁶ S. I. Cooper,¹³⁶ C. Henderson,¹³⁶ P. Rumerio,¹³⁶ C. West,¹³⁶ D. Arcaro,¹³⁷ T. Bose,¹³⁷ D. Gastler,¹³⁷ D. Pinna,¹³⁷ D. Rankin,¹³⁷ C. Richardson,¹³⁷ J. Rohlf,¹³⁷ L. Sulak,¹³⁷ D. Zou,¹³⁷ G. Benelli,¹³⁸ X. Coubez,¹³⁸ D. Cutts,¹³⁸ M. Hadley,¹³⁸ J. Hakala,¹³⁸ U. Heintz,¹³⁸ J. M. Hogan,^{138,mmm} K. H. M. Kwok,¹³⁸ E. Laird,¹³⁸ G. Landsberg,¹³⁸ J. Lee,¹³⁸ Z. Mao,¹³⁸ M. Narain,¹³⁸ S. Sagir,^{138,nnn} R. Syarif,¹³⁸ E. Usai,¹³⁸ D. Yu,¹³⁸ R. Band,¹³⁹ C. Brainerd,¹³⁹ R. Breedon,¹³⁹ D. Burns,¹³⁹ M. Calderon De La Barca Sanchez,¹³⁹ M. Chertok,¹³⁹ J. Conway,¹³⁹ R. Conway,¹³⁹ P. T. Cox,¹³⁹ R. Erbacher,¹³⁹ C. Flores,¹³⁹ G. Funk,¹³⁹ W. Ko,¹³⁹ O. Kukral,¹³⁹ R. Lander,¹³⁹ M. Mulhearn,¹³⁹ D. Pellett,¹³⁹ J. Pilot,¹³⁹ S. Shalhout,¹³⁹ M. Shi,¹³⁹ D. Stolp,¹³⁹ D. Taylor,¹³⁹ K. Tos,¹³⁹ M. Tripathi,¹³⁹ Z. Wang,¹³⁹ F. Zhang,¹³⁹ M. Bachtis,¹⁴⁰ C. Bravo,¹⁴⁰ R. Cousins,¹⁴⁰ A. Dasgupta,¹⁴⁰ A. Florent,¹⁴⁰ J. Hauser,¹⁴⁰ M. Ignatenko,¹⁴⁰ N. Mccoll,¹⁴⁰ S. Regnard,¹⁴⁰ D. Saltzberg,¹⁴⁰ C. Schnaible,¹⁴⁰ V. Valuev,¹⁴⁰ E. Bouvier,¹⁴¹ K. Burt,¹⁴¹ R. Clare,¹⁴¹ J. W. Gary,¹⁴¹ S. M. A. Ghiasi Shirazi,¹⁴¹ G. Hanson,¹⁴¹ G. Karapostoli,¹⁴¹ E. Kennedy,¹⁴¹ F. Lacroix,¹⁴¹ O. R. Long,¹⁴¹ M. Olmedo Negrete,¹⁴¹ M. I. Paneva,¹⁴¹ W. Si,¹⁴¹

L. Wang,¹⁴¹ H. Wei,¹⁴¹ S. Wimpenny,¹⁴¹ B. R. Yates,¹⁴¹ J. G. Branson,¹⁴² P. Chang,¹⁴² S. Cittolin,¹⁴² M. Derdzinski,¹⁴² R. Gerosa,¹⁴² D. Gilbert,¹⁴² B. Hashemi,¹⁴² A. Holzner,¹⁴² D. Klein,¹⁴² G. Kole,¹⁴² V. Krutelyov,¹⁴² J. Letts,¹⁴² M. Masciovecchio,¹⁴² D. Olivito,¹⁴² S. Padhi,¹⁴² M. Pieri,¹⁴² M. Sani,¹⁴² V. Sharma,¹⁴² S. Simon,¹⁴² M. Tadel,¹⁴² A. Vartak,¹⁴² S. Wasserbaech,^{142,ooo} J. Wood,¹⁴² F. Würthwein,¹⁴² A. Yagil,¹⁴² G. Zevi Della Porta,¹⁴² N. Amin,¹⁴³ R. Bhandari,¹⁴³ C. Campagnari,¹⁴³ M. Citron,¹⁴³ V. Dutta,¹⁴³ M. Franco Sevilla,¹⁴³ L. Gouskos,¹⁴³ R. Heller,¹⁴³ J. Incandela,¹⁴³ A. Ovcharova,¹⁴³ H. Qu,¹⁴³ J. Richman,¹⁴³ D. Stuart,¹⁴³ I. Suarez,¹⁴³ S. Wang,¹⁴³ J. Yoo,¹⁴³ D. Anderson,¹⁴⁴ A. Bornheim,¹⁴⁴ J. M. Lawhorn,¹⁴⁴ N. Lu,¹⁴⁴ H. B. Newman,¹⁴⁴ T. Q. Nguyen,¹⁴⁴ M. Spiropulu,¹⁴⁴ J. R. Vlimant,¹⁴⁴ R. Wilkinson,¹⁴⁴ S. Xie,¹⁴⁴ Z. Zhang,¹⁴⁴ R. Y. Zhu,¹⁴⁴ M. B. Andrews,¹⁴⁵ T. Ferguson,¹⁴⁵ T. Mudholkar,¹⁴⁵ M. Paulini,¹⁴⁵ M. Sun,¹⁴⁵ I. Vorobiev,¹⁴⁵ M. Weinberg,¹⁴⁵ J. P. Cumalat,¹⁴⁶ W. T. Ford,¹⁴⁶ F. Jensen,¹⁴⁶ A. Johnson,¹⁴⁶ E. MacDonald,¹⁴⁶ T. Mulholland,¹⁴⁶ R. Patel,¹⁴⁶ A. Perloff,¹⁴⁶ K. Stenson,¹⁴⁶ K. A. Ulmer,¹⁴⁶ S. R. Wagner,¹⁴⁶ J. Alexander,¹⁴⁷ J. Chaves,¹⁴⁷ Y. Cheng,¹⁴⁷ J. Chu,¹⁴⁷ A. Datta,¹⁴⁷ K. McDermott,¹⁴⁷ N. Mirman,¹⁴⁷ J. R. Patterson,¹⁴⁷ D. Quach,¹⁴⁷ A. Rinkevicius,¹⁴⁷ A. Ryd,¹⁴⁷ L. Skinnari,¹⁴⁷ L. Soffi,¹⁴⁷ S. M. Tan,¹⁴⁷ Z. Tao,¹⁴⁷ J. Thom,¹⁴⁷ J. Tucker,¹⁴⁷ P. Wittich,¹⁴⁷ M. Zientek,¹⁴⁷ S. Abdullin,¹⁴⁸ M. Albrow,¹⁴⁸ M. Alyari,¹⁴⁸ G. Apollinari,¹⁴⁸ A. Apresyan,¹⁴⁸ A. Apyan,¹⁴⁸ S. Banerjee,¹⁴⁸ L. A. T. Bauerdick,¹⁴⁸ A. Beretvas,¹⁴⁸ J. Berryhill,¹⁴⁸ P. C. Bhat,¹⁴⁸ K. Burkett,¹⁴⁸ J. N. Butler,¹⁴⁸ A. Canepa,¹⁴⁸ G. B. Cerati,¹⁴⁸ H. W. K. Cheung,¹⁴⁸ F. Chlebana,¹⁴⁸ M. Cremonesi,¹⁴⁸ J. Duarte,¹⁴⁸ V. D. Elvira,¹⁴⁸ J. Freeman,¹⁴⁸ Z. Gecse,¹⁴⁸ E. Gottschalk,¹⁴⁸ L. Gray,¹⁴⁸ D. Green,¹⁴⁸ S. Grünendahl,¹⁴⁸ O. Gutsche,¹⁴⁸ J. Hanlon,¹⁴⁸ R. M. Harris,¹⁴⁸ S. Hasegawa,¹⁴⁸ J. Hirschauer,¹⁴⁸ Z. Hu,¹⁴⁸ B. Jayatilaka,¹⁴⁸ S. Jindariani,¹⁴⁸ M. Johnson,¹⁴⁸ U. Joshi,¹⁴⁸ B. Klima,¹⁴⁸ M. J. Kortelainen,¹⁴⁸ B. Kreis,¹⁴⁸ S. Lammel,¹⁴⁸ D. Lincoln,¹⁴⁸ R. Lipton,¹⁴⁸ M. Liu,¹⁴⁸ T. Liu,¹⁴⁸ J. Lykken,¹⁴⁸ K. Maeshima,¹⁴⁸ J. M. Marraffino,¹⁴⁸ D. Mason,¹⁴⁸ P. McBride,¹⁴⁸ P. Merkel,¹⁴⁸ S. Mrenna,¹⁴⁸ S. Nahn,¹⁴⁸ V. O'Dell,¹⁴⁸ K. Pedro,¹⁴⁸ C. Pena,¹⁴⁸ O. Prokofyev,¹⁴⁸ G. Rakness,¹⁴⁸ L. Ristori,¹⁴⁸ A. Savoy-Navarro,^{148,ppp} B. Schneider,¹⁴⁸ E. Sexton-Kennedy,¹⁴⁸ A. Soha,¹⁴⁸ W. J. Spalding,¹⁴⁸ L. Spiegel,¹⁴⁸ S. Stoynev,¹⁴⁸ J. Strait,¹⁴⁸ N. Strobbe,¹⁴⁸ L. Taylor,¹⁴⁸ S. Tkaczyk,¹⁴⁸ N. V. Tran,¹⁴⁸ L. Uplegger,¹⁴⁸ E. W. Vaandering,¹⁴⁸ C. Vernieri,¹⁴⁸ M. Verzocchi,¹⁴⁸ R. Vidal,¹⁴⁸ M. Wang,¹⁴⁸ H. A. Weber,¹⁴⁸ A. Whitbeck,¹⁴⁸ D. Acosta,¹⁴⁹ P. Avery,¹⁴⁹ P. Bortignon,¹⁴⁹ D. Bourilkov,¹⁴⁹ A. Brinkerhoff,¹⁴⁹ L. Cadamuro,¹⁴⁹ A. Carnes,¹⁴⁹ D. Curry,¹⁴⁹ R. D. Field,¹⁴⁹ S. V. Gleyzer,¹⁴⁹ B. M. Joshi,¹⁴⁹ J. Konigsberg,¹⁴⁹ A. Korytov,¹⁴⁹ K. H. Lo,¹⁴⁹ P. Ma,¹⁴⁹ K. Matchev,¹⁴⁹ H. Mei,¹⁴⁹ G. Mitselmakher,¹⁴⁹ D. Rosenzweig,¹⁴⁹ K. Shi,¹⁴⁹ D. Sperka,¹⁴⁹ J. Wang,¹⁴⁹ S. Wang,¹⁴⁹ X. Zuo,¹⁴⁹ Y. R. Joshi,¹⁵⁰ S. Linn,¹⁵⁰ A. Ackert,¹⁵¹ T. Adams,¹⁵¹ A. Askew,¹⁵¹ S. Hagopian,¹⁵¹ V. Hagopian,¹⁵¹ K. F. Johnson,¹⁵¹ T. Kolberg,¹⁵¹ G. Martinez,¹⁵¹ T. Perry,¹⁵¹ H. Prosper,¹⁵¹ A. Saha,¹⁵¹ C. Schiber,¹⁵¹ R. Yohay,¹⁵¹ M. M. Baarmand,¹⁵² V. Bhopatkar,¹⁵² S. Colafranceschi,¹⁵² M. Hohmann,¹⁵² D. Noonan,¹⁵² M. Rahmani,¹⁵² T. Roy,¹⁵² F. Yumiceva,¹⁵² M. R. Adams,¹⁵³ L. Apanasevich,¹⁵³ D. Berry,¹⁵³ R. R. Betts,¹⁵³ R. Cavanaugh,¹⁵³ X. Chen,¹⁵³ S. Dittmer,¹⁵³ O. Evdokimov,¹⁵³ C. E. Gerber,¹⁵³ D. A. Hangal,¹⁵³ D. J. Hofman,¹⁵³ K. Jung,¹⁵³ J. Kamin,¹⁵³ C. Mills,¹⁵³ I. D. Sandoval Gonzalez,¹⁵³ M. B. Tonjes,¹⁵³ H. Trauger,¹⁵³ N. Varelas,¹⁵³ H. Wang,¹⁵³ X. Wang,¹⁵³ Z. Wu,¹⁵³ J. Zhang,¹⁵³ M. Alhusseini,¹⁵⁴ B. Bilki,^{154,qqq} W. Clarida,¹⁵⁴ K. Dilsiz,^{154,rrr} S. Durgut,¹⁵⁴ R. P. Gandrajula,¹⁵⁴ M. Haytmyradov,¹⁵⁴ V. Khristenko,¹⁵⁴ J.-P. Merlo,¹⁵⁴ A. Mestvirishvili,¹⁵⁴ A. Moeller,¹⁵⁴ J. Nachtman,¹⁵⁴ H. Ogul,^{154,sss} Y. Onel,¹⁵⁴ F. Ozok,^{154,ttt} A. Penzo,¹⁵⁴ C. Snyder,¹⁵⁴ E. Tiras,¹⁵⁴ J. Wetzel,¹⁵⁴ B. Blumenfeld,¹⁵⁵ A. Cocoros,¹⁵⁵ N. Eminizer,¹⁵⁵ D. Fehling,¹⁵⁵ L. Feng,¹⁵⁵ A. V. Gritsan,¹⁵⁵ W. T. Hung,¹⁵⁵ P. Maksimovic,¹⁵⁵ J. Roskes,¹⁵⁵ U. Sarica,¹⁵⁵ M. Swartz,¹⁵⁵ M. Xiao,¹⁵⁵ C. You,¹⁵⁵ A. Al-bataineh,¹⁵⁶ P. Baringer,¹⁵⁶ A. Bean,¹⁵⁶ S. Boren,¹⁵⁶ J. Bowen,¹⁵⁶ A. Bylinkin,¹⁵⁶ J. Castle,¹⁵⁶ S. Khalil,¹⁵⁶ A. Kropivnitskaya,¹⁵⁶ D. Majumder,¹⁵⁶ W. Mcbrayer,¹⁵⁶ M. Murray,¹⁵⁶ C. Rogan,¹⁵⁶ S. Sanders,¹⁵⁶ E. Schmitz,¹⁵⁶ J. D. Tapia Takaki,¹⁵⁶ Q. Wang,¹⁵⁶ S. Duric,¹⁵⁷ A. Ivanov,¹⁵⁷ K. Kaadze,¹⁵⁷ D. Kim,¹⁵⁷ Y. Maravin,¹⁵⁷ D. R. Mendis,¹⁵⁷ T. Mitchell,¹⁵⁷ A. Modak,¹⁵⁷ A. Mohammadi,¹⁵⁷ L. K. Saini,¹⁵⁷ F. Rebassoo,¹⁵⁸ D. Wright,¹⁵⁸ A. Baden,¹⁵⁹ O. Baron,¹⁵⁹ A. Belloni,¹⁵⁹ S. C. Eno,¹⁵⁹ Y. Feng,¹⁵⁹ C. Ferraioli,¹⁵⁹ N. J. Hadley,¹⁵⁹ S. Jabeen,¹⁵⁹ G. Y. Jeng,¹⁵⁹ R. G. Kellogg,¹⁵⁹ J. Kunkle,¹⁵⁹ A. C. Mignerey,¹⁵⁹ S. Nabili,¹⁵⁹ F. Ricci-Tam,¹⁵⁹ Y. H. Shin,¹⁵⁹ A. Skuja,¹⁵⁹ S. C. Tonwar,¹⁵⁹ K. Wong,¹⁵⁹ D. Abercrombie,¹⁶⁰ B. Allen,¹⁶⁰ V. Azzolini,¹⁶⁰ A. Baty,¹⁶⁰ G. Bauer,¹⁶⁰ R. Bi,¹⁶⁰ S. Brandt,¹⁶⁰ W. Busza,¹⁶⁰ I. A. Cali,¹⁶⁰ M. D'Alfonso,¹⁶⁰ Z. Demiragli,¹⁶⁰ G. Gomez Ceballos,¹⁶⁰ M. Goncharov,¹⁶⁰ P. Harris,¹⁶⁰ D. Hsu,¹⁶⁰ M. Hu,¹⁶⁰ Y. Iiyama,¹⁶⁰ G. M. Innocenti,¹⁶⁰ M. Klute,¹⁶⁰ D. Kovalskyi,¹⁶⁰ Y.-J. Lee,¹⁶⁰ P. D. Luckey,¹⁶⁰ B. Maier,¹⁶⁰ A. C. Marini,¹⁶⁰ C. Mcginn,¹⁶⁰ C. Mironov,¹⁶⁰ S. Narayanan,¹⁶⁰ X. Niu,¹⁶⁰ C. Paus,¹⁶⁰ C. Roland,¹⁶⁰ G. Roland,¹⁶⁰ Z. Shi,¹⁶⁰ G. S. F. Stephans,¹⁶⁰ K. Sumorok,¹⁶⁰ K. Tatar,¹⁶⁰ D. Velicanu,¹⁶⁰ J. Wang,¹⁶⁰ T. W. Wang,¹⁶⁰ B. Wyslouch,¹⁶⁰ A. C. Benvenuti,^{161,†} R. M. Chatterjee,¹⁶¹ A. Evans,¹⁶¹ P. Hansen,¹⁶¹ J. Hiltbrand,¹⁶¹ Sh. Jain,¹⁶¹ S. Kalafut,¹⁶¹ M. Krohn,¹⁶¹ Y. Kubota,¹⁶¹ Z. Lesko,¹⁶¹ J. Mans,¹⁶¹ N. Ruckstuhl,¹⁶¹ R. Rusack,¹⁶¹ M. A. Wadud,¹⁶¹ J. G. Acosta,¹⁶² S. Oliveros,¹⁶² E. Avdeeva,¹⁶³ K. Bloom,¹⁶³ D. R. Claes,¹⁶³ C. Fangmeier,¹⁶³ F. Golf,¹⁶³ R. Gonzalez Suarez,¹⁶³ R. Kamalieddin,¹⁶³ I. Kravchenko,¹⁶³

J. Monroy,¹⁶³ J. E. Siado,¹⁶³ G. R. Snow,¹⁶³ B. Stieger,¹⁶³ A. Godshalk,¹⁶⁴ C. Harrington,¹⁶⁴ I. Iashvili,¹⁶⁴ A. Kharchilava,¹⁶⁴ C. Mclean,¹⁶⁴ D. Nguyen,¹⁶⁴ A. Parker,¹⁶⁴ S. Rappoccio,¹⁶⁴ B. Roozbahani,¹⁶⁴ G. Alverson,¹⁶⁵ E. Barberis,¹⁶⁵ C. Freer,¹⁶⁵ Y. Haddad,¹⁶⁵ A. Hortiangtham,¹⁶⁵ D. M. Morse,¹⁶⁵ T. Orimoto,¹⁶⁵ R. Teixeira De Lima,¹⁶⁵ T. Wamorkar,¹⁶⁵ B. Wang,¹⁶⁵ A. Wisecarver,¹⁶⁵ D. Wood,¹⁶⁵ S. Bhattacharya,¹⁶⁶ J. Bueghly,¹⁶⁶ O. Charaf,¹⁶⁶ K. A. Hahn,¹⁶⁶ N. Mucia,¹⁶⁶ N. Odell,¹⁶⁶ M. H. Schmitt,¹⁶⁶ K. Sung,¹⁶⁶ M. Trovato,¹⁶⁶ M. Velasco,¹⁶⁶ R. Bucci,¹⁶⁷ N. Dev,¹⁶⁷ M. Hildreth,¹⁶⁷ K. Hurtado Anampa,¹⁶⁷ C. Jessop,¹⁶⁷ D. J. Karmgard,¹⁶⁷ N. Kellams,¹⁶⁷ K. Lannon,¹⁶⁷ W. Li,¹⁶⁷ N. Loukas,¹⁶⁷ N. Marinelli,¹⁶⁷ F. Meng,¹⁶⁷ C. Mueller,¹⁶⁷ Y. Musienko,^{167,ij} M. Planer,¹⁶⁷ A. Reinsvold,¹⁶⁷ R. Ruchti,¹⁶⁷ P. Siddireddy,¹⁶⁷ G. Smith,¹⁶⁷ S. Taroni,¹⁶⁷ M. Wayne,¹⁶⁷ A. Wightman,¹⁶⁷ M. Wolf,¹⁶⁷ A. Woodard,¹⁶⁷ J. Alimena,¹⁶⁸ L. Antonelli,¹⁶⁸ B. Bylsma,¹⁶⁸ L. S. Durkin,¹⁶⁸ S. Flowers,¹⁶⁸ B. Francis,¹⁶⁸ C. Hill,¹⁶⁸ W. Ji,¹⁶⁸ T. Y. Ling,¹⁶⁸ W. Luo,¹⁶⁸ B. L. Winer,¹⁶⁸ S. Cooperstein,¹⁶⁹ P. Elmer,¹⁶⁹ J. Hardenbrook,¹⁶⁹ S. Higginbotham,¹⁶⁹ A. Kalogeropoulos,¹⁶⁹ D. Lange,¹⁶⁹ M. T. Lucchini,¹⁶⁹ J. Luo,¹⁶⁹ D. Marlow,¹⁶⁹ K. Mei,¹⁶⁹ I. Ojalvo,¹⁶⁹ J. Olsen,¹⁶⁹ C. Palmer,¹⁶⁹ P. Piroué,¹⁶⁹ J. Salfeld-Nebgen,¹⁶⁹ D. Stickland,¹⁶⁹ C. Tully,¹⁶⁹ Z. Wang,¹⁶⁹ S. Malik,¹⁷⁰ S. Norberg,¹⁷⁰ A. Barker,¹⁷¹ V. E. Barnes,¹⁷¹ S. Das,¹⁷¹ L. Gutay,¹⁷¹ M. Jones,¹⁷¹ A. W. Jung,¹⁷¹ A. Khatiwada,¹⁷¹ B. Mahakud,¹⁷¹ D. H. Miller,¹⁷¹ N. Neumeister,¹⁷¹ C. C. Peng,¹⁷¹ S. Piperov,¹⁷¹ H. Qiu,¹⁷¹ J. F. Schulte,¹⁷¹ J. Sun,¹⁷¹ F. Wang,¹⁷¹ R. Xiao,¹⁷¹ W. Xie,¹⁷¹ T. Cheng,¹⁷² J. Dolen,¹⁷² N. Parashar,¹⁷² Z. Chen,¹⁷³ K. M. Ecklund,¹⁷³ S. Freed,¹⁷³ F. J. M. Geurts,¹⁷³ M. Kilpatrick,¹⁷³ W. Li,¹⁷³ B. P. Padley,¹⁷³ R. Redjimi,¹⁷³ J. Roberts,¹⁷³ J. Rorie,¹⁷³ W. Shi,¹⁷³ Z. Tu,¹⁷³ A. Zhang,¹⁷³ A. Bodek,¹⁷⁴ P. de Barbaro,¹⁷⁴ R. Demina,¹⁷⁴ Y. t. Duh,¹⁷⁴ J. L. Dulemba,¹⁷⁴ C. Fallon,¹⁷⁴ T. Ferbel,¹⁷⁴ M. Galanti,¹⁷⁴ A. Garcia-Bellido,¹⁷⁴ J. Han,¹⁷⁴ O. Hindrichs,¹⁷⁴ A. Khukhunaishvili,¹⁷⁴ E. Ranken,¹⁷⁴ P. Tan,¹⁷⁴ R. Taus,¹⁷⁴ A. Agapitos,¹⁷⁵ J. P. Chou,¹⁷⁵ Y. Gershtein,¹⁷⁵ E. Halkiadakis,¹⁷⁵ A. Hart,¹⁷⁵ M. Heindl,¹⁷⁵ E. Hughes,¹⁷⁵ S. Kaplan,¹⁷⁵ R. Kunnawalkam Elayavalli,¹⁷⁵ S. Kyriacou,¹⁷⁵ A. Lath,¹⁷⁵ R. Montalvo,¹⁷⁵ K. Nash,¹⁷⁵ M. Osherson,¹⁷⁵ H. Saka,¹⁷⁵ S. Salur,¹⁷⁵ S. Schnetzer,¹⁷⁵ D. Sheffield,¹⁷⁵ S. Somalwar,¹⁷⁵ R. Stone,¹⁷⁵ S. Thomas,¹⁷⁵ P. Thomassen,¹⁷⁵ M. Walker,¹⁷⁵ A. G. Delannoy,¹⁷⁶ J. Heideman,¹⁷⁶ G. Riley,¹⁷⁶ S. Spanier,¹⁷⁶ O. Bouhali,^{177,uuu} A. Celik,¹⁷⁷ M. Dalchenko,¹⁷⁷ M. De Mattia,¹⁷⁷ A. Delgado,¹⁷⁷ S. Dildick,¹⁷⁷ R. Eusebi,¹⁷⁷ J. Gilmore,¹⁷⁷ T. Huang,¹⁷⁷ T. Kamon,^{177,vvv} S. Luo,¹⁷⁷ R. Mueller,¹⁷⁷ D. Overton,¹⁷⁷ L. Perniè,¹⁷⁷ D. Rathjens,¹⁷⁷ A. Safonov,¹⁷⁷ N. Akchurin,¹⁷⁸ J. Damgov,¹⁷⁸ F. De Guio,¹⁷⁸ P. R. Duerdo,¹⁷⁸ S. Kunori,¹⁷⁸ K. Lamichhane,¹⁷⁸ S. W. Lee,¹⁷⁸ T. Mengke,¹⁷⁸ S. Muthumuni,¹⁷⁸ T. Peltola,¹⁷⁸ S. Undleeb,¹⁷⁸ I. Volobouev,¹⁷⁸ Z. Wang,¹⁷⁸ S. Greene,¹⁷⁹ A. Gurrola,¹⁷⁹ R. Janjam,¹⁷⁹ W. Johns,¹⁷⁹ C. Maguire,¹⁷⁹ A. Melo,¹⁷⁹ H. Ni,¹⁷⁹ K. Padeken,¹⁷⁹ J. D. Ruiz Alvarez,¹⁷⁹ P. Sheldon,¹⁷⁹ S. Tuo,¹⁷⁹ J. Velkovska,¹⁷⁹ M. Verweij,¹⁷⁹ Q. Xu,¹⁷⁹ M. W. Arenton,¹⁸⁰ P. Barria,¹⁸⁰ B. Cox,¹⁸⁰ R. Hirosky,¹⁸⁰ M. Joyce,¹⁸⁰ A. Ledovskoy,¹⁸⁰ H. Li,¹⁸⁰ C. Neu,¹⁸⁰ T. Sinthuprasith,¹⁸⁰ Y. Wang,¹⁸⁰ E. Wolfe,¹⁸⁰ F. Xia,¹⁸⁰ R. Harr,¹⁸¹ P. E. Karchin,¹⁸¹ N. Poudyal,¹⁸¹ J. Sturdy,¹⁸¹ P. Thapa,¹⁸¹ S. Zaleski,¹⁸¹ M. Brodski,¹⁸² J. Buchanan,¹⁸² C. Caillol,¹⁸² D. Carlsmith,¹⁸² S. Dasu,¹⁸² I. De Bruyn,¹⁸² L. Dodd,¹⁸² B. Gomber,¹⁸² M. Grothe,¹⁸² M. Herndon,¹⁸² A. Hervé,¹⁸² U. Hussain,¹⁸² P. Klabbers,¹⁸² A. Lanaro,¹⁸² K. Long,¹⁸² R. Loveless,¹⁸² T. Ruggles,¹⁸² A. Savin,¹⁸² V. Sharma,¹⁸² N. Smith,¹⁸² W. H. Smith,¹⁸² and N. Woods¹⁸²

(CMS Collaboration)

¹*Yerevan Physics Institute, Yerevan, Armenia*

²*Institut für Hochenergiephysik, Wien, Austria*

³*Institute for Nuclear Problems, Minsk, Belarus*

⁴*Universiteit Antwerpen, Antwerpen, Belgium*

⁵*Vrije Universiteit Brussel, Brussel, Belgium*

⁶*Université Libre de Bruxelles, Bruxelles, Belgium*

⁷*Ghent University, Ghent, Belgium*

⁸*Université Catholique de Louvain, Louvain-la-Neuve, Belgium*

⁹*Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*

¹⁰*Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*

^{11a}*Universidade Estadual Paulista, São Paulo, Brazil*

^{11b}*Universidade Federal do ABC, São Paulo, Brazil*

¹²*Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria*

¹³*University of Sofia, Sofia, Bulgaria*

¹⁴*Beihang University, Beijing, China*

¹⁵*Institute of High Energy Physics, Beijing, China*

¹⁶*State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

¹⁷*Tsinghua University, Beijing, China*

¹⁸*Universidad de Los Andes, Bogota, Colombia*

- ¹⁹*University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia*
- ²⁰*University of Split, Faculty of Science, Split, Croatia*
- ²¹*Institute Rudjer Boskovic, Zagreb, Croatia*
- ²²*University of Cyprus, Nicosia, Cyprus*
- ²³*Charles University, Prague, Czech Republic*
- ²⁴*Escuela Politecnica Nacional, Quito, Ecuador*
- ²⁵*Universidad San Francisco de Quito, Quito, Ecuador*
- ²⁶*Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt*
- ²⁷*National Institute of Chemical Physics and Biophysics, Tallinn, Estonia*
- ²⁸*Department of Physics, University of Helsinki, Helsinki, Finland*
- ²⁹*Helsinki Institute of Physics, Helsinki, Finland*
- ³⁰*Lappeenranta University of Technology, Lappeenranta, Finland*
- ³¹*IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France*
- ³²*Laboratoire Leprince-Ringuet, Ecole polytechnique, CNRS/IN2P3, Université Paris-Saclay, Palaiseau, France*
- ³³*Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France*
- ³⁴*Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France*
- ³⁵*Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France*
- ³⁶*Georgian Technical University, Tbilisi, Georgia*
- ³⁷*Tbilisi State University, Tbilisi, Georgia*
- ³⁸*RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany*
- ³⁹*RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
- ⁴⁰*RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany*
- ⁴¹*Deutsches Elektronen-Synchrotron, Hamburg, Germany*
- ⁴²*University of Hamburg, Hamburg, Germany*
- ⁴³*Karlsruher Institut fuer Technologie, Karlsruhe, Germany*
- ⁴⁴*Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece*
- ⁴⁵*National and Kapodistrian University of Athens, Athens, Greece*
- ⁴⁶*National Technical University of Athens, Athens, Greece*
- ⁴⁷*University of Ioánnina, Ioánnina, Greece*
- ⁴⁸*MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary*
- ⁴⁹*Wigner Research Centre for Physics, Budapest, Hungary*
- ⁵⁰*Institute of Nuclear Research ATOMKI, Debrecen, Hungary*
- ⁵¹*Institute of Physics, University of Debrecen, Debrecen, Hungary*
- ⁵²*Indian Institute of Science (IISc), Bangalore, India*
- ⁵³*National Institute of Science Education and Research, HBNI, Bhubaneswar, India*
- ⁵⁴*Panjab University, Chandigarh, India*
- ⁵⁵*University of Delhi, Delhi, India*
- ⁵⁶*Saha Institute of Nuclear Physics, HBNI, Kolkata, India*
- ⁵⁷*Indian Institute of Technology Madras, Madras, India*
- ⁵⁸*Bhabha Atomic Research Centre, Mumbai, India*
- ⁵⁹*Tata Institute of Fundamental Research-A, Mumbai, India*
- ⁶⁰*Tata Institute of Fundamental Research-B, Mumbai, India*
- ⁶¹*Indian Institute of Science Education and Research (IISER), Pune, India*
- ⁶²*Institute for Research in Fundamental Sciences (IPM), Tehran, Iran*
- ⁶³*University College Dublin, Dublin, Ireland*
- ^{64a}*INFN Sezione di Bari, Bari, Italy*
- ^{64b}*Università di Bari, Bari, Italy*
- ^{64c}*Politecnico di Bari, Bari, Italy*
- ^{65a}*INFN Sezione di Bologna, Bologna, Italy*
- ^{65b}*Università di Bologna, Bologna, Italy*
- ^{66a}*INFN Sezione di Catania, Catania, Italy*
- ^{66b}*Università di Catania, Catania, Italy*
- ^{67a}*INFN Sezione di Firenze, Firenze, Italy*
- ^{67b}*Università di Firenze, Firenze, Italy*

- ⁶⁸*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
^{69a}*INFN Sezione di Genova, Genova, Italy*
^{69b}*Università di Genova, Genova, Italy*
^{70a}*INFN Sezione di Milano-Bicocca, Milano, Italy*
^{70b}*Università di Milano-Bicocca, Milano, Italy*
^{71a}*INFN Sezione di Napoli, Napoli, Italy*
^{71b}*Università di Napoli 'Federico II', Napoli, Italy*
^{71c}*Università della Basilicata, Potenza, Italy*
^{71d}*Università G. Marconi, Roma, Italy*
^{72a}*INFN Sezione di Padova, Padova, Italy*
^{72b}*Università di Padova, Padova, Italy*
^{72c}*Università di Trento, Trento, Italy*
^{73a}*INFN Sezione di Pavia, Pavia, Italy*
^{73b}*Università di Pavia, Pavia, Italy*
^{74a}*INFN Sezione di Perugia, Perugia, Italy*
^{74b}*Università di Perugia, Perugia, Italy*
^{75a}*INFN Sezione di Pisa, Pisa, Italy*
^{75b}*Università di Pisa, Pisa, Italy*
^{75c}*Scuola Normale Superiore di Pisa, Pisa, Italy*
^{76a}*INFN Sezione di Roma, Rome, Italy*
^{76b}*Sapienza Università di Roma, Rome, Italy*
^{77a}*INFN Sezione di Torino, Torino, Italy*
^{77b}*Università di Torino, Torino, Italy*
^{77c}*Università del Piemonte Orientale, Novara, Italy*
^{78a}*INFN Sezione di Trieste, Trieste, Italy*
^{78b}*Università di Trieste, Trieste, Italy*
⁷⁹*Kyungpook National University, Daegu, Korea*
⁸⁰*Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea*
⁸¹*Hanyang University, Seoul, Korea*
⁸²*Korea University, Seoul, Korea*
⁸³*Sejong University, Seoul, Korea*
⁸⁴*Seoul National University, Seoul, Korea*
⁸⁵*University of Seoul, Seoul, Korea*
⁸⁶*Sungkyunkwan University, Suwon, Korea*
⁸⁷*Vilnius University, Vilnius, Lithuania*
⁸⁸*National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia*
⁸⁹*Universidad de Sonora (UNISON), Hermosillo, Mexico*
⁹⁰*Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico*
⁹¹*Universidad Iberoamericana, Mexico City, Mexico*
⁹²*Benemerita Universidad Autonoma de Puebla, Puebla, Mexico*
⁹³*Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico*
⁹⁴*University of Auckland, Auckland, New Zealand*
⁹⁵*University of Canterbury, Christchurch, New Zealand*
⁹⁶*National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan*
⁹⁷*National Centre for Nuclear Research, Swierk, Poland*
⁹⁸*Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland*
⁹⁹*Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal*
¹⁰⁰*Joint Institute for Nuclear Research, Dubna, Russia*
¹⁰¹*Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia*
¹⁰²*Institute for Nuclear Research, Moscow, Russia*
¹⁰³*Institute for Theoretical and Experimental Physics, Moscow, Russia*
¹⁰⁴*Moscow Institute of Physics and Technology, Moscow, Russia*
¹⁰⁵*National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia*
¹⁰⁶*P.N. Lebedev Physical Institute, Moscow, Russia*
¹⁰⁷*Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia*
¹⁰⁸*Novosibirsk State University (NSU), Novosibirsk, Russia*
¹⁰⁹*Institute for High Energy Physics of National Research Centre 'Kurchatov Institute', Protvino, Russia*
¹¹⁰*National Research Tomsk Polytechnic University, Tomsk, Russia*
¹¹¹*University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia*

- ¹¹²*Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain*
¹¹³*Universidad Autónoma de Madrid, Madrid, Spain*
¹¹⁴*Universidad de Oviedo, Oviedo, Spain*
¹¹⁵*Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain*
¹¹⁶*University of Ruhuna, Department of Physics, Matarara, Sri Lanka*
¹¹⁷*CERN, European Organization for Nuclear Research, Geneva, Switzerland*
¹¹⁸*Paul Scherrer Institut, Villigen, Switzerland*
¹¹⁹*ETH Zurich—Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland*
¹²⁰*Universität Zürich, Zurich, Switzerland*
¹²¹*National Central University, Chung-Li, Taiwan*
¹²²*National Taiwan University (NTU), Taipei, Taiwan*
¹²³*Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand*
¹²⁴*Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey*
¹²⁵*Middle East Technical University, Physics Department, Ankara, Turkey*
¹²⁶*Bogazici University, Istanbul, Turkey*
¹²⁷*Istanbul Technical University, Istanbul, Turkey*
¹²⁸*Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkov, Ukraine*
¹²⁹*National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine*
¹³⁰*University of Bristol, Bristol, United Kingdom*
¹³¹*Rutherford Appleton Laboratory, Didcot, United Kingdom*
¹³²*Imperial College, London, United Kingdom*
¹³³*Brunel University, Uxbridge, United Kingdom*
¹³⁴*Baylor University, Waco, Texas, USA*
¹³⁵*Catholic University of America, Washington, DC, USA*
¹³⁶*The University of Alabama, Tuscaloosa, Alabama, USA*
¹³⁷*Boston University, Boston, Massachusetts, USA*
¹³⁸*Brown University, Providence, Rhode Island, USA*
¹³⁹*University of California, Davis, Davis, California, USA*
¹⁴⁰*University of California, Los Angeles, California, USA*
¹⁴¹*University of California, Riverside, Riverside, California, USA*
¹⁴²*University of California, San Diego, La Jolla, California, USA*
¹⁴³*University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA*
¹⁴⁴*California Institute of Technology, Pasadena, California, USA*
¹⁴⁵*Carnegie Mellon University, Pittsburgh, Pennsylvania, USA*
¹⁴⁶*University of Colorado Boulder, Boulder, Colorado, USA*
¹⁴⁷*Cornell University, Ithaca, New York, USA*
¹⁴⁸*Fermi National Accelerator Laboratory, Batavia, Illinois, USA*
¹⁴⁹*University of Florida, Gainesville, Florida, USA*
¹⁵⁰*Florida International University, Miami, Florida, USA*
¹⁵¹*Florida State University, Tallahassee, Florida, USA*
¹⁵²*Florida Institute of Technology, Melbourne, Florida, USA*
¹⁵³*University of Illinois at Chicago (UIC), Chicago, Illinois, USA*
¹⁵⁴*The University of Iowa, Iowa City, Iowa, USA*
¹⁵⁵*Johns Hopkins University, Baltimore, Maryland, USA*
¹⁵⁶*The University of Kansas, Lawrence, Kansas, USA*
¹⁵⁷*Kansas State University, Manhattan, Kansas, USA*
¹⁵⁸*Lawrence Livermore National Laboratory, Livermore, California, USA*
¹⁵⁹*University of Maryland, College Park, Maryland, USA*
¹⁶⁰*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
¹⁶¹*University of Minnesota, Minneapolis, Minnesota, USA*
¹⁶²*University of Mississippi, Oxford, Mississippi, USA*
¹⁶³*University of Nebraska-Lincoln, Lincoln, Nebraska, USA*
¹⁶⁴*State University of New York at Buffalo, Buffalo, New York, USA*
¹⁶⁵*Northeastern University, Boston, Massachusetts, USA*
¹⁶⁶*Northwestern University, Evanston, Illinois, USA*
¹⁶⁷*University of Notre Dame, Notre Dame, Indiana, USA*
¹⁶⁸*The Ohio State University, Columbus, Ohio, USA*
¹⁶⁹*Princeton University, Princeton, New Jersey, USA*
¹⁷⁰*University of Puerto Rico, Mayaguez, Puerto Rico*
¹⁷¹*Purdue University, West Lafayette, Indiana, USA*

- ¹⁷²*Purdue University Northwest, Hammond, Indiana, USA*
¹⁷³*Rice University, Houston, Texas, USA*
¹⁷⁴*University of Rochester, Rochester, New York, USA*
¹⁷⁵*Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA*
¹⁷⁶*University of Tennessee, Knoxville, Tennessee, USA*
¹⁷⁷*Texas A&M University, College Station, Texas, USA*
¹⁷⁸*Texas Tech University, Lubbock, Texas, USA*
¹⁷⁹*Vanderbilt University, Nashville, Tennessee, USA*
¹⁸⁰*University of Virginia, Charlottesville, Virginia, USA*
¹⁸¹*Wayne State University, Detroit, Michigan, USA*
¹⁸²*University of Wisconsin - Madison, Madison, Wisconsin, USA*

[†]Deceased.

^aAlso at Vienna University of Technology, Vienna, Austria.

^bAlso at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France.

^cAlso at Universidade Estadual de Campinas, Campinas, Brazil.

^dAlso at Federal University of Rio Grande do Sul, Porto Alegre, Brazil.

^eAlso at Université Libre de Bruxelles, Bruxelles, Belgium.

^fAlso at University of Chinese Academy of Sciences, Beijing, China.

^gAlso at Institute for Theoretical and Experimental Physics, Moscow, Russia.

^hAlso at Joint Institute for Nuclear Research, Dubna, Russia.

ⁱAlso at British University in Egypt, Cairo, Egypt.

^jAlso at Suez University, Suez, Egypt.

^kAlso at Zewail City of Science and Technology, Zewail, Egypt.

^lAlso at Department of Physics, King Abdulaziz University, Jeddah, Saudi Arabia.

^mAlso at Université de Haute Alsace, Mulhouse, France.

ⁿAlso at Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow, Russia.

^oAlso at Tbilisi State University, Tbilisi, Georgia.

^pAlso at CERN, European Organization for Nuclear Research, Geneva, Switzerland.

^qAlso at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany.

^rAlso at University of Hamburg, Hamburg, Germany.

^sAlso at Brandenburg University of Technology, Cottbus, Germany.

^tAlso at Institute of Physics, University of Debrecen, Debrecen, Hungary.

^uAlso at Institute of Nuclear Research ATOMKI, Debrecen, Hungary.

^vAlso at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary.

^wAlso at IIT Bhubaneswar, Bhubaneswar, India.

^xAlso at Institute of Physics, Bhubaneswar, India.

^yAlso at Shoolini University, Solan, India.

^zAlso at University of Visva-Bharati, Santiniketan, India.

^{aa}Also at Isfahan University of Technology, Isfahan, Iran.

^{bb}Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran.

^{cc}Also at Università degli Studi di Siena, Siena, Italy.

^{dd}Also at Scuola Normale e Sezione dell'INFN, Pisa, Italy.

^{ee}Also at Kyunghee University, Seoul, Korea.

^{ff}Also at International Islamic University of Malaysia, Kuala Lumpur, Malaysia.

^{gg}Also at Malaysian Nuclear Agency, MOSTI, Kajang, Malaysia.

^{hh}Also at Consejo Nacional de Ciencia y Tecnología, Mexico city, Mexico.

ⁱⁱAlso at Warsaw University of Technology, Institute of Electronic Systems, Warsaw, Poland.

^{jj}Also at Institute for Nuclear Research, Moscow, Russia.

^{kk}Also at National Research Nuclear University 'Moscow Engineering Physics Institute' (MEPhI), Moscow, Russia.

^{ll}Also at St. Petersburg State Polytechnical University, St. Petersburg, Russia.

^{mm}Also at University of Florida, Gainesville, Florida, USA.

ⁿⁿAlso at P.N. Lebedev Physical Institute, Moscow, Russia.

^{oo}Also at California Institute of Technology, Pasadena, California, USA.

^{pp}Also at Budker Institute of Nuclear Physics, Novosibirsk, Russia.

^{qq}Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia.

^{rr}Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy.

^{ss}Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia.

^{tt}Also at National and Kapodistrian University of Athens, Athens, Greece.

^{uu}Also at Riga Technical University, Riga, Latvia.

- ^{vv} Also at Universität Zürich, Zurich, Switzerland.
- ^{ww} Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria.
- ^{xx} Also at Gaziosmanpasa University, Tokat, Turkey.
- ^{yy} Also at Istanbul Aydin University, Istanbul, Turkey.
- ^{zz} Also at Mersin University, Mersin, Turkey.
- ^{aaa} Also at Piri Reis University, Istanbul, Turkey.
- ^{bbb} Also at Adiyaman University, Adiyaman, Turkey.
- ^{ccc} Also at Ozyegin University, Istanbul, Turkey.
- ^{ddd} Also at Izmir Institute of Technology, Izmir, Turkey.
- ^{eee} Also at Marmara University, Istanbul, Turkey.
- ^{fff} Also at Kafkas University, Kars, Turkey.
- ^{ggg} Also at Istanbul University, Faculty of Science, Istanbul, Turkey.
- ^{hhh} Also at Istanbul Bilgi University, Istanbul, Turkey.
- ⁱⁱⁱ Also at Hacettepe University, Ankara, Turkey.
- ^{jjj} Also at Rutherford Appleton Laboratory, Didcot, United Kingdom.
- ^{kkk} Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.
- ^{lll} Also at Monash University, Faculty of Science, Clayton, Australia.
- ^{mmm} Also at Bethel University, St. Paul, Minnesota, USA.
- ⁿⁿⁿ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey.
- ^{ooo} Also at Utah Valley University, Orem, Utah, USA.
- ^{ppp} Also at Purdue University, West Lafayette, Indiana, USA.
- ^{qqq} Also at Beykent University, Istanbul, Turkey.
- ^{rrr} Also at Bingol University, Bingol, Turkey.
- ^{sss} Also at Sinop University, Sinop, Turkey.
- ^{ttt} Also at Mimar Sinan University, Istanbul, Istanbul, Turkey.
- ^{uuu} Also at Texas A&M University at Qatar, Doha, Qatar.
- ^{vvv} Also at Kyungpook National University, Daegu, Korea.