

## Influence of soil properties on N<sub>2</sub>O and CO<sub>2</sub> emissions from excreta deposited on tropical pastures in Kenya

Yuhao Zhu<sup>a,b</sup>, Lutz Merbold<sup>b</sup>, Sonja Leitner<sup>b</sup>, Longlong Xia<sup>a</sup>, David E. Pelster<sup>b,1</sup>, Eugenio Diaz-Pines<sup>c</sup>, Sheila Abwanda<sup>b</sup>, Paul M. Mutuo<sup>b</sup>, Klaus Butterbach-Bahl<sup>a,b,\*</sup>

<sup>a</sup> Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Karlsruhe Institute of Technology (KIT), Kreuzteckbahnstr. 19, 82467, Garmisch-Partenkirchen, Germany

<sup>b</sup> Mazingira Centre, International Livestock Research Institute (ILRI), P.O. Box 30709, Nairobi, 00100, Kenya

<sup>c</sup> Institute of Soil Research, University of Natural Resources and Life Sciences (BOKU), Peter-Jordan-Straße 82, 1190, Vienna, Austria

### ARTICLE INFO

#### Keywords:

Soil properties  
Excreta  
N<sub>2</sub>O emission factor  
Dung  
Urine  
Seasonality  
Sub-Saharan Africa (SSA)

### ABSTRACT

Urine and dung patches deposited by grazing cattle on grassland are an important source of nitrous oxide (N<sub>2</sub>O). While a number of studies have investigated the effects of excreta on soil N<sub>2</sub>O fluxes in developed economies and in China, observations in sub-Saharan Africa (SSA) are scarce. Moreover, the effects of soil properties (e.g. pH or texture) on N<sub>2</sub>O emissions from excreta patches have hardly been studied. In this study we investigated the importance of soil properties on N<sub>2</sub>O and carbon dioxide (CO<sub>2</sub>) emissions from cattle excreta (dung, urine, and manure [dung + urine]) for five typical tropical soils in Kenya. For this, intact soil cores were translocated from Western Kenya (Nandi county) to Nairobi, where N<sub>2</sub>O and CO<sub>2</sub> fluxes were measured over four individual periods (two during dry seasons and two during wet seasons). Fluxes were measured for between 25 and 73 days following surface application of excreta, depending on how quickly emissions returned to baseline. Both dung and manure applications led to increased CO<sub>2</sub> and N<sub>2</sub>O fluxes during both dry and wet seasons. On average, the N<sub>2</sub>O emission factor (EF) for manure was higher than for dung. The EFs during the wet season were higher for both the dung (0.12%) and urine (0.50%) compared to the dry season EFs (0.01% and 0.07% for dung and urine respectively). Soil type had no measurable effect on N<sub>2</sub>O and CO<sub>2</sub> emissions for either dung or manure application. In contrast, soil clay content was negatively ( $P < 0.05$ ) and pH positively ( $P < 0.05$ ) correlated with N<sub>2</sub>O emissions after urine application. Assuming an excreta-N ratio of dung to urine of 66:34, as evidenced in earlier studies for SSA, and averaging across all treatments and soils, we calculated a cattle excreta N<sub>2</sub>O EF of 0.14%, which is one magnitude lower than the IPCC default N<sub>2</sub>O EF of 2%. Our results call for a revision of the IPCC guidelines for calculating N<sub>2</sub>O emissions from excreta deposition on tropical rangelands.

### 1. Introduction

Nitrous oxide (N<sub>2</sub>O) is a potent greenhouse gas with a global warming potential 265 times greater than that of carbon dioxide (CO<sub>2</sub>) on a per mass basis over 100-years (IPCC, 2014), and it is estimated to account for 6% of total anthropogenic global warming (Davidson, 2009). Current atmospheric concentrations of N<sub>2</sub>O are approximately 18% higher than pre-industrial levels and concentrations were projected to further increase by 35–60% between 2007 and 2030 (Smith et al., 2007). More specifically, annual N<sub>2</sub>O emissions from excreta (i.e. dung

and urine) deposited by grazing livestock were estimated to be 1.5 Tg N<sub>2</sub>O–N yr<sup>-1</sup>, which is equivalent to 41% of all N<sub>2</sub>O emissions derived from global livestock production systems and 22% of total anthropogenic N<sub>2</sub>O emissions (Oenema et al., 2005; Taghizadeh-Toosi et al., 2011). Smallholder mixed-crop and pastoral livestock systems, which rely on native pasture grazing are widespread across sub-Saharan Africa (SSA), and support an estimated 123 million cattle (Thornton and Herrero, 2014). Due to the free grazing in daytime, it is estimated that in SSA more than 40% of total cattle excreta are deposited on rangelands (Rufino et al., 2006). As a result, direct N<sub>2</sub>O emissions from urine and

\* Corresponding author. Institute of Meteorology and Climate Research, Atmospheric Environmental Research (IMK-IFU), Karlsruhe Institute of Technology (KIT), Kreuzteckbahnstr. 19, 82467, Garmisch-Partenkirchen, Germany.

E-mail address: [klaus.butterbach-bahl@kit.edu](mailto:klaus.butterbach-bahl@kit.edu) (K. Butterbach-Bahl).

<sup>1</sup> Now at Agriculture and Agri-Food Canada, 2560 Blvd Hochelaga, Quebec, QC, Canada.

<https://doi.org/10.1016/j.soilbio.2019.107636>

Received 7 May 2019; Received in revised form 18 October 2019; Accepted 20 October 2019

Available online 23 October 2019

0038-0717/© 2019 The Authors.

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

dung excreta on SSA pastures were estimated to cause 62% of total annual N<sub>2</sub>O emissions in the region (Tully et al., 2017).

Up to 75–90% of nitrogen (N) ingested by grass-fed animals is returned to the soil through dung and urine excretion (Saggar et al., 2013; Bell et al., 2015). As adult cattle can excrete as much as 23 kg dung and 21 L urine over 13 dung and ten urine patches per day (Haynes and Williams, 1993), the amount of N contained in individual urine or dung patches by far exceeds plant N demands in the deposited area (Di and Cameron, 2007; Chadwick et al., 2018). In fact, the amount of N in a dung patch was calculated to be equivalent to up to 1130 kg N ha<sup>-1</sup> (Saarijärvi et al., 2006), while N loading rates for urine patches were found to be equivalent to 613 kg N ha<sup>-1</sup> for dairy cattle and 345 kg N ha<sup>-1</sup> for beef cattle (Selbie et al., 2015). Thus, high environmental N losses due to nitrate (NO<sub>3</sub><sup>-</sup>) leaching, ammonia (NH<sub>3</sub>) volatilization, or other gaseous emissions (N<sub>2</sub>O, nitric oxide [NO] and dinitrogen [N<sub>2</sub>]) are often associated with N excretion on rangeland (Chadwick et al., 2018).

Although there are many studies focusing on N<sub>2</sub>O emissions from excreta patches, most of these have focused either on timing (Lessa et al., 2014; Rochette et al., 2014; Bell et al., 2015) or application rate (Van Groenigen et al., 2005a; Sordi et al., 2014; Zhu et al., 2018) of excreta, while neglecting the influence of soil properties on N<sub>2</sub>O emissions. However, the magnitude and temporal dynamics of soil N<sub>2</sub>O emissions are determined not only by soil N availability but also by key soil properties (Butterbach-Bahl et al., 2013; Neira et al., 2015; Samad et al., 2016; Wang et al., 2018a,b; Ghezzehei et al., 2019). For example, soil texture and soil organic carbon content (SOC) affect the water holding capacity and therefore the gas diffusivity of soils. Both factors are crucial parameters controlling the availability of oxygen (O<sub>2</sub>) in soils and consequently soil microbial processes, thus N<sub>2</sub>O production (Davidson et al., 2000; Butterbach-Bahl et al., 2013; Balaine et al., 2016). Consistent with this, two other previous studies suggested that certain soil properties such as soil texture, pH and bulk density may affect denitrification rates and soil N<sub>2</sub>O emissions from livestock excreta applied to pasture soils (Cai and Akiyama, 2016; Wang et al., 2018a,b). However, to our knowledge, no previous study has systematically explored how soil properties may affect N<sub>2</sub>O emissions from excreta patches.

In general, N<sub>2</sub>O emissions from animal excreta on grasslands have been found to scale linearly with the dung mass (Zhu et al., 2018) or urine volume (Sordi et al., 2014), with higher N<sub>2</sub>O emissions from cattle urine compared to cattle dung due to higher N availability in urine and greater interactions of urine with the soil microbial communities as it infiltrates into the soil (Cai et al., 2017).

Depending on diet, N partitioning between cattle dung and urine is thought to range from 50:50 to 25:75 (Valk, 1994; Webb and Misselbrook, 2004; Van der Weerden et al., 2011; Chadwick et al., 2018). The split depends on the crude protein (CP) intake and concentration in feedstuffs. In Western Kenya, CP of feed ranges from 3.2 to 14% (Onyango et al., 2019), which is much less than that in intensive production systems such as in the USA where CP concentrations range from 17 to 23% (Council, 2015; Korir et al., 2016). The low CP concentration of feeds in SSA therefore results in an average N partitioning between cattle dung and urine of 66:34 (Rufino et al., 2006).

Given the differences in N<sub>2</sub>O emission factors (EF) between dung and urine (Cai et al., 2017), disregarding excreta-N partitioning might cause large uncertainties when estimating regional N<sub>2</sub>O emission inventories. However, the Intergovernmental Panel on Climate Change (IPCC) guidelines do not disaggregate the EF for urine and dung, rather they propose a default N<sub>2</sub>O EF of 2% excreta-N (IPCC, 2006). As such, many recent studies have suggested that disentangling N<sub>2</sub>O emissions from urine and dung is critical to improve our understanding of N<sub>2</sub>O emissions from grazed pastures, better assess this key source for atmospheric N<sub>2</sub>O and identify potential mitigation options (Van der Weerden et al., 2011; Krol et al., 2016; Chadwick et al., 2018). Despite differences in climate and soils, most countries in SSA, due to the scarcity of local and regional studies, still use the IPCC default value to estimate their

country-level greenhouse gas (GHG) inventories, even though this may not accurately reflect the specific conditions of SSA (Ogle et al., 2014). The objectives of our study were to: a) examine the influence of soil type on the CO<sub>2</sub> and N<sub>2</sub>O emissions from single dung, urine or manure patches after deposition on tropical pasture; b) quantify the cumulative CO<sub>2</sub> and N<sub>2</sub>O emissions from urine and dung applications to the five soil types; and c) examine whether CO<sub>2</sub> and N<sub>2</sub>O emission magnitudes from dung, urine or manure patches would differ between dry and wet seasons. We hypothesized that: a) soil properties such as pH or texture have minimal effects on N<sub>2</sub>O emissions from dung but significant effects on N<sub>2</sub>O emissions from urine deposited onto pasture; b) soils with a high SOC would have higher N<sub>2</sub>O emissions from urine deposited onto pasture than soils with low SOC; c) N<sub>2</sub>O emissions would be higher and effects of soil properties would be stronger during the wet than during the dry season; and d) soil CO<sub>2</sub> fluxes would be stimulated by all additions of excreta.

## 2. Materials and methods

### 2.1. Soil and site description

For our study, five soil types including (i) poorly-drained Gleysols, (ii) well-drained Nitisols, (iii) well-drained Acrisols, (iv) well-drained Cambisols, and (v) well-drained Ferralsols (IUSS Working Group WRB, 2014) differing in SOC (34–45 g C kg<sup>-1</sup> dry matter), clay content (29–53%) and pH (5.3–6.4) were selected in Nandi County, Western Kenya (Table 1). Sampling locations were selected based on a soil map for Kenya (Jaetzold et al., 2010) and the willingness of smallholder farmers to participate in the study. In total, 14 farms with grazing pastures or rangelands were chosen. There were four farms each for Nitisols and Gleysols, while two farms each were allocated for the other soil types. Before taking the cores, grass was cut down to 2 cm above the soil surface, after which intact soil cores with a diameter of 26 cm and a depth of 12 cm were collected with spades. In the area immediately adjacent to the soil cores, we took soil samples using 100 cm<sup>3</sup> soil cylinders to measure soil bulk density (BD) and with a 4.5 cm diameter soil auger to 5 cm depth to measure soil carbon (C), N and pH (for a detailed description see Saiz and Albrecht, 2016).

The intact soil cores were carefully wrapped in plastic bags and put into 50 L-buckets. Small holes were made to allow for gas exchange, and the cores were immediately transported to the Mazingira Centre of the International Livestock Research Institute (ILRI) in Nairobi, Kenya (S 1°16'13"; E 36°43'23"; altitude 1809 m asl). At the study site, soil cores were embedded into a flat grassland dominated by a mixture of Kikuyu grass (*Pennisetum clandestinum* Hochst. ex Chiov.) and Rhodes grass (*Chloris gayana* Kunth), i.e. grass species that were also found in the swards of the sampled rangelands. In addition to the five different soil types, dried and sieved (mesh width 2 mm) sand was included in this study as a control. Unfortunately, the sand contained high amount of soil, which still supported microbial activities. The soil cores were placed into holes (30 cm diameter, 15 cm deep) in an existing grassland immediately adjacent to the Mazingira Centre that had been lined with the same sand at the bottom. The sand was also used to fill the gaps on the sides of the cores in order to separate the soil cores from the adjacent soil. This procedure did not likely significantly affect N movement in the soil as the main rooting depth of the sward was generally less than 15 cm and as vertical N transport with the soil water movement was not influenced. After installation, soil cores were left to settle for three weeks to allow for equilibration with environmental conditions at the ILRI site. Thereafter, the grass was cut down to 2 cm again and excreta applied in the following days. Note that before the start of each trial, new soil cores were obtained from the field sites in Nandi County, transferred to ILRI, Nairobi, and placed in newly opened holes to avoid any legacy effects from the previous trial(s). A meteorological station was installed to record precipitation (tipping rain gauge, ECRN-100 high-resolution, Decagon, Pullman, WA; USA) and air temperature

**Table 1**

Soil pH, bulk density, carbon and nitrogen concentration, C/N ratio, soil sand and clay content of the different soils used in the experiment.


Soil type	pH	Bulk density (g cm <sup>-3</sup> )	C content (g kg <sup>-1</sup> dry matter)	N content (g kg <sup>-1</sup> dry matter)	CEC (cmol kg <sup>-1</sup> )	C/N ratio	Sand (%)	Clay (%)
Gleysols	6.0 ± 0.3 <sup>bc</sup>	0.94 ± 0.13 <sup>b</sup>	45.6 ± 9.3 <sup>a</sup>	3.91 ± 0.92 <sup>a</sup>	151 ± 16 <sup>a</sup>	12 ± 0.5 <sup>bc</sup>	27.2 ± 14.4 <sup>bc</sup>	46.6 ± 2.9 <sup>a</sup>
Nitisols	5.9 ± 0.5 <sup>c</sup>	0.95 ± 0.11 <sup>b</sup>	42.9 ± 8.7 <sup>a</sup>	3.63 ± 0.93 <sup>ab</sup>	147 ± 62 <sup>ab</sup>	12 ± 0.9 <sup>b</sup>	25.2 ± 5.0 <sup>c</sup>	50.1 ± 5.3 <sup>a</sup>
Acrisols	6.4 ± 0.6 <sup>b</sup>	1.17 ± 0.14 <sup>a</sup>	35.2 ± 6.1 <sup>b</sup>	3.12 ± 0.64 <sup>b</sup>	137 ± 37 <sup>ab</sup>	11 ± 0.6 <sup>c</sup>	52.7 ± 1.4 <sup>a</sup>	28.6 ± 21.4 <sup>b</sup>
Cambisols	5.6 ± 0.4 <sup>d</sup>	1.10 ± 0.13 <sup>a</sup>	34.7 ± 6.5 <sup>b</sup>	2.89 ± 0.69 <sup>c</sup>	85 ± 43 <sup>bc</sup>	12 ± 0.8 <sup>b</sup>	37.7 ± 2.8 <sup>b</sup>	42.6 ± 4.2 <sup>a</sup>
Ferralsols	5.3 ± 0.5 <sup>e</sup>	0.98 ± 0.08 <sup>b</sup>	37.0 ± 5.1 <sup>b</sup>	2.80 ± 0.61 <sup>c</sup>	67 ± 7 <sup>c</sup>	14 ± 1.7 <sup>a</sup>	24.5 ± 1.4 <sup>c</sup>	52.6 ± 4.3 <sup>a</sup>
Sand	8.4 ± 0.0 <sup>a</sup>	1.07 ± 0.11 <sup>ab</sup>	0.3 ± 0.1 <sup>c</sup>	0.00 ± 0.00 <sup>d</sup>	–	–	–	–

Values are mean ± standard deviation (n = 3). Different lowercase letters indicate significant differences within columns (P < 0.05).


and humidity (ATMOS 14, Decagon, Pullman, WA; USA) at a 5 min resolution. Soil moisture and temperature at 0.05 m depth were also measured during gas sampling (Decagon 5TM sensors, Decagon, Pullman, WA; USA).

## 2.2. Experimental design


In this experiment, four separate trials were conducted; two during the wet season and two during the dry season. Trial 1 was conducted from 26 July 2016 to 25 August 2016 (dry season) with eight cores of six different soils, resulting in a total of 48 soil cores. We applied 0.5 kg of fresh dung to the soil surface of four cores of each soil type, while the other four served as controls (i.e. no application, Fig. 1). For the following three trials, 12 new soil cores of the same six soils were installed, giving a total of 72 soil cores for each of the remaining three trials. Trial 2 was conducted from 16 October 2017 to 01 December 2017 (wet season). The 12 soil cores of each soil type were divided evenly into three groups that received either no application, or a surface application of either 0.5 kg fresh dung or 0.5 kg fresh manure (dung + urine) (Fig. 2). Trial 3 was conducted from 25 March 2018 to 14 May 2018 (wet season, Fig. 3), while the fourth trial was conducted from 02 July 2018 to 04 October 2018 (dry season, Fig. 4). In these two trials, 12 soil cores of each soil type were divided evenly into four groups with either no application, or with 0.5 L distilled water, 0.5 L urine or a surface application of 0.5 kg fresh dung. Excreta application was done following a few days of background measurements (i.e. prior to excreta application) at the beginning of each trial (see Figs. 1–4). Excreta application rates were scaled to the chamber area (0.04 m<sup>2</sup>) based on previous studies that found that emissions scale linearly with mass/volume of applied excreta (Sordi et al., 2014; Zhu et al., 2018). Due to an



**Fig. 1.** Observation time and treatments in Trial 1. The lower panel shows air temperature (°C) and precipitation (mm). Red arrows indicate application dates for dung. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)




**Fig. 2.** Observation time and treatments in Trial 2. The lower panel shows air temperature (°C) and precipitation (mm). Red arrows indicate application dates for dung or manure. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)



**Fig. 3.** Observation time and treatments in Trial 3. The lower panel shows air temperature (°C) and precipitation (mm). Red arrows indicate application dates for excreta (or water). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

initial lack of urine-collection harnesses, urine collection was only possible in 2018. Therefore, urine and soil interaction effects on soil N<sub>2</sub>O fluxes could only be studied in trials 3 and 4.

Annual rainfall distribution at Nairobi, Kenya (mean 1982–2012: 869 mm) is bimodal with a long rainy season from the end of March to



**Fig. 4.** Observation time and treatments in Trial 4. The lower panel shows air temperature ( $^{\circ}\text{C}$ ) and precipitation (mm). Red arrows indicate application dates for excreta (or water). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

the end of June and a short rainy season from October to December. Trials 1 and 4 were conducted during the dry season, with 34.8 mm precipitation in 25 days (trial 1) and 41.6 mm precipitation in 73 days (trial 4); trial 2 was conducted during the short rainy season with 83.2 mm precipitation in 43 days and trial 3 was conducted during the long rainy season with 596.4 mm precipitation in 39 days. Across our four observation periods, the mean temperatures were 16.0, 17.8, 17.8 and 16.5  $^{\circ}\text{C}$  during trials 1, 2, 3 and 4, respectively (Figs. 1–4).

Fresh dung and manure were collected at the ILRI farm from a cattle herd that was grazed on pasture dominated by a mixture of Kikuyu grass and Rhodes grass during the day and then housed in single-animal pens at night, where they had access to hay from Kikuyu and Rhodes grass. Excreta were collected from the concrete floors of each pen in the mornings, after the animals had been taken outside for grazing. To avoid contamination of dung with urine, only the upper half of dung cakes was collected from the floor. To collect manure (dung + urine), a waterproof plastic sheet was placed on the concrete floor, and the edges were raised upward to capture all the urine and dung from individual animals overnight, which was then homogenized in a bucket. Dung and manure were collected immediately before application to prevent nutrient losses during storage. Fresh urine was collected from three steers, each fitted with collection harnesses. The three urine samples were pooled after collection and the pH was measured immediately. Urine was acidified with 50 ml 20% HCl to a pH  $\leq 3$  to minimize N loss during collection and storage and then frozen at  $-20^{\circ}\text{C}$  until application. Before application, urine was thawed and the pH re-adjusted to the original pH with 2 M NaOH.

collected via a sampling port in the centre of the lid at times 0, 10, 20, and 30 min. The airtight lid was removed between samplings. Gas samples were immediately analyzed for  $\text{CO}_2$  and  $\text{N}_2\text{O}$  concentrations using a gas chromatograph (SRI 8610C, SRI Instruments, Torrance, CA, USA) equipped with a flame ionization detector (FID) for  $\text{CO}_2$  and an electron capture detector (ECD) for  $\text{N}_2\text{O}$ . The oven was operated at  $70^{\circ}\text{C}$ , and the ECD and FID were heated to 340 and  $350^{\circ}\text{C}$ . The carrier gas (pure  $\text{N}_2$ ) had a flow rate of  $25\text{ mL min}^{-1}$ .  $\text{CO}_2$  (5%) in 95%  $\text{N}_2$  was used as purge gas ( $3\text{ mL min}^{-1}$ ) for the ECD to avoid cross-interferences of the  $\text{N}_2\text{O}$  signal with  $\text{CO}_2$  (Zheng et al., 2004). Greenhouse gas fluxes were calculated on basis of temporal changes of  $\text{CO}_2$  and  $\text{N}_2\text{O}$  concentrations in the headspace of closed chambers. Slopes were calculated either using linear or non-linear regression analysis, using  $R^2$  values as decision criteria (Yao et al., 2015). Corrections for air pressure and headspace temperature were applied as described by Wolf et al. (2010). The flux detection limits were  $2.3\ \mu\text{g N}_2\text{O-N m}^{-2}\text{ h}^{-1}$  and  $1.3\text{ mg CO}_2\text{-C m}^{-2}\text{ h}^{-1}$  (Parkin et al., 2012). The fluxes were calculated by the following formula:

$$F = (b \times M_W \times V_{Ch} \times 60 \times 10^6) / (A_{Ch} \times V_m \times 10^9)$$

Where  $F$  is the flux rate ( $\text{mg CO}_2\text{-C m}^{-2}\text{ h}^{-1}$  or  $\mu\text{g N}_2\text{O-N m}^{-2}\text{ h}^{-1}$ ),  $b$  is the slope of concentration change ( $\text{ppm min}^{-1}$  or  $\text{ppb min}^{-1}$ ),  $M_W$  is the molecular weight of component ( $12\text{ g CO}_2\text{-C mol}^{-1}$  or  $28\text{ g N}_2\text{O-N mol}^{-1}$ ),  $V_{Ch}$  is the chamber volume ( $\text{m}^3$ ),  $A_{Ch}$  is the chamber area ( $\text{m}^2$ ), and  $V_m$  is the corrected standard gaseous molar volume ( $\text{m}^3\text{ mol}^{-1}$ ).

The sampling scheme was as follows: Before excreta application  $\text{CO}_2$  and  $\text{N}_2\text{O}$  fluxes were measured for at least three individual days. Following excreta application for the first trial, gas fluxes were measured twice per day for three days, then every two days for the next two weeks, and every three days for the last eight days. For the other three trials, fluxes were measured after excreta application daily for the first week, then every two days for the next two weeks, and three or four days per time for the remaining days. To capture the entire  $\text{CO}_2$  and  $\text{N}_2\text{O}$  emissions of the applied excreta, samples were collected for a minimum of one additional week (sometimes longer) after  $\text{CO}_2$  and  $\text{N}_2\text{O}$  fluxes had returned to background in all treatments, i.e. to levels as observed for the unamended soil cores. In the majority of cases fluxes returned to background levels within two or three weeks after application of dung or manure. Still, it needs to be noted that during the third trial (wet season) fluxes in the urine treatment only returned to background levels after five weeks. Accordingly, we extended the observation time for the fourth trial to make sure that the entire period with elevated fluxes was captured. In total,  $\text{CO}_2$  and  $\text{N}_2\text{O}$  fluxes were measured on 17 occasions/14 days (trial 1), 19 days (trial 2), 18 days (trial 3), and 30 days (trial 4).

Cumulative emissions were calculated using trapezoidal integration. Net cumulative emissions were calculated by subtracting cumulative emissions from control plots, i.e. plots not receiving excreta, from cumulative emissions of plots with excreta. The  $\text{N}_2\text{O}$  EF (i.e. the proportion of applied N emitted as  $\text{N}_2\text{O}$ ) was calculated by dividing the net cumulative emissions by the amount of added excreta-N according to the equation:

$$\text{N}_2\text{O EF (\%)} = \frac{\text{Cumulative N}_2\text{O emission (g N}_2\text{O - N) from excreta application} - \text{Cumulative N}_2\text{O emission (g N}_2\text{O - N) from control}}{\text{Nitrogen content applied as excreta (g N)}} \times 100$$

### 2.3. Measurements of $\text{CO}_2$ and $\text{N}_2\text{O}$ fluxes

For  $\text{CO}_2$  and  $\text{N}_2\text{O}$  fluxes measurements, we used closed static chambers (Norman et al., 1997). Polyvinyl chloride collars with 22.5 cm inner diameter and 13 cm height were inserted 8 cm into the soil on the plots with the soil cores. During chamber deployment, the collars were covered with an opaque, airtight lid and four 20 ml gas samples were

### 2.4. Soil and excreta analysis

For soil BD determination soil was sampled with a  $100\text{ cm}^3$  ring, the oven-dried at  $105^{\circ}\text{C}$  for 24 h, and weighed. Soil pH was measured using

a glass electrode in a water:soil suspension of 2.5:1 (weight:weight). Soil CEC was measured by ETH Zurich using BaCl<sub>2</sub>-TEA method (Hendershot et al., 2007). Water content of dung and manure was determined by drying fresh samples in the oven at 105 °C until constant weight. For total C and N, soil, dung or manure samples were dried at 50 °C for three days, ground and analyzed with an elemental combustion system (Costech International S. p.A., Milano, Italy). Total urine N concentration was analyzed by the Kjeldahl method (Kirk, 1950). Soil texture was estimated based on the soil particle size distribution as analyzed by the hydrometer technique (van Reeuwijk, 2002; Scrimgeour, 2008).

### 2.5. Data analysis

Two-way ANOVA with Tukey's HSD test was used to test for differences of cumulative emissions from unamended control plots and excreta amended plots and for testing differences between soils during the same trial. The differences of N<sub>2</sub>O EF for urine, dung and manure among soils in the same trial were also tested using two-way ANOVA with Tukey's HSD test. All statistical calculations were done in R v3.5.3 (R core team, 2019). The linear regression between urine EF and soil clay content or pH was conducted by Sigmaplot 12.5 (Systat Software, Inc. SigmaPlot for Windows).

## 3. Results

### 3.1. Soil properties

The pH of the different soils sampled from individual farms ranged from 5.3 ± 0.5 (Ferralsols) to 8.4 ± 0.0 (Sand) (Table 1). Soil BD varied from 0.94 ± 0.13 g cm<sup>-3</sup> (Gleysols) to 1.17 ± 0.14 g cm<sup>-3</sup> (Acrisols). The Gleysols had the highest C and N concentrations (45.6 g ± 9.3 C kg<sup>-1</sup> dry matter [DM] and 3.91 ± 0.92 g N kg<sup>-1</sup> DM), while the Sand had the lowest C and N concentrations (0.3 ± 0.1 g C kg<sup>-1</sup> DM and 0.0 ± 0.0 g N kg<sup>-1</sup> DM). There were large differences in the proportion of sand and clay in the different soils, with the sand content ranging from 24.5% (Ferralsols) to 52.7% (Acrisols) and clay content ranging from 28.6% (Acrisols) to 52.6% (Ferralsols) (Table 1).

### 3.2. Excreta properties

Dung properties used in the different trials varied substantially (Table 2) because the quality of the grasses in the pasture differed between dry and wet season. As a result, the dung used in trial 1 (dry season) had the highest N content (18.10 ± 0.12 g N kg<sup>-1</sup> DM) and lowest C content (397.2 ± 0.2 g C kg<sup>-1</sup> DM), while the lowest N concentration and highest C concentration was measured in the dung used for the third trial (wet season), which had 11.75 ± 0.09 g N kg<sup>-1</sup> DM and 422.7 ± 0.3 g C kg<sup>-1</sup> DM (Table 2). Consequently, the dung C/N ratio also varied significantly from 22 (trial 1, dry season) to 36 (trial 3, wet season). Compared to dung, manure had a higher water content, and lower C and N concentrations. Urine N concentration also varied between the two trials and ranged from 6.42 ± 0.23 g N L<sup>-1</sup> (trial 3, wet season) to 5.69 ± 0.14 g N L<sup>-1</sup> (trial 4, dry season) (Table 2).

**Table 2**

Water content, carbon and nitrogen concentrations and C/N ratio of dung and nitrogen concentrations of urine applied to soil cores during the four trials.

Period	Season	Excreta type	Water content (%)	C concentration (g C kg <sup>-1</sup> DM)	N concentration (g N kg <sup>-1</sup> DM or g N L <sup>-1</sup> )	C/N ratio
26-Jul — 25-Aug-16	Dry season	dung	82.7 ± 0.1 <sup>a</sup>	397.2 ± 0.2 <sup>d</sup>	18.10 ± 0.12 <sup>a</sup>	22 ± 0.1 <sup>e</sup>
16-Oct — 01-Dec-17	Wet season	dung	80.6 ± 0.2 <sup>c</sup>	412.0 ± 0.2 <sup>b</sup>	13.60 ± 0.04 <sup>c</sup>	30 ± 0.1 <sup>c</sup>
		manure	81.8 ± 0.1 <sup>b</sup>	399.9 ± 0.5 <sup>c</sup>	11.93 ± 0.03 <sup>d</sup>	34 ± 0.1 <sup>b</sup>
25-Mar — 14-May-18	Wet season	dung	81.7 ± 0.2 <sup>b</sup>	422.7 ± 0.3 <sup>a</sup>	11.75 ± 0.09 <sup>d</sup>	36 ± 0.3 <sup>a</sup>
		urine	—	—	6.42 ± 0.23 <sup>A</sup>	—
02-Jul — 04-Oct-18	Dry season	dung	78.9 ± 0.4 <sup>d</sup>	410.3 ± 1.5 <sup>b</sup>	14.19 ± 0.17 <sup>b</sup>	29 ± 0.4 <sup>d</sup>
		urine	—	—	5.69 ± 0.14 <sup>B</sup>	—

Values are mean ± standard deviation (n = 3). Different lowercase letters indicate significant differences among dung and manure property and the uppercase letters indicate the significant difference of urine N concentration (P < 0.05).

### 3.3. Background N<sub>2</sub>O and CO<sub>2</sub> emissions from different soils

Cumulative CO<sub>2</sub> emissions from the unamended soils only differed significantly between soil types in trial 3 (wet season, 2018) with the lowest emissions from the Sand (1186 ± 235 kg CO<sub>2</sub>-C ha<sup>-1</sup> 39 days<sup>-1</sup>) and the highest from the Acrisols (3234 ± 968 kg CO<sub>2</sub>-C ha<sup>-1</sup> 39 days<sup>-1</sup>) (P < 0.05). Cumulative N<sub>2</sub>O emissions from the different unamended soils were similar in all of the trials (Tables 3–6). Soil N<sub>2</sub>O fluxes were largely similar or below the detection limit (2.3 μg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup>) during the dry season (trial 1 and 4; Figs. 5 and 8) and the only notable fluxes were following a re-wetting by a 28 mm rainfall at the end of trial 1. As expected, rainfall did not cause elevated N<sub>2</sub>O fluxes for the sand plots (Fig. 5). During the wet season, soil CO<sub>2</sub> and N<sub>2</sub>O fluxes were higher than during the dry season (see Figs. 5–8).

### 3.4. Influence of soil type and type of excreta on CO<sub>2</sub> and N<sub>2</sub>O emissions

In all trials, dung and manure application increased CO<sub>2</sub> fluxes from all soils, with higher fluxes observed during wet seasons than during dry seasons (Figs. 5–8). The highest CO<sub>2</sub> flux of 890 ± 70 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> was measured in Acrisols amended with manure in trial 2 (Fig. 6). Urine application increased CO<sub>2</sub> fluxes more than either dung or water application within 1–2 days in all soils. However, the stimulative effect of urine additions differed between soils, ranging from 456 ± 35 (Gleysols) to 1464 ± 175 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> (Ferralsols) in trial 3 (Fig. 7), and from 71 ± 3 (Acrisols) to 1026 ± 81 mg CO<sub>2</sub>-C m<sup>-2</sup> h<sup>-1</sup> (Gleysols) in trial 4 (Fig. 8).

The N<sub>2</sub>O fluxes also increased following either dung or manure application. Among dung and manure applications, the highest N<sub>2</sub>O flux of 461 ± 136 μg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup> was measured for manure amended Gleysols in trial 2 (Fig. 6). Compared to unamended control soils, cumulative N<sub>2</sub>O emissions from dung application were similar across all soils during all trials (Tables 3–6), while during trial 2 the manure application resulted in higher cumulative N<sub>2</sub>O emissions compared to the unamended controls only in the Gleysols and Ferralsols (Table 4; P < 0.05).

Neither the water nor the dung application significantly increased cumulative N<sub>2</sub>O emissions relative to the unamended control across all soil types and all trials. However, the response to urine additions was highly variable across soils (Tables 5 and 6), with fluxes ranging from 33 ± 38 (Ferralsols – trial 4; Fig. 8) to 8760 ± 1322 μg N<sub>2</sub>O-N m<sup>-2</sup> h<sup>-1</sup> (Acrisols – trial 3; Fig. 7). Soil N<sub>2</sub>O fluxes were not only larger but also lasted for longer (several weeks) for urine as compared to dung additions (Figs. 7 and 8). As a result, cumulative N<sub>2</sub>O emissions were higher for urine application compared to dung application in the Acrisols (Trials 3 and 4) and Sand (only trial 4) (Tables 5 and 6). The urine EF was negatively correlated with soil clay content in both dry (EF = 0.5936–0.0118 × soil clay content, n = 15, R<sup>2</sup> = 0.53, P < 0.05) and wet season (EF = 2.5526–0.0466 × soil clay content, n = 15, R<sup>2</sup> = 0.51, P < 0.05), while positively correlated with soil pH in both dry (EF = -1.3850 + 0.2367 × soil pH, n = 15, R<sup>2</sup> = 0.36, P < 0.05) and wet season (EF = -6.5575 + 1.2290 × soil pH, n = 15, R<sup>2</sup> = 0.60, P < 0.05).

**Table 3**

Cumulative CO<sub>2</sub> and N<sub>2</sub>O emission and dung N<sub>2</sub>O EF (% applied excreta-N) over 25 days as affected by addition of 0.5 kg cattle dung to different soil cores in Trial 1 (01–25 August 2016, dry season).

Soil type	Cumulative CO <sub>2</sub> emissions (kg CO <sub>2</sub> -C ha <sup>-1</sup> )		Cumulative N <sub>2</sub> O emissions (g N <sub>2</sub> O-N ha <sup>-1</sup> )		N <sub>2</sub> O EF (%)
	None	Dung	None	Dung	
Gleysols	472 ± 100 <sup>Aa</sup>	754 ± 148 <sup>Aa</sup>	22.3 ± 11.0 <sup>Aa</sup>	49.7 ± 44.1 <sup>Aab</sup>	0.01 ± 0.01 <sup>a</sup>
Nitisols	353 ± 117 <sup>Ba</sup>	684 ± 75 <sup>Aa</sup>	11.6 ± 9.3 <sup>Aa</sup>	31.3 ± 17.4 <sup>Aab</sup>	0.00 ± 0.00 <sup>a</sup>
Acrisols	518 ± 149 <sup>Aa</sup>	743 ± 113 <sup>Aa</sup>	71.8 ± 71.4 <sup>Aa</sup>	108.5 ± 77.6 <sup>Aa</sup>	0.01 ± 0.02 <sup>a</sup>
Cambisols	441 ± 69 <sup>Ba</sup>	753 ± 125 <sup>Aa</sup>	8.9 ± 5.4 <sup>Aa</sup>	29.3 ± 7.3 <sup>Aab</sup>	0.00 ± 0.00 <sup>a</sup>
Ferralsols	386 ± 82 <sup>Ba</sup>	747 ± 132 <sup>Aa</sup>	12.7 ± 13.3 <sup>Aa</sup>	21.4 ± 15.1 <sup>Ab</sup>	0.00 ± 0.00 <sup>a</sup>
Sand	505 ± 140 <sup>Aa</sup>	748 ± 121 <sup>Aa</sup>	15.5 ± 15.4 <sup>Aa</sup>	20.9 ± 21.2 <sup>Ab</sup>	0.00 ± 0.00 <sup>a</sup>

Values are mean ± standard deviation (n = 4); different lowercase letters indicate significant differences among soil types within the same treatment and different uppercase letters indicate significant differences among treatment within the same soil type (P < 0.05).

**Table 4**

Cumulative CO<sub>2</sub> and N<sub>2</sub>O emission and N<sub>2</sub>O EF (% applied excreta-N) over 43 days as affected by addition of 0.5 kg cattle dung or 0.5 kg cattle manure to different soil cores in Trial 2 (20 October — 01 December 2017, wet season).

Soil type	Cumulative CO <sub>2</sub> emissions (kg CO <sub>2</sub> -C ha <sup>-1</sup> )			Cumulative N <sub>2</sub> O emissions (g N <sub>2</sub> O-N ha <sup>-1</sup> )			N <sub>2</sub> O EF (%)	
	None	Dung	Manure	None	Dung	Manure	Dung	Manure
Gleysols	2028 ± 226 <sup>Aa</sup>	2635 ± 520 <sup>Aa</sup>	2951 ± 573 <sup>Aa</sup>	127 ± 37 <sup>Ba</sup>	579 ± 296 <sup>ABa</sup>	916 ± 86 <sup>Aa</sup>	0.13 ± 0.08 <sup>Aa</sup>	0.28 ± 0.03 <sup>Aa</sup>
Nitisols	1890 ± 455 <sup>Aa</sup>	2555 ± 101 <sup>Aa</sup>	2705 ± 232 <sup>Aa</sup>	150 ± 38 <sup>Aa</sup>	644 ± 448 <sup>Aa</sup>	765 ± 56 <sup>Aa</sup>	0.14 ± 0.13 <sup>Aa</sup>	0.22 ± 0.03 <sup>Aa</sup>
Acrisols	1997 ± 134 <sup>Aa</sup>	3034 ± 661 <sup>Aa</sup>	3024 ± 526 <sup>Aa</sup>	218 ± 47 <sup>Aa</sup>	571 ± 253 <sup>Aa</sup>	709 ± 479 <sup>Aa</sup>	0.10 ± 0.06 <sup>Aa</sup>	0.17 ± 0.18 <sup>Aa</sup>
Cambisols	1945 ± 405 <sup>Aa</sup>	3093 ± 544 <sup>Aa</sup>	3163 ± 1041 <sup>Aa</sup>	110 ± 30 <sup>Aa</sup>	818 ± 616 <sup>Aa</sup>	814 ± 538 <sup>Aa</sup>	0.20 ± 0.17 <sup>Aa</sup>	0.25 ± 0.18 <sup>Aa</sup>
Ferralsols	1540 ± 286 <sup>Ba</sup>	3092 ± 833 <sup>Aa</sup>	2949 ± 711 <sup>ABa</sup>	158 ± 148 <sup>Ba</sup>	620 ± 154 <sup>ABa</sup>	916 ± 403 <sup>Aa</sup>	0.13 ± 0.07 <sup>Aa</sup>	0.27 ± 0.09 <sup>Aa</sup>
Sand	1959 ± 1273 <sup>Aa</sup>	2273 ± 328 <sup>Aa</sup>	2501 ± 729 <sup>Aa</sup>	179 ± 112 <sup>Aa</sup>	335 ± 203 <sup>Aa</sup>	264 ± 96 <sup>Aa</sup>	0.05 ± 0.07 <sup>Aa</sup>	0.03 ± 0.05 <sup>Aa</sup>

Values are mean ± standard deviation (n = 4); there was no significant differences among soil types within the same treatment; different uppercase letters indicate significant differences among treatment within the same soil type (P < 0.05).

### 3.5. N<sub>2</sub>O emission factors for dung, manure and urine

The N<sub>2</sub>O EF for dung in trial 1 (dry season) was negligible and similar among all soil types, varying from 0.00 to 0.01% (Table 3), while the N<sub>2</sub>O EF for dung was in the range of 0.00–0.20% in all other trial periods (Tables 4–6). Despite a lower dung N concentration, the dung EF was higher during the wet seasons (Trials 2 and 3) than during the dry seasons (Trials 1 and 4; P < 0.05). However, the N<sub>2</sub>O EF for dung did not differ across soils. Manure was only applied in the second trial (wet season), and during this period the manure N<sub>2</sub>O EF ranged from 0.03 to 0.28% with no differences between the soils tested (Table 4).

In contrast to dung and manure, the urine N<sub>2</sub>O EF varied largely among the different soils during both wet and dry seasons. The highest urine N<sub>2</sub>O EF was observed for Acrisols both in the wet season (1.36%) and dry season (0.29%), while the lowest urine N<sub>2</sub>O EF was measured in the Ferralsols (wet season: 0.12%; dry season: 0.01%). Furthermore, in both wet and dry seasons, the urine N<sub>2</sub>O EF was markedly higher for Acrisols than EFs for all other soils (Tables 5 and 6; P < 0.05).

## 4. Discussion

### 4.1. Influence of soil properties on CO<sub>2</sub> and N<sub>2</sub>O emissions after excreta application

In our study, the CO<sub>2</sub> fluxes after dung or manure applications were similar and no soil type effect was detected indicating that the majority of CO<sub>2</sub> releases was from the fresh dung or manure itself. This was similar to observations by Lin et al. (2009) after yak dung application under laboratory conditions. The short-lived CO<sub>2</sub> fluxes after urine application were likely associated with the rapid hydrolysis of urine urea (Ambus et al., 2007; Cai et al., 2017), during which CO<sub>2</sub> is released, and did not cause any significant difference in cumulative CO<sub>2</sub> emissions among soils in any of the trials.

Consistent with our original hypothesis, the N<sub>2</sub>O EF from both manure and dung application to pasture was not affected by the underlying soil properties such as e.g. soil C/N ratio, SOC or pH. This was likely because the low dung-N concentrations limited N availability for

N<sub>2</sub>O formation in the dung itself, and because the rapid crusting of the dung hampered incorporation of excreta-N into the soil (Zhu et al., 2018). However, the presence of termites in some of the soil cores was noted shortly after application, which might have altered soil aeration and thus influenced N<sub>2</sub>O production and emission. Also, the termites may have transferred some of the dung/manure away from the area covered by the chambers. Therefore, it is possible that the presence of the termites may have contributed to our observation that soil texture and pH did not have measurable effects on soil N<sub>2</sub>O production after dung application. The lack of measurable soil effects was consistent with earlier findings by Van der Weerden et al. (2011) who also did not find differences in the N<sub>2</sub>O EF for dung in three contrasting regions with differing soil types.

In line with a recent global meta-analysis (Wang et al., 2018a,b), the N<sub>2</sub>O EF for urine application to soils was negatively correlated with clay content. Clay particles hold water tightly in soil aggregates. Moreover, clay soils mostly have a low gas diffusivity due to small mean pore sizes (Weitz et al., 2001; Gu et al., 2013). Both factors favor anaerobic conditions in soils and, thus, might promote complete denitrification towards the end product N<sub>2</sub>, causing lower N<sub>2</sub>O emissions from clay soils (Weitz et al., 2001). In addition, high clay content generally correlates with a high cation exchange capacity. As a result, the mineralized NH<sub>4</sub><sup>+</sup> ion can be adsorbed due to the high CEC or even fixed to the clay minerals (Chantigny et al., 2004), which decreases soil NH<sub>4</sub><sup>+</sup> ion availability for nitrification and, thus, NO<sub>3</sub><sup>-</sup> ion production. Thus, the soil ammonium sorption capacity has been found to affect N<sub>2</sub>O production in soils which otherwise showed largely similar physical and chemical properties (Venterea et al., 2015). Consequently, a number of studies have found limited N<sub>2</sub>O production following urine application to clay-rich soils (Jarecki et al., 2008; Zhou et al., 2017). Contradicting our findings, a study in Canada reported higher urine EFs in a clay-rich soil in comparison with a sandy loam soil and argued that the good soil aeration in the sandy loam soil limited the N<sub>2</sub>O production due to denitrification (Rochette et al., 2014).

In addition to soil clay content, soil pH is also an important factor controlling the magnitude of N<sub>2</sub>O emissions after N additions (Wang et al., 2018a,b). Wang et al., 2018a,b found in a meta-data analysis on

**Table 5**  
Cumulative CO<sub>2</sub> and N<sub>2</sub>O emission and N<sub>2</sub>O EF over 39 days as affected by addition of 0.5 L cattle urine or 0.5 kg cattle dung to different soil cores in Trial 3 (06 April –14 May 2018, wet season).


Soil type	Cumulative CO <sub>2</sub> emissions (kg CO <sub>2</sub> -C ha <sup>-1</sup> )			Cumulative N <sub>2</sub> O emissions (g N <sub>2</sub> O-N ha <sup>-1</sup> )			N <sub>2</sub> O EF (%)	
	None	Water	Dung	None	Water	Dung	Urine	Dung
	Gleysols	2890 ± 737 <sup>Aa</sup>	2610 ± 136 <sup>Aa</sup>	3881 ± 1125 <sup>Aa</sup>	219 ± 69 <sup>Aa</sup>	177 ± 73 <sup>Aa</sup>	640 ± 332 <sup>Aa</sup>	0.22 ± 0.12 <sup>Ab</sup>
Nitisols	2540 ± 939 <sup>Ab</sup>	2499 ± 526 <sup>Aa</sup>	3168 ± 498 <sup>Aa</sup>	186 ± 120 <sup>Aa</sup>	448 ± 440 <sup>Aa</sup>	533 ± 365 <sup>Aa</sup>	0.54 ± 0.49 <sup>Ab</sup>	0.12 ± 0.09 <sup>Aa</sup>
Acrisols	3234 ± 968 <sup>Aa</sup>	2158 ± 327 <sup>Aa</sup>	3244 ± 958 <sup>Aa</sup>	186 ± 210 <sup>Ba</sup>	194 ± 187 <sup>Ba</sup>	241 ± 273 <sup>Ba</sup>	1.36 ± 0.70 <sup>Aa</sup>	0.02 ± 0.02 <sup>Ba</sup>
Cambisols	2457 ± 123 <sup>Ab</sup>	2398 ± 370 <sup>Aa</sup>	3578 ± 32 <sup>Aa</sup>	133 ± 119 <sup>Aa</sup>	101 ± 93 <sup>Aa</sup>	377 ± 153 <sup>Aa</sup>	0.25 ± 0.08 <sup>Ab</sup>	0.09 ± 0.06 <sup>Aa</sup>
Ferralsols	2078 ± 172 <sup>Ab</sup>	2503 ± 671 <sup>Aa</sup>	3300 ± 572 <sup>Aa</sup>	87 ± 35 <sup>Aa</sup>	240 ± 67 <sup>Aa</sup>	218 ± 267 <sup>Aa</sup>	0.12 ± 0.12 <sup>Ab</sup>	0.05 ± 0.10 <sup>Aa</sup>
Sand	1186 ± 235 <sup>Bb</sup>	1397 ± 140 <sup>Ba</sup>	3484 ± 145 <sup>Aa</sup>	52 ± 18 <sup>Aa</sup>	88 ± 39 <sup>Aa</sup>	178 ± 110 <sup>Aa</sup>	0.15 ± 0.03 <sup>Ab</sup>	0.04 ± 0.04 <sup>Aa</sup>

Values are mean ± standard deviation (n = 3); different lowercase letters indicate significant differences among soil types within the same treatment and different uppercase letters indicate significant differences among treatment within the same soil type (P < 0.05).

**Table 6**  
Cumulative CO<sub>2</sub> and N<sub>2</sub>O emission and N<sub>2</sub>O EF over 73 days as affected by addition of 0.5 L cattle urine or 0.5 kg cattle dung to different soil cores in Trial 4 (24 July — 04 October 2018, dry season).

Soil type	Cumulative CO <sub>2</sub> emissions (kg CO <sub>2</sub> -C ha <sup>-1</sup> )			Cumulative N <sub>2</sub> O emissions (g N <sub>2</sub> O-N ha <sup>-1</sup> )			N <sub>2</sub> O EF (%)	
	None	Water	Dung	None	Water	Dung	Urine	Dung
	Gleysols	1589 ± 136 <sup>Aa</sup>	1989 ± 564 <sup>Aa</sup>	1716 ± 530 <sup>Aa</sup>	18 ± 25 <sup>Aa</sup>	75 ± 63 <sup>Aa</sup>	60 ± 52 <sup>Aa</sup>	0.01 ± 0.01 <sup>Ab</sup>
Nitisols	1937 ± 435 <sup>Aa</sup>	1635 ± 201 <sup>Aa</sup>	2286 ± 629 <sup>Aa</sup>	65 ± 60 <sup>Aa</sup>	77 ± 14 <sup>Aa</sup>	210 ± 81 <sup>Aa</sup>	0.03 ± 0.03 <sup>Ab</sup>	0.04 ± 0.02 <sup>Aa</sup>
Acrisols	2430 ± 370 <sup>Aa</sup>	2256 ± 307 <sup>Aa</sup>	1826 ± 133 <sup>Aa</sup>	84 ± 57 <sup>Ba</sup>	107 ± 202 <sup>Ba</sup>	100 ± 37 <sup>Ba</sup>	0.29 ± 0.23 <sup>Aa</sup>	0.00 ± 0.02 <sup>Ba</sup>
Cambisols	1939 ± 438 <sup>Aa</sup>	2403 ± 119 <sup>Aa</sup>	2399 ± 1040 <sup>Aa</sup>	41 ± 42 <sup>Aa</sup>	82 ± 33 <sup>Aa</sup>	117 ± 64 <sup>Aa</sup>	0.02 ± 0.00 <sup>Ab</sup>	0.02 ± 0.01 <sup>Aa</sup>
Ferralsols	1922 ± 398 <sup>Aa</sup>	1965 ± 285 <sup>Aa</sup>	2144 ± 485 <sup>Aa</sup>	36 ± 20 <sup>Aa</sup>	40 ± 0 <sup>Aa</sup>	162 ± 156 <sup>Aa</sup>	0.01 ± 0.01 <sup>Ab</sup>	0.03 ± 0.04 <sup>Aa</sup>
Sand	1927 ± 568 <sup>Aa</sup>	1887 ± 264 <sup>Aa</sup>	2208 ± 476 <sup>Aa</sup>	77 ± 27 <sup>Aa</sup>	98 ± 57 <sup>Ba</sup>	31 ± 36 <sup>Ba</sup>	0.15 ± 0.01 <sup>Ab</sup>	-0.01 ± 0.01 <sup>Aa</sup>

Values are mean ± standard deviation (n = 3); different lowercase letters indicate significant differences among soil types within the same treatment and different uppercase letters indicate significant differences among treatment within the same soil type (P < 0.05).




**Fig. 5.** Dynamics of (a)  $\text{CO}_2$  and (b)  $\text{N}_2\text{O}$  daily fluxes from different soil cores to which dung was added (left panel) compared to the same soils without excreta application (right panel) (Trial 1). The lower panels show the observed temporal dynamics of (c) mean daily soil moisture (0.05 m depth), (d) soil temperature (0.05 m depth) measured during gas sampling. Each flux value represents the mean of three chambers ( $\pm$ SE).

$\text{N}_2\text{O}$  emissions from excreta applied to grasslands, that the magnitude of  $\text{N}_2\text{O}$  emissions is negatively correlated with pH, i.e. that emissions increase at lower soil pHs. However, Khan et al., (2011) reported higher cumulative  $\text{N}_2\text{O}$  fluxes from bovine urine applied onto limed acid soils than on non-limed acid soils. The fact that all the soils except for the sand used in our study (pH 8.4, not included in the analysis) were slightly acidic and fairly similar in pH (pH range: 5.3–6.4) might explain why we found a positive correlation between urine EF and soil pH in both dry and wet season.

Contrary to our expectations, soil C and soil C/N ratio did not have any effect on urine  $\text{N}_2\text{O}$  EFs in our study, contrary to Pelster et al. (2012) who reported a potential C-limitation of soil  $\text{N}_2\text{O}$  emissions after N fertilizer application to temperate soils in Canada. However, because of the high C content in our soils ( $35 \text{ g kg}^{-1}$  compared to  $19 \text{ g kg}^{-1}$  in the previous study), soil  $\text{N}_2\text{O}$  production was likely N- and not C-limited. This would also explain why urine, but not dung or manure additions significantly stimulated  $\text{N}_2\text{O}$  emissions as urine N is quickly available as urease is ubiquitous in all soils (Van Groenigen et al., 2005b). Moreover, not only N but also significant amounts of water was added with urine applications, which increases soil moisture and soil microbial activity

(Marsden et al., 2016). It is noteworthy that in our study water-only additions did not stimulate  $\text{N}_2\text{O}$  emissions while additions of the same amounts of urine did (Figs. 7 and 8), indicating that although significant changes in soil moisture were observed after water application, the effect of the water was minor compared to the urine-N addition effect. As the soil moisture was generally below 60% water filled pore space (WFPS), it is likely that at least in the lighter textured soils nitrification was the dominant process of  $\text{N}_2\text{O}$  production after excreta application (Bell et al., 2015). Generally, high soil moisture promotes  $\text{N}_2\text{O}$  production, as shown in our study when comparing dry and wet season fluxes, but soil moisture can also be related to soil properties causing differences in  $\text{N}_2\text{O}$  fluxes between different soil types. For instance, De Klein et al. (2003) in a study in New Zealand grasslands reported an urine EF of 0.5% for a well-drained stony silt loam soil and 3.7% for a moderately-drained silt loam soil. However, Van der Weerden et al. (2011) reported that  $\text{N}_2\text{O}$  emissions did not always relate to drainage class, when summarizing experimental findings obtained for well- and poor-drained silt loam soils at three sites in New Zealand. Balaine et al. (2013) highlighted that the relative soil gas diffusivity is a key indicator to rank soil  $\text{N}_2\text{O}$  emission potentials.



**Fig. 6.** Dynamics of (a) CO<sub>2</sub>-C and (b) N<sub>2</sub>O-N daily fluxes from different soil cores to which manure (left panel) or dung (middle panel) was added compared to soils without excreta application (right panel) (Trial 2). The lower panels show temporal dynamics of (c) mean daily soil moisture (0.05 m depth) and (d) soil temperature (0.05 m depth) measured during gas sampling. Each flux value represents the mean of three chambers (±SE).


#### 4.2. Influence of season (wet vs dry) and time elapsed since excreta application on CO<sub>2</sub> and N<sub>2</sub>O emissions

The higher soil CO<sub>2</sub> fluxes after excreta application in the wet season compared to the dry season was likely caused by the higher soil moisture, since temperature was similar during all four trials, as both soil moisture and temperature are controlling factors for CO<sub>2</sub> production (Rochette and Gregorich, 1998; Ginting et al., 2003). Similar results had been observed by Zhu et al. (2018) after cattle dung application to an East African grassland.

A one year field observation in Canada showed that most of the emissions occur within a few weeks after excreta application, although smaller peaks may occur later (Rochette et al., 2014). In our study, the flux patterns after dung application were consistent with a previous study for tropical rangeland soils amended with dung by Zhu et al. (2018). They reported that N<sub>2</sub>O fluxes stayed elevated for around 14 days after dung application. The rather restricted period of elevated N<sub>2</sub>O emissions after dung application can likely be attributed to the relatively low mineral N content of the dung, fast crust formation and the wide dung C/N ratio (22–36, Table 2), which likely promoted N immobilization during C decomposition (Van der Weerden et al., 2011; Zhu et al., 2018).

In the present study, N<sub>2</sub>O fluxes returned to background levels within 39–73 observation days after urine application, which is consistent with Sordi et al. (2014) who reported  $41 \pm 10$  days of elevated emissions following cattle urine application to a subtropical pasture of Brazil during three different seasons, after which the N<sub>2</sub>O fluxes rapidly diminished and returned to background levels. This was also consistent with another study in New Zealand, which showed that soil NH<sub>4</sub><sup>+</sup>-N concentrations returned to background concentrations after 27 days, while elevated soil NO<sub>3</sub><sup>-</sup> concentrations were observed for around 40 days after cattle urine application (Clough et al., 2009). Similarly, a study carried out in the UK reported that the majority of N<sub>2</sub>O emissions occurred during the first 20 days after sheep urine application onto Eutric Cambisol mesocosms (Marsden et al., 2016). Thus, the frequent gas sampling and the length of the observation period in our study should have been sufficient to capture most of the N<sub>2</sub>O emissions caused by excreta application.

The higher N concentration and lower C/N ratio of the dung used during the dry seasons (C/N ratio: 22–29, Table 2) compared to the dung used during the rainy seasons (C/N ratio: 30–36, Table 2) should have translated into more substrate for denitrification and generally greater N<sub>2</sub>O production during the dry season; however, the dung N<sub>2</sub>O EFs were in fact lower during the dry seasons (range 0.00–0.04%) than during the



**Fig. 7.** Dynamics of (a) CO<sub>2</sub>-C and (b) N<sub>2</sub>O-N daily fluxes from different soil cores to which dung (left panel), urine (second-left panel), water (second-right panel), or nothing (right panel) was added (Trial 3). The lower panels show temporal dynamics of (c) mean daily soil moisture (0.05 m depth) and (d) soil temperature (0.05 m depth) measured during gas sampling. Each flux value represents the mean of three chambers ( $\pm$ SE).


wet seasons (0.02–0.20%). This was likely due to delayed dung crust formation due to rainfall (Mazzetto et al., 2014) and higher dung mineralization activity during the wet seasons with more favorable soil environmental conditions for N<sub>2</sub>O production (see soil moisture levels in Figs. 5–8), which is in agreement with our hypothesis. The magnitude of total rainfall may cause differences in the amount of anaerobic microsites in the soil/dung, which could provide suitable conditions for further reduction of the microbially-produced N<sub>2</sub>O to N<sub>2</sub> (Sordi et al., 2014; Zhu et al., 2018). This may explain why the dung N<sub>2</sub>O EFs during trial 3 (range: 0.02–0.15%), which had 596 mm rain, were lower than for the other wet season period (trial 2; range: 0.10–0.20%), during which only 83 mm of rainfall. Furthermore, this lower EF might also have been related to increased N leaching as a consequence of the heavy rainfalls, although this remains speculation as we did not measure N leaching.

The higher urine N<sub>2</sub>O EF in wet season than in dry season was similar to other studies in Brazil and Kenya (Sordi et al., 2014; Tully et al., 2017). Seasonal variations for urine N<sub>2</sub>O EFs was also measured under a temperate climate (Van Groenigen et al., 2005b), although the authors attributed this seasonality to the seasonality of background N<sub>2</sub>O emissions. Nevertheless, it should be noted that temperate climates show large intra-annual changes in temperature with less seasonality in

precipitation rates. In Kenya, air temperature only changes minimally during the year, whereas rainfalls have a distinct seasonality (i.e. a clear separation between dry and wet season). Soil moisture, which exerts significant effects on soil N<sub>2</sub>O emissions through modulating soil O<sub>2</sub> concentration and nutrient availability (Butterbach-Bahl et al., 2013), typically mirrors rainfall distribution. Therefore, our observation that excreta effects on soil N<sub>2</sub>O emissions were generally stronger in the wet season is not surprising, as soil moisture content was generally higher during the wet season soils favouring coupled nitrification denitrification processes, i.e. the main N<sub>2</sub>O production pathway after urine application (Monaghan and Barraclough, 1993). During the dry season, N mineralization and nitrification may be moisture limited, while denitrification would be unlikely as it is limited by the low soil moisture values (Linn and Doran, 1984).

#### 4.3. Influence of excreta type on CO<sub>2</sub> and N<sub>2</sub>O emissions

We confirmed our fourth hypothesis that both cattle urine and dung applications would stimulate CO<sub>2</sub> fluxes. The cumulative CO<sub>2</sub> emissions after dung application were similar to or slightly higher than after urine application in our study depending on the amount of C applied (Bertora et al., 2008), which is consistent with Lin et al. (2009) who also reported



**Fig. 8.** Dynamics of (a)  $\text{CO}_2\text{-C}$  and (b)  $\text{N}_2\text{O-N}$  daily fluxes from different soil cores to which water, urine or dung was added (Trial 4). The lower panels show the observed temporal dynamics of (c) mean daily soil moisture (0.05 m depth), (d) soil temperature (0.05 m depth) measured during gas sampling. Each flux value represents the mean of three chambers ( $\pm$ SE).

higher cumulative  $\text{CO}_2$  emissions from yak dung than from yak urine.

The  $\text{N}_2\text{O}$  EF for dung application (range: 0.00–0.20%, mean = 0.06%) was consistent with another study in Kenya that reported dung  $\text{N}_2\text{O}$  EF ranged from 0.00 to 0.04% (Tully et al., 2017). However, our mean EF was lower than the value of 0.28% synthesized for cattle dung patches through a global meta-analysis by Cai and Akiyama (2016). We suggest that the lower  $\text{N}_2\text{O}$  EF for cattle dung in Kenya in the present study compared to the global mean was caused by low quality feeds (i.e. low protein content) (Wassie et al., 2019) that resulted in lower N excretion and a higher C/N ratio in dung patches. In our study the dung and manure C/N ratios ranged from 22 to 36, much higher than the C/N ratios reported for dung/manure from cattle in temperate regions (<20, see Zhu et al., 2018). Moreover, the soils have low N concentrations (<0.39 g N  $\text{kg}^{-1}$  DM, Table 1), which could result in rapid immobilization of the applied N causing low  $\text{NO}_3^-$  availability for denitrification. Furthermore, higher solar radiation and high vapor pressure deficits in the Kenyan highlands leads to fast crust formation on the manure and dung and thus less incorporation of excreta N into the soil (Zhu et al., 2018).

The manure (i.e. a mixture of dung and urine) EF of 0.23% was also similar to a previous study in Kenya that reported a manure EF for the wet season of 0.15% (Tully et al., 2017). The higher manure EF in the present study compared to the dung EF might be explained by

synergistic effects of labile C from the dung and inorganic N from the urine in manure, which combined with higher water contents likely promoted  $\text{N}_2\text{O}$  formation in soils and the manure itself (Hyde et al., 2016).

The observed range of urine EFs in our study (0.01–1.36%, with a mean of 0.29%) were within the range of EFs determined by studies carried out in Brazil (0.19–0.33%) (Barneze et al., 2014; Sordi et al., 2014) and Kenya (0.05–0.21%) (Tully et al., 2017). Cattle urine EFs of more than 3% were reported for an incubation experiment conducted at 25 °C and 80% relative humidity in Brazil (Cardoso et al., 2017). However, this previous study might be misleading, as the high temperature and humidity conditions likely favored  $\text{N}_2\text{O}$  production while the incubation system used by Cardoso et al. (2017) did not allow for N leaching; a major N loss pathway of urine N. In their review, Cai and Akiyama (2016) estimated that at least 17.8% of applied urine N is leached under field conditions. Another recent study from Kenya that simulated a 20 mm rainfall event after application, reported a cattle urine EF of 1.2%, which can be ascribed to more favorable soil condition to  $\text{N}_2\text{O}$  production after rainfall (Pelster et al., 2016). As synthesized by Cai and Akiyama (2016) the average  $\text{N}_2\text{O}$  EF for cattle urine was 0.76%, i.e. approx. three times higher than in our study. Although the N application rate of 845 and 749  $\text{kg N ha}^{-1}$  applied as urine in our study was in the range of most studies analyzed in the global meta-analysis

(Cai and Akiyama, 2016), the soil properties and climate conditions in our study differed from most of the studies in the Cai and Akiyama review, which were predominantly data from temperate regions.

As the IPCC default value does not disaggregate the EF for urine and dung, we cannot compare the separate EFs for urine and dung in our study with the default value of 2%. However, the N partitioning between dung and urine for cattle excreta in SSA has been reported to be around 2:1 (Rufino et al., 2006), which is consistent with a recent study in Kenya by Wassie et al. (2019) who found that the N partitioning between dung and urine varied from ~1:1 to ~2:1. Those ratios differ significantly from estimates for western European countries (40:60) (Chadwick et al., 2018). If we apply the split as determined by Rufino et al. (2006) (i.e. an excreta-N ratio of dung to urine of 66:34), the overall N<sub>2</sub>O EF for cattle excreta on grassland ranges from 0.06 to 0.30% depending on the soil type, with an average of 0.14%. Thus, the EFs found here are approximately one order of magnitude lower than the IPCC default value or the EF determined for livestock excreta in the tropics and subtropics by Albanito et al. (2017) of 1.2%.

## 5. Conclusion

Here we showed that while soil type had no influence on CO<sub>2</sub> emissions after excreta application it did affect N<sub>2</sub>O emissions after urine, but not manure or dung, application. The differences between soils in N<sub>2</sub>O emissions following manure application were likely related to soil properties such as texture and soil pH. Environmental conditions (wet and dry season), and the N content of excreta did affect N<sub>2</sub>O losses, with a higher N<sub>2</sub>O EF (0.29%) from urine application versus application of cattle dung (EF = 0.06%). Based on the excreta N split estimated by existing literature in SSA, the overall mean N<sub>2</sub>O EF for cattle excreta deposited onto grassland was 0.14%, less than 10% of the IPCC default value (2%). Our findings suggest that current IPCC methodology provides a substantial overestimation of the N<sub>2</sub>O EF for cattle excreta on tropical rangeland. In order to improve estimates and to identify adequate, region specific mitigation options to reduce emissions from livestock excreta, environmental conditions should be taken into account when estimating N<sub>2</sub>O losses.

## Declaration of competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The authors would like to thank the farmers for granting access to their farms, George Nandhoka Wanyama, Jesse Owino, Phyllis Ndungu for helping with soil core collection, Rodgers Nyamosi for gas sampling and analysis, Daniel Korir and Alice Onyango for urine analysis, and other colleagues in the Mazingira Center. Yuhao Zhu acknowledges financial support from the China Scholarship Council (201506350087). Major funding was provided by the Research Program ATMO within the Earth and Environment Program of the German Helmholtz Association, by the German Federal Ministry for Economic Cooperation and development (BMZ), and the German Technical Cooperation (GIZ) under the project “In situ assessment of GHG emissions from two livestock systems in East Africa— determining current status and quantifying mitigation options” as well as under the International Fund for Agricultural Development (IFAD) funds project “Greening livestock: Incentive-based interventions for reducing the climate impact of livestock in East Africa”. Lutz Merbold, David Pelster, Sheila Abwanda, and Paul M. Mutuo further acknowledge the support received for CGIAR Fund Council, Australia (ACIAR), Irish Aid, the European Union, the Netherlands, New Zealand, Switzerland, UK, USAID and Thailand for funding to the CGIAR Research Program on Climate Change, Agriculture and Food Security

(CCAFS) as well as for the CGIAR Research Program on Livestock.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.soilbio.2019.107636>.

## References

- Davidson, E.A., Keller, M., Erickson, H.E., Verchot, L.V., Veldkamp, E., 2000. Testing a conceptual model of soil emissions of nitrous and nitric oxides. *BioScience* 50, 667–680. [https://doi.org/10.1641/0006-3568\(2000\)050\[0667:TACMOS\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2000)050[0667:TACMOS]2.0.CO;2).
- Albanito, F., Lebender, U., Cornulier, T., Sapkota, T.B., Bentrup, F., Stirling, C., Hillier, J., 2017. Direct nitrous oxide emissions from tropical and sub-tropical agricultural systems - A review and modelling of emission factors. *Scientific Reports* 7, Article number 44235.
- Ambus, P., Petersen, S.O., Soussana, J.F., 2007. Short-term carbon and nitrogen cycling in urine patches assessed by combined carbon-13 and nitrogen-15 labelling. *Agriculture, Ecosystems & Environment* 121, 84–92. <https://doi.org/10.1016/j.agee.2006.12.007>.
- Balaine, N., Clough, T.J., Beare, M.H., Thomas, S.M., Meenken, E.D., Ross, J.G., 2013. Changes in relative gas diffusivity explain soil nitrous oxide flux dynamics. *Soil Science Society of America Journal* 77, 1496–1505. <https://doi.org/10.2136/sssaj2013.04.0141>.
- Balaine, N., Clough, T.J., Beare, M.H., Thomas, S.M., Meenken, E.D., 2016. Soil gas diffusivity controls N<sub>2</sub>O and N<sub>2</sub> emissions and their ratio. *Soil Science Society of America Journal* 80, 529–540. <https://doi.org/10.2136/sssaj2015.09.0350>.
- Barneze, A.S., Mazzetto, A.M., Zani, C.F., Misselbrook, T., Cerri, C.C., 2014. Nitrous oxide emissions from soil due to urine deposition by grazing cattle in Brazil. *Atmospheric Environment* 92, 394–397. <https://doi.org/10.1016/j.atmosenv.2014.04.046>.
- Bell, M.J., Rees, R.M., Cloy, J.M., Topp, C.F.E., Bagnall, A., Chadwick, D.R., 2015. Nitrous oxide emissions from cattle excreta applied to a Scottish grassland: effects of soil and climatic conditions and a nitrification inhibitor. *The Science of the Total Environment* 508, 343–353. <https://doi.org/10.1016/j.scitotenv.2014.12.008>.
- Bertora, C., Alluvione, F., Zavattaro, L., van Groenigen, J.W., Velthof, G., Grignani, C., 2008. Pig slurry treatment modifies slurry composition, N<sub>2</sub>O, and CO<sub>2</sub> emissions after soil incorporation. *Soil Biology and Biochemistry* 40, 1999–2006. <https://doi.org/10.1016/j.soilbio.2008.03.021>.
- Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S., 2013. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society of London Series B Biological Sciences* 368, 20130122. <https://doi.org/10.1098/rstb.2013.0122>.
- Cai, Y., Akiyama, H., 2016. Nitrogen loss factors of nitrogen trace gas emissions and leaching from excreta patches in grassland ecosystems: a summary of available data. *The Science of the Total Environment* 572, 185–195. <https://doi.org/10.1016/j.scitotenv.2016.07.222>.
- Cai, Y., Chang, S.X., Cheng, Y., 2017. Greenhouse gas emissions from excreta patches of grazing animals and their mitigation strategies. *Earth-Science Reviews* 171, 44–57. <https://doi.org/10.1016/j.earscirev.2017.05.013>.
- Cardoso Abmael da, S., Quintana, B.G., Januskiewicz, E.R., Brito Liziane de, F., Morgado Elian da, S., Reis, R.A., Ruggieri, A.C., 2017. N<sub>2</sub>O emissions from urine-treated tropical soil: effects of soil moisture and compaction, urine composition, and dung addition. *Catena* 157, 325–332. <https://doi.org/10.1016/j.catena.2017.05.036>.
- Chadwick, D.R., Cardenas, L.M., Dhanoa, M.S., Donovan, N., Misselbrook, T., Williams, J.R., Thorman, R.E., McGeough, K.L., Watson, C.J., Bell, M., Anthony, S.G., Rees, R.M., 2018. The contribution of cattle urine and dung to nitrous oxide emissions: quantification of country specific emission factors and implications for national inventories. *The Science of the Total Environment* 635, 607–617. <https://doi.org/10.1016/j.scitotenv.2018.04.152>.
- Chantigny, M.H., Angers, D.A., Morvan, T., Pomar, C., 2004. Dynamics of pig slurry nitrogen in soil and plant as determined with <sup>15</sup>N. *Soil Science Society of America Journal* 68, 637–643. <https://doi.org/10.2136/sssaj2004.6370>.
- Clough, T.J., Ray, J.L., Buckthought, L.E., Calder, J., Baird, D., O'Callaghan, M., Sherlock, R.R., Condron, L.M., 2009. The mitigation potential of hippuric acid on N<sub>2</sub>O emissions from urine patches: an in situ determination of its effect. *Soil Biology and Biochemistry* 41, 2222–2229. <https://doi.org/10.1016/j.soilbio.2009.07.032>.
- Council, N.R., 2015. *Nutrient Requirements of Beef Cattle*, 8th Revised Edition. National Academies Press, Washington, D.C. <https://doi.org/10.17226/19014>.
- Davidson, E.A., 2009. The contribution of manure and fertilizer nitrogen to atmospheric nitrous oxide since 1860. *Nature Geoscience* 2, 659–662. <https://doi.org/10.1038/ngeo608>.
- De Klein, C.A.M.M., Barton, L., Sherlock, R.R., Li, Z., Littlejohn, R.P., 2003. Estimating a nitrous oxide emission factor for animal urine from some New Zealand pastoral soils. *Australian Journal of Soil Research* 41, 381–399. <https://doi.org/10.1071/SR02128>.
- Di, H.J., Cameron, K.C., 2007. Nitrate leaching losses and pasture yields as affected by different rates of animal urine nitrogen returns and application of a nitrification inhibitor - a lysimeter study. *Nutrient Cycling in Agroecosystems* 79, 281–290. <https://doi.org/10.1007/s10705-007-9115-5>.

- Ghezzehei, T.A., Sulman, B., Arnold, C.L., Bogie, N.A., Asefaw Berhe, A., 2019. On the role of soil water retention characteristic on aerobic microbial respiration. *Biogeosciences* 16, 1187–1209. <https://doi.org/10.5194/bg-16-1187-2019>.
- Ginting, D., Kessavalou, A., Eghball, B., Doran, J.W., 2003. Greenhouse gas emissions and soil indicators four years after manure and compost applications. *Journal of Environmental Quality* 32, 23–32. <https://doi.org/10.2134/jeq2003.2300>.
- Gu, J., Nicoulaud, B., Rochette, P., Gossel, A., Hénault, C., Cellier, P., Richard, G., 2013. A regional experiment suggests that soil texture is a major control of N<sub>2</sub>O emissions from tile-drained winter wheat fields during the fertilization period. *Soil Biology and Biochemistry* 60, 134–141. <https://doi.org/10.1016/j.soilbio.2013.01.029>.
- Haynes, R.J., Williams, P.H., 1993. Nutrient cycling and soil fertility in the grazed pasture ecosystem. *Advances in Agronomy* 49, 119–199. [https://doi.org/10.1016/S0065-2113\(08\)60794-4](https://doi.org/10.1016/S0065-2113(08)60794-4).
- Hendershot, W., Lalonde, H., Duquette, M., 2007. Ion exchange and exchangeable cations. In: Carter, M.R., Gregorich, E.G. (Eds.), *Soil Sampling and Methods of Analysis*, second ed. Canadian Society of Soil Science, CRC Press and Taylor & Francis Group, Oxford, UK. <https://doi.org/10.1201/9781420005271.ch18>.
- Hyde, B.P., Forrester, P.J., Jahangir, M.M.R., Ryan, M., Fanning, A.F., Carton, O.T., Lanigan, G.J., Richards, K.G., 2016. The interactive effects of fertilizer nitrogen with dung and urine on nitrous oxide emissions in grassland. *Irish Journal of Agricultural & Food Research* 0, 1–9. <https://doi.org/10.1515/ijaf-2016-0001>.
- IPCC, 2006. *IPCC Guide- Lines for National Greenhouse Gas Inventories*. IPCC Geneva, Switzerland.
- IPCC, 2014. *Summary for Policymakers. Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, pp. 1–33. <https://doi.org/10.1017/CBO9781107415324>.
- IUSS Working Group WRB, 2014. *World Reference Base for Soil Resources 2014, Update 2015. International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*. World Soil Resources Reports, vol 106. <https://doi.org/10.1017/S0014479706394902>. World Soil Resources Reports No. 106.
- Jaetzold, R., Schmidt, H., Hornetz, B., Shisanya, C., 2010. *Farm Management Handbook of Kenya: Subpart B1a, Southern Rift Valley Province, second ed.* Ministry of Agriculture, Kenya. in *Cooperation with the German Agency for Technical Cooperation (GTZ)*.
- Jarecki, M.K., Parkin, T.B., Chan, A.S.K., Hatfield, J.L., Jones, R., 2008. Greenhouse gas emissions from two soils receiving nitrogen fertilizer and swine manure slurry. *Journal of Environmental Quality* 37, 1432–1438. <https://doi.org/10.2134/jeq2007.0427>.
- Khan, S., Clough, T.J., Goh, K.M., Sherlock, R.R., 2011. Influence of soil pH on NO<sub>x</sub> and N<sub>2</sub>O emissions from bovine urine applied to soil columns. *New Zealand Journal of Agricultural Research* 54, 285–301. <https://doi.org/10.1080/00288233.2011.607831>.
- Kirk, P.L., 1950. Kjeldahl method for total nitrogen. *Analytical Chemistry* 22, 354–358. <https://doi.org/10.1021/ac60038a038>.
- Korir, D., Goopy, J.P., Gachuiri, C., Butterbach-Bahl, K., 2016. Supplementation with *Calliandra calothyrsus* improves nitrogen retention in cattle fed low-protein diets. *Animal Production Science* 56, 619–626. <https://doi.org/10.1071/AN15569>.
- Krol, D.J., Carolan, R., Minet, E., McGeough, K.L., Watson, C.J., Forrester, P.J., Lanigan, G.J., Richards, K.G., 2016. Improving and disaggregating N<sub>2</sub>O emission factors for ruminant excreta on temperate pasture soils. *The Science of the Total Environment* 568, 327–338. <https://doi.org/10.1016/j.scitotenv.2016.06.016>.
- Lessa, A.C.R., Madari, B.E., Paredes, D.S., Boddey, R.M., Urquiaga, S., Jantalia, C.P., Alves, B.J.R., 2014. Bovine urine and dung deposited on Brazilian savannah pastures contribute differently to direct and indirect soil nitrous oxide emissions. *Agriculture, Ecosystems & Environment* 190, 104–111. <https://doi.org/10.1016/j.agee.2014.01.010>.
- Lin, X., Wang, S., Ma, X., Xu, G., Luo, C., Li, Y., Jiang, G., Xie, Z., 2009. Fluxes of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O in an alpine meadow affected by yak excreta on the Qinghai-Tibetan plateau during summer grazing periods. *Soil Biology and Biochemistry* 41, 718–725. <https://doi.org/10.1016/j.soilbio.2009.01.007>.
- Linn, D.M., Doran, J.W., 1984. Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and nontilled soils 1. *Soil Science Society of America Journal* 48, 1267–1272. <https://doi.org/10.2136/sssaj1984.03615995004800060013x>.
- Marsden, K.A., Jones, D.L., Chadwick, D.R., 2016. The urine patch diffusional area: an important N<sub>2</sub>O source? *Soil Biology and Biochemistry* 92, 161–170. <https://doi.org/10.1016/j.soilbio.2015.10.011>.
- Mazzetto, A.M., Barneze, A.S., Feigl, B.J., Van Groeningen, J.W., Oenema, O., Cerri, C.C., 2014. Temperature and moisture affect methane and nitrous oxide emission from bovine manure patches in tropical conditions. *Soil Biology & Biochemistry* 76, 242–248.
- Monaghan, R.M., Barraclough, D., 1993. Nitrous oxide and dinitrogen emissions from urine-affected soil under controlled conditions. *Plant and Soil* 151, 127–138. <https://doi.org/10.1007/BF00010793>.
- Neira, J., Ortiz, M., Morales, L., Acevedo, E., 2015. Oxygen diffusion in soils: understanding the factors and processes needed for modeling. *Chilean Journal of Agricultural Research* 75, 35–44. <https://doi.org/10.4067/s0718-58392015000300005>.
- Norman, J.M., Kucharik, C.J., Gower, S.T., Baldocchi, D.D., Crill, P.M., Rayment, M., Savage, K., Striegl, R.G., 1997. A comparison of six methods for measuring soil-surface carbon dioxide fluxes. *Journal of Geophysical Research: Atmosphere* 102, 28771–28777. <https://doi.org/10.1029/97JD01440>.
- Oenema, O., Wrage, N., Velthof, G.L., Van Groeningen, J.W., Dolfin, J., Kuikman, P.J., 2005. Trends in global nitrous oxide emissions from animal production systems. *Nutrient Cycling in Agroecosystems* 72, 51–65. <https://doi.org/10.1007/s10705-004-7354-2>.
- Ogle, S.M., Olander, L., Wollenberg, L., Rosenstock, T., Tubiello, F., Paustian, K., Buendia, L., Nihart, A., Smith, P., 2014. Reducing greenhouse gas emissions and adapting agricultural management for climate change in developing countries: providing the basis for action. *Global Change Biology* 20, 1–6. <https://doi.org/10.1111/gcb.12361>.
- Onyango, A.A., Dickhoefer, U., Rufino, M.C., Butterbach-Bahl, K., Goopy, J.P., 2019. Temporal and spatial variability in the nutritive value of pasture vegetation and supplement feedstuffs for domestic ruminants in Western Kenya. *Asian-Australasian Journal of Animal Sciences* 32, 637–647. <https://doi.org/10.5713/ajas.18.0114>.
- Parkin, T.B., Venterea, R.T., Hargreaves, S.K., 2012. Calculating the detection limits of chamber-based soil greenhouse gas flux measurements. *Journal of Environmental Quality* 41, 705–715. <https://doi.org/10.2134/jeq2011.0394>.
- Pelster, D.E., Chantigny, M.H., Rochette, P., Angers, D.A., Rieux, C., Vanasse, A., 2012. Nitrous Oxide emissions respond differently to mineral and organic nitrogen sources in contrasting soil types. *Journal of Environmental Quality* 41, 427–435. <https://doi.org/10.2134/jeq2011.0261>.
- Pelster, D.E., Gisore, B., Goopy, J., Korir, D., Koske, J.K., Rufino, M.C., Butterbach-Bahl, K., 2016. Methane and nitrous oxide emissions from cattle excreta on an East African grassland. *Journal of Environmental Quality* 45, 1531–1539. <https://doi.org/10.2134/jeq2016.02.0050>.
- Reeuwijk, L., 2002. *Procedures for Soil Analysis. Technical Paper. International Soil Reference and Information Centre (ISIRC), Wageningen, the Netherlands*.
- Rochette, P., Gregorich, E.G., 1998. Dynamics of soil microbial biomass C, soluble organic C and CO<sub>2</sub> evolution after three years of manure application. *Canadian Journal of Soil Science* 78, 283–290. <https://doi.org/10.4141/S97-066>.
- Rochette, P., Chantigny, M.H., Ziadi, N., Angers, D.A., Bélanger, G., Charbonneau, É., Pellerin, D., Liang, C., Bertrand, N., 2014. Soil nitrous oxide emissions after deposition of dairy cow excreta in eastern Canada. *Journal of Environmental Quality* 43, 829–841. <https://doi.org/10.2134/jeq2013.11.0474>.
- Rufino, M.C., Rowe, E.C., Delve, R.J., Giller, K.E., 2006. Nitrogen cycling efficiencies through resource-poor African crop-livestock systems. *Agriculture, Ecosystems & Environment* 112, 261–282. <https://doi.org/10.1016/j.agee.2005.08.028>.
- Saarjärvi, K., Mattila, P.K., Virkajärvi, P., 2006. Ammonia volatilization from artificial dung and urine patches measured by the equilibrium concentration technique (JTI method). *Atmospheric Environment* 40, 5137–5145. <https://doi.org/10.1016/j.atmosenv.2006.03.052>.
- Sagar, S., Jha, N., Deslippe, J., Bolan, N.S., Luo, J., Giltrap, D.L., Kim, D.G., Zaman, M., Tillman, R.W., 2013. Denitrification and N<sub>2</sub>O:N<sub>2</sub> production in temperate grasslands: processes, measurements, modelling and mitigating negative impacts. *The Science of the Total Environment* 465, 173–195. <https://doi.org/10.1016/j.scitotenv.2012.11.050>.
- Saiz, G., Albrecht, A., 2016. Methods for smallholder quantification of soil carbon stocks and stock changes. In: Rosenstock, T., Rufino, M., Butterbach-Bahl, K., Wollenberg, L., Richards, M. (Eds.), *Methods for Measuring Greenhouse Gas Balances and Evaluating Mitigation Options in Smallholder Agriculture*. Springer International Publishing, Cham, pp. 135–162. [https://doi.org/10.1007/978-3-319-29794-1\\_7](https://doi.org/10.1007/978-3-319-29794-1_7).
- Samad, M.S., Bakken, L.R., Nadeem, S., Clough, T.J., de Klein, C.A.M., Richards, K.G., Lanigan, G.J., Morales, S.E., 2016. High-resolution denitrification kinetics in pasture soils link N<sub>2</sub>O Emissions to pH, and denitrification to C mineralization. *PLOS ONE* 11, e0151713. <https://doi.org/10.1371/journal.pone.0151713>.
- Scrimgeour, C., 2008. In: second ed. Carter, M.R., Gregorich, E.G. (Eds.), *Soil Sampling and Methods of Analysis*. CRC Press (2008), Boca Raton, FL, USA, p. 1224. <https://doi.org/10.1017/S0014479708006546>. £85.00. ISBN-13: 978-0-8593-3586-0. *Experimental Agriculture* 44, 437–437.
- Selbie, D.R., Buckthought, L.E., Shepherd, M.A., 2015. The challenge of the urine patch for managing nitrogen in grazed pasture systems. *Advances in Agronomy* 129, 229–292. <https://doi.org/10.1016/b.s.agron.2014.09.004>.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., S. O., 2007. *Agriculture. In: Climate change 2007: mitigation*. In: Metz, B., Davidson, O.R., Bosch, P.R., Dave, R., M, L.A. (Eds.), *Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 498–540.
- Sordi, A., Dieckow, J., Bayer, C., Albuquerque, M.A., Piva, J.T., Zanatta, J.A., Tomazi, M., da Rosa, C.M., de Moraes, A., 2014. Nitrous oxide emission factors for urine and dung patches in a subtropical Brazilian pastureland. *Agriculture, Ecosystems & Environment* 190, 94–103. <https://doi.org/10.1016/j.agee.2013.09.004>.
- Taghizadeh-Toosi, A., Clough, T.J., Condon, L.M., Sherlock, R.R., Anderson, C.R., Craigie, R.A., 2011. Biochar incorporation into pasture soil suppresses in situ nitrous oxide emissions from ruminant urine patches. *Journal of Environmental Quality* 40, 468–476. <https://doi.org/10.2134/jeq2010.0419>.
- Thornton, P.K., Herrero, M., 2014. Climate change adaptation in mixed crop-livestock systems in developing countries. *Global Food Security* 3, 99–107. <https://doi.org/10.1016/j.gfs.2014.02.002>.
- Tully, K.L., Abwanda, S., Thiong'o, M., Mutuo, P.M., Rosenstock, T.S., 2017. Nitrous oxide and methane fluxes from urine and dung deposited on Kenyan pastures. *Journal of Environmental Quality* 46, 921–929. <https://doi.org/10.2134/jeq2017.01.0040>.
- Valk, H., 1994. Effects of partial replacement of herbage by maize silage on N utilization and milk production of dairy cows. *Livestock Production Science* 40, 241–250. [https://doi.org/10.1016/0301-6226\(94\)90092-2](https://doi.org/10.1016/0301-6226(94)90092-2).

- Van der Weerden, T.J., Luo, J., de Klein, C.A.M., Hoogendoorn, C.J., Littlejohn, R.P., Rys, G.J., 2011. Disaggregating nitrous oxide emission factors for ruminant urine and dung deposited onto pastoral soils. *Agriculture, Ecosystems & Environment* 141, 426–436. <https://doi.org/10.1016/j.agee.2011.04.007>.
- Van Groenigen, J.W., Kuikman, P.J., De Groot, W.J.M., Velthof, G.L., 2005. Nitrous oxide emission from urine-treated soil as influenced by urine composition and soil physical conditions. *Soil Biology and Biochemistry* 37, 463–473. <https://doi.org/10.1016/j.soilbio.2004.08.009>.
- Van Groenigen, J.W., Velthof, G.L., Van Der Bolt, F.J.E., Vos, A., Kuikman, P.J., 2005. Seasonal variation in N<sub>2</sub>O emissions from urine patches: effects of urine concentration, soil compaction and dung. *Plant and Soil* 273, 15–27. <https://doi.org/10.1007/s11104-004-6261-2>.
- Venterea, R.T., Clough, T.J., Coulter, J.A., Breuillin-Sessoms, F., 2015. Ammonium sorption and ammonia inhibition of nitrite-oxidizing bacteria explain contrasting soil N<sub>2</sub>O production. *Scientific Reports* 5, 1–15. <https://doi.org/10.1038/srep12153>.
- Wang, J., Chadwick, D.R., Cheng, Y., Yan, X., 2018. Global analysis of agricultural soil denitrification in response to fertilizer nitrogen. *The Science of the Total Environment* 616–617, 908–917. <https://doi.org/10.1016/j.scitotenv.2017.10.229>.
- Wang, Y., Guo, J., Vogt, R.D., Mulder, J., Wang, J., Zhang, X., 2018. Soil pH as the chief modifier for regional nitrous oxide emissions: new evidence and implications for global estimates and mitigation. *Global Change Biology* 24, e617–e626. <https://doi.org/10.1111/gcb.13966>.
- Wassie, S.E., Ali, A.I.M., Korir, D., Butterbach-Bahl, K., Goopy, J., Merbold, L., Schlecht, E., Dickhoefer, U., 2019. Effects of feed intake level on efficiency of microbial protein synthesis and nitrogen balance in Boran steers consuming tropical poor-quality forage. *Archives of Animal Nutrition* 73, 140–157. <https://doi.org/10.1080/1745039X.2019.1572343>.
- Webb, J., Misselbrook, T.H., 2004. A mass-flow model of ammonia emissions from UK livestock production. *Atmospheric Environment* 38, 2163–2176. <https://doi.org/10.1016/j.atmosenv.2004.01.023>.
- Weitz, A.M., Linder, E., Frolking, S., Crill, P.M., Keller, M., 2001. N<sub>2</sub>O emissions from humid tropical agricultural soils: effects of soil moisture, texture and nitrogen availability. *Soil Biology and Biochemistry* 33, 1077–1093. [https://doi.org/10.1016/S0038-0717\(01\)00013-X](https://doi.org/10.1016/S0038-0717(01)00013-X).
- Wolf, B., Zheng, X., Brüggemann, N., Chen, W., Dannenmann, M., Han, X., Sutton, M.A., Wu, H., Yao, Z., Butterbach-Bahl, K., 2010. Grazing-induced reduction of natural nitrous oxide release from continental steppe. *Nature* 464, 881–884. <https://doi.org/10.1038/nature08931>.
- Yao, Z., Wei, Y., Liu, C., Zheng, X., Xie, B., 2015. Organically fertilized tea plantation stimulates N<sub>2</sub>O emissions and lowers NO fluxes in subtropical China. *Biogeosciences* 12, 5915–5928. <https://doi.org/10.5194/bg-12-5915-2015>.
- Zheng, X., Han, S., Huang, Y., Wang, Y., Wang, M., 2004. Re-quantifying the emission factors based on field measurements and estimating the direct N<sub>2</sub>O emission from Chinese croplands. *Global Biogeochemical Cycles* 18, 1–19. <https://doi.org/10.1029/2003GB002167>.
- Zhou, M., Zhu, B., Wang, S., Zhu, X., Vereecken, H., Brüggemann, N., 2017. Stimulation of N<sub>2</sub>O emission by manure application to agricultural soils may largely offset carbon benefits: a global meta-analysis. *Global Change Biology* 23, 4068–4083. <https://doi.org/10.1111/gcb.13648>.
- Zhu, Y., Merbold, L., Pelster, D., Diaz-Pines, E., Wanyama, G.N., Butterbach-Bahl, K., 2018. Effect of dung quantity and quality on greenhouse gas fluxes from tropical pastures in Kenya. *Global Biogeochemical Cycles* 32, 1589–1604. <https://doi.org/10.1029/2018GB005949>.