

Sharp constant in the curl inequality and ground states for curl-curl problem with critical exponent

Jarosław Mederski and Andrzej Szulkin

CRC Preprint 2020/3, January 2020

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

SHARP CONSTANT IN THE CURL INEQUALITY AND GROUND STATES FOR CURL-CURL PROBLEM WITH CRITICAL EXPONENT

JAROSŁAW MEDERSKI AND ANDRZEJ SZULKIN

ABSTRACT. Let $\Omega \subset \mathbb{R}^3$ be a Lipschitz domain and let $S_{\text{curl}}(\Omega)$ be the largest constant such that

$$\int_{\mathbb{R}^3} |\nabla \times u|^2 dx \geq S_{\text{curl}}(\Omega) \inf_{\substack{w \in W_0^6(\text{curl}; \mathbb{R}^3) \\ \nabla \times w = 0}} \left(\int_{\mathbb{R}^3} |u + w|^6 dx \right)^{\frac{1}{3}}$$

for any u in $W_0^6(\text{curl}; \Omega) \subset W_0^6(\text{curl}; \mathbb{R}^3)$ where $W_0^6(\text{curl}; \Omega)$ is the closure of $C_0^\infty(\Omega, \mathbb{R}^3)$ in $\{u \in L^6(\Omega, \mathbb{R}^3) : \nabla \times u \in L^2(\Omega, \mathbb{R}^3)\}$ with respect to the norm $(|u|_6^2 + |\nabla \times u|_2^2)^{1/2}$. We show that $S_{\text{curl}}(\Omega)$ is strictly larger than the classical Sobolev constant S in \mathbb{R}^3 . Moreover, $S_{\text{curl}}(\Omega)$ is independent of Ω and is attained by a ground state solution to the curl-curl problem

$$\nabla \times (\nabla \times u) = |u|^4 u$$

if $\Omega = \mathbb{R}^3$. With the aid of those results, we also investigate ground states of the Brezis-Nirenberg-type problem for the curl-curl operator in a bounded domain Ω

$$\nabla \times (\nabla \times u) + \lambda u = |u|^4 u \quad \text{in } \Omega$$

with the so-called metallic boundary condition $\nu \times u = 0$ on $\partial\Omega$, where ν is the exterior normal to $\partial\Omega$.

1. INTRODUCTION

Sharp Sobolev-type inequalities have been widely studied by a large number of authors and the best Sobolev constants play an important role e.g. in the theory of partial differential equations, differential geometry, isoperimetric inequalities as well as in mathematical physics, see e.g. [4, 20, 33]. In particular, if Ω is a domain in \mathbb{R}^3 , then the best constant S in the Sobolev inequality

$$(1.1) \quad \int_{\Omega} |\nabla u|^2 dx \geq S \left(\int_{\Omega} |u|^6 dx \right)^{\frac{1}{3}} \quad \text{for } u \in \mathcal{D}^{1,2}(\Omega)$$

has been computed explicitly by Talenti [33] and as is well-known, it is achieved (i.e., equality holds) if and only if $\Omega = \mathbb{R}^3$ and u is the Aubin-Talenti instanton $U_{\varepsilon, y}(x) := 3^{1/4}(\varepsilon^2 + |x - y|^2)^{-1/2}$, see [4, 33]. When $\varepsilon = 1$, this is the unique (up to translations in \mathbb{R}^3) positive solution to the equation $-\Delta u = |u|^4 u$ in $\mathcal{D}^{1,2}(\mathbb{R}^3)$ and a ground state, i.e. a minimizer for the energy functional among all nontrivial solutions.

The aim of this work is to perform a similar analysis for the curl operator $\nabla \times (\cdot)$. This is challenging from the mathematical point of view and important in mathematical physics; such operator appears e.g. in Maxwell equations as well as in Navier-Stokes problems [13,

2010 *Mathematics Subject Classification.* Primary: 35Q60; Secondary: 35Q61, 35J20.

Key words and phrases. Sharp constant, Sobolev inequality, time-harmonic Maxwell equations, ground state, variational methods, strongly indefinite functional, curl-curl problem.

[17, 26]. Finding a formulation in the spirit of (1.1), but involving the curl operator, is not straightforward and there are several essential difficulties as we shall see later.

For instance, the kernel of $\nabla \times (\cdot)$ is of infinite dimension since $\nabla \times (\nabla \varphi) = 0$ for all $\varphi \in \mathcal{C}^2(\Omega)$. Hence the inequality (1.1) with ∇u replaced by $\nabla \times u$ would hold for all $u \in \mathcal{C}_0^\infty(\mathbb{R}^3, \mathbb{R}^3)$ only if $S = 0$. This makes it necessary to introduce a Sobolev-like constant in a different way which we now proceed to do.

Let Ω be a Lipschitz domain in \mathbb{R}^3 and for $2 \leq p \leq 6$, let

$$W^p(\text{curl}; \Omega) := \{u \in L^p(\Omega, \mathbb{R}^3) : \nabla \times u \in L^2(\Omega, \mathbb{R}^3)\}.$$

This is a Banach space if provided with the norm

$$\|u\|_{W^p(\text{curl}; \Omega)} := (|u|_p^2 + |\nabla \times u|_2^2)^{1/2}.$$

Here and in the sequel $|\cdot|_q$ denotes the L^q -norm for $q \in [1, \infty]$. We also define

$$(1.2) \quad W_0^p(\text{curl}; \Omega) := \text{closure of } \mathcal{C}_0^\infty(\Omega, \mathbb{R}^3) \text{ in } W^p(\text{curl}; \Omega).$$

If $\Omega = \mathbb{R}^3$, these two spaces coincide, see Lemma 2.1. Although results of this kind are well known, we provide a proof for the reader's convenience. The spaces $W^2(\text{curl}; \Omega)$ and $W_0^2(\text{curl}; \Omega)$ are studied in detail in [13, 18, 26]. Extending $u \in W_0^p(\text{curl}; \Omega)$ by 0 outside Ω we may assume $W_0^p(\text{curl}; \Omega) \subset W_0^p(\text{curl}; \mathbb{R}^3)$. Denote the kernel of $\nabla \times (\cdot)$ in $W_0^6(\text{curl}; \mathbb{R}^3)$ by

$$\mathcal{W} := \{w \in W_0^6(\text{curl}; \mathbb{R}^3) : \nabla \times w = 0\}.$$

Let $S_{\text{curl}}(\Omega)$ be the largest possible constant such that the inequality

$$(1.3) \quad \int_{\mathbb{R}^3} |\nabla \times u|^2 dx \geq S_{\text{curl}}(\Omega) \inf_{w \in \mathcal{W}} \left(\int_{\mathbb{R}^3} |u + w|^6 dx \right)^{\frac{1}{3}}$$

holds for any $u \in W_0^6(\text{curl}; \Omega)$. Note that here u but not necessarily w is supported in Ω . It is not a priori clear that $S_{\text{curl}}(\Omega)$ is positive or that it is independent of Ω . That this is the case follows from Theorems 1.1 and 1.2(a) below:

Theorem 1.1. $S_{\text{curl}}(\Omega) = S_{\text{curl}}$ where $S_{\text{curl}} := S_{\text{curl}}(\mathbb{R}^3)$.

In the next result we show that S_{curl} is attained provided $\Omega = \mathbb{R}^3$ and the optimal function is (up to rescaling) a ground state solution to the curl-curl problem with critical exponent. Existence of a ground state in this case has been an open question for some time. Let

$$(1.4) \quad J(u) := \frac{1}{2} \int_{\mathbb{R}^3} |\nabla \times u|^2 dx - \frac{1}{6} \int_{\mathbb{R}^3} |u|^6 dx$$

and introduce the following constraint:

$$(1.5) \quad \mathcal{N} := \left\{ u \in W_0^6(\text{curl}; \mathbb{R}^3) \setminus \mathcal{W} : \int_{\mathbb{R}^3} |\nabla \times u|^2 = \int_{\mathbb{R}^3} |u|^6 dx \text{ and } \text{div}(|u|^4 u) = 0 \right\}.$$

As we shall see later, this set is a variant of a generalization of the Nehari manifold [27] which may be found in [28] for a Schrödinger equation.

Theorem 1.2. (a) $S_{\text{curl}} > S$.

(b) $\inf_{\mathcal{N}} J = \frac{1}{3} S_{\text{curl}}^{3/2}$ and is attained. Moreover, if $u \in \mathcal{N}$ and $J(u) = \inf_{\mathcal{N}} J$, then u is a ground state solution to the equation

$$(1.6) \quad \nabla \times (\nabla \times u) = |u|^4 u \quad \text{in } \mathbb{R}^3$$

and equality holds in (1.3) for this u . If u satisfies equality in (1.3), then there are unique $t > 0$ and $w \in \mathcal{W}$ such that $t(u + w) \in \mathcal{N}$ and $J(t(u + w)) = \inf_{\mathcal{N}} J$.

A natural question arises whether ground states must have some symmetry properties. It follows from Theorem 1.1 in [5] that any $\mathcal{O}(3)$ -equivariant (weak) solution to (1.6) is trivial, hence a ground state cannot be radially symmetric.

The curl-curl problem $\nabla \times (\nabla \times u) = f(x, u)$ in a bounded domain or in \mathbb{R}^3 has been recently studied e.g. in [5–8, 22, 24] under different hypotheses on f but always assuming f is subcritical, i.e. $f(x, u)/|u|^5 \rightarrow 0$ as $|u| \rightarrow \infty$. However, the occurrence of ground states to (1.6) (i.e., in the critical exponent case) has been an open problem as we have already mentioned. In view of the existence of Aubin-Talenti instantons, this is a very natural question. While the instantons are given explicitly, we have no such explicit formula for ground states in the curl-curl case. Since the instantons are radially symmetric up to translations, one can find them by ODE methods. In view of the above remark concerning $\mathcal{O}(3)$ -equivariant solutions, such methods do not seem available for the curl-curl problem and a different approach is needed. Note further that there is no maximum principle for the curl-curl operator and, to our knowledge, no unique continuation principle applicable to our case. An approach different than for (1.1) is also required for the proof of Ω -independence of S_{curl} , see Section 5. Moreover concentration-compactness analysis for the curl operator is considerably different from that in [16, 21, 36] – see our approach in Section 3.

We would like to emphasize an important role of the analysis of nonlinear curl-curl problems from the physical point of view. Solutions u to nonlinear curl-curl equations describe the profiles of time-harmonic solutions $E(x, t) = u(x) \cos(\omega t)$ to the time-dependent nonlinear electromagnetic wave equation, which together with material constitutive laws and Maxwell equations, describes the *exact* propagation of electromagnetic waves in a nonlinear medium [1, 6, 31]. Since finding propagation exactly may be very difficult, there are several simplifications in the literature which rely on approximations of the nonlinear electromagnetic wave equation. The most prominent one is the scalar or vector nonlinear Schrödinger equation. For instance, one assumes that the term $\nabla(\text{div}(u))$ in $\nabla \times (\nabla \times u) = \nabla(\text{div}(u)) - \Delta u$ is negligible and can be dropped, or one uses the so-called *slowly varying envelope approximation*. However, such simplifications may produce *non-physical* solutions; see [2, 11] and the references therein.

We also point out that the term $|u|^4 u$ in (1.6) as well as in (1.7) below allows to consider the so-called quintic effect in nonlinear optics modelled by Maxwell equations. See for instance [1, 6, 14, 15, 23, 25, 31] and the references therein. We hope that our results will prompt further analytical studies of physical phenomena involving the quintic nonlinearity, e.g. the well-known cubic-quintic effect in nonlinear optics [14, 25].

Using our concentration-compactness result we are also able to treat the Brezis-Nirenberg problem [10] for the curl-curl operator

$$(1.7) \quad \nabla \times (\nabla \times u) + \lambda u = |u|^4 u \quad \text{in } \Omega$$

together with the so-called metallic boundary condition

$$(1.8) \quad \nu \times u = 0 \quad \text{on } \partial\Omega.$$

Here $\nu : \partial\Omega \rightarrow \mathbb{R}^3$ is the exterior normal and $\Omega \subset \mathbb{R}^3$ is a bounded domain. This boundary condition is natural in the theory of Maxwell equations and it holds when Ω is surrounded

by a perfect conductor. If the boundary of Ω is not of class \mathcal{C}^1 , then we assume (1.8) is satisfied in a generalized sense by which we mean u is in the space $W_0^6(\text{curl}; \Omega)$ defined in (1.2). Weak solutions to (1.7)–(1.8) correspond to critical points of the associated energy functional $J_\lambda : W_0^6(\text{curl}; \Omega) \rightarrow \mathbb{R}$ given by

$$(1.9) \quad J_\lambda(u) := \frac{1}{2} \int_\Omega |\nabla \times u|^2 dx + \frac{\lambda}{2} \int_\Omega |u|^2 dx - \frac{1}{6} \int_\Omega |u|^6 dx.$$

Recall from [7, 23] that the spectrum of the curl-curl operator in $H_0(\text{curl}; \Omega) := W_0^2(\text{curl}; \Omega)$ consists of the eigenvalue $\lambda_0 = 0$ with infinite multiplicity and of a sequence of eigenvalues

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \rightarrow \infty$$

with corresponding finite multiplicities $m(\lambda_k) \in \mathbb{N}$. Let \mathcal{N}_λ be the generalized Nehari manifold for J_λ (see (6.1) for the definition), and for $\lambda \leq 0$ let

$$c_\lambda := \inf_{\mathcal{N}_\lambda} J_\lambda.$$

Denote the Lebesgue measure of Ω by $|\Omega|$. We introduce the following condition:

(Ω) Ω is a bounded domain, either convex or with $\mathcal{C}^{1,1}$ -boundary.

The reason for this assumption will be explained in the next section.

In domains $\Omega \neq \mathbb{R}^3$ we also introduce another constant, $\bar{S}_{\text{curl}}(\Omega)$, such that the inequality

$$(1.10) \quad \int_\Omega |\nabla \times u|^2 dx \geq \bar{S}_{\text{curl}}(\Omega) \inf_{w \in \mathcal{W}_\Omega} \left(\int_\Omega |u + w|^6 dx \right)^{\frac{1}{3}}$$

holds for any $u \in W_0^6(\text{curl}; \Omega)$, where $\mathcal{W}_\Omega := \{w \in W_0^6(\text{curl}; \Omega) : \nabla \times w = 0\}$, and $\bar{S}_{\text{curl}}(\Omega)$ is largest with this property. Although $\bar{S}_{\text{curl}}(\Omega)$ seems to be more natural than $S_{\text{curl}}(\Omega)$, we do not know whether it equals S_{curl} . We are only able to prove the following result:

Theorem 1.3. *Let Ω be a Lipschitz domain in \mathbb{R}^3 , possibly unbounded, $\Omega \neq \mathbb{R}^3$. Then $S_{\text{curl}} \geq \bar{S}_{\text{curl}}(\Omega)$. If Ω satisfies (Ω), then $\bar{S}_{\text{curl}}(\Omega) \geq S$.*

Finally, the main result concerning the Brezis-Nirenberg problem for the curl-curl operator (1.7) reads as follows.

Theorem 1.4. *Suppose Ω satisfies (Ω). Let $\lambda \in (-\lambda_\nu, -\lambda_{\nu-1}]$ for some $\nu \geq 1$. Then $c_\lambda > 0$ and the following statements hold:*

(a) *If $c_\lambda < c_0$, then there is a ground state solution to (1.7)–(1.8), i.e. c_λ is attained by a critical point of J_λ . A sufficient condition for this inequality to hold is $\lambda \in (-\lambda_\nu, -\lambda_\nu + \bar{S}_{\text{curl}}(\Omega)|\Omega|^{-2/3})$.*

(b) *There exists $\varepsilon_\nu \geq \bar{S}_{\text{curl}}(\Omega)|\Omega|^{-2/3}$ such that c_λ is not attained for $\lambda \in (-\lambda_\nu + \varepsilon_\nu, -\lambda_{\nu-1}]$, and $c_\lambda = c_0$ for $\lambda \in [-\lambda_\nu + \varepsilon_\nu, -\lambda_{\nu-1}]$. We do not exclude that $\varepsilon_\nu > \lambda_\nu - \lambda_{\nu-1}$, so these intervals may be empty.*

(c) *$c_\lambda \rightarrow 0$ as $\lambda \rightarrow -\lambda_\nu^-$, and the function*

$$(-\lambda_\nu, -\lambda_\nu + \varepsilon_\nu] \cap (-\lambda_\nu, -\lambda_{\nu-1}] \ni \lambda \mapsto c_\lambda \in (0, +\infty)$$

is continuous and strictly increasing.

(d) *There are at least $\#\{k : -\lambda_k < \lambda < -\lambda_k + \frac{1}{3}\bar{S}_{\text{curl}}(\Omega)|\Omega|^{-\frac{2}{3}}\}$ pairs of solutions $\pm u$ to (1.7)–(1.8).*

Note that if λ is as in (a), then the relation $-\lambda_k < \lambda < -\lambda_k + \frac{1}{3}\overline{S}_{\text{curl}}(\Omega)|\Omega|^{-\frac{2}{3}}$ holds for $k = \nu, \dots, \nu + m - 1$ where m is the multiplicity of λ_ν but it may also hold for some k with $\lambda_k > \lambda_\nu$.

The above result is known for cylindrically symmetric domains where it is possible to reduce the curl-curl operator to a positive definite one, see [23]. However, the solution obtained there is a ground state in a subspace of functions having cylindric symmetry and we do not know whether it is a ground state in the full space.

Let us recall from earlier work that the main difficulties when treating J and J_λ , also in the subcritical case, are that these functionals are strongly indefinite, i.e., they are unbounded from above and from below, even on subspaces of finite codimension. Moreover, the quadratic part of J has infinite-dimensional kernel and J', J'_λ are not (sequentially) weak-to-weak* continuous, i.e. $u_n \rightharpoonup u$ does not imply that $J'_\lambda(u_n)\varphi \rightarrow J'_\lambda(u)\varphi$ for all $\varphi \in \mathcal{C}_0^\infty(\Omega, \mathbb{R}^3)$. This lack of continuity is caused by the fact that $W_0^p(\text{curl}; \Omega)$ is not (locally) compactly embedded in any Lebesgue space and we do not know whether necessarily $u_n \rightarrow u$ a.e. in Ω . A consequence of this is that for a Palais-Smale sequence $u_n \rightharpoonup u$ it is not clear whether u is a critical point. In the subcritical case one can overcome these difficulties since either a variant of the Palais-Smale condition is satisfied or some compactness can be recovered on a suitable topological manifold, see e.g. [6, 22, 24]. In the critical case however, there are additional difficulties. In Section 3 we introduce a general concentration-compactness analysis for this case. We show that the topological manifold

$$\left\{ u \in W_0^6(\text{curl}; \mathbb{R}^3) : \text{div}(|u|^4 u) = 0 \right\}$$

is locally compactly embedded in $L^p(\mathbb{R}^3, \mathbb{R}^3)$ for $1 \leq p < 6$ and that if a sequence (u_n) is contained in this manifold and $u_n \rightharpoonup u$, then $u_n \rightarrow u$ a.e. after passing to a subsequence. This result will play a crucial role because it implies that if such (u_n) is a Palais-Smale sequence, then u is a solution for our equation. If the condition $\text{div}(|u|^4 u) = 0$ is violated, the embedding need not be locally compact.

The paper is organized as follows. In Section 2 we introduce the functional setting and some notation. Section 3 concerns the concentration-compactness analysis as we have already mentioned. In Section 4 we prove Theorem 1.2, and in Section 5 we show that $S_{\text{curl}}(\Omega)$ is independent of Ω and we also prove Theorem 1.3. The proof of Theorem 1.4 is contained in Section 6 whereas in Section 7 we state some open problems.

2. FUNCTIONAL SETTING AND PRELIMINARIES

Throughout the paper we assume that Ω is a Lipschitz domain in \mathbb{R}^3 and $2 \leq p \leq 2^* = 6$. The curl of u , $\nabla \times u$, should be understood in the distributional sense. We shall look for solutions to (1.6) and (1.7)–(1.8) in the space $W_0^6(\text{curl}; \mathbb{R}^3)$ and $W_0^6(\text{curl}; \Omega)$ respectively. We introduce the subspaces

$$\begin{aligned} \mathcal{V}_\Omega &:= \left\{ v \in W_0^6(\text{curl}; \Omega) : \int_\Omega \langle v, \varphi \rangle dx = 0 \text{ for every } \varphi \in \mathcal{C}_0^\infty(\Omega, \mathbb{R}^3) \text{ with } \nabla \times \varphi = 0 \right\}, \\ \mathcal{W}_\Omega &:= \left\{ w \in W_0^6(\text{curl}; \Omega) : \int_\Omega \langle w, \nabla \times \varphi \rangle dx = 0 \text{ for all } \varphi \in \mathcal{C}_0^\infty(\Omega, \mathbb{R}^3) \right\} \\ &= \{ w \in W_0^6(\text{curl}; \Omega) : \nabla \times w = 0 \text{ in the sense of distributions} \}. \end{aligned}$$

The second one has already been defined in Section 1. Here and below $\langle \cdot, \cdot \rangle$ denotes the inner product in \mathbb{R}^3 . If $\Omega = \mathbb{R}^3$, we shall usually write \mathcal{V} and \mathcal{W} for $\mathcal{V}_{\mathbb{R}^3}$ and $\mathcal{W}_{\mathbb{R}^3}$.

In the sequel Ω is always a Lipschitz domain and C denotes a generic positive constant which may vary from one equation to another.

In the following subsections we consider two cases.

2.1. $\Omega = \mathbb{R}^3$.

Lemma 2.1. $W^p(\text{curl}; \mathbb{R}^3) = W_0^p(\text{curl}; \mathbb{R}^3)$ for each $p \in [2, 6]$.

Proof. We show $\mathcal{C}_0^\infty(\mathbb{R}^3, \mathbb{R}^3)$ is dense in $W^p(\text{curl}; \mathbb{R}^3)$. Let $\chi_R \in \mathcal{C}_0^\infty(\mathbb{R}^3)$ be such that $|\nabla \chi_R| \leq 2/R$, $\chi_R = 1$ for $|x| \leq R$ and $\chi_R = 0$ for $|x| \geq 2R$. Take $u = (u_1, u_2, u_3) \in W^p(\text{curl}; \mathbb{R}^3)$. Then $\chi_R u \rightarrow u$ in $L^p(\mathbb{R}^3, \mathbb{R}^3)$ as $R \rightarrow \infty$. We have

$$(2.1) \quad \partial_i(\chi_R u_j) - \partial_j(\chi_R u_i) = (\partial_i \chi_R) u_j - (\partial_j \chi_R) u_i + \chi_R (\partial_i u_j - \partial_j u_i), \quad i \neq j.$$

If $p = 2$, it is clear that $(\partial_i \chi_R) u_j \rightarrow 0$ in $L^2(\mathbb{R}^3)$. If $2 < p \leq 6$, then

$$\int_{\mathbb{R}^3} (\partial_i \chi_R)^2 u_j^2 dx \leq \left(\int_{R \leq |x| \leq 2R} |\partial_i \chi_R|^q dx \right)^{2/q} \left(\int_{R \leq |x| \leq 2R} |u_j|^p dx \right)^{2/p}$$

where $q = 2p/(p-2) \geq 3$. Since

$$\int_{R \leq |x| \leq 2R} |\partial_i \chi_R|^q dx \leq CR^{3-q} < +\infty,$$

also here $(\partial_i \chi_R) u_j \rightarrow 0$ in $L^2(\mathbb{R}^3)$. As $\partial_i u_j - \partial_j u_i \in L^2(\mathbb{R}^3)$, it follows that the left-hand side in (2.1) tends to $\partial_i u_j - \partial_j u_i$ in $L^2(\mathbb{R}^3)$ as $R \rightarrow \infty$. Hence $\chi_R u \rightarrow u$ in $W^p(\text{curl}; \mathbb{R}^3)$ and functions of compact support are dense in $W^p(\text{curl}; \mathbb{R}^3)$.

Suppose now $u \in W^p(\text{curl}; \mathbb{R}^3)$ has a compact support. Clearly, $j_\varepsilon * u \rightarrow u$ in $L^p(\mathbb{R}^3, \mathbb{R}^3)$ as $\varepsilon \rightarrow 0$ where j_ε is the standard mollifier. Since

$$(2.2) \quad \partial_i(j_\varepsilon * u_j) - \partial_j(j_\varepsilon * u_i) = j_\varepsilon * (\partial_i u_j - \partial_j u_i)$$

and $\partial_i u_j - \partial_j u_i \in L^2(\mathbb{R}^3)$, the right-hand side above tends to $\partial_i u_j - \partial_j u_i$ in $L^2(\mathbb{R}^3)$ as $\varepsilon \rightarrow 0$. This completes the proof. \square

As usual, let $\mathcal{D}^{1,2}(\mathbb{R}^3, \mathbb{R}^3)$ denote the completion of $\mathcal{C}_0^\infty(\mathbb{R}^3, \mathbb{R}^3)$ with respect to the norm $|\nabla \cdot|_2$. The following Helmholtz decomposition holds, see [22, 24]:

Lemma 2.2. \mathcal{V} and \mathcal{W} are closed subspaces of $W_0^6(\text{curl}; \mathbb{R}^3)$ and

$$(2.3) \quad W_0^6(\text{curl}; \mathbb{R}^3) = \mathcal{V} \oplus \mathcal{W}.$$

Moreover, $\mathcal{V} \subset \mathcal{D}^{1,2}(\mathbb{R}^3, \mathbb{R}^3)$ and the norms $|\nabla \cdot|_2$ and $\|\cdot\|_{W^6(\text{curl}; \mathbb{R}^3)}$ are equivalent in \mathcal{V} .

We note that \mathcal{W} is the closure of $\{\nabla \varphi : \varphi \in \mathcal{C}_0^\infty(\mathbb{R}^3)\}$. Indeed, if $w \in \mathcal{W}$, then $\nabla \times w = 0$, hence we can find φ_n such that $\nabla \varphi_n \rightarrow w$ and $\nabla \varphi_n \in \mathcal{C}_0^\infty(\mathbb{R}^3, \mathbb{R}^3)$ [22, 24]. Since $\nabla \varphi_n = 0$ outside of some ball, φ_n is constant there and we may assume this constant is 0.

2.2. Ω **bounded.** Recall $H_0(\text{curl}; \Omega) := W_0^2(\text{curl}; \Omega)$ and note that

$$\mathcal{V}_\Omega \subset \{u \in H_0(\text{curl}; \Omega) : \text{div}(u) \in L^2(\Omega, \mathbb{R}^3)\}.$$

Here we have used the fact that if φ in the definition of \mathcal{V}_Ω is supported in a ball, then $\varphi = \nabla\psi$ for some ψ and hence $u \in \mathcal{V}_\Omega$ implies $\text{div}(u) = 0$. It follows from [3, 12] that \mathcal{V}_Ω is continuously embedded in $H^s(\Omega, \mathbb{R}^3)$ for some $s \in [1/2, 1]$, hence compactly in $L^2(\Omega, \mathbb{R}^3)$. If, in addition Ω satisfies (Ω) , then \mathcal{V}_Ω is continuously embedded in $H^1(\Omega, \mathbb{R}^3)$, hence compactly in $L^p(\Omega, \mathbb{R}^3)$ for $1 \leq p < 6$ and continuously in $L^6(\Omega, \mathbb{R}^3)$. This implies in particular that

$$(2.4) \quad \mathcal{V}_\Omega = \left\{ v \in H_0(\text{curl}; \Omega) : \int_\Omega \langle v, \varphi \rangle dx = 0 \text{ for every } \varphi \in \mathcal{C}_0^\infty(\Omega, \mathbb{R}^3) \text{ with } \nabla \times \varphi = 0 \right\}$$

is a Hilbert space with inner product

$$(v, z) = \int_\Omega \langle \nabla \times v, \nabla \times z \rangle dx \equiv \int_\Omega \langle \nabla v, \nabla z \rangle dx.$$

Observe that the right-hand side of (2.4) is a closed linear subspace of $W_0^6(\text{curl}; \Omega)$ as a consequence of (Ω) . Using this, it follows from the decomposition in [18, Theorem 4.21(c)] that also here there is a Helmholtz type decomposition

$$W_0^6(\text{curl}; \Omega) = \mathcal{V}_\Omega \oplus \mathcal{W}_\Omega$$

and that

$$\int_\Omega \langle v, w \rangle dx = 0 \quad \text{if } v \in \mathcal{V}_\Omega, w \in \mathcal{W}_\Omega$$

which means that \mathcal{V}_Ω and \mathcal{W}_Ω are orthogonal in $L^2(\Omega, \mathbb{R}^3)$. In $W_0^6(\text{curl}; \Omega) = \mathcal{V}_\Omega \oplus \mathcal{W}_\Omega$ we can use the norm

$$\|v + w\| := \left((v, v) + |w|_6^2 \right)^{\frac{1}{2}}, \quad v \in \mathcal{V}_\Omega, w \in \mathcal{W}_\Omega$$

which is equivalent to $\|\cdot\|_{W_0^6(\text{curl}; \Omega)}$ if (Ω) is satisfied.

According to [13, Theorem IX.2] or [26, Theorem 3.33], there is a continuous tangential trace operator $\gamma_t : H(\text{curl}; \Omega) := W^2(\text{curl}; \Omega) \rightarrow H^{-1/2}(\partial\Omega)$ such that

$$\gamma_t(u) = \nu \times u|_{\partial\Omega} \quad \text{for any } u \in \mathcal{C}^\infty(\overline{\Omega}, \mathbb{R}^3)$$

and

$$H_0(\text{curl}; \Omega) = \{u \in H(\text{curl}; \Omega) : \gamma_t(u) = 0\}.$$

Hence any vector field $u \in W_0^6(\text{curl}; \Omega) = \mathcal{V}_\Omega \oplus \mathcal{W}_\Omega \subset H_0(\text{curl}; \Omega)$ satisfies the metallic boundary condition (1.8).

Denote the subspace of all gradient vector fields in $W_0^{1,6}(\Omega)$ by $\nabla W_0^{1,6}(\Omega)$. Clearly, $\nabla W_0^{1,6}(\Omega) \subset \mathcal{W}_\Omega$. However, for general domains $\{w \in \mathcal{W}_\Omega : \text{div}(w) = 0\}$ may be nontrivial and hence $\nabla W_0^{1,6}(\Omega) \subsetneq \mathcal{W}_\Omega$, see [7, pp. 4314 and 4315] and [26, Theorem 3.42].

Lemma 2.3. *There holds $\mathcal{W}_\Omega = W_0^6(\text{curl}; \Omega) \cap \mathcal{W} = W_0^6(\text{curl}; \Omega) \cap \nabla W^{1,6}(\Omega)$. If $\partial\Omega$ is connected, then $\mathcal{W}_\Omega = \nabla W_0^{1,6}(\Omega)$. If Ω is unbounded, $\mathcal{W}_\Omega = W_0^6(\text{curl}; \Omega) \cap \mathcal{W}$ still holds.*

Proof. Let $w \in \mathcal{W}_\Omega$ and take a sequence $(\varphi_n) \subset \mathcal{C}_0^\infty(\Omega, \mathbb{R}^3)$ such that $\varphi_n \rightarrow w$ in $W_0^6(\text{curl}; \Omega)$. Extend φ_n by 0 in $\mathbb{R}^3 \setminus \Omega$ and note that (φ_n) is a Cauchy sequence, so $\varphi_n \rightarrow \tilde{w}$ in $W_0^6(\mathbb{R}^3, \mathbb{R}^3)$ where $\tilde{w}|_\Omega = w$ and $\tilde{w} = 0$ in $\mathbb{R}^3 \setminus \Omega$. As

$$\int_{\mathbb{R}^3} \langle \tilde{w}, \nabla \times \psi \rangle dx = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^3} \langle \varphi_n, \nabla \times \psi \rangle dx = \lim_{n \rightarrow \infty} \int_{\mathbb{R}^3} \langle \nabla \times \varphi_n, \psi \rangle dx \leq \lim_{n \rightarrow \infty} |\nabla \times \varphi_n|_2 |\psi|_2 = 0$$

for any $\psi \in C_0^\infty(\mathbb{R}^3, \mathbb{R}^3)$, it follows that $\tilde{w} \in \mathcal{W}$. Moreover, since $\tilde{w} \in L^6(\mathbb{R}^3, \mathbb{R}^3)$ and $\nabla \times \tilde{w} = 0$, in view of [19, Lemma 1.1] we obtain $\tilde{w} = \nabla \psi$ for some $\psi \in W_{loc}^{1,6}(\mathbb{R}^3)$. Therefore $w = \nabla \psi|_\Omega \in \nabla W^{1,6}(\Omega)$. Clearly, $W_0^6(\text{curl}; \Omega) \cap \mathcal{W}$ and $W_0^6(\text{curl}; \Omega) \cap \nabla W^{1,6}(\Omega)$ are contained in \mathcal{W}_Ω .

Suppose that $\partial\Omega$ is connected. Similarly as above, we obtain $w = \nabla \psi$ for some $\psi \in W^{1,6}(\Omega)$ and the surface gradient

$$\nabla_S \psi = (\nu \times \nabla \psi) \times \nu = 0.$$

Therefore we may assume that $\psi \in W_0^{1,6}(\Omega)$, cf. [26, Theorem 4.3 and Remark 4.4]. \square

3. GENERAL CONCENTRATION-COMPACTNESS ANALYSIS IN \mathbb{R}^N

In this, self-contained, section we have $N \geq 3$ and we work in subspaces of $L^{2^*}(\mathbb{R}^N, \mathbb{R}^N)$ where $2^* := 2N/(N-2)$.

Let Ω be a domain in \mathbb{R}^N , \mathcal{V} a closed subspace of $\mathcal{D}^{1,2}(\mathbb{R}^N, \mathbb{R}^N)$ and

$$(3.1) \quad \mathcal{W} := \{w = (w_1, \dots, w_N) \in L^{2^*}(\Omega, \mathbb{R}^N) : \nabla \times w = 0\}$$

where $\nabla \times w$ denotes the skew-symmetric, matrix-valued distribution having $\partial_k w_l - \partial_l w_k \in \mathcal{D}'(\Omega)$ as matrix elements. So for $N = 3$, \mathcal{W} corresponds to \mathcal{W}_Ω in Section 2 but \mathcal{V} may be a more general subspace. Note that $\nabla \times$ is the usual curl operator if $N = 3$. Let Z be a finite-dimensional subspace of $L^{2^*}(\Omega, \mathbb{R}^N)$ such that $Z \cap \mathcal{W} = \{0\}$ and put

$$\widetilde{\mathcal{W}} := \mathcal{W} \oplus Z.$$

Assume

- (F1) $F : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}$ is differentiable with respect to the second variable $u \in \mathbb{R}^N$ for a.e. $x \in \Omega$, $F(x, 0) = 0$ and $f = \partial_u F : \Omega \times \mathbb{R}^N \rightarrow \mathbb{R}^N$ is a Carathéodory function (i.e., f is measurable in $x \in \Omega$ for all $u \in \mathbb{R}^N$ and continuous in $u \in \mathbb{R}^N$ for a.e. $x \in \Omega$);
- (F2) F is uniformly strictly convex with respect to $u \in \mathbb{R}^N$, i.e. for any compact $A \subset (\mathbb{R}^N \times \mathbb{R}^N) \setminus \{(u, u) : u \in \mathbb{R}^N\}$

$$\inf_{\substack{x \in \Omega \\ (u_1, u_2) \in A}} \left(\frac{1}{2} (F(x, u_1) + F(x, u_2)) - F\left(x, \frac{u_1 + u_2}{2}\right) \right) > 0;$$

- (F3) There are $c_1, c_2 > 0$ and $a \in L^{N/2}(\Omega)$, $a \geq 0$, such that

$$c_1 |u|^{2^*} \leq F(x, u) \quad \text{and} \quad |f(x, u)| \leq a(x)|u| + c_2 |u|^{2^*-1}$$

for every $u \in \mathbb{R}^N$ and a.e. $x \in \Omega$.

In view of (F2) and (F3), for any $v \in \mathcal{V}$ we find a unique $\tilde{w}_\Omega(v) \in \widetilde{\mathcal{W}}$ such that

$$(3.2) \quad \int_\Omega F(x, v + \tilde{w}_\Omega(v)) dx \leq \int_\Omega F(x, v + \tilde{w}) dx \quad \text{for all } \tilde{w} \in \widetilde{\mathcal{W}}.$$

This implies that

$$(3.3) \quad \int_\Omega \langle f(x, v + \tilde{w}), \zeta \rangle dx = 0 \quad \text{for all } \zeta \in \widetilde{\mathcal{W}} \text{ if and only if } \tilde{w} = \tilde{w}_\Omega(v).$$

Denote the space of finite measures in \mathbb{R}^N by $\mathcal{M}(\mathbb{R}^N)$.

Theorem 3.1. *Assume that (F1)–(F3) are satisfied. Suppose $(v_n) \subset \mathcal{V}$, $v_n \rightharpoonup v_0$ in \mathcal{V} , $v_n \rightarrow v_0$ a.e. in \mathbb{R}^N , $|\nabla v_n|^2 \rightharpoonup \mu$ and $|v_n|^{2^*} \rightharpoonup \rho$ in $\mathcal{M}(\mathbb{R}^N)$. Then there exists an at most countable set $I \subset \mathbb{R}^N$ and nonnegative weights $\{\mu_x\}_{x \in I}$, $\{\rho_x\}_{x \in I}$ such that*

$$\mu \geq |\nabla v_0|^2 + \sum_{x \in I} \mu_x \delta_x, \quad \rho = |v_0|^{2^*} + \sum_{x \in I} \rho_x \delta_x,$$

and passing to a subsequence, $\tilde{w}_\Omega(v_n) \rightharpoonup \tilde{w}_\Omega(v_0)$ in $\widetilde{\mathcal{W}}$, $\tilde{w}_\Omega(v_n) \rightarrow \tilde{w}_\Omega(v_0)$ a.e. in Ω and in $L^p_{loc}(\Omega)$ for any $1 \leq p < 2^*$.

Remark 3.2. We shall use this theorem in Sections 4 and 6. In Section 4 we have $\Omega = \mathbb{R}^3$ and $Z = \{0\}$, so $\tilde{w} = w$ and we will write $w(v)$ for $w_{\mathbb{R}^3}(v)$. In Section 6, where we treat a Brezis-Nirenberg problem, Ω will be bounded and Z the subspace of \mathcal{V}_Ω on which the quadratic part of J_λ (see (1.9)) is negative semidefinite.

Proof of Theorem 3.1. Step 1. Let $\varphi \in C_0^\infty(\mathbb{R}^N)$. By the Sobolev inequality,

$$(3.4) \quad \left(\int_{\mathbb{R}^N} |\varphi|^{2^*} |v_n - v_0|^{2^*} dx \right)^{1/2^*} \leq S^{-1/2} \left(\int_{\mathbb{R}^N} |\nabla[\varphi(v_n - v_0)]|^2 dx \right)^{1/2} \\ = S^{-1/2} \left(\int_{\mathbb{R}^N} |\varphi|^2 |\nabla(v_n - v_0)|^2 dx \right)^{1/2} + o(1).$$

Passing to the limit and using the Brezis-Lieb lemma [9, 36] on the left-hand side above we obtain

$$(3.5) \quad \left(\int_{\mathbb{R}^N} |\varphi|^{2^*} d\bar{\rho} \right)^{1/2^*} \leq S^{-1/2} \left(\int_{\mathbb{R}^N} |\varphi|^2 d\bar{\mu} \right)^{1/2}$$

where $\bar{\mu} := \mu - |\nabla v_0|^2$ and $\bar{\rho} := \rho - |v_0|^{2^*}$. Set $I = \{x \in \mathbb{R}^N : \mu_x := \mu(\{x\}) > 0\}$. Since μ is finite and $\mu, \bar{\mu}$ have the same singular set, I is at most countable and $\mu \geq |\nabla v_0|^2 + \sum_{x \in I} \mu_x \delta_x$. As in the proof of Theorem 1.9 in [16] it follows from (3.5) that $\bar{\rho} = \sum_{x \in I} \rho_x \delta_x$, see also Proposition 4.2 in [35]. So μ and ρ are as claimed.

Step 2. Using (F3) and (3.2) we infer that

$$c_1 |v_n + \tilde{w}_\Omega(v_n)|_{2^*}^{2^*} \leq \int_{\Omega} F(x, v_n + \tilde{w}_\Omega(v_n)) \leq \int_{\Omega} F(x, v_n) dx \\ \leq c_2 |v_n|_{2^*}^{2^*} + |a|_{N/2} |v_n|_{2^*}^2,$$

and since the right-hand side above is bounded, so is $(|\tilde{w}_\Omega(v_n)|_{2^*})$. Hence, up to a subsequence, $\tilde{w}_\Omega(v_n) \rightharpoonup \tilde{w}_0$ for some \tilde{w}_0 . Write $\tilde{w}_\Omega(v_n) = w_n + z_n$, $\tilde{w}_0 = w_0 + z_0$ where $w_n, w_0 \in \mathcal{W}$ and $z_n, z_0 \in Z$. We shall show that $\tilde{w}_\Omega(v_n) \rightarrow \tilde{w}_0$ a.e. in Ω after taking subsequences. Obviously, we may assume $z_n \rightarrow z_0$ in Z and a.e. in Ω .

We can find a sequence of open balls $(B_l)_{l=1}^\infty$ such that $\Omega = \bigcup_{l=1}^\infty B_l$. Fix $l \geq 1$. In view of [19, Lemma 1.1] there exists $\xi_n \in W^{1,2^*}(B_l)$ such that $w_n = \nabla \xi_n$ and we may assume without loss of generality that $\int_{B_l} \xi_n dx = 0$. Then by the Poincaré inequality,

$$\|\xi_n\|_{W^{1,2^*}(B_l)} \leq C |w_n|_{L^{2^*}(B_l, \mathbb{R}^N)} \leq C |w_n|_{2^*}$$

and passing to a subsequence, $\xi_n \rightarrow \xi$ for some $\xi \in W^{1,2^*}(B_l)$. So $\xi_n \rightarrow \xi$ in $L^{2^*}(B_l)$. Now take any $\varphi \in C_0^\infty(B_l)$. Since $\nabla(|\varphi|^{2^*}(\xi_n - \xi)) \in \mathcal{W}$, in view of (3.3) we get

$$\int_{\Omega} \langle f(x, v_n + \tilde{w}_\Omega(v_n)), \nabla(|\varphi|^{2^*}(\xi_n - \xi)) \rangle dx = 0,$$

that is,

$$\int_{\Omega} |\varphi|^{2^*} \langle f(x, v_n + \tilde{w}_{\Omega}(v_n)), w_n - \nabla \xi \rangle dx = \int_{\Omega} \langle f(x, v_n + \tilde{w}_{\Omega}(v_n)), \nabla(|\varphi|^{2^*})(\xi - \xi_n) \rangle dx,$$

where the right-hand side tends to 0 as $n \rightarrow \infty$. Since $w_n \rightharpoonup \nabla \xi$ in $L^{2^*}(B_l)$,

$$\int_{\Omega} |\varphi|^{2^*} \langle f(x, v_0 + \nabla \xi + z_0), w_n - \nabla \xi \rangle dx = o(1),$$

hence, recalling that $\tilde{w}_{\Omega}(v_n) = w_n + z_n$ and $z_n \rightarrow z_0$, we obtain

$$(3.6) \quad \int_{\Omega} |\varphi|^{2^*} \langle f(x, v_n + \tilde{w}_{\Omega}(v_n)) - f(x, v_0 + \nabla \xi + z_0), \tilde{w}_{\Omega}(v_n) - \nabla \xi - z_0 \rangle dx = o(1).$$

The convexity of F in u implies that

$$F\left(x, \frac{u_1 + u_2}{2}\right) \geq F(x, u_1) + \left\langle f(x, u_1), \frac{u_2 - u_1}{2} \right\rangle$$

and

$$F\left(x, \frac{u_1 + u_2}{2}\right) \geq F(x, u_2) + \left\langle f(x, u_2), \frac{u_1 - u_2}{2} \right\rangle.$$

Adding these inequalities and using (F2), we obtain for any $k \geq 1$ and $|u_1 - u_2| \geq \frac{1}{k}$, $|u_1|, |u_2| \leq k$ that

$$(3.7) \quad m_k \leq \frac{1}{2}(F(x, u_1) + F(x, u_2)) - F\left(x, \frac{u_1 + u_2}{2}\right) \leq \frac{1}{4} \langle f(x, u_1) - f(x, u_2), u_1 - u_2 \rangle$$

where

$$(3.8) \quad m_k := \inf_{\substack{x \in \Omega, u_1, u_2 \in \mathbb{R}^N \\ \frac{1}{k} \leq |u_1 - u_2|, \\ |u_1|, |u_2| \leq k}} \frac{1}{2}(F(x, u_1) + F(x, u_2)) - F\left(x, \frac{u_1 + u_2}{2}\right) > 0.$$

Let

$$\Omega_{n,k} := \left\{ x \in \Omega : |v_n + \tilde{w}_{\Omega}(v_n) - v_0 - \nabla \xi - z_0| \geq \frac{1}{k} \text{ and } |v_n + \tilde{w}_{\Omega}(v_n)|, |v_0 + \nabla \xi + z_0| \leq k \right\}.$$

Taking into account (3.6) and using (F3), (3.7) and Hölder's inequality, we get

$$\begin{aligned} & 4m_k \int_{\Omega_{n,k}} |\varphi|^{2^*} dx \\ & \leq \int_{\Omega} |\varphi|^{2^*} \langle f(x, v_n + \tilde{w}_{\Omega}(v_n)) - f(x, v_0 + \nabla \xi + z_0), v_n + \tilde{w}_{\Omega}(v_n) - v_0 - \nabla \xi - z_0 \rangle dx \\ & \leq \int_{\Omega} |\varphi|^{2^*} \langle f(x, v_n + \tilde{w}_{\Omega}(v_n)) - f(x, v_0 + \nabla \xi + z_0), v_n - v_0 \rangle dx + o(1) \\ & \leq C \left(\int_{\Omega} |\varphi|^{2^*} |v_n - v_0|^{2^*} dx \right)^{1/2^*} + o(1) = C \left(\int_{\Omega} |\varphi|^{2^*} d\bar{\rho} \right)^{1/2^*} + o(1), \end{aligned}$$

where k is fixed. Here we have used the fact that $\int_{\Omega} a(x) |v_n - v_0|^2 dx \rightarrow 0$ if $v_n \rightharpoonup v_0$ in $L^{2^*}(\Omega, \mathbb{R}^N)$. Since $\varphi \in \mathcal{C}_0^\infty(B_l)$ is arbitrary,

$$(3.9) \quad 4m_k |\Omega_{n,k} \cap E| \leq (\bar{\rho}(E))^{1/2^*} + o(1)$$

for any Borel set $E \subset B_l$. We find an open set $E_k \supset I$ such that $|E_k| < 1/2^{k+1}$. Then, taking $E = B_l \setminus E_k$ in (3.9), we have $4m_k|\Omega_{n_k,k} \cap (B_l \setminus E_k)| = o(1)$ as $n \rightarrow \infty$ because $\text{supp}(\bar{\rho}) \subset I$; hence we can find a sufficiently large n_k such that $|\Omega_{n_k,k} \cap B_l| < 1/2^k$ and we obtain

$$\left| \bigcap_{j=1}^{\infty} \bigcup_{k=j}^{\infty} \Omega_{n_k,k} \cap B_l \right| \leq \lim_{j \rightarrow \infty} \sum_{k=j}^{\infty} |\Omega_{n_k,k} \cap B_l| \leq \lim_{j \rightarrow \infty} \frac{1}{2^{j-1}} = 0.$$

If $x \notin \bigcap_{j=1}^{\infty} \bigcup_{k=j}^{\infty} \Omega_{n_k,k}$ and $x \in B_l$, then

$$\begin{aligned} |v_{n_k}(x) + \tilde{w}_{\Omega}(v_{n_k})(x) - v_0(x) - \nabla \xi(x) - z_0(x)| &< \frac{1}{k}, \text{ or } |v_{n_k}(x) + \tilde{w}_{\Omega}(v_{n_k})(x)| > k, \\ \text{or } |v_0(x) + \nabla \xi(x) + z_0(x)| &> k \end{aligned}$$

for all sufficiently large k . Since $v_{n_k} + \tilde{w}_{\Omega}(v_{n_k})$ is bounded in $L^{2^*}(\Omega, \mathbb{R}^N)$, the second and the third inequality above cannot hold on a set of positive measure for all large k . We infer that $v_{n_k} + \tilde{w}_{\Omega}(v_{n_k}) \rightarrow v_0 + \nabla \xi + z_0$, hence $\tilde{w}_{\Omega}(v_{n_k}) \rightarrow \nabla \xi + z_0$ a.e. in B_l . Since $\tilde{w}_{\Omega}(v_n) \rightharpoonup \tilde{w}_0$, $\tilde{w}_0 = \nabla \xi + z_0$ a.e. in B_l . Now employing the diagonal procedure, we find a subsequence of $\tilde{w}_{\Omega}(v_n)$ which converges to \tilde{w}_0 a.e. in $\Omega = \bigcup_{l=1}^{\infty} B_l$.

Let $p \in [1, 2^*)$. For $\Omega' \subset \Omega$ such that $|\Omega'| < +\infty$ we have

$$\int_{\Omega'} |v_n - v_0 + \tilde{w}_{\Omega}(v_n) - \tilde{w}_0|^p dx \leq |\Omega'|^{1-\frac{p}{2^*}} \left(\int_{\Omega} |v_n - v_0 + \tilde{w}_{\Omega}(v_n) - \tilde{w}_0|^{2^*} dx \right)^{\frac{p}{2^*}},$$

hence by the Vitali convergence theorem, $v_n - v_0 + \tilde{w}_{\Omega}(v_n) - \tilde{w}_0 \rightarrow 0$ in $L_{loc}^p(\Omega)$ after passing to a subsequence.

Step 3. We show that $\tilde{w}_{\Omega}(v_0) = \tilde{w}_0$. Take any $\tilde{w} \in \tilde{\mathcal{W}}$ and observe that by the Vitali convergence theorem,

$$0 = \int_{\Omega} \langle f(x, v_n + \tilde{w}_{\Omega}(v_n)), \tilde{w} \rangle dx \rightarrow \int_{\Omega} \langle f(x, v_0 + \tilde{w}_0), \tilde{w} \rangle dx$$

up to a subsequence. Now (3.3) implies that $w_0 = \tilde{w}_{\Omega}(v_0)$ which completes the proof. \square

4. PROBLEM IN $\Omega = \mathbb{R}^3$ AND PROOF OF THEOREM 1.2

Let S be the best Sobolev constant for the embedding of $\mathcal{D}^{1,2}(\mathbb{R}^3)$ into $L^6(\mathbb{R}^3)$, see (1.1). It is clear that a minimizer $w(u)$ in (3.2) exists uniquely for any $u \in W_0^6(\text{curl}; \Omega)$, not only for $u \in \mathcal{V}$. Here we have $F(x, u) = \frac{1}{6}|u|^6$ and $Z = \{0\}$. So by Lemma 2.3, $u + w(u) = v + w(v) \in \mathcal{V} \oplus \mathcal{W}$ for some $v \in \mathcal{V}$ and therefore

$$(4.1) \quad \inf_{w \in \mathcal{W}} \int_{\mathbb{R}^3} |u + w|^6 dx = \int_{\mathbb{R}^3} |u + w(u)|^6 dx = \int_{\mathbb{R}^3} |v + w(v)|^6 dx.$$

Since $\text{div}(v) = 0$,

$$(4.2) \quad S_{\text{curl}} = \inf_{\substack{u \in W_0^6(\text{curl}; \mathbb{R}^3) \\ \nabla \times u \neq 0}} \frac{|\nabla \times u|_2^2}{|u + w(u)|_6^2} = \inf_{v \in \mathcal{V} \setminus \{0\}} \frac{|\nabla v|_2^2}{|v + w(v)|_6^2}.$$

Lemma 4.1. $S_{\text{curl}} \geq S$.

Proof. Given $\varepsilon > 0$, by (4.2) we can find $v \neq 0$ such that

$$(4.3) \quad \int_{\mathbb{R}^3} |\nabla v|^2 dx \leq (S_{\text{curl}} + \varepsilon) \left(\int_{\mathbb{R}^3} |v + w(v)|^6 dx \right)^{\frac{1}{3}}.$$

Let $v = (v_1, v_2, v_3)$. By the Hölder inequality,

$$(4.4) \quad \int_{\mathbb{R}^3} v_1^2 v_2^2 v_3^2 dx \leq \left(\int_{\mathbb{R}^3} v_1^6 dx \int_{\mathbb{R}^3} v_2^6 dx \int_{\mathbb{R}^3} v_3^6 dx \right)^{\frac{1}{3}}$$

and

$$(4.5) \quad \int_{\mathbb{R}^3} v_i^4 v_j^2 dx \leq \left(\int_{\mathbb{R}^3} v_i^6 dx \right)^{\frac{2}{3}} \left(\int_{\mathbb{R}^3} v_j^6 dx \right)^{\frac{1}{3}}, \quad i \neq j.$$

Using this and the Sobolev inequality gives

$$(4.6) \quad \int_{\mathbb{R}^3} |\nabla v|^2 dx \geq S \sum_{i=1}^3 \left(\int_{\mathbb{R}^3} |v_i|^6 dx \right)^{1/3} \geq S \left(\int_{\mathbb{R}^3} |v|^6 dx \right)^{1/3},$$

and since $w(v)$ is a minimizer, we obtain using (4.3) and (4.6)

$$(4.7) \quad \begin{aligned} \int_{\mathbb{R}^3} |\nabla v|^2 dx &\leq (S_{\text{curl}} + \varepsilon) \left(\int_{\mathbb{R}^3} |v + w(v)|^6 dx \right)^{\frac{1}{3}} \leq (S_{\text{curl}} + \varepsilon) \left(\int_{\mathbb{R}^3} |v|^6 dx \right)^{\frac{1}{3}} \\ &\leq (S_{\text{curl}} + \varepsilon)/S \int_{\mathbb{R}^3} |\nabla v|^2 dx. \end{aligned}$$

Hence $S_{\text{curl}} + \varepsilon \geq S$ for all $\varepsilon > 0$ and the conclusion follows. \square

Next we look for ground states for the curl-curl problem (1.6), i.e. nontrivial solutions with least possible associated energy J given by (1.4). Throughout the rest of the paper we shall make repeated use of the following fact:

Lemma 4.2. *Let $\lambda > 0$. Then $w(\lambda u) = \lambda w(u)$. Similarly, if Ω is a proper subset of \mathbb{R}^3 , then $w_\Omega(\lambda u) = \lambda w_\Omega(u)$.*

Proof. We prove this for w_Ω . Using the minimizing property of $w_\Omega(u)$ we obtain

$$\begin{aligned} \lambda^6 \int_{\Omega} |u + w_\Omega(u)|^6 dx &= \int_{\Omega} |\lambda u + \lambda w_\Omega(u)|^6 dx \geq \int_{\Omega} |\lambda u + w_\Omega(\lambda u)|^6 dx \\ &= \lambda^6 \int_{\Omega} |u + w_\Omega(\lambda u)/\lambda|^6 dx \geq \lambda^6 \int_{\Omega} |u + w_\Omega(u)|^6 dx. \end{aligned}$$

Since the minimizer is unique, $w_\Omega(u) = w_\Omega(\lambda u)/\lambda$ as claimed. \square

Lemma 4.3. *Let \mathcal{N} be the set defined in (1.5). Then*

$$(4.8) \quad \mathcal{N} = \{u \in W_0^6(\text{curl}; \mathbb{R}^3) \setminus \mathcal{W} : J'(u)u = 0 \text{ and } J'(u)|_{\mathcal{W}} = 0\}.$$

Proof. The first condition in (1.5) is equivalent to $J'(u)u = 0$. The second condition is satisfied because $\text{div}(|u|^4 u) = 0$ if and only if $\int_{\mathbb{R}^3} \langle |u|^4 u, \nabla \varphi \rangle dx = 0$ for all $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^3)$ and each element of \mathcal{W} can be approximated by such φ , see the comment preceding Section 2.2. \square

By Lemma 2.2, $W_0^6(\text{curl}; \mathbb{R}^3) = \mathcal{V} \oplus \mathcal{W}$. It follows from (3.2) and (3.3) that if $v \in \mathcal{V}$, then $J'(v + w(v))|_{\mathcal{W}} = 0$, and as

$$(4.9) \quad J(t(v + w(v))) = \frac{t^2}{2} \int_{\mathbb{R}^3} |\nabla v|^2 dx - \frac{t^6}{6} \int_{\mathbb{R}^3} |v + w(v)|^6 dx,$$

there is a unique $t(v) > 0$ such that

$$(4.10) \quad m(v) := t(v)(v + w(v)) \in \mathcal{N} \quad \text{for } v \in \mathcal{V} \setminus \{0\}.$$

We note that

$$(4.11) \quad J(m(v)) \geq J(t(v+w)) \quad \text{for all } t > 0 \text{ and } w \in \mathcal{W}.$$

Since $J(m(v)) \geq J(v)$ and there exist $a, r > 0$ such that $J(v) \geq a$ if $\|v\| = r$, \mathcal{N} is bounded away from \mathcal{W} and hence closed.

Lemma 4.4. *The mapping $m : \mathcal{V} \setminus \{0\} \rightarrow \mathcal{N}$ given by (4.10) is continuous.*

Proof. Let $v_n \rightarrow v_0 \neq 0$ in \mathcal{V} . Since

$$(4.12) \quad \int_{\mathbb{R}^3} |v_n + w(v_n)|^6 dx \leq \int_{\mathbb{R}^3} |v_n|^6 dx,$$

it follows that $(w(v_n))$ is bounded and it is then clear from (4.9) that so is $(t(v_n))$. Hence we may assume $t(v_n) \rightarrow t_0$ and $w(v_n) \rightharpoonup w_0$ in $L^6(\mathbb{R}^3, \mathbb{R}^3)$. By the weak sequential lower semicontinuity of the second integral in (4.9) and by (4.11),

$$J(t_0(v_0 + w_0)) \geq \limsup_{n \rightarrow \infty} J(t(v_n)(v_n + w(v_n))) \geq \limsup_{n \rightarrow \infty} J(t_0(v_n + w_0)) = J(t_0(v_0 + w_0)).$$

So $w(v_n) \rightarrow w_0$ and since \mathcal{N} is closed, $t_0(v_0 + w_0) = t(v_0)(v_0 + w(v_0)) = m(v_0)$. \square

Now it is easily seen that $m|_{\mathcal{S}} : \mathcal{S} := \{v \in \mathcal{V} : \|v\| = 1\} \rightarrow \mathcal{N}$ is a homeomorphism with the inverse $u = v + w(v) \mapsto v/\|v\|$. Note that \mathcal{N} is an infinite-dimensional topological manifold of infinite codimension. Although J is of class \mathcal{C}^2 , we do not know whether \mathcal{N} is of class \mathcal{C}^1 . However, repeating the argument in [22, Proposition 4.4(b)] or [32, Proposition 2.9] we see that $J \circ m|_{\mathcal{S}} : \mathcal{S} \rightarrow \mathbb{R}$ is of class \mathcal{C}^1 and is bounded from below by the constant $a > 0$ introduced above. By the Ekeland variational principle [36, Theorem 8.5], there is a Palais-Smale sequence $(v_n) \subset \mathcal{S}$ such that

$$(4.13) \quad (J \circ m)(v_n) \rightarrow \inf_{\mathcal{S}} J \circ m = \inf_{\mathcal{N}} J \geq a > 0.$$

It follows from [22, Proposition 4.4(b)] again or from [32, Corollary 2.10] that $(m(v_n))$ is a Palais-Smale sequence for J on \mathcal{N} , so in particular, $J'(m(v_n)) \rightarrow 0$ as $n \rightarrow \infty$. See also an abstract critical point theory on the generalized Nehari manifold in [6, Section 4] and in [7, Section 4].

For $s > 0$, $y \in \mathbb{R}^N$ and $u : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ we denote $T_{s,y}(u) := s^{1/2}u(s \cdot + y)$. The following lemma is a special case of [29, Theorem 1], see also [34, Lemma 5.3].

Lemma 4.5. *Suppose that $(v_n) \subset \mathcal{D}^{1,2}(\mathbb{R}^3, \mathbb{R}^3)$ is bounded. Then $v_n \rightarrow 0$ in $L^6(\mathbb{R}^3, \mathbb{R}^3)$ if and only if $T_{s_n, y_n}(v_n) \rightarrow 0$ in $\mathcal{D}^{1,2}(\mathbb{R}^3, \mathbb{R}^3)$ for all $(s_n) \subset \mathbb{R}^+$ and $(y_n) \subset \mathbb{R}^3$.*

Lemma 4.6. *$T_{s,y}$ is an isometric isomorphism of $W_0^6(\text{curl}; \mathbb{R}^3)$ which leaves the functional J and the subspaces \mathcal{V}, \mathcal{W} invariant. In particular, $w(T_{s,y}u) = T_{s,y}w(u)$.*

The proof is by an explicit (and simple) computation.

Lemma 4.7. *Suppose $u + w(u) \in \mathcal{N}$. Then*

$$\frac{|\nabla \times u|_2^2}{|u + w(u)|_6^2} = A \quad \text{if and only if} \quad J(u + w(u)) = \frac{1}{3}A^{3/2}.$$

In particular, $\inf_{\mathcal{N}} J = \frac{1}{3}S_{\text{curl}}^{3/2}$.

Proof. Since $u + w(u) \in \mathcal{N}$, $J'(u)u = 0$, i.e. $|\nabla \times u|_2^2 = |u + w(u)|_6^6$. Hence

$$\frac{|\nabla \times u|_2^2}{|u + w(u)|_6^2} = |u + w(u)|_6^4 \quad \text{and} \quad J(u + w(u)) = \frac{1}{3}|u + w(u)|_6^6.$$

□

Proof of Theorem 1.2. We prove part (b) first. Take a minimizing sequence $(u_n) = (m(v_n)) \subset \mathcal{N}$ constructed above and write $u_n = t(v_n)(v_n + w(v_n)) = v'_n + w(v'_n) \in \mathcal{V} \oplus \mathcal{W}$. As

$$(4.14) \quad J(u_n) = J(u_n) - \frac{1}{6}J'(u_n)u_n = \frac{1}{3}|\nabla \times u_n|_2^2 = \frac{1}{3}|\nabla v'_n|_2^2$$

and $|\nabla \cdot|_2$ is an equivalent norm in \mathcal{V} , (v'_n) is bounded. We also have

$$(4.15) \quad J(u_n) = J(u_n) - \frac{1}{2}J'(u_n)u_n = \frac{1}{3}|u_n|_6^6.$$

Since $J(u_n)$ is bounded away from 0, $|u_n|_6 \not\rightarrow 0$ and hence by (4.12), $|v'_n|_6 \not\rightarrow 0$. Therefore, passing to a subsequence and using Lemma 4.5, $\tilde{v}_n := T_{s_n, y_n}(v'_n) \rightharpoonup v_0$ for some $v_0 \neq 0$, $(s_n) \subset \mathbb{R}^+$ and $(y_n) \subset \mathbb{R}^3$. Taking subsequences again we also have that $\tilde{v}_n \rightarrow v_0$ a.e. in \mathbb{R}^3 and in view of Theorem 3.1, $w(\tilde{v}_n) \rightharpoonup w(v_0)$ and $w(\tilde{v}_n) \rightarrow w(v_0)$ a.e. in \mathbb{R}^3 . We set $u := v_0 + w(v_0)$ and by Lemma 4.6 we may assume without loss of generality that $s_n = 1$ and $y_n = 0$. So if $z \in W_0^6(\text{curl}; \mathbb{R}^3)$, then using weak and a.e. convergence,

$$J'(u_n)z = \int_{\mathbb{R}^3} \langle \nabla \times u_n, z \rangle dx - \int_{\mathbb{R}^3} \langle |u_n|^4 u_n, z \rangle dx \rightarrow J'(u)z.$$

Here we have used that $|u_n|^4 u_n \rightharpoonup \zeta$ in $L^{6/5}(\mathbb{R}^3, \mathbb{R}^3)$ for some ζ but since $|u_n|^4 u_n \rightarrow |u|^4 u$ a.e., $\zeta = |u|^4 u$. So u is a solution to (1.6). To show it is a ground state, we note that using Fatou's lemma,

$$\begin{aligned} \inf_{\mathcal{N}} J &= J(u_n) + o(1) = J(u_n) - \frac{1}{2}J'(u_n)u_n + o(1) = \frac{1}{3}|u_n|_6^6 + o(1) \\ &\geq \frac{1}{3}|u|_6^6 + o(1) = J(u) - \frac{1}{2}J'(u)u + o(1) = J(u) + o(1). \end{aligned}$$

Hence $J(u) \leq \inf_{\mathcal{N}} J$ and as a solution, $u \in \mathcal{N}$. It follows using Lemma 4.7 that $J(u) = \inf_{\mathcal{N}} J = \frac{1}{3}S_{\text{curl}}^{3/2}$.

If u satisfies equality in (1.3), then $t(u)(u + w(u)) \in \mathcal{N}$ and is a minimizer for $J|_{\mathcal{N}}$. But then the corresponding point v in \mathcal{S} is a minimizer for $J \circ m|_{\mathcal{S}}$, see (4.13). So v is a critical point of $J \circ m|_{\mathcal{S}}$ and $m(v) = u$ is a critical point of J . This completes the proof of (b).

(a) By Lemma 4.1, $S_{\text{curl}} \geq S$ and by part (b), there exists $u = v + w(v)$ for which S_{curl} is attained. Suppose $S_{\text{curl}} = S$. Then all inequalities become equalities in (4.7) with $\varepsilon = 0$, and therefore also in (4.6). But then $\int_{\mathbb{R}^3} |\nabla v_i|^2 dx = S|v_i|_6^2$ for $i = 1, 2, 3$ and hence all v_i are instantons, up to multiplicative constants. Since $v \neq 0$ and $\text{div}(v) = 0$, this is impossible. It follows that $S_{\text{curl}} > S$. □

5. PROOF OF THEOREMS 1.1 AND 1.3

Let Ω be a Lipschitz domain in \mathbb{R}^3 . Recall from Section 2 that we have the Helmholtz decompositions

$$(5.1) \quad W_0^6(\text{curl}; \mathbb{R}^3) = \mathcal{V} \oplus \mathcal{W} \quad \text{and} \quad W_0^6(\text{curl}; \Omega) = \mathcal{V}_\Omega \oplus \mathcal{W}_\Omega$$

where the second one holds if condition (Ω) from Introduction is satisfied. For $u \in W_0^6(\text{curl}; \Omega)$, denote the minimizer of

$$\int_{\Omega} |u + w|^6 dx, \quad w \in \mathcal{W}_\Omega$$

by $w_\Omega(u)$ (cf. (4.1)) and, according to our notational convention, write $w(u)$ for $w_{\mathbb{R}^3}(u)$. Recall from (1.3) the definition of $S_{\text{curl}}(\Omega)$:

$$\int_{\mathbb{R}^3} |\nabla \times u|^2 dx \geq S_{\text{curl}}(\Omega) \inf_{w \in \mathcal{W}} \left(\int_{\mathbb{R}^3} |u + w|^6 dx \right)^{1/3}$$

where $u \in W_0^6(\text{curl}; \Omega)$ and $S_{\text{curl}}(\Omega)$ is the largest constant with this property. By (5.1) we have $u = v + w \in \mathcal{V} \oplus \mathcal{W}$. We emphasize that although $u = 0$ in $\mathbb{R}^3 \setminus \overline{\Omega}$, v and w need not be 0 there. Note that $S_{\text{curl}}(\Omega)$ can be characterized as

$$(5.2) \quad S_{\text{curl}}(\Omega) = \inf_{\substack{u \in W_0^6(\text{curl}; \Omega) \\ \nabla \times u \neq 0}} \sup_{w \in \mathcal{W}} \frac{|\nabla \times u|_2^2}{|u + w|_6^2} = \inf_{\substack{u \in W_0^6(\text{curl}; \Omega) \\ \nabla \times u \neq 0}} \frac{|\nabla \times u|_2^2}{|u + w(u)|_6^2}$$

(cf. (4.2)). In domains $\Omega \neq \mathbb{R}^3$ there is also another constant, $\overline{S}_{\text{curl}}(\Omega)$, introduced in (1.10). Similarly as in (5.2), it can be characterized as

$$(5.3) \quad \overline{S}_{\text{curl}}(\Omega) = \inf_{\substack{u \in W_0^6(\text{curl}; \Omega) \\ \nabla \times u \neq 0}} \sup_{w \in \mathcal{W}_\Omega} \frac{|\nabla \times u|_2^2}{|u + w|_6^2} = \inf_{\substack{u \in W_0^6(\text{curl}; \Omega) \\ \nabla \times u \neq 0}} \frac{|\nabla \times u|_2^2}{|u + w_\Omega(u)|_6^2}.$$

As we have noticed in Introduction, although this constant seems more natural, we do not know whether it equals S_{curl} .

Lemma 5.1. *The mapping $u \mapsto w_\Omega(u) : L^6(\Omega, \mathbb{R}^3) \rightarrow L^6(\Omega, \mathbb{R}^3)$ is continuous ($\Omega = \mathbb{R}^3$ is admitted).*

Proof. Let $u_n \rightarrow u_0$. Since $(w_\Omega(u_n))$ is bounded, $w_\Omega(u_n) \rightharpoonup w_0$ after passing to a subsequence. By the maximality and uniqueness of $w_\Omega(\cdot)$,

$$\begin{aligned} \int_{\Omega} |u_0 + w_\Omega(u_0)|^6 dx &\leq \int_{\Omega} |u_0 + w_0|^6 dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |u_n + w_\Omega(u_n)|^6 dx \\ &\leq \liminf_{n \rightarrow \infty} \int_{\Omega} |u_n + w_\Omega(u_0)|^6 dx = \int_{\Omega} |u_0 + w_\Omega(u_0)|^6 dx. \end{aligned}$$

Hence all inequalities above must be equalities and it follows that $w_0 = w_\Omega(u_0)$ and $w_\Omega(u_n) \rightarrow w_\Omega(u_0)$. \square

We shall need the following inequality:

Lemma 5.2. *If $u \in W_0^6(\text{curl}; \Omega) \setminus \{0\}$, $w \in \mathcal{W}_\Omega$ and $t \geq 0$, then*

$$(5.4) \quad J(u) \geq J(tu + w) - J'(u) \left[\frac{t^2 - 1}{2} u + tw \right].$$

Moreover, strict inequality holds unless $t = 1$ and $w = 0$. ($\Omega = \mathbb{R}^3$ admitted.)

Proof. The proof follows a similar argument as in [22, Proposition 4.1] and [23, Lemma 4.1]. We include it for the reader's convenience. We show that

$$(5.5) \quad J(u) - J(tu + w) + J'(u) \left[\frac{t^2 - 1}{2}u + tw \right] = \int_{\mathbb{R}^3} \varphi(t, x) dx \geq 0,$$

where

$$\varphi(t, x) := - \left\langle |u|^4 u, \frac{t^2 - 1}{2}u + tw \right\rangle - \frac{1}{6}|u|^6 + \frac{1}{6}|tu + w|^6.$$

An explicit computation using $\nabla \times w = 0$ shows that both sides of (5.5) are equal. Clearly, $\varphi(t, x) \geq 0$ if $u(x) = 0$. So let $u(x) \neq 0$. It is easy to check that $\varphi(0, x) > 0$ and $\varphi(t, x) \rightarrow \infty$ as $t \rightarrow \infty$. Note that if $\partial_t \varphi(t_0, x) = 0$ for some $t_0 > 0$, then either $\langle u, t_0 u + w \rangle = 0$ or $|u| = |t_0 u + w|$. In the first case, substituting $-\langle u, w \rangle = t_0 |u|^2$, we obtain $\varphi(t_0, x) = \left(\frac{t_0^2}{2} + \frac{1}{3}\right)|u|^6 + \frac{1}{6}|t_0 u + w|^6 > 0$. In the second case we have, using $-t_0 \langle u, w \rangle = \frac{t_0^2 - 1}{2}|u|^2 + \frac{1}{2}|w|^2$, that $\varphi(t_0, x) = \frac{1}{2}|u|^4 |w|^2 \geq 0$. Hence $\varphi(t, x) \geq 0$ for all $t \geq 0$ and the inequality is strict if $w \neq 0$. If $w = 0$, then $\varphi(t, x) = \left(\frac{t^6}{6} - \frac{t^2}{2} + \frac{1}{3}\right)|u|^6 > 0$ provided $t \neq 1$. \square

Similarly as in (4.8) we introduce the set

$$(5.6) \quad \mathcal{N}_\Omega := \left\{ u \in W_0^6(\text{curl}; \Omega) \setminus \mathcal{W}_\Omega : J'(u)u = 0 \text{ and } J'(u)|_{\mathcal{W}_\Omega} = 0 \right\}.$$

Proof of Theorems 1.1 and 1.3. Since $tu + w(tu) = t(u + w(u))$ according to Lemma 4.2, we may assume without loss of generality that $u + w(u) \in \mathcal{N}$ in (5.2) and similarly, $u + w_\Omega(u) \in \mathcal{N}_\Omega$ in (5.3). According to Lemma 4.7,

$$\inf_{\mathcal{N}} J|_{W_0^6(\text{curl}; \Omega)} = \frac{1}{3} S_{\text{curl}}(\Omega)^{\frac{3}{2}}, \quad \inf_{\mathcal{N}_\Omega} J = \frac{1}{3} \bar{S}_{\text{curl}}(\Omega)^{\frac{3}{2}}, \quad \inf_{\mathcal{N}} J = \frac{1}{3} S_{\text{curl}}^{\frac{3}{2}}.$$

In view of Lemma 2.3, $\mathcal{W}_\Omega \subset \mathcal{W}$, hence we easily infer from (5.2), (5.3) that $S_{\text{curl}}(\Omega) \geq \bar{S}_{\text{curl}}(\Omega)$. As $W_0^6(\text{curl}; \Omega) \subset W_0^6(\text{curl}; \mathbb{R}^3)$, it follows that $S_{\text{curl}} \leq S_{\text{curl}}(\Omega)$.

Next we show that $S_{\text{curl}}(\Omega) \leq S_{\text{curl}}$. Let u_0 be a minimizer for J on \mathcal{N} provided by Theorem 1.2(b) and find a sequence $(u_n) \subset \mathcal{C}_0^\infty(\mathbb{R}^3, \mathbb{R}^3)$ such that $u_n \rightarrow u_0$. We can decompose u_n as $u_n = v_n + w_n$, $v_n \in \mathcal{V}$, $w_n \in \mathcal{W}$. Since $u_0 = v_0 + w(v_0)$ (recall $u_0 \in \mathcal{N}$), $u_n = v_n + w_n \rightarrow u_0 = v_0 + w(v_0)$ and therefore $v_n \rightarrow v_0$, $w_n \rightarrow w(v_0)$. So $v_0 \neq 0$ and v_n are bounded away from 0 in $L^6(\mathbb{R}^3, \mathbb{R}^3)$. Assume without loss of generality that $0 \in \Omega$. There exist λ_n such that \tilde{u}_n given by $\tilde{u}_n(x) := \lambda_n^{1/2} u_n(\lambda_n x)$ are supported in Ω . Set $\tilde{w}_n := w(\tilde{u}_n) \in \mathcal{W}$ and choose t_n so that $t_n(\tilde{u}_n + \tilde{w}_n) \in \mathcal{N}$. Then

$$(5.7) \quad t_n^2 = \frac{|\nabla \times \tilde{u}_n|_2}{|\tilde{u}_n + \tilde{w}_n|_6^3}.$$

According to Lemma 4.6, $\|\tilde{u}_n\| = \|u_n\|$ and $|\tilde{u}_n + \tilde{w}_n|_6 = |u_n + w(u_n)|_6 = |v_n + w(v_n)|_6$. As (u_n) is bounded, so is (\tilde{u}_n) and as $|v_n + w(v_n)|_6 \rightarrow |v_0 + w(v_0)|_6$, $|\tilde{u}_n + \tilde{w}_n|_6$ is bounded away from 0. So (t_n) is bounded. Moreover, $|\tilde{w}_n|_6 = |w(u_n)|_6$ and therefore (\tilde{w}_n) is bounded. Since

$J(\tilde{u}_n) = J(u_n) \rightarrow \frac{1}{3}S_{\text{curl}}^{3/2}$ and $\|J'(\tilde{u}_n)\| = \|J'(u_n)\| \rightarrow 0$, it follows from Lemma 5.2 that

$$\begin{aligned} \frac{1}{3}S_{\text{curl}}^{3/2} &= \lim_{n \rightarrow \infty} J(\tilde{u}_n) \geq \lim_{n \rightarrow \infty} \left(J(t_n(\tilde{u}_n + \tilde{w}_n)) - J'(\tilde{u}_n) \left[\frac{t_n^2 - 1}{2} \tilde{u}_n + t_n^2 \tilde{w}_n \right] \right) \\ &= \lim_{n \rightarrow \infty} J(t_n(\tilde{u}_n + \tilde{w}_n)) \geq \frac{1}{3}S_{\text{curl}}(\Omega)^{3/2}. \end{aligned}$$

The last inequality follows from Lemma 4.7 and the fact that \tilde{u}_n are as in (5.2), i.e. $\tilde{u}_n \in W_0^6(\text{curl}; \Omega)$.

It remains to show that $\bar{S}_{\text{curl}}(\Omega) \geq S$ if (Ω) is satisfied. But this follows by repeating the argument of Lemma 4.1 with obvious changes: S_{curl} should be replaced by $\bar{S}_{\text{curl}}(\Omega)$, $w(v)$ by $w_\Omega(v)$ and the domain of integration should be Ω . \square

Remark 5.3. Let $\Omega \neq \mathbb{R}^3$ and suppose $S_{\text{curl}}(\Omega)$ is attained by some u . Extend u by 0 outside Ω . As $S_{\text{curl}}(\Omega) = S_{\text{curl}}$, u also solves (1.6) in \mathbb{R}^3 , possibly after replacing u with αu for an appropriate $\alpha > 0$. In particular, if $S_{\text{curl}}(\Omega)$ were attained in a bounded Ω , this would imply the existence of ground states in \mathbb{R}^3 which have compact support. To our knowledge, there is no unique continuation principle which could rule out this possibility.

In view of this remark we expect that similarly as is the case for the Sobolev constant, S_{curl} is attained if and only if $\Omega = \mathbb{R}^3$. We leave this problem as a conjecture.

6. THE BREZIS-NIRENBERG-TYPE PROBLEM AND PROOF OF THEOREM 1.4

Let $\lambda \leq 0$. In this section $\Omega \subset \mathbb{R}^3$ is a fixed bounded domain satisfying (Ω) but λ will be varying. Therefore we drop the subscript Ω from notation and replace it by λ (J_λ , \mathcal{N}_λ etc.). We also write \mathcal{V}, \mathcal{W} for $\mathcal{V}_\Omega, \mathcal{W}_\Omega$.

Recall from Introduction and Subsection 2.2 that the spectrum of the curl-curl operator in $H_0(\text{curl}; \Omega)$ consists of the eigenvalue $\lambda_0 = 0$ whose eigenspace is \mathcal{W} and of a sequence of eigenvalues

$$0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \rightarrow \infty$$

with finite multiplicities $m(\lambda_k) \in \mathbb{N}$. The eigenfunctions corresponding to different eigenvalues are L^2 -orthogonal and those corresponding to $\lambda_k > 0$ are in \mathcal{V} .

For $\lambda \leq 0$ we find two closed and orthogonal subspaces \mathcal{V}^+ and $\tilde{\mathcal{V}}$ of \mathcal{V} such that the quadratic form $Q : \mathcal{V} \rightarrow \mathbb{R}$ given by

$$Q(v) := \int_{\Omega} (|\nabla \times v|^2 + \lambda|v|^2) dx \equiv \int_{\Omega} (|\nabla v|^2 + \lambda|v|^2) dx$$

is positive definite on \mathcal{V}^+ and negative semidefinite on $\tilde{\mathcal{V}}$ where $\dim \tilde{\mathcal{V}} < \infty$. Writing $u = v + w = v^+ + \tilde{v} + w \in \mathcal{V}^+ \oplus \tilde{\mathcal{V}} \oplus \mathcal{W}$, we have

$$Q(v) = Q(v^+) + Q(\tilde{v})$$

and our functional J_λ (see (1.9)) can be expressed as

$$J_\lambda(u) = \frac{1}{2}Q(v^+) + \frac{1}{2}Q(\tilde{v}) + \frac{\lambda}{2} \int_{\Omega} |w|^2 dx - \frac{1}{6} \int_{\Omega} |u|^6 dx.$$

We shall use Theorem 3.1 with

$$F(x, u) = \frac{1}{6}|u|^6 - \frac{\lambda}{2}|u|^2.$$

Here $\widetilde{\mathcal{W}} := \widetilde{\mathcal{V}} \oplus \mathcal{W}$ (so $Z = \widetilde{\mathcal{V}}$ in the notation of Section 3) and $\widetilde{w} = \widetilde{v} + w$. \mathcal{V} , and hence \mathcal{V}^+ , may be considered, after a proper extension, as closed subspaces of $\mathcal{D}^{1,2}(\mathbb{R}^3, \mathbb{R}^3)$. Indeed, let U be a bounded domain in \mathbb{R}^3 , $U \supset \overline{\Omega}$. Since $\mathcal{V} \subset H^1(\Omega, \mathbb{R}^3)$, each $v \in \mathcal{V}$ may be extended to $v' \in H_0^1(U, \mathbb{R}^3)$ such that $v'|_{\Omega} = v$. This extension is bounded as a mapping from \mathcal{V} to $H_0^1(U, \mathbb{R}^3)$. Since

$$\mathcal{V}' := \{v' \in H_0^1(U, \mathbb{R}^3) : v'|_{\Omega} \in \mathcal{V}\}$$

is a closed subspace of $H_0^1(U, \mathbb{R}^3)$, and hence of $\mathcal{D}^{1,2}(\mathbb{R}^3, \mathbb{R}^3)$, we can apply Theorem 3.1 with F as above and \mathcal{V}^+ replacing \mathcal{V} . The generalized Nehari manifold is now given by

$$(6.1) \quad \mathcal{N}_{\lambda} := \{u \in W_0^6(\text{curl}; \Omega) \setminus (\widetilde{\mathcal{V}} \oplus \mathcal{W}) : J'_{\lambda}(u)|_{\mathbb{R}u \oplus \widetilde{\mathcal{V}} \oplus \mathcal{W}} = 0\}.$$

As in Section 4, also here it is not clear whether \mathcal{N}_{λ} is of class \mathcal{C}^1 . Setting $m_{\lambda}(v^+) := v^+ + \widetilde{w}(v^+)$ where $v^+ \in \mathcal{V}^+$ and $\widetilde{w}(v^+) \equiv \widetilde{w}_{\Omega}(v^+)$ is the minimizer as in (3.2), we have

$$m_{\lambda}(v^+) := t(v^+)(v^+ + \widetilde{w}(v^+)) \in \mathcal{N}_{\lambda}, \quad v^+ \in \mathcal{V}^+ \setminus \{0\}$$

(cf. (4.10)) and $J_{\lambda} \circ m_{\lambda}$ is of class \mathcal{C}^1 on \mathcal{S}^+ . Moreover, $m_{\lambda}|_{\mathcal{S}^+}$ is a homeomorphism between \mathcal{S}^+ and \mathcal{N}_{λ} . As in (4.13), we may also find a Palais-Smale sequence $(v_n^+) \subset \mathcal{S}^+$ such that

$$(6.2) \quad (J_{\lambda} \circ m_{\lambda})(v_n^+) \rightarrow \inf_{\mathcal{S}^+} J_{\lambda} \circ m_{\lambda} = c_{\lambda} \text{ and } J'_{\lambda}(m_{\lambda}(v_n^+)) \rightarrow 0$$

where

$$c_{\lambda} := \inf_{\mathcal{N}_{\lambda}} J_{\lambda}.$$

Note that $c_0 = \frac{1}{3}\overline{S}_{\text{curl}}(\Omega)^{3/2} \geq \frac{1}{3}S^{3/2}$.

Lemma 6.1. *Let $\lambda \in (-\lambda_{\nu}, -\lambda_{\nu-1}]$ for some $\nu \geq 1$. There holds*

$$c_{\lambda} \leq \frac{1}{3}(\lambda + \lambda_{\nu})^{3/2}|\Omega| \quad \text{and} \quad c_{\lambda} < c_0 \text{ if } \lambda < -\lambda_{\nu} + \overline{S}_{\text{curl}}(\Omega)|\Omega|^{-2/3}.$$

Proof. The first inequality has been established in [23, Lemma 4.7]. However, for the reader's convenience we include the argument. Let e_{ν} be an eigenvector corresponding to λ_{ν} . Then $e_{\nu} \in \mathcal{V}^+$. Choose $t > 0$, $\widetilde{v} \in \widetilde{\mathcal{V}}$ and $w \in \mathcal{W}$ so that $u = v + w = te_{\nu} + \widetilde{v} + w \in \mathcal{N}_{\lambda}$. Since $\lambda_k \leq \lambda_{\nu}$ for $k < \nu$,

$$\begin{aligned} c_{\lambda} &\leq J_{\lambda}(u) = \frac{1}{2} \int_{\Omega} |\nabla \times v|^2 dx + \frac{\lambda}{2} \int_{\Omega} |u|^2 dx - \frac{1}{6} \int_{\Omega} |u|^6 dx \\ &\leq \frac{\lambda_{\nu}}{2} \int_{\Omega} |v|^2 dx + \frac{\lambda}{2} \int_{\Omega} |u|^2 dx - \frac{1}{6} \int_{\Omega} |u|^6 dx \leq \frac{\lambda + \lambda_{\nu}}{2} \int_{\Omega} |u|^2 dx - \frac{1}{6} \int_{\Omega} |u|^6 dx \\ &\leq \frac{\lambda + \lambda_{\nu}}{2} |\Omega|^{2/3} \left(\int_{\Omega} |u|^6 dx \right)^{1/3} - \frac{1}{6} \int_{\Omega} |u|^6 dx \leq \frac{1}{3}(\lambda + \lambda_{\nu})^{3/2}|\Omega|. \end{aligned}$$

In the last step we have used the elementary inequality $\frac{A}{2}t^2 - \frac{1}{6}t^6 \leq \frac{1}{3}A^{3/2}$ ($A > 0$).

Since $c_0 = \frac{1}{3}\overline{S}_{\text{curl}}(\Omega)^{3/2}$, the second inequality follows immediately. \square

If $c_\lambda < c_0$, then in view of [23, Theorem 2.2 (a)] there is a Palais-Smale sequence $(u_n) \subset \mathcal{N}_\lambda$ such that $J_\lambda(u_n) \rightarrow c_\lambda > 0$ and $u_n \rightharpoonup u_0 \neq 0$ in $W_0^6(\text{curl}; \Omega)$. It has been unclear so far whether u_0 is a critical point of J_λ . Now we shall show using the concentration-compactness analysis from Section 3 that u_0 is not only a solution but even a ground state for (1.7). The following lemma plays a crucial role.

Lemma 6.2. *If $(u_n) \subset \mathcal{N}_\lambda$ is bounded, then, passing to a subsequence, $u_n \rightarrow u_0$ in $L^2(\Omega, \mathbb{R}^3)$ for some u_0 .*

Proof. Let $u_n = m_\lambda(v_n^+) = v_n^+ + \tilde{w}(v_n^+)$. Since \mathcal{V}^+ and $\tilde{\mathcal{W}}$ are complementary subspaces, (v_n^+) is bounded in \mathcal{V}^+ . So passing to a subsequence, $v_n^+ \rightharpoonup v_0^+$ in \mathcal{V}^+ , and $v_n^+ \rightarrow v_0^+$ in $L^2(\Omega, \mathbb{R}^3)$ and a.e. in Ω . Hence by Theorem 3.1, $\tilde{w}(v_n^+) \rightarrow \tilde{w}(v_0^+)$ in $L^2(\Omega, \mathbb{R}^3)$, and therefore also $u_n \rightarrow u_0$ there. \square

Lemma 6.3. *(cf. [23, Lemma 4.6]) J_λ is coercive on \mathcal{N}_λ .*

Proof. Let (u_n) be a sequence in \mathcal{N}_λ such that $J_\lambda(u_n) \leq d$. Then

$$d \geq J_\lambda(u_n) = J_\lambda(u_n) - \frac{1}{2}J'_\lambda(u_n)u_n = \frac{1}{3} \int_\Omega |u_n|^6 dx,$$

hence (u_n) is bounded in $L^6(\Omega, \mathbb{R}^3)$, and therefore also in $L^2(\Omega, \mathbb{R}^3)$. It follows that

$$d \geq J_\lambda(u_n) = \frac{1}{2}Q(v_n^+) + \frac{1}{2}Q(\tilde{v}_n) + \frac{\lambda}{2} \int_\Omega |w_n|^2 dx - \frac{1}{6} \int_\Omega |u_n|^6 dx$$

where the last three terms are bounded (recall $\dim \tilde{\mathcal{V}} < \infty$). Hence also (v_n^+) is bounded. \square

Let

$$N(u) := |u|^4 u.$$

It is clear that $N : L^6(\Omega, \mathbb{R}^3) \rightarrow L^{6/5}(\Omega, \mathbb{R}^3)$. We shall need the following version of the Brezis-Lieb lemma:

Lemma 6.4. *Suppose (u_n) is bounded in $L^6(\Omega, \mathbb{R}^3)$ and $u_n \rightarrow u$ a.e. in Ω . Then*

$$N(u_n) - N(u_n - u) \rightarrow N(u) \quad \text{in } L^{6/5}(\Omega, \mathbb{R}^3) \text{ as } n \rightarrow \infty.$$

Proof. Since $N(u_n) - N(u_n - u) \rightarrow N(u)$ a.e. in Ω and $N(u_n) - N(u_n - u)$ is bounded in $L^{6/5}(\Omega, \mathbb{R}^3)$, $N(u_n) - N(u_n - u) \rightharpoonup N(u)$. We claim that $|N(u_n) - N(u_n - u)|_{6/5} \rightarrow |N(u)|_{6/5}$. Using Vitali's convergence theorem we obtain

$$\begin{aligned} \int_\Omega \left| |u_n|^4 u_n - |u_n - u|^4 (u_n - u) \right|^{6/5} dx &= \int_\Omega \int_0^1 \frac{d}{dt} \left| |u_n + (t-1)u|^4 (u_n + (t-1)u) \right|^{6/5} dt dx \\ &= \int_\Omega \int_0^1 \frac{d}{dt} |u_n + (t-1)u|^6 dt dx = 6 \int_0^1 \int_\Omega \langle |u_n + (t-1)u|^4 (u_n + (t-1)u), u \rangle dx dt \\ &\rightarrow 6 \int_0^1 \int_\Omega t^5 |u|^6 dx dt = \int_\Omega |u|^6 dx. \end{aligned}$$

Hence $N(u_n) - N(u_n - u)$ converges strongly to $N(u)$. \square

Lemma 6.5. *Let $\beta < c_0$. Then J_λ satisfies the $(PS)_\beta$ -condition in \mathcal{N}_λ , i.e. if $(u_n) \subset \mathcal{N}_\lambda$, $J_\lambda(u_n) \rightarrow \beta$ and $J'_\lambda(u_n) \rightarrow 0$ as $n \rightarrow \infty$, then $u_n \rightarrow u_0 \neq 0$ in $W_0^6(\text{curl}; \Omega)$ along a subsequence. In particular, u_0 is a nontrivial solution for (1.7)–(1.8).*

Proof. Let (u_n) be a $(PS)_\beta$ -sequence such that $(u_n) \subset \mathcal{N}_\lambda$. According to Lemma 6.3, (u_n) is bounded and we may assume $u_n \rightharpoonup u_0$ in $W_0^6(\text{curl}; \Omega)$. By Lemma 6.2, $u_n \rightarrow u_0$ in $L^2(\Omega, \mathbb{R}^3)$ and hence also a.e. in Ω after passing to a subsequence if necessary. As in the proof of Theorem 1.2 in Section 4 we see that $J'_\lambda(u_0) = 0$, i.e. u_0 is a solution for (1.7)–(1.8). According to the Brezis-Lieb lemma [9],

$$\lim_{n \rightarrow \infty} \left(\int_{\Omega} |u_n|^6 dx - \int_{\Omega} |u_n - u_0|^6 dx \right) = \int_{\Omega} |u_0|^6 dx,$$

hence

$$(6.3) \quad \lim_{n \rightarrow \infty} (J_\lambda(u_n) - J_\lambda(u_n - u_0)) = J_\lambda(u_0) \geq 0,$$

and by Lemma 6.4,

$$(6.4) \quad \lim_{n \rightarrow \infty} (J'_\lambda(u_n) - J'_\lambda(u_n - u_0)) = J'_\lambda(u_0) = 0.$$

Since $J'_\lambda(u_n) \rightarrow 0$ and $u_n \rightarrow u_0$ in $L^2(\Omega, \mathbb{R}^3)$,

$$(6.5) \quad \lim_{n \rightarrow \infty} J'_0(u_n - u_0) = 0.$$

Suppose $\liminf_{n \rightarrow \infty} \|u_n - u_0\| > 0$. Since $\lim_{n \rightarrow \infty} J'_0(u_n - u_0)(u_n - u_0) = 0$, we infer that

$$\liminf_{n \rightarrow \infty} |\nabla \times (u_n - u_0)|_2 > 0.$$

Let $u_n - u_0 = v_n + \tilde{w}_n \in \mathcal{V} \oplus \mathcal{W}$ according to the Helmholtz decomposition in $W_0^6(\text{curl}; \Omega)$. If $v_n \rightarrow 0$ in $L^6(\Omega, \mathbb{R}^3)$, then by (6.5) we have $J'_0(u_n - u_0)v_n \rightarrow 0$, thus

$$|\nabla \times (u_n - u_0)|_2^2 = |\nabla \times v_n|_2^2 = J'_0(u_n - u_0)v_n + \int_{\Omega} |u_n - u_0|^4 (u_n - u_0), v_n dx \rightarrow 0$$

as $n \rightarrow \infty$ which is a contradiction. Therefore $|v_n|_6$ is bounded away from 0. If $w_n := w(u_n - u_0) \in \mathcal{W}$, then (w_n) is bounded and since $u_n - u_0 + w_n = v_n + w(v_n) \in \mathcal{V} \oplus \mathcal{W}$, $|u_n - u_0 + w_n|_6$ is bounded away from 0. Choose t_n so that $t_n(u_n - u_0 + w_n) \in \mathcal{N}_0$ ($\mathcal{N}_0 \equiv \mathcal{N}_\Omega$ in the notation of Section 5). As in (5.7) we have

$$t_n^2 = \frac{|\nabla \times (u_n - u_0)|_2}{|u_n - u_0 + w_n|_6^3},$$

so (t_n) is bounded. Using Lemma 5.2, as in the proof of Theorems 1.1 and 1.3 we get

$$J_0(u_n - u_0) \geq J_0(t_n(u_n - u_0 + w_n)) - J'_0(u_n - u_0) \left[\frac{t_n^2 - 1}{2} (u_n - u_0) + t_n^2 w_n \right],$$

so by (6.5) and since $u_n \rightarrow u_0$ in $L^2(\Omega, \mathbb{R}^3)$,

$$\beta = \lim_{n \rightarrow \infty} J_\lambda(u_n - u_0) = \lim_{n \rightarrow \infty} J_0(u_n - u_0) \geq \lim_{n \rightarrow \infty} J_0(t_n(u_n - u_0 + w_n)) \geq c_0,$$

a contradiction. Therefore, passing to a subsequence, $u_n \rightarrow u_0$. Since $u_0 \in \mathcal{N}_\lambda$, $u_0 \neq 0$. \square

Proof of Theorem 1.4. (a) It follows from (6.2) and Lemma 6.5 that if $c_\lambda < c_0$, then c_λ is attained and hence there exists a ground state solution. By Lemma 6.1, this inequality is satisfied whenever $\lambda \leq \lambda_{\nu-1}$ and $\lambda \in (-\lambda_\nu, -\lambda_\nu + \bar{S}_{\text{curl}}(\Omega)|\Omega|^{-2/3})$.

In view of [23, Theorem 2.2(b)], the function $(-\lambda_\nu, -\lambda_{\nu-1}] \ni \lambda \mapsto c_\lambda \in (0, +\infty)$ is non-decreasing, continuous and $c_\lambda \rightarrow 0$ as $\lambda \rightarrow -\lambda_\nu^-$, and if $c_{\mu_1} = c_{\mu_2}$ for some $-\lambda_\nu < \mu_1 < \mu_2 \leq -\lambda_{\nu-1}$, then c_λ is not attained for $\lambda \in (\mu_1, \mu_2]$. Hence (b) and (c) follow.

(d) Since J_λ is even and, by Lemma 6.5, satisfies the Palais-Smale condition in \mathcal{N}_λ at any level below c_0 , then, in view of [23, Theorem 3.2(c)], J_λ has at least $m(\mathcal{N}_\lambda, c_0)$ pairs of critical points $\pm u$ such that $u \neq 0$ and $c_\lambda \leq J_\lambda(u) < c_0$ where

$$(6.6) \quad m(\mathcal{N}_\lambda, c_0) := \sup\{\gamma(J_\lambda^{-1}((0, \beta]) \cap \mathcal{N}_\lambda) : \beta < c_0\}$$

and γ is the Krasnoselskii genus [30]. This is a consequence of the standard fact that if

$$\beta_k := \inf\{\beta \in \mathbb{R} : \gamma(J_\lambda^{-1}((0, \beta]) \cap \mathcal{N}_\lambda) \geq k\},$$

then there are at least as many pairs of critical points as the number of k for which $(PS)_{\beta_k}$ holds, see e.g. [30].

In order to complete the proof we show that

$$m(\mathcal{N}_\lambda, c_0) \geq \tilde{m}_\lambda := \#\left\{k : -\lambda_k < \lambda < -\lambda_k + \frac{1}{3}\overline{S}_{\text{curl}}(\Omega)|\Omega|^{-\frac{2}{3}}\right\}.$$

Let

$$A(\lambda) := \left\{k \geq 1 : -\lambda_k < \lambda < -\lambda_k + \frac{1}{3}\overline{S}_{\text{curl}}(\Omega)|\Omega|^{-\frac{2}{3}} \text{ and } \lambda_k > \lambda_{k-1}\right\}$$

and observe that

$$\tilde{m}_\lambda = \sum_{k \in A(\lambda)} m(\lambda_k),$$

where $m(\lambda_k)$ stands for the multiplicity of λ_k . For $k \in A(\lambda)$, let $\mathcal{V}(\lambda_k)$ denote the eigenspace corresponding to λ_k . Then $\dim \mathcal{V}(\lambda_k) = m(\lambda_k)$. Let $S(\lambda)$ be the unit sphere in $\bigoplus_{k \in A(\lambda)} \mathcal{V}(\lambda_k) \subset \mathcal{V}^+$. Recall that $m_\lambda|_{\mathcal{S}^+}$ is a homeomorphism from \mathcal{S}^+ to \mathcal{N}_λ . Since J_λ is even, m_λ is odd. Similarly as in Lemma 6.1 we show that for $u \in S(\lambda)$

$$J_\lambda(m_\lambda(u)) \leq \max_{k \in A(\lambda)} \frac{1}{3}(\lambda + \lambda_k)^{\frac{3}{2}}|\Omega| =: \beta$$

and thus $m_\lambda(S(\lambda)) \subset J_\lambda^{-1}((0, \beta]) \cap \mathcal{N}_\lambda$. Hence

$$\gamma(J_\lambda^{-1}(0, \beta]) \cap \mathcal{N}_\lambda \geq \gamma(S(\lambda)) = \tilde{m}_\lambda.$$

Since $\lambda < -\lambda_k + \frac{1}{3}\overline{S}_{\text{curl}}(\Omega)|\Omega|^{-\frac{2}{3}}$, we have $\beta < c_0$ and it follows that $m(\mathcal{N}_\lambda, c_0) \geq \tilde{m}_\lambda$ which completes the proof. \square

7. OPEN PROBLEMS

In this section we state some open problems. Some of them have already been mentioned earlier.

- (P1) Does there exist a ground state solution u whose support is a proper subset of \mathbb{R}^3 ? In particular, can a ground state have compact support?
- (P2) Can one find an explicit expression for a ground state? Or at least, what can be said about the decaying properties of ground states? If they are the same as for the Aubin-Talenti instantons, then one could hopefully retrieve the formulas in the middle of p. 35 in [36] which could be useful when looking for ground states for (1.6) with the right-hand side $|u|^4u + g(x, u)$ where g is a monotone lower order term.
- (P3) Do the solutions to (1.6) have any symmetry properties? How regular are they?

- (P4) If Ω is a bounded domain which is neither convex nor has $\mathcal{C}^{1,1}$ boundary, then $\mathcal{V} \subset H^s(\Omega, \mathbb{R}^3)$ where $s \in [1/2, 1]$ and s may be strictly less than 1, see Subsection 2.2 and [12]. Note that the critical exponent for H^s is $6/(3 - 2s) < 6$ if $s < 1$. Do the results of Theorem 1.4 remain valid (with the same right-hand side)? Here the boundary condition (1.8) should be understood in the generalized sense, i.e. u should be in $W_0^6(\text{curl}; \Omega)$.
- (P5) Can the inequality $S_{\text{curl}} \geq \bar{S}_{\text{curl}}(\Omega) \geq S$ be sharpened? Do there exist domains as in (P4) for which $\bar{S}_{\text{curl}}(\Omega) < S$?

Acknowledgements. J. Mederski was partially supported by the National Science Centre, Poland (Grant No. 2017/26/E/ST1/00817). He was also partially supported by the Alexander von Humboldt Foundation (Germany) and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 258734477 – SFB 1173 during the stay at Karlsruhe Institute of Technology.

REFERENCES

- [1] G. Agrawal: *Nonlinear Fiber Optics*, Academic Press 2013, 5-th Edition.
- [2] N.N. Akhmediev, A. Ankiewicz, J.M. Soto-Crespo: *Does the nonlinear Schrödinger equation correctly describe beam propagation?*, Opt. Lett. **18** (1993), 411.
- [3] C. Amrouche, C. Bernardi, M. Dauge, V. Girault: *Vector potentials in three-dimensional non-smooth domains*, Math. Methods Appl. Sci. **21** (1998), no. 9, 823–864.
- [4] T. Aubin: *Équations différentielles non linéaires et problème de Yamabe concernant la courbure scalaire*, J. Math. Pures Appl. (9) **55** (1976), no. 3, 269–296.
- [5] T. Bartsch, T. Dohnal, M. Plum, W. Reichel: *Ground states of a nonlinear curl-curl problem in cylindrically symmetric media*, Nonlinear Diff. Eq. Appl. **23:52** (2016), 34 pp.
- [6] T. Bartsch, J. Mederski: *Ground and bound state solutions of semilinear time-harmonic Maxwell equations in a bounded domain*, Arch. Rational Mech. Anal., 215 (1), (2015), 283–306.
- [7] T. Bartsch, J. Mederski: *Nonlinear time-harmonic Maxwell equations in an anisotropic bounded medium*, J. Funct. Anal. **272** (2017), no. 10, 4304–4333.
- [8] V. Benci, D. Fortunato: *Towards a unified field theory for classical electrodynamics*, Arch. Rat. Mech. Anal. **173** (2004), 379–414.
- [9] H. Brézis, E. Lieb: *A relation between pointwise convergence of functions and convergence of functionals*, Proc. Amer. Math. Soc. **88** (1983), no. 3, 486–490.
- [10] H. Brezis, L. Nirenberg: *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983), 437–477.
- [11] A. Ciattoni, B. Crossignani, P. Di Porto, A. Yariv: *Perfect optical solitons: spatial Kerr solitons as exact solutions of Maxwell's equations*, J. Opt. Soc. Am. B **22** (2005), 1384–94.
- [12] M. Costabel: *A remark on the regularity of solutions of Maxwell's equations on Lipschitz domains*, Math. Methods Appl. Sci. **12**, (1990), 365–368.
- [13] R. Dautray, J.L. Lions: *Mathematical Analysis and Numerical Methods for Science and Technology, Vol. 3*, Springer 1990.
- [14] A. Desyatnikov, A. Maimistov, B. Malomed: *Three-dimensional spinning solitons in dispersive media with the cubic-quintic nonlinearity*, Phys. Rev. E. **61**(3), (2000), 3107–3113.
- [15] W. Dörfler, A. Lechleiter, M. Plum, G. Schneider, C. Wieners: *Photonic Crystals: Mathematical Analysis and Numerical Approximation*, Springer 2012.
- [16] L. C. Evans: *Weak convergence methods for nonlinear partial differential equations*, CBMS Regional Conference Series in Mathematics # 74, AMS, Providence, RI 1990.
- [17] V. Girault, P.-A. Raviart: *Finite Element Methods for Navier-Stokes Equations: Theory and Algorithms*, Springer 1986.
- [18] A. Kirsch, F. Hettlich: *The Mathematical Theory of Time-Harmonic Maxwell's Equations: Expansion-, Integral-, and Variational Methods*, Springer 2015.
- [19] H. Leinfelder: *Gauge invariance of Schrödinger operators and related spectral properties*, J. Operator Theory 9 (1983), 163–179.
- [20] E.H. Lieb: *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*, Ann. of Math. (2) **118** (1983), no. 2, 349–374.

- [21] P.L. Lions: *The concentration-compactness principle in the calculus of variations. The limit case. Part I and II.*, Rev. Mat. Iberoamer. **1** (1985), no. 1, 145–201 and no. 2, 45–121.
- [22] J. Mederski: *Ground states of time-harmonic semilinear Maxwell equations in \mathbb{R}^3 with vanishing permittivity*, Arch. Rational Mech. Anal. **218** (2), (2015), 825–861.
- [23] J. Mederski: *The Brezis-Nirenberg problem for the curl-curl operator*, J. Funct. Anal. **274** (5) (2018), 1345–1380.
- [24] J. Mederski, J. Schino, A. Szulkin: *Multiple solutions to a nonlinear curl-curl problem in \mathbb{R}^3* , Arch. Rational Mech. Anal., DOI: 10.1007/s00205-019-01469-3, Online First.
- [25] D. Mihalache, D. Mazilu, L.-C. Crasovan, I. Towers, A.V. Buryak, B.A. Malomed, L. Torner: *Stable spinning solitons in three dimensions*, Phys. Rev. Lett. **88**(7), 4 (2002).
- [26] P. Monk: *Finite Element Methods for Maxwell's Equations*, Oxford University Press 2003.
- [27] Z. Nehari: *Characteristic values associated with a class of non-linear second-order differential equations*, Acta Math. **105** (1961), 141–175.
- [28] A. Pankov: *Periodic Nonlinear Schrödinger Equation with Application to Photonic Crystals*, Milan J. Math. **73** (2005), 259–287.
- [29] S. Solimini: *A note on compactness-type properties with respect to Lorentz norms of bounded subsets of a Sobolev space*, Ann. Inst. H. Poincaré Anal. Non Linéaire **12** (1995), no. 3, 319–337.
- [30] M. Struwe: *Variational Methods*, Springer 2008.
- [31] C. A. Stuart: *Guidance Properties of Nonlinear Planar Waveguides*, Arch. Rational Mech. Anal. **125** (1993), no. 1, 145–200.
- [32] A. Szulkin, T. Weth: *Ground state solutions for some indefinite variational problems*, J. Funct. Anal. **257** (2009), no. 12, 3802–3822.
- [33] G. Talenti: *Best constant in Sobolev inequality*, Ann. Mat. Pura Appl. (4) **110** (1976), 353–372.
- [34] K. Tintarev, K.-H. Fieseler: *Concentration Compactness: Functional-analytic Grounds And Applications*, Imperial College Press 2007.
- [35] S. Waliullah, *Minimizers and symmetric minimizers for problems with critical Sobolev exponent*, Topol. Meth. Nonl. Anal. **34** (2009), 291–326.
- [36] M. Willem: *Minimax Theorems*, Birkhäuser 1996.

(J. Mederski)

INSTITUTE OF MATHEMATICS,
POLISH ACADEMY OF SCIENCES,
UL. ŚNIADECKICH 8, 00-656 WARSAW, POLAND
AND

CRC 1173 WAVE PHENOMENA: ANALYSIS AND NUMERICS,
DEPARTEMENT OF MATHEMATICS, KARLSRUHE INSTITUTE OF TECHNOLOGY (KIT),
D-76128 KARLSRUHE, GERMANY
Email address: jmederski@impan.pl

(A. Szulkin)

DEPARTMENT OF MATHEMATICS,
STOCKHOLM UNIVERSITY,
106 91 STOCKHOLM, SWEDEN
Email address: andrzej@math.su.se