

CONDENSED MATTER PHYSICS

Enhanced quantum coherence in exchange coupled spins via singlet-triplet transitions

Y. Bae^{1,2,3*}, K. Yang^{2*}, P. Willke^{1,2,3}, T. Choi^{1,3}, A. J. Heinrich^{1,3†}, C. P. Lutz^{2†}

Manipulation of spin states at the single-atom scale underlies spin-based quantum information processing and spintronic devices. These applications require protection of the spin states against quantum decoherence due to interactions with the environment. While a single spin is easily disrupted, a coupled-spin system can resist decoherence by using a subspace of states that is immune to magnetic field fluctuations. Here, we engineered the magnetic interactions between the electron spins of two spin-1/2 atoms to create a “clock transition” and thus enhance their spin coherence. To construct and electrically access the desired spin structures, we use atom manipulation combined with electron spin resonance (ESR) in a scanning tunneling microscope. We show that a two-level system composed of a singlet state and a triplet state is insensitive to local and global magnetic field noise, resulting in much longer spin coherence times compared with individual atoms. Moreover, the spin decoherence resulting from the interaction with tunneling electrons is markedly reduced by a homodyne readout of ESR. These results demonstrate that atomically precise spin structures can be designed and assembled to yield enhanced quantum coherence.

INTRODUCTION

The coherent control of spin states is a prerequisite for the use of spins in quantum information technologies (1–3). However, the quantum properties of spin states in solid-state nanostructures are easily disrupted by interactions with the environment such as electric or magnetic field noise (4), as well as unwanted coupling to nearby spins (5, 6). To protect the spin states against decoherence, ion traps (7, 8), silicon-based qubits (9), and quantum dots (10–12) adopted particular spin transitions called “clock transitions,” which are inherently robust against magnetic field fluctuations (7). By carefully tuning the parameters in the spin Hamiltonian of a coupled electron-nuclear (7, 8) or electron-electron system (9), these clock transition-based spin qubits have been created and shown to be insensitive to magnetic field noise, at least to first order.

To experimentally address sources of decoherence, well-controlled studies of individual spin centers are critical (13). Scanning tunneling microscopy (STM) has been intensively used to construct and characterize spin structures (14, 15). While the spin relaxation time (T_1) of individual atoms (16, 17), molecules (18), and nanostructures (19–21) has been studied using STM, the spin coherence time (T_2) of surface atoms is mostly discussed for individual atoms (17, 22) and in theoretical works (23–25). Recently, electron spin resonance (ESR) in STM has been applied to electrically sense and control individual magnetic atoms on the surface (26) as well as interactions between them (27–29). Combining the high-energy resolution of ESR and the capability of STM to position individual spin centers with atomic precision, ESR-STM now enables the exploration of decoherence in assembled nanostructures.

In this work, we create a two-level system using magnetic field-independent spin states of two magnetically coupled spin-1/2 titanium (Ti) atoms. The spacing between the atoms is precisely chosen to create a relatively strong magnetic coupling (~30 GHz) that protects the spin states from fluctuating magnetic fields. The two-level system consists of

the singlet and triplet states having magnetic quantum number $m = 0$, and thus, it is not sensitive to magnetic field fluctuations to first order (3). This gives a spin coherence time that is more than one order of magnitude longer compared with other states in this system of coupled atoms and with individual Ti atoms. We further improve the coherence time by setting the DC bias voltage to zero to reduce decoherence induced by tunneling electrons (22). This is achieved by using homodyne detection, a mechanism previously used in electrical detection of ferromagnetic resonance (30, 31) and here applied to ESR-STM.

RESULTS

Spin resonance of singlet and triplet states

We used a low-temperature STM that allows imaging, atom manipulation, and single-atom ESR (Fig. 1A) (26–29). One or a few Fe atoms were transferred to the tip apex to create a magnetic tip for ESR driving and sensing. We deposited Ti atoms on two monolayers (ML) of MgO grown on Ag(001) (see Materials and Methods and section S1). On this surface, Ti atoms have two binding sites: on top of the oxygen atom (Ti_O) and at the bridge site between two oxygen atoms (Ti_B). Both species have a spin of 1/2, most likely due to an attached H atom (29), and show negligible magnetic anisotropy; thus, to good approximation, the spins align to the uniform external magnetic field \mathbf{B}_ext (fig. S2).

We positioned two Ti atoms to form Ti_O - Ti_B dimers (Fig. 1A) and characterized the magnetic interactions between Ti atoms using ESR. During ESR data acquisition, the STM tip is positioned over the Ti_O atom because it yields a better ESR signal amplitude than Ti_B (fig. S10). Furthermore, this position compensates for the subtle difference of the gyromagnetic ratios of Ti_O and Ti_B atoms (figs. S2 and S6). When two spin-1/2 atoms are magnetically coupled, the eigenstates are given by the singlet ($|S\rangle$) and triplet ($|T_0\rangle$, $|T_-\rangle$, $|T_+\rangle$) states. While two of the triplet states are the Zeeman product states ($|T_-\rangle = |00\rangle$ and $|T_+\rangle = |11\rangle$), the spin-spin interaction causes the superposition of $|01\rangle$ and $|10\rangle$ states and results in the remaining two eigenstates: $|S\rangle = (|01\rangle - |10\rangle)/\sqrt{2}$ and $|T_0\rangle = (|01\rangle + |10\rangle)/\sqrt{2}$. Here, 0 and 1 designate the spin-up and spin-down states of the constituting spins, respectively.

Figure 1B shows an ESR spectrum obtained from a Ti_O - Ti_B dimer with the atomic separation $r = 0.92$ nm. Four ESR peaks arise from the four transitions that change the total magnetic quantum

Copyright © 2018
The Authors, some
rights reserved;
exclusive licensee
American Association
for the Advancement
of Science. No claim to
original U.S. Government
Works. Distributed
under a Creative
Commons Attribution
NonCommercial
License 4.0 (CC BY-NC).

Downloaded from <http://advances.sciencemag.org/> on June 4, 2020

¹Center for Quantum Nanoscience, Institute for Basic Science, Seoul 03760, Republic of Korea. ²IBM Almaden Research Center, San Jose, CA 95120, USA. ³Department of Physics, Ewha Womans University, Seoul 03760, Republic of Korea.

*These authors contributed equally to this work.

†Corresponding author. Email: heinrich.andreas@qns.science (A.J.H.); cplutz@us.ibm.com (C.P.L.)




Fig. 1. Spin resonance for two coupled spin-1/2 Ti atoms. (A) Schematic of the ESR-STM setup with the topographic image of a pair of Ti atoms on 2 ML MgO, where the Ti atoms are separated by $r = 0.92$ nm. The two species appear with different apparent heights in the STM image: ~ 1 Å for Ti at the O binding site of MgO (Ti_O) and ~ 1.8 Å for Ti at a bridge site (Ti_B) ($V_\text{DC} = 40$ mV, $I = 10$ pA, $T = 1.2$ K). The external magnetic field is applied almost parallel to the surface. (B and C) ESR spectrum measured on Ti_O in a Ti_O - Ti_B dimer with (B) $r = 0.92$ nm and (C) $r = 0.72$ nm [$V_\text{DC} = 40$ mV, $T = 1.2$ K, $B_\text{ext} = 0.9$ T; (B) $I = 10$ pA, $V_\text{RF} = 30$ mV; (C) $I = 20$ pA, $V_\text{RF} = 15$ mV]. Insets: STM images of the Ti_O - Ti_B dimer used to measure each ESR spectrum. The grid intersections indicate the positions of oxygen atoms of the MgO lattice. The mark “x” shows the position of the tip in the ESR measurement. (D and E) Schematic energy level diagrams for two coupled spin-1/2 atoms. In (D), the Zeeman energy is larger than the interaction energy J between two atoms, leading to the triplet state as the ground state. In (E), the singlet state becomes the ground state when J is larger than the Zeeman energy. The resonance peaks in (B) and (C) are marked by the same colors as transition labels in (D) and (E), respectively.

number m by ± 1 , as given in the schematic energy diagram (Fig. 1D). The peak heights are proportional to the difference in thermal occupation of the initial and final states (29); thus, the two tallest peaks correspond to transitions out of the ground state. The difference between resonance frequencies ($f_{T_0T_+} - f_{ST_+}$ or, equivalently, $f_{ST_+} - f_{T_0T_+}$) directly gives the magnetic interaction energy J between the two spins (29), which is 0.77 ± 0.02 GHz for this dimer spacing (fig. S5). Because the interaction energy J is smaller than the Zeeman energy, the $|T_- \rangle$ state becomes the ground state.

We find that the Ti_O - Ti_B dimer can be positioned close enough to yield coupling strengths sufficiently strong to shift the singlet state down in energy to become the ground state (Fig. 1E). This interaction strength was not accessible for the Ti_O - Ti_O dimers (29). Decreasing the spacing of the atoms in the Ti_O - Ti_B dimer to $r = 0.72$ nm (Fig. 1C) results in three ESR peaks in our measurement range (5 to 30 GHz). In addition to the two triplet-triplet transitions ($f_{T_0T_+}$ and $f_{T_0T_-}$), the singlet-triplet (S - T_0) transition is now visible as resonance at $f_{ST_0} \approx 29$ GHz. Here, the resonance frequency f_{ST_0} directly gives the value of J when the detuning (see below) is negligible.

While a traditional ESR measurement applies a radio frequency (RF) magnetic field, in the ESR-STM technique used here, the RF magnetic field at the Ti atom arises from the modulation of the atom’s position (32) in the nonuniform magnetic field B_tip (29, 33) generated by the STM tip. This enables the ESR transition of $\Delta m = 0, \pm 1$. The triplet-triplet transitions in the dimer are driven by a gradient in B_tip^x , the component perpendicular to B_ext . This is the same component required for driving the $|0\rangle$ to $|1\rangle$ transition for an individual Ti atom ($\Delta m = \pm 1$; section S6). In contrast, driving the S - T_0 transition ($\Delta m = 0$) requires a gradient in B_tip^z , the component of B_tip parallel to B_ext (12). In this work, we chose a tip having both spatial components of B_tip , which therefore can drive the transitions of $\Delta m = 0, \pm 1$ in the dimer.

Heisenberg exchange coupled spin-1/2 Ti atoms

From the ESR peak splitting, we determined the magnetic interaction energy J for 30 dimers with different separations and orientations (section S3). To exclude the effects of B_tip on the resonance frequencies, we measured the ESR spectra as a function of B_tip and determined the value of J from the fit based on the spin Hamiltonian (section S4). The measured values of J are given in Fig. 2A as a function of atomic separations (r , ranging from 0.72 to 1.3 nm). We find that for atomic distances of less than 1 nm, the Ti_O - Ti_B dimers are dominantly coupled by the Heisenberg exchange interaction $J\mathbf{S}_1 \cdot \mathbf{S}_2$, where \mathbf{S}_1 and \mathbf{S}_2 are the spin operators. Moreover, the interaction is found to be isotropic (fig. S3).

The exchange interaction generally shows exponential dependence on the separation between spins (34). Given the isotropic interaction energy $J = J_0 \exp[-(r - r_0)/d]$ (34) and taking $r_0 = 0.72$ nm, we obtain for Ti_O - Ti_B dimers a decay constant $d = 64.6 \pm 4.9$ pm and a prefactor $J_0 = 28.9 \pm 1.3$ GHz. The decay constant matches well with reported values for exchange interactions across a vacuum gap (29, 33, 35). For Ti_O - Ti_O and Ti_B - Ti_B dimers, we obtain $d = 40.0 \pm 2.0$ pm (29) and 94.0 ± 0.3 pm (fig. S3D), respectively. This difference in decay constant between the dimer types indicates the sensitivity of the exchange interaction to either the orbitals being involved in the interaction or the spatial distribution of spin density (36), resulting from the different interaction potentials (34) and the different magnetic ground states (29). As determined from the intensity of peaks in the ESR spectra (fig. S3) (27, 29), J is positive, and thus, the coupling between Ti atom spins is antiferromagnetic.

Energy detuning of superposition states

While the nonuniform magnetic field arising from the STM tip is necessary to drive the spin resonance, this tip magnetic field also provides a means to control the quantum states by applying a local




Fig. 2. Singlet-triplet energy detuning of Ti_O - Ti_B dimers with different interaction energies. (A) Magnetic interaction energy determined from ESR measurements for Ti_O - Ti_B dimers with different atomic separations. The red line shows the exponential fit, indicative of Heisenberg exchange interaction. The slight deviation of the Ti_O - Ti_B interaction energy from the exponential fit is due to the contribution from the dipole-dipole interaction at larger distances. (B) The ESR frequency shift of the S- T_0 transition (Δf_{ST_0}) for dimers with different J as a function of the magnitude of the tip field (B_{tip}). B_{tip} is calibrated for each tip that we used (section S4). For the dimers with $J = 0.5, 0.8,$ and 3 GHz, the resonance frequencies are obtained by $f_{\text{ST}_0} = f_{\text{T}_0\text{T}_1} - f_{\text{ST}_1}$; for the dimers with $J = 9$ and 30 GHz, f_{ST_0} is directly obtained from ESR spectra. Strengthening the exchange interaction between Ti atoms protects the $|S\rangle$ and $|T_0\rangle$ states from detuning by B_{tip} , reducing Δf_{ST_0} . (C) First-order tip field dependence of f_{ST_0} for the dimers in (B). The clock transitions appear at $B_{\text{tip}} = 38 \pm 12$ mT, where $df_{\text{ST}_0}/dB_{\text{tip}} = 0$.

magnetic field to one atom in the dimer. The eigenstates deviate from ideal singlet and triplet states because of an energy detuning ϵ , which is the difference in Zeeman energies between the two atoms. The detuning arises from two sources: (i) the tip magnetic field that is applied only on one of the atoms (29) and (ii) a slight difference in the gyromagnetic ratios γ_1 and γ_2 for the two atoms at different binding sites (fig. S2). The Hamiltonian (in units of angular frequency) describing the two spins dominantly coupled by the exchange interaction is then given by

$$H = \gamma_1 \mathbf{S}_1 \cdot (\mathbf{B}_{\text{ext}} + \mathbf{B}_{\text{tip}}) + \gamma_2 \mathbf{S}_2 \cdot \mathbf{B}_{\text{ext}} + 2\pi J \mathbf{S}_1 \cdot \mathbf{S}_2 \quad (1)$$

Under the approximation that \mathbf{B}_{tip} is parallel to \mathbf{B}_{ext} , the energy detuning is given by $\epsilon = [(\gamma_1 - \gamma_2)B_{\text{ext}} + \gamma_1 B_{\text{tip}}]/2\pi$ (29), resulting in the quantum eigenstates

$$\begin{aligned} |T_0(\xi)\rangle &= \sin\frac{\xi}{2}|01\rangle + \cos\frac{\xi}{2}|10\rangle \\ |S(\xi)\rangle &= \cos\frac{\xi}{2}|01\rangle - \sin\frac{\xi}{2}|10\rangle \end{aligned}$$

where ξ is a mixing parameter given by $\tan \xi = J/\epsilon$. When the energy detuning is negligible ($J \gg \epsilon$), the eigenstates are the ideal singlet and triplet states: $|S\rangle = (|01\rangle - |10\rangle)/\sqrt{2}$ and $|T_0\rangle = (|01\rangle + |10\rangle)/\sqrt{2}$. In contrast, increasing the energy detuning leads to the Zeeman product states: $|01\rangle$ and $|10\rangle$.

The effect of this energy detuning on the eigenstates is a shift of their energy levels, which results in a corresponding ESR frequency shift of the S- T_0 transition (Δf_{ST_0}) from the minimum value of f_{ST_0} . Figure 2B shows the measured Δf_{ST_0} for Ti_O - Ti_B dimers as a function of B_{tip} for different values of J . The minimum in f_{ST_0} is reached at $B_{\text{tip}} = 38 \pm 12$ mT (Fig. 2C), where the transition constitutes a magnetic field-independent clock transition to first order. At this field, the detuning is absent,

that is, $\epsilon = 0$, because the tip field fully compensates for the subtle difference in magnetic moments of the Ti_O and Ti_B atoms (fig. S2).

We calculated the eigenvalues and eigenstates (fig. S4) using the Hamiltonian in Eq. 1 to fit the experimental results. When \mathbf{B}_{tip} is parallel to \mathbf{B}_{ext} , the singlet-triplet energy detuning is given by $\Delta f_{\text{ST}_0} = \sqrt{J^2 + \epsilon^2} - J$. However, it is necessary to account for the tilting of \mathbf{B}_{tip} with respect to \mathbf{B}_{ext} to obtain adequate fits to the ESR frequencies for all allowed transitions in each dimer of Fig. 2. For the STM tips used in Fig. 2, we find that the angles between \mathbf{B}_{tip} and \mathbf{B}_{ext} fall in the range of 21° to 51° depending on the tip apex (section S4).

For weakly coupled dimers ($J < 1$ GHz), a slight increase of energy detuning due to B_{tip} shifts f_{ST_0} considerably from its minimum value. Using the model Hamiltonian of Eq. 1, at the typical tip field of 110 mT, the eigenstates of the dimer with $J = 0.8$ GHz are $|S(\xi)\rangle = 0.92|01\rangle - 0.39|10\rangle$ and $|T_0(\xi)\rangle = 0.39|01\rangle + 0.92|10\rangle$, which more closely resemble the Zeeman product states.

We find that the effects of Zeeman energy detuning (ϵ) on Δf_{ST_0} can be regulated by the coupling strength of two atoms. As shown in Fig. 2C, increasing J to 30 GHz ($J \gg \epsilon$) markedly reduces the sensitivity of f_{ST_0} on the magnetic field variation. At the same tip field ($B_{\text{tip}} = 110$ mT), the eigenstates now remain almost in the ideal singlet and triplet states ($|S(\xi)\rangle = 0.71|01\rangle - 0.70|10\rangle$ and $|T_0(\xi)\rangle = 0.70|01\rangle + 0.71|10\rangle$). Thus, in the following, we ensure that $J \gg \epsilon$ by keeping B_{tip} small (< 150 mT) and by using large J (30 GHz) and show that this choice results in a decoherence-free subspace.

Enhanced spin coherence using magnetic field-independent states

On the basis of the results from the previous sections, we now focus on the spin coherence times of strongly coupled Ti_O - Ti_B dimers ($r = 0.72$ nm, $J \approx 30$ GHz). The spin coherence for the singlet-triplet transition and its sensitivity to the external and local magnetic fields are compared in Fig. 3 with (i) the triplet-triplet transition of the same dimer and (ii) the $|0\rangle$ to $|1\rangle$ transition of an individual Ti_O atom. We obtained the spin coherence time of each transition by fitting the




Fig. 3. Spin coherence of ESR transitions and their sensitivity to external and local magnetic fields. (A) ESR peak width as a function of V_{RF} for the $S-T_0$ and T_0-T_- transitions measured on TiO in a strongly coupled dimer ($r = 0.72$ nm, $J \approx 30$ GHz), and the $|0\rangle$ to $|1\rangle$ transition of an individual TiO atom ($V_{DC} = 40$ mV, $I = 10$ pA, $B_{tip} = 110$ mT, $B_{ext} = 0.9$ T, $T = 1.2$ K). Solid lines are fits to $\Gamma = \sqrt{1 + AV_{RF}^2/\pi T_2^*}$, derived from the Bloch equation model, where the spin coherence time T_2^* is determined by the intercept at the y axis and A is a constant. (B) ESR frequencies as a function of the external magnetic field B_{ext} . For the $S-T_0$ transition, the frequency f_{ST_0} remains almost constant, characteristic of a clock transition. Inset: Energy diagram for the four eigenstates at different B_{ext} ($V_{DC} = 40$ mV, $I = 10$ pA, $B_{tip} = 110$ mT, $T = 1.2$ K). (C) ESR frequencies as a function of the tip magnetic field B_{tip} . B_{tip} is set by the junction impedance (V_{DC}/I) and calibrated from the fit (red curves; see also section S4). For the $S-T_0$ transition, the frequency f_{ST_0} remains almost constant and measurably increases when B_{tip} is larger than 150 mT, which reflects the change of eigenstates from the ideal singlet and triplet states. Inset: Energy diagram at different B_{tip} ($V_{DC} = 40$ mV, $I = 10$ pA, $B_{ext} = 0.9$ T, $T = 1.2$ K).




Fig. 4. Homodyne detection and enhanced spin coherence of the $S-T_0$ transition. (A) The tip field effects on ESR line shape of the $S-T_0$ transition. The ESR spectra are normalized and vertically offset. (B) DC bias dependence of the ESR signals for the T_0-T_- transition (top) and $S-T_0$ transition (bottom). For the $S-T_0$ transition, homodyne detection allows V_{DC} to be decreased without losing signal intensity ($B_{tip} = 110$ mT, $V_{RF} = 20$ mV, $T = 1.2$ K). (C) Spin coherence time, T_2^* , as a function of V_{DC} . Red curves are reciprocal fits. At fixed junction impedance ($V_{DC} = 40$ mV, $I = 10$ pA), T_2^* increases with lowering V_{DC} because of the reduction of tunneling electrons per unit time. For the $S-T_0$ transition, setting V_{DC} to zero provides further improvement in the spin coherence time by reducing the DC tunneling current. Labels #1 and #2 indicate different dimers (measured with different tips) with the same separation ($r = 0.72$ nm) to confirm the reproducibility of T_2^* .

ESR linewidth Γ to the Bloch equation model (26) (section S5), as a function of RF voltage (V_{RF}) (Fig. 3A). In the limit of small V_{RF} , the coherence time is given by $1/\pi\Gamma$. This coherence time includes the effect of inhomogeneous line broadening and is designated T_2^* to distinguish it from the intrinsic spin coherence time T_2 . In our single-spin experiment, inhomogeneous broadening may be due to any time-varying magnetic fields that are present to give temporal ensemble broadening (37).

For typical ESR conditions and $B_{tip} = 110$ mT, we find $T_2^* = 99.0 \pm 9.7$ ns for the $S-T_0$ transition (Fig. 3A). Under the same conditions, T_2^* of the triplet-triplet (T_0-T_-) transition and of the individual TiO is ~ 8 and ~ 20 times smaller, at only 13.0 ± 0.3 ns and 5.3 ± 0.7 ns, respectively.

The spin coherence time is closely related to the sensitivity of spin states to the time-varying external and local magnetic fields. The sensitivity to uniform external magnetic fields was characterized by varying the external field magnitude B_{ext} from 0.5 to 1.1 T (Fig. 3B). ESR frequencies $f_{T_0T_-}$ and $f_{T_0T_+}$ ($\Delta m = \pm 1$ transition) shift linearly with B_{ext}

due to the Zeeman effect on the states $|T_- \rangle$ and $|T_+ \rangle$. In contrast, f_{ST_0} ($\Delta m = 0$ transition) shows no Zeeman shift and remains nearly independent of B_{ext} , an essential property of a clock transition (7–9).

We now investigate the effect of a local magnetic field by varying B_{tip} over a large range, extending from 10 mT to 0.45 T (Fig. 3C). For the transitions between triplet states, the resonance frequencies $f_{T_0T_-}$ and $f_{T_0T_+}$ again increase steadily by the Zeeman energy owing to B_{tip} applied to one atom in the dimer. In contrast, f_{ST_0} remains essentially constant when B_{tip} is lower than ~ 150 mT. The results in Fig. 3 show that the singlet-triplet transition is insensitive to both external and local magnetic field fluctuations, which results in the measured increase of its spin coherence time.

Homodyne detection as a means to spin decoherence reduction

In addition to magnetic field fluctuations, tunneling electrons are a major source of decoherence of the surface atom's spin in magnetoresistively

sensed ESR (22). Here, we show how to achieve further improvements in T_2^* based on the ESR detection mechanism.

In the ESR spectrum of the dimer (Fig. 1C), a notable difference between the singlet-triplet (S- T_0) and triplet-triplet (T_0 - T_+) transitions is the line shape of ESR peaks. For the individual Ti atoms (fig. S2) or triplet-triplet transitions, the ESR signal is nearly symmetric. In contrast, the S- T_0 transition appears antisymmetric for low B_{tip} (Fig. 4A). As the tip field increases, the ESR line shape becomes more symmetric. Since the ESR detection mechanism depends on the nature of the spin states, these changes in the ESR line shape for the S- T_0 transition in Fig. 4A are a direct consequence of the states changing from the $|S\rangle$ and $|T_0\rangle$ states toward the Zeeman product states ($|01\rangle$ and $|10\rangle$) as B_{tip} increases (section S5).

Figure 4B shows the ESR spectra for the T_0 - T_+ and S- T_0 transitions at $B_{\text{tip}} = 110$ mT, where the superposition states more closely approximate the $|S\rangle$ and $|T_0\rangle$ states. For the T_0 - T_+ transition, the nearly symmetric ESR signal results from the change in time-average population of spin states for the atom under the tip (26), as detected by V_{DC} . Thus, the peak amplitude decreases with decreasing V_{DC} (Fig. 4B, top). For the S- T_0 transition, the time-average population of spin states of the atom under the tip does not vary; thus, it cannot be detected by DC conductance changes. However, the magnetization of the atom along the quantization axis oscillates in time during ESR. The oscillating magnetoconductance at the frequency of the driving voltage V_{RF} is multiplied by V_{RF} to produce a DC tunnel current, which can thus be detected. This rectification is known as a homodyne detection (for a full description of the ESR line shape, see section S5) (30, 31). Thus, in the case of the S- T_0 transition, both driving and sensing the spin resonance signal can be achieved by using V_{RF} only, enabling us to set V_{DC} to zero. In Fig. 4B, we find that, for the S- T_0 resonance signal, the peak width is narrower for lower V_{DC} . As a result, we find that at $V_{\text{DC}} = 0$, the ESR signal of the S- T_0 transition is the sharpest because the tunneling current due to V_{DC} is absent.

As seen in Fig. 4C, the coherence times T_2^* for all transitions observed increase rapidly with decreasing V_{DC} . Since nearly every tunneling electron induces decoherence of the surface spin (22), reducing the number of tunneling electrons improves the spin coherence significantly. At $V_{\text{DC}} = 0$, we obtain $T_2^* = 257 \pm 80$ ns for the S- T_0 transition. Note that the ESR measurement at $V_{\text{DC}} = 0$ is only possible for the S- T_0 transition (Fig. 4B). Although we set V_{DC} to zero, the remaining tunneling current generated by V_{RF} , the finite temperature (22), and the relatively short spin relaxation time T_1 (29, 38) limit the spin coherence time of the S- T_0 transition, resulting in the deviation of T_2^* from the reciprocal curve in Fig. 4C.

DISCUSSION

By controlling the magnetic coupling between electron spins of two atoms, we have demonstrated robust singlet and triplet states and achieved a significantly enhanced spin coherence time. Both driving and sensing the singlet-triplet transition do not require a DC voltage, providing an additional way to improve the spin coherence. As a result, we achieved a large improvement of spin coherence by a factor of about 10 compared with the triplet-triplet transition in the same dimer. Moreover, this exceeds the spin coherence time previously determined for individual Fe atoms (26), despite the much shorter spin relaxation time T_1 for individual Ti atoms (29). These engineered atomic-scale magnetic structures may serve as the smallest component for assembling custom spin chains and arrays with enhanced quantum coherence times. The ability of ESR-STM to construct desired multispin

systems and to electrically access their many-body states might enable the exploration of quantum phases, spintronic information processing, and quantum simulation.

MATERIALS AND METHODS

Experiments were performed using a home-built STM system at the IBM Almaden Research Center. We evaporated Ti atoms onto a cold (< 10 K) 2 ML MgO grown on Ag(001). The MgO layer was used to decouple the spin of Ti atoms from the underlying substrate electrons (39). Previous works showed that the Ti atoms are likely hydrogenated because of residual hydrogen gas in the vacuum chamber (29, 40), and here, we denote the hydrogenated Ti atoms simply as Ti. An external magnetic field (\mathbf{B}_{ext}) was applied nearly in-plane. An RF voltage V_{RF} was applied across the tunnel junction for driving spin resonance, and a DC bias voltage V_{DC} was applied for the DC magnetoresistive sensing of the spin states (Fig. 1A) (26).

SUPPLEMENTARY MATERIALS

Supplementary material for this article is available at <http://advances.sciencemag.org/cgi/content/full/4/11/eaau4159/DC1>

Section S1. Binding site of Ti atoms on 2 ML MgO/Ag(001)

Section S2. ESR of individual Ti atoms

Section S3. Magnetic interaction of Ti dimers

Section S4. Spin Hamiltonian of two coupled Ti spins

Section S5. Detection mechanism of ESR

Section S6. Driving mechanism of ESR

Section S7. Determination of spin coherence time

Fig. S1. Tunneling current as a function of the tip-atom distance for T_{Ib} (blue) and T_{Io} (red) on 2 ML MgO/Ag(001) at $V_{\text{DC}} = 10$ mV.

Fig. S2. ESR of individual Ti atoms.

Fig. S3. Characterization of magnetic interaction between Ti atoms.

Fig. S4. Tip field effects on eigenenergies and eigenstates of T_{Io} - T_{Ib} dimers.

Fig. S5. Resonance frequencies measured on each atom of a weakly coupled dimer ($r = 0.92$ nm, $J \approx 0.8$ GHz) at different tip fields.

Fig. S6. ESR frequencies measured on each atom of strongly coupled dimers ($r = 0.72$ nm, $J \approx 30$ GHz) at different tip fields.

Fig. S7. ESR signals for the S- T_0 transition with different DC biases.

Fig. S8. Determination of spin coherence time at different DC biases and temperatures.

Fig. S9. Spin coherence time with different DC biases (or different tunneling currents) for individual T_{Io} and T_{Ib} atoms.

Fig. S10. Comparison of ESR signals for the singlet-triplet transition measured on the T_{Io} (black) and T_{Ib} (green) atoms in the T_{Io} - T_{Ib} dimer.

REFERENCES AND NOTES

1. P. W. Shor, Scheme for reducing decoherence in quantum computer memory. *Phys. Rev. A* **52**, R2493 (1995).
2. A. M. Steane, Error correcting codes in quantum theory. *Phys. Rev. Lett.* **77**, 793–797 (1996).
3. D. A. Lidar, I. L. Chuang, K. B. Whaley, Decoherence-free subspaces for quantum computation. *Phys. Rev. Lett.* **81**, 2594–2597 (1998).
4. P. Jaminon, M. Lesik, J. P. Tetienne, I. Alvizu, L. Mayer, A. Dréau, S. Kosen, J.-F. Roch, S. Pezzagna, J. Meijer, T. Teraji, Y. Kubo, P. Bertet, J. R. Maze, V. Jacques, Competition between electric field and magnetic field noise in the decoherence of a single spin in diamond. *Phys. Rev. B* **93**, 024305 (2016).
5. A. M. Tyryshkin, S. Tojo, J. J. L. Morton, H. Riemann, N. V. Abrosimov, P. Becker, H.-J. Pohl, T. Schenkel, M. L. W. Thewalt, K. M. Itoh, S. A. Lyon, Electron spin coherence exceeding seconds in high-purity silicon. *Nat. Mater.* **11**, 143–147 (2012).
6. A. V. Khaetskii, D. Loss, L. Glazman, Electron spin decoherence in quantum dots due to interaction with nuclei. *Phys. Rev. Lett.* **88**, 186802 (2002).
7. J. Bollinger, J. D. Prestage, W. M. Itano, D. J. Wineland, Laser-cooled-atomic frequency standard. *Phys. Rev. Lett.* **54**, 1000–1003 (1985).
8. P. C. Haljan, P. J. Lee, K.-A. Brickman, M. Acton, L. Deslauriers, C. Monroe, Entanglement of trapped-ion clock states. *Phys. Rev. A* **72**, 062316 (2005).

9. G. Wolfowicz, A. M. Tyryshkin, R. E. George, H. Riemann, N. V. Abrosimov, P. Becker, H.-J. Pohl, M. L. W. Thewalt, S. A. Lyon, J. J. L. Morton, Atomic clock transitions in silicon-based spin qubits. *Nat. Nanotechnol.* **8**, 561–564 (2013).
10. J. R. Petta, A. C. Johnson, J. M. Taylor, E. A. Laird, A. Yacoby, M. D. Lukin, C. M. Marcus, M. P. Hanson, A. C. Gossard, Coherent manipulation of coupled electron spins in semiconductor quantum dots. *Science* **309**, 2180–2184 (2005).
11. F. H. L. Koppens, C. Buizert, K. J. Tielrooij, I. T. Vink, K. C. Nowack, T. Meunier, L. P. Kouwenhoven, L. M. Vandersypen, Driven coherent oscillations of a single electron spin in a quantum dot. *Nature* **442**, 766–771 (2006).
12. S. Foletti, H. Bluhm, D. Mahalu, V. Umansky, A. Yacoby, Universal quantum control of two-electron spin quantum bits using dynamic nuclear polarization. *Nat. Phys.* **5**, 903–908 (2009).
13. N. Zhao, J. Honert, B. Schmid, M. Klas, J. Isoya, M. Markham, D. Twitchen, F. Jelezko, R. B. Liu, H. Fedder, J. Wrachtrup, Sensing single remote nuclear spins. *Nat. Nanotechnol.* **7**, 657–662 (2012).
14. R. Wiesendanger, Spin mapping at the nanoscale and atomic scale. *Rev. Mod. Phys.* **81**, 1495–1550 (2009).
15. C. F. Hirjibehedin, C. P. Lutz, A. J. Heinrich, Spin coupling in engineered atomic structures. *Science* **312**, 1021–1024 (2006).
16. S. Loth, M. Etzkorn, C. P. Lutz, D. M. Eigler, A. J. Heinrich, Measurement of fast electron spin relaxation times with atomic resolution. *Science* **329**, 1628–1630 (2010).
17. J. Hermenau, M. Ternes, M. Steinbrecher, R. Wiesendanger, J. Wiebe, Long spin relaxation times in a transition metal atom in direct contact to a metal substrate. *Nano Lett.* **18**, 1978–1983 (2018).
18. B. W. Heinrich, L. Braun, J. I. Pascual, K. J. Franke, Protection of excited spin states by a superconducting energy gap. *Nat. Phys.* **9**, 765–768 (2013).
19. S. Loth, S. Baumann, C. P. Lutz, D. M. Eigler, A. J. Heinrich, Bistability in atomic-scale antiferromagnets. *Science* **335**, 196–199 (2012).
20. A. A. Khajetoorians, B. Baxevanis, C. Hübner, T. Schlenk, S. Krause, T. O. Wehling, S. Lounis, A. Lichtenstein, D. Pfannkuche, J. Wiebe, R. Wiesendanger, Current-driven spin dynamics of artificially constructed quantum magnets. *Science* **339**, 55–59 (2013).
21. S. Yoshida, Y. Aizawa, Z.-h. Wang, R. Oshima, Y. Mera, E. Matsuyama, H. Oigawa, O. Takeuchi, H. Shigekawa, Probing ultrafast spin dynamics with optical pump–probe scanning tunnelling microscopy. *Nat. Nanotechnol.* **9**, 588–593 (2014).
22. P. Willke, W. Paul, F. D. Natterer, K. Yang, Y. Bae, T. Choi, J. Fernández-Rossier, A. J. Heinrich, C. P. Lutz, Probing quantum coherence in single-atom electron spin resonance. *Sci. Adv.* **4**, eaag1543 (2018).
23. J.-P. Gauyacq, N. Lorente, Decoherence-governed magnetic-moment dynamics of supported atomic objects. *J. Phys. Condens. Matter* **27**, 455301 (2015).
24. F. Delgado, J. Fernández-Rossier, Spin decoherence of magnetic atoms on surfaces. *Prog. Surf. Sci.* **92**, 40–82 (2017).
25. A. M. Shalikhov, A. N. Rubtsov, A. I. Lichtenstein, P. Ribeiro, Relaxation and decoherence of qubits encoded in collective states of engineered magnetic structures. *Phys. Rev. B* **96**, 094410 (2017).
26. S. Baumann, W. Paul, T. Choi, C. P. Lutz, A. Ardavan, A. J. Heinrich, Electron paramagnetic resonance of individual atoms on a surface. *Science* **350**, 417–420 (2015).
27. T. Choi, W. Paul, S. Rolf-Pissarczyk, A. J. Macdonald, F. D. Natterer, K. Yang, P. Willke, C. P. Lutz, A. J. Heinrich, Atomic-scale sensing of the magnetic dipolar field from single atoms. *Nat. Nanotechnol.* **12**, 420–424 (2017).
28. F. D. Natterer, K. Yang, W. Paul, P. Willke, T. Choi, T. Greber, A. J. Heinrich, C. P. Lutz, Reading and writing single-atom magnets. *Nature* **543**, 226–228 (2017).
29. K. Yang, Y. Bae, W. Paul, F. D. Natterer, P. Willke, J. L. Lado, A. Ferrón, T. Choi, J. Fernández-Rossier, A. J. Heinrich, C. P. Lutz, Engineering the eigenstates of coupled spin-1/2 atoms on a surface. *Phys. Rev. Lett.* **119**, 227206 (2017).
30. H. Kubota, A. Fukushima, K. Yakushiji, T. Nagahama, S. Yuasa, K. Ando, H. Maehara, Y. Nagamine, K. Tsunekawa, D. D. Djayaprawira, N. Watanabe, Y. Suzuki, Quantitative measurement of voltage dependence of spin-transfer torque in MgO-based magnetic tunnel junctions. *Nat. Phys.* **4**, 37–41 (2008).
31. T. Nozaki, Y. Shiota, S. Miwa, S. Murakami, F. Bonell, S. Ishibashi, H. Kubota, K. Yakushiji, T. Saruya, A. Fukushima, S. Yuasa, T. Shinjo, Y. Suzuki, Electric-field-induced ferromagnetic resonance excitation in an ultrathin ferromagnetic metal layer. *Nat. Phys.* **8**, 491–496 (2012).
32. J. L. Lado, A. Ferrón, J. Fernández-Rossier, Exchange mechanism for electron paramagnetic resonance of individual atoms. *Phys. Rev. B* **96**, 205420 (2017).
33. S. Yan, D.-J. Choi, J. A. J. Burgess, S. Rolf-Pissarczyk, S. Loth, Control of quantum magnets by atomic exchange bias. *Nat. Nanotechnol.* **10**, 40–45 (2015).
34. R. Ahlrichs, H. J. Böhm, S. Brode, Interaction potentials for alkali ion-rare gas and halogen ion-rare gas systems. *J. Chem. Phys.* **88**, 6290 (1988).
35. R. Schmidt, C. Lazo, U. Kaiser, A. Schwarz, S. Heinze, R. Wiesendanger, Quantitative measurement of the magnetic exchange interaction across a vacuum gap. *Phys. Rev. Lett.* **106**, 257202 (2011).
36. J. H. Rodríguez, D. E. Wheeler, J. K. McCusker, Density functional studies of a Heisenberg spin coupled chromium–semiquinone complex and its chromium–catechol analog. *J. Am. Chem. Soc.* **120**, 12051–12068 (1998).
37. A. Abragam, B. Bleaney, *Electron Paramagnetic Resonance of Transition Ions* (Oxford Univ. Press, 2012).
38. X. R. Wang, Y. S. Zheng, S. Yin, Spin relaxation and decoherence of two-level systems. *Phys. Rev. B* **72**, 121303(R) (2005).
39. W. Paul, K. Yang, S. Baumann, N. Romming, T. Choi, C. P. Lutz, A. J. Heinrich, Control of the millisecond spin lifetime of an electrically probed atom. *Nat. Phys.* **13**, 403–407 (2017).
40. F. D. Natterer, F. Patthey, H. Brune, Quantifying residual hydrogen adsorption in low-temperature STMs. *Surf. Sci.* **615**, 80–87 (2013).

Acknowledgments: We thank B. Melior for expert technical assistance. **Funding:** We acknowledge financial support from the Office of Naval Research. Y.B., P.W., T.C., and A.J.H. acknowledge support from the Institute for Basic Science under grant IBS-R027-D1. P.W. acknowledges support from the Alexander von Humboldt Foundation. **Author contributions:** Y.B. and K.Y. conceived the experiment. Y.B. wrote the manuscript with support from all authors. Y.B., K.Y., and P.W. carried out the measurements and analyzed the data. All authors interpreted the results. A.J.H. and C.P.L. supervised the project. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials. Additional data related to this paper may be requested from the authors.

Submitted 8 June 2018
 Accepted 10 October 2018
 Published 9 November 2018
 10.1126/sciadv.aau4159

Citation: Y. Bae, K. Yang, P. Willke, T. Choi, A. J. Heinrich, C. P. Lutz, Enhanced quantum coherence in exchange coupled spins via singlet-triplet transitions. *Sci. Adv.* **4**, eaau4159 (2018).

Enhanced quantum coherence in exchange coupled spins via singlet-triplet transitions

Y. Bae, K. Yang, P. Willke, T. Choi, A. J. Heinrich and C. P. Lutz

Sci Adv 4 (11), eaau4159.
DOI: 10.1126/sciadv.aau4159

ARTICLE TOOLS

<http://advances.sciencemag.org/content/4/11/eaau4159>

SUPPLEMENTARY MATERIALS

<http://advances.sciencemag.org/content/suppl/2018/11/05/4.11.eaau4159.DC1>

REFERENCES

This article cites 39 articles, 7 of which you can access for free
<http://advances.sciencemag.org/content/4/11/eaau4159#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)

Science Advances (ISSN 2375-2548) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science Advances* is a registered trademark of AAAS.

Copyright © 2018 The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works. Distributed under a Creative Commons Attribution NonCommercial License 4.0 (CC BY-NC).