

Review

Advanced 3D Cell Culture Techniques in Micro-Bioreactors, Part I: A Systematic Analysis of the Literature Published between 2000 and 2020

Christoph Grün ¹, **Brigitte Altmann** ^{2,3,§} and **Eric Gottwald** ^{1,*,\$} ¹ Institute of Functional Interfaces, Group 3D Cell Culture Systems, Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany; christoph.gruen@kit.edu² Department of Prosthetic Dentistry, Center for Dental Medicine, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, 79106 Freiburg, Germany; brigitte.altmann@uniklinik-freiburg.de³ G.E.R.N Center for Tissue Replacement, Regeneration & Neogenesis, Department of Prosthetic Dentistry, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, 79108 Freiburg, Germany

* Correspondence: eric.gottwald@kit.edu; Tel.: +49-721-608-26803

§ The authors contributed equally to this work.

Received: 31 October 2020; Accepted: 12 December 2020; Published: 15 December 2020



Abstract: Bioreactors have proven useful for a vast amount of applications. Besides classical large-scale bioreactors and fermenters for prokaryotic and eukaryotic organisms, micro-bioreactors, as specialized bioreactor systems, have become an invaluable tool for mammalian 3D cell cultures. In this systematic review we analyze the literature in the field of eukaryotic 3D cell culture in micro-bioreactors within the last 20 years. For this, we define complexity levels with regard to the cellular 3D microenvironment concerning cell–matrix-contact, cell–cell-contact and the number of different cell types present at the same time. Moreover, we examine the data with regard to the micro-bioreactor design including mode of cell stimulation/nutrient supply and materials used for the micro-bioreactors, the corresponding 3D cell culture techniques and the related cellular microenvironment, the cell types and in vitro models used. As a data source we used the National Library of Medicine and analyzed the studies published from 2000 to 2020.

Keywords: micro-bioreactor; 3D cell culture; scaffolds; microfluidics; material; cellular microenvironment; tissue engineering; HTS; in vitro models

1. Introduction

Micro-bioreactors (MBRs) represent specialized bioreactor systems that were, unlike their big brothers, namely classical bioreactors for biotechnological applications or industrial production, developed for growing mammalian cells and/or tissues *in vitro* on a smaller scale. In fact, MBRs may be considered as an intermediate stage towards more complex organ-on-a-chip (OoC) systems as they incorporate design elements of classical bioreactors, e.g., stirred or perfused flasks/chambers instead of large-scale tanks, and novel technologies including microfluidic components, cell-scaffolds, and setups enabling microscopic analysis and cell-based assays which are also part of OoC systems. Despite the technological advances in the field of OoC systems, MBRs are still prevalent in cell-based biomedical research and have become invaluable tools in eukaryotic 3D cell culture research because they (i) provide controllable 3D cell culture conditions usually including an active fluid flow supply, (ii) enable, unlike large-scale bioreactors, the usage of small amounts of chemical entities and low cell numbers when primary cell and/or tissue availability is limited, and (iii) depending on the design, are frequently amenable to microscopic imaging, or other, even more sophisticated analysis techniques,

which facilitates cell culture evaluation during bioreactor operation. This has led to an ever-increasing amount of applications which provide insights that have not been possible before. In this context, a multitude of in vitro studies using 3D cell culture techniques individually or in combination with MBRs identified the following parameters which are relevant for the modulation of cell behavior: (i) spatial and temporal gradients of signaling molecules (e.g., growth factors, cytokines, and hormones), (ii) spatial distribution of cell–extracellular matrix (ECM) interactions, which are also inevitably coupled with the molecular and mechanical ECM properties, (iii) the spatial distribution of homologous and heterologous cell–cell contacts, and (iv) biomechanical forces emerging from interstitial fluid flow and tissue deformation.

Against this background and within the scope of this special issue of *Processes* bearing the same title, we would like to give an overview of the work published in the last 20 years in the area of 3D cell culture in micro-bioreactors and to thereby address the following questions:

- What has been done in the last 20 years in the field of micro-bioreactor design and what are the applications of such systems?
- Which systems have prevailed up till now?
- What are currently the most common techniques of 3D cell culture-based MBRs?
- Where is the journey taking us?

To answer these questions, we reviewed the studies in the field of 3D cell culture in MBRs between the years 2000 and 2020 (9 July 2020) systematically with respect to MBR design, the corresponding 3D cell culture techniques and the related cellular microenvironment, the mode of cell stimulation and/or nutrient supply, the materials used for MBRs and scaffold fabrication, the applications of the systems and the used cell type or in vitro model.

In this first part of the review, we analyze the published papers in terms of MBR design; this includes the used 3D cell culture technique, the mode of fluid flow and MBR materials, the applications of such systems and the origin and type of the cells in the last 20 years. On the basis of the literature research results we have structured the different approaches for better clarity by means of the complexity of the corresponding 3D cell culture techniques used in the systems, hereafter referred to as “complexity level”, defined the main fluid flow mode and classified the micro-bioreactor types according to their principal design. Furthermore, as organ-on-a-chip (OoC) systems are sometimes very hard to distinguish from MBRs with incorporated microfluidics we try to elucidate what distinguishes MBRs from OoC systems, although barriers are fluent.

In review part two, we provide a more detailed description of the results of our literature research structured according to the aforementioned complexity levels and with focus on the existing MBR types and their applications, as well as on common cell/tissue types cultured in such systems. We then disseminate the different types of MBR applications including simulation studies and finally give an outlook to future developments in the field of 3D cell culture in MBR. With this review, we hope to be able to adequately introduce the reader into the great contributions of this special issue of *Processes* on Advanced 3D-Cell Culture Techniques in Micro-Bioreactors.

2. Methods and Definitions

For the detection and analysis of the relevant literature, the PubMed® database of the National Library of Medicine of the National Center for Biotechnology Information (NCBI) at the U.S. National Library of Medicine (<https://pubmed.ncbi.nlm.nih.gov/>) was used. The following search terms were defined to include different spellings of the keywords as well as their synonyms:

1. “3D cell culture” AND “microbioreactor”;
2. (bioreactor OR microbioreactor OR micro-bioreactor) AND (“three-dimensional cell culture” OR “3D cell culture” OR “3-D cell culture”);
3. (microbioreactor OR micro-bioreactor) AND “tissue engineering”.

Moreover, secondary literature meeting the same search term results were included (Figure 1). Based on the results of our literature research we could extract the following parameters which are characteristic for MBRs and were used to define MBRs in the present review:

- culture volumes of less than 500 mL;
- provision of an active fluid flow, being realized by a pump, a stirring mechanism, piston movements or mechanical movement of the cell-based constructs in culture medium;
- 3D cell culture approach with or without scaffold.




Figure 1. Number of publications yielded by the search terms in the period from 2000 to 2020. Term 1: “3D cell culture” AND “microbioreactor”; Term 2: (bioreactor OR microbioreactor OR micro-bioreactor) AND (“three-dimensional cell culture” OR “3D cell culture” OR “3-D cell culture”); Term 3: (microbioreactor OR micro-bioreactor) AND “tissue engineering”.

By using this search strategy, 326 papers were retrieved of which 92 were excluded because they did not fulfil above mentioned MBR-definitions or because information on the used bioreactor type was insufficient. In addition to 192 relevant publications, we found 26 reviews and 10 publications which were dedicated to the simulation of flow rates, shear stress, or also oxygen distribution (Figure 1). These studies are discussed in the second part of the review. A total of 192 publications was the basis of the analysis with the corresponding figures and all papers were analyzed with regard to the following parameters: complexity level, bioreactor type, materials used for fabrication, scaffold- or non-scaffold-based culture technique, species origin of the cells used, type of organ, type of cells, and application.

Since 3D cell culture comprises many different techniques, we defined as mentioned before, complexity levels with regard to the culture configuration to structure the different strategies. Cellular complexity levels of the 3D cell culture configurations were defined as follows (Figure 2A):

1. Complexity level 1: cells immobilized in (hydro-)gels as monoculture (one cell type) or coculture (at least two cell types);
2. Complexity level 2: multicellular aggregates consisting of one cell type in 3D scaffolds or in scaffold-free cultures;
3. Complexity level 3: multicellular aggregates consisting of at least two cell types in 3D scaffold-based or in scaffold-free cultures.




Figure 2. (A) Complexity levels defined by means of the applied 3D cell culture technique and with respect to the existing 3D microenvironment. (B) Differentiation between superfusion and perfusion of different 3D cell cultures. (C) Frequently used micro-bioreactor (MBR) types: 1. based on spinner flasks or stirred tank systems, 2. rotating wall vessels (RWVs), 3. fluidic MBR, 4. hollow fiber bioreactors, and 5. MBR for mechanical stimulation. The sketches show examples of the structure of the respective MBRs. Details are discussed in review part II.

With respect to the fluid flow mode in the context of microfluidic bioreactors, the term perfusion is now used almost inflationary for all fluidic models. In the medical context, perfusion is defined as the flushing of a blood vessel or the lymphatic system with a liquid. In a broader meaning, perfusion also refers to the fluid flow through three-dimensional cell constructs or scaffolds. Systems in which only cell layers are flushed, or tissues or scaffolds are flushed around them, do not fulfil these definitions. This is generally called superfusion or perifusion [1]. The perfusion of tissues/scaffolds can be clearly differentiated from superfusion. However, the distinction between superfusion and perfusion of (blood) vessels is difficult to determine in the current *in vitro* models. Therefore, in the context of this review, systems in which the medium flows through a channel surrounded by cells are considered to be part of a perfusion system (Figure 2B).

In order to realize a medium flow around or through the 3D cell-constructs/scaffolds or tissues in the MBRs, several bioreactor types have been established in the last years including (1) stirred systems or spinner flasks, (2) rotating wall vessels (RWVs), (3) microfluidic MBRs, and (4) hollow fiber bioreactors. Especially in the case of microfluidic MBR there are various designs enabling perfusion of the cell-based constructs/scaffolds in addition to superfusion (Figure 2C). Furthermore, in MBRs designed for mechanical stimulation ((5) in Figure 2C) the medium flow or mass transport through the 3D cell culture emerges by stretching or compressing the cell-based construct/scaffold or tissue. These are mostly in-house developments of the labs and details are presented in the second part of this review.

3. Results

In order to give the reader an overview of the MBR configurations, the materials used for fabrication, and 3D cell culture techniques employed in the available literature, we analyzed the following parameters in 5-year periods: (i) the complexity level of the underlying 3D cell culture technique (Figure 3), (ii) the frequencies of the MBR types (Figure 4), (iii) the materials used for MBR fabrication (Figure 5), (iv) the complexity level for scaffold-based and scaffold-free applications in relation to the MBR type (Figure 6), (v) the application field of the MBR types (Figure 7), and (vi) the type and origin of cells that are used for the different approaches (Figure 8).




Figure 3. Distribution of studies in the period from 2000 to 2020 with regard to complexity level (for definitions, see “Section 2”). The number of the respective publications is shown in brackets.

* status as per 9 July 2020.

Regarding the distribution of the studies over time with regard to publication date and complexity level, it can be seen from Figure 3 that applications of complexity level 2 contribute a considerable portion to the total number of publications (123 out of 192). Complexity levels 1 and 3 are less represented in this area and share a similar proportion with 29 and 40 publications, respectively. This distribution reflects the notion that 3D aggregates of the same cell type are the most popular model systems in the field of MBR culture.

Of the 192 publications falling in the time frame between 2000 and 2020 (9 July 2020), an increase in the publication number from 25 in the 5-year period of 2000–2004 to 43 in the period of 2005–2009, to 66 in the period of 2010–2014 can be observed. In the period of 2015–2019 a slight decrease to 57 publications in total can be detected. With respect to the temporal distribution of the study numbers within the complexity level groups, our analysis further demonstrates that the number of studies using cell culture configurations with complexity levels 1 and 2 in MBRs peaked from 2005 to 2014 and from 2010 to 2014, respectively. In contrast, the number of published studies on complexity level 3 approaches in MBRs increased continuously during the evaluation period. This trend indicates that cell culture models made of different cell types gain in importance within the area of 3D cell culture in MBRs.




Figure 4. Distribution of micro-bioreactor types from 2000 to 2020 in the literature under study.

If the analysis is done with regard to the used bioreactor type in the same 5-year periods, it is clearly demonstrated that MBR systems with integrated microfluidics are the most prominent ones, followed by RWV type bioreactors (Figure 4). The use of MBRs for mechanical stimulation and hollow fiber MBRs take an intermediate position and vary between the specified periods. Furthermore, it is interesting to note that MBR types, developed at a very early stage of 3D-bioreactor culture, such as spinner flasks and hollow fiber systems, are still used nowadays.

Materials used for micro-bioreactor fabrication




Figure 5. Distribution of studies in the period from 2000 to 2020 with regard to the materials used for MBR fabrication. The materials used for the most common types of MBR, namely fluidic MBR, RWVs and spinner flasks, MBR for mechanical stimulation and hollow fiber MBR. Abbreviations: PDMS (polydimethylsiloxane), PC (polycarbonate), PMMA (Poly(methylmethacrylate)), PS (Polystyrene), PU (polyurethane), PES (polyethersulfone), PVDF (polyvinylidene fluoride); “other polymers” include the publications with the less frequently applied materials stereolithography resin, polyethylene terephthalate, polylactic acid, polytetrafluoroethylene, PU (individually), polyethylene, polyoxymethylene, polypropylene, polyether ether ketone; the category “unknown materials” includes publications in which the applied MBR materials were not further specified.

When looking closer to the materials used for bioreactor manufacturing, it can be seen that polymer-based materials represent the major part. Polydimethylsiloxane (PDMS), individually or in combination with other polymers or glass, is thereby the most commonly used material and mainly

applied when microfluidics is asked to be incorporated (Figure 5). This may be due to numerous in-house developments and optimizations in the field of fluidic MBR as PDMS is comparatively easy to process and to shape. Occasionally, 3D printed systems are also used [2,3]. Another large group of widely used MBRs, the RWV bioreactor type, is usually based on commercially available systems, e.g., from Synthecon. These are available as plastic or glass versions. However, the RWV version used in the corresponding publications were not always specified and therefore summarized under unknown materials in Figure 5. In addition to the polymer-based materials, stainless steel, besides glass, is often used in microfluidic systems and MBRs for mechanical stimulation. As the latter MBR type is primarily used in bone or cartilage tissue reconstruction to simulate the *in vivo* situation by applying mechanical forces, tougher materials like stainless steel are used for construction to meet the biomechanical requirements. In the group of hollow fiber MBRs, polyethersulfone (PU) and polyvinylidenefluoride (PVDF) predominate as these materials are mainly used for the capillaries or hollow fibers of the MBR.

Cell culture techniques described in the analyzed literature




Figure 6. 3D cell culture techniques and their application described in the literature under study. The techniques were organized with regard to their complexity level, the use and nature of the scaffolds and their application in the various MBR types. The numbers in parentheses refer to the number of corresponding publications.




Figure 7. Application fields of the MBRs described in the analyzed literature with respect to the complexity levels. The dimensions of the pie charts reflect the number of publications in the respective application group and complexity level subgroup. Number of publications: 192.

As described earlier, parameters of the 3D microenvironment, such as gradients of signaling molecules, cell/cell- and cell/ECM-interactions, determine the cellular behavior *in vitro*. Hence, in order to induce or maintain tissue specific cell differentiation and functions, various cell culture techniques have been established which were classified in three complexity levels in the present review as described in Section 2 (Figure 2A). The techniques can further be divided in two groups, namely techniques using scaffolds for the immobilization and organization of the cells and scaffold-free techniques (Figure 6). Both techniques utilize the self-organizing, polarizing, and adhesion capabilities of the cells to form multicellular aggregates within scaffolds or without supporting structures. To give an overview of the distribution of the different techniques in the literature under study, we analyzed the described cell culture techniques with regard to complexity level, the use and nature of the scaffolds and their application in the various MBR types (Figure 6). The results from our literature analysis reveal that in the scaffold-based group the majority of studies used cell culture approaches with a complexity level of 2, i.e., multicellular aggregates of one single cell type (Figure 6). In the scaffold-free group, cell aggregates with one cell type (complexity level 2) also comprise a greater proportion when compared to complexity level 3 but is less strongly represented than compared to the scaffold-based group. If we take a look at the application of the specific cell culture techniques in the different MBR systems, our analysis demonstrates that gel-based scaffolds, composed, amongst others, of collagen, hyaluronan, polyethylene glycol, or Matrigel®, are most frequently used for 3D culture

in microfluidic MBR systems, followed by sponge-like and fibrous scaffolds, and microstructured substrates. Microbead-based scaffolds were exclusively described in studies using RWVs and thus could only be assigned to this type of MBR, whereas spheroids are commonly applied in RWVs and spinner flask/stirred tank systems. This distribution can be explained by the immobilizing and structuring function of the scaffolds and hollow fibers, which is a prerequisite for proper perfusion or superfusion in microfluidic and hollow fiber MBRs. In contrast, the concept of RWV systems is to provide the nutrient and gas supply of the cells by stirring or rotating movement of the 3D-cell constructs such as microbeads and spheroids. Regarding the techniques used frequently in MBRs for mechanical stimulation, gel-based, fibrous, and sponge-like scaffolds, as well as decellularized tissues play a relevant role (Figure 6) because they allow for transmission of the applied mechanical forces to the cells.




Figure 8. Distribution of the cells used in the publications under study with respect to the origin of the cells (A), the target tissues/organs (B), and the type of cells (C). Others: bladder, placenta, muscle, endometrium, skin, vaginal epithelium, retina, kidney, pancreas, nerves, connective tissue, and brain. Number of publications: 192.

Since 3D culture techniques have proven useful in many applications we grouped the papers into the three most common fields of application that could be extracted, namely “*in vitro* models”,

“tissue engineering”, and “high-throughput-screening (HTS)”, split them according to the complexity level in subgroups and examined the MBR distribution within the respective subgroups. For this, we deduced the main application area indicated in the papers or categorized them by the soundness and impact of the results when no main application was specified, or more than one application possibility was given. The presented results may therefore not be considered as strictly as Figure 7 proposes since there are smooth transitions possible, especially between the groups “in vitro models” and “tissue engineering”.

As presented in Figure 7, most of the analyzed studies could be assigned to the “in vitro models” group and were mainly done with cell cultures of complexity level 2, i.e., cell aggregates of one cell type. In this group, approaches that used two or more cell types for the in vitro model, i.e., complexity level 3, represented the second largest proportion or subgroup which was followed by complexity level 1 techniques using immobilized cells in (hydro-) gels. In the “tissue engineering” group again publications using complexity level 2 cell cultures predominate and the number of studies with complexity level 3 was lowest. The same trend holds true for the HTS group. The pie charts further reveal that microfluidic MBRs are most strongly represented throughout all groups, and prevail particularly in the HTS group. The latter might be due to the possibility to integrate analysis features into microfluidic MBRs which support data evaluation in HTS systems. The second largest share among the MBR systems are represented by RWVs and spinner flasks as they support the generation of uniform and complex cell aggregates which are often used for in vitro models. Table 1 lists publications that are included in our analysis.

Table 1. List of the 192 publications used for the analysis, broken down by complexity level and type of micro-bioreactor.

Complexity Level	Type of Micro-Bioreactor				
	Spinner Flask or Stirred Tank	RWVs and Derivatives	Fluidic Micro-Bioreactor	Hollow Fiber Micro-Bioreactor	MBR for Mechanical Stimulation
1		[4–5]	[6–24]		[25–32]
2	[33–35]	[36–71]	[3,24,72–132]	[133–144]	[145–153]
3	[154–156]	[157–165]	[2,166–190]	[191]	[192]

Since our literature analysis revealed that most publications under study involve “in vitro models” we were next interested in the distribution of the applied cell types and target tissues/organs in the MBR field (Figure 8). With respect to the origin of the cells, the examination of the published literature between 2000 and 2019 revealed that at the beginning of the 2000s, human and rodent cells were the most common models, applied in 41% (human cells and combinations) and 45% of the studies, respectively (Figure 8A). By comparison, the share of publications using cells from other mammals, mainly bovine for bone and cartilage models, was at 14% and thus relatively low. During the period under review, the usage of human cells individually or in combination with cells of other origin increased to about 80% by 2019, whereas publications using rodent and other mammalian models declined to 13% and 4%, and thus play a minor role in this research field by now. The reason for this trend becomes apparent when the distribution of the target tissues/organs is analyzed (Figure 8B). From this follows that most published studies address the liver, with the primary objective to establish liver models for metabolism and toxicity testing (data not shown). Since meanwhile it is commonly agreed upon that rodent hepatocytes do not adequately reflect human hepatocyte functions with respect to the panel of detoxification enzymes and proteins, in vitro models based on rodent primary cells lost most of their importance. A similar situation may prevail for other tissues displayed in Figure 8B. Regarding the type of cells cultured in the MBR systems, our literature research demonstrates that in the early 2000s a higher number of studies used primary cells, i.e., differentiated and/or mature cells, followed by stem and progenitor cells, and to a lesser extent cell lines (Figure 8C). Interestingly,

the application of primary cells considerably increased until 2014 and sharply decreased from 2015 to 2019 to the same level as cell lines which also increased to a lesser extent in the same period. The progress in the field of stem cell research however may lead to a moderate but constant increase of the publications using stem and progenitor cells, and thereby is gradually replacing primary cells and cell lines.

As mentioned at the beginning, it is not always easy to distinguish MBR approaches and organ-on-a-chip (OoC) systems, as definitions are not consistent and transitions are fluid. OoCs are defined as microengineered devices with microfluidic channels with at least one cell culture compartment in which functional units of organs are modeled [193–195]. Organ-on-a-chip systems do not only mimic human organs' in vivo physiology, for instance by providing an active flow, they also integrate actuators or sensors for further analysis.

To give the reader an idea of the number of published studies using OoC in comparison to MBR systems we conducted a second literature research with the term [("organ on chip" OR "organ-on-chip" OR "organ on a chip" OR "organ-on-a-chip") AND ("tissue engineering" OR "three dimensional cell culture" OR "3d cell culture" OR "3-d cell culture")]. As depicted in Figure 9 the number of recently published studies using 3D cell cultures in MBR has been almost constant in the last years, whereas the number of OoC-related publications has increased dramatically in the last 5 years. At second glance, however, the published systems do not always reach the complexity defined for OoCs, such as integrated sensors, respectively, actors or modular combination of different organ models. Although MBRs are often referred to as organ-on-chip systems [81], a distinction was made between OoCs and MBRs as described above. Since OoC systems are repeatedly considered to be among the top emerging technologies [196], it could be speculated that the use of this keyword is to increase the popularity of the results.




Figure 9. Search results in the database PubMed between 2000 and 2020 with the search terms 1–3 used in this review for micro-bioreactors (Term 1: 3D cell culture" AND "microbioreactor; Term 2: (bioreactor OR microbioreactor OR micro-bioreactor) AND ("three-dimensional cell culture" OR "3D cell culture" OR "3-D cell culture"); Term 3: (microbioreactor OR micro-bioreactor) AND "tissue engineering" compared to organs-on-a-chip (search term: ("organ on chip" OR "organ-on-chip" OR "organ on a chip" OR "organ-on-a-chip") AND ("tissue engineering" OR "three dimensional cell culture" OR "3d cell culture" OR "3-d cell culture")), status as per 27 October 2020).

4. Conclusions

The results of our systematic review revealed that MBRs became very popular between 2000 and 2014, with microfluidic MBRs being most strongly represented in the fields of fundamental research with in vitro models, tissue engineering, and HTS. With respect to the cell culture configuration used in the MBR systems during this period, 3D aggregates of the same cell type, consisting mainly of human primary cells or cell lines are the most frequently used technique in this period, irrespective of whether scaffold-based techniques are applied or scaffold-free ones. Our analysis further demonstrates that MBR systems established very early in bioreactor-based 3D culture and tissue engineering are still used until today, possibly due to the commercialization of the systems. In the last 5 years, however, a new trend can be observed towards more complex 3D culture systems with an increasing number of publications using 3D coculture models, i.e., cell cultures consisting of at least two cell types, and human cells. Within the group of human cells, stem and progenitor cells have become very popular. This trend towards more complex human stem/progenitor cell-based in vitro models requires however further miniaturization of the systems as cell/tissue sources, especially with respect to personalized medicine, are often limited or yield low cell amounts. Consequently, further developments in 3D cell culture techniques, such as unique media formulations for coculture models, and in cell function analysis of low cell numbers or even on a single-cell level will also advance in the next years. With respect to the bioreactor systems, this trend manifests by a decrease in the number of MBR publications which is accompanied by a pronounced increase in the organ-on-a-chip literature. This most likely reflects the progress made in manufacturing and/or cell culturing techniques as well as the increase in experience made by using MBRs.

In part II of the review (Systems and Applications) we will have a closer look at the micro-bioreactors and the specific applications that they have been used for.

Author Contributions: Conceptualization, B.A., C.G. and E.G.; methodology, C.G. and E.G.; formal analysis, C.G. and E.G.; investigation, B.A., C.G. and E.G.; resources, National Library of Medicine; data curation, C.G.; writing—original draft preparation, B.A., C.G. and E.G.; writing—review and editing, B.A., C.G. and E.G.; visualization, C.G.; supervision, E.G. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: We thank all colleagues that sent us those papers that we had no access to.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gaddum, J.H. The technique of superfusion. 1953. *Br. J. Pharmacol.* **1997**, *120*, 82–87. [CrossRef] [PubMed]
2. Daneshgar, A.; Tang, P.; Remde, C.; Lommel, M.; Feufel, M.A.; Kertzscher, U.; Klein, O.; Weinhart, M.; Pratschke, J.; Sauer, I.M.; et al. Teburu—Open source 3D printable bioreactor for tissue slices as dynamic three-dimensional cell culture models. *Artif. Organs* **2019**, *43*, 1035–1041. [CrossRef] [PubMed]
3. Schmid, J.; Schwarz, S.; Meier-Staude, R.; Sudhop, S.; Clausen-Schaumann, H.; Schieker, M.; Huber, R. A Perfusion Bioreactor System for Cell Seeding and Oxygen-Controlled Cultivation of Three-Dimensional Cell Cultures. *Tissue Eng. Part C Methods* **2018**, *24*, 585–595. [CrossRef] [PubMed]
4. Hwang, Y.S.; Cho, J.; Tay, F.; Heng, J.Y.; Ho, R.; Kazarian, S.G.; Williams, D.R.; Boccaccini, A.R.; Polak, J.M.; Mantalaris, A. The use of murine embryonic stem cells, alginate encapsulation, and rotary microgravity bioreactor in bone tissue engineering. *Biomaterials* **2009**, *30*, 499–507. [CrossRef] [PubMed]
5. Vecchiatini, R.; Penolazzi, L.; Lambertini, E.; Angelozzi, M.; Morganti, C.; Mazzitelli, S.; Trombelli, L.; Nastruzzi, C.; Piva, R. Effect of dynamic three-dimensional culture on osteogenic potential of human periodontal ligament-derived mesenchymal stem cells entrapped in alginate microbeads. *J. Periodontal Res.* **2015**, *50*, 544–553. [CrossRef] [PubMed]
6. Chang, R.; Nam, Y.; Sun, W. Direct cell writing of 3D microorgan for in vitro pharmacokinetic model. *Tissue Eng. Part. C-Me* **2008**, *14*, 157–166. [CrossRef] [PubMed]

7. Cui, Z.F.; Xu, X.; Trainor, N.; Triffitt, J.T.; Urban, J.P.; Tirlapur, U.K. Application of multiple parallel perfused microbioreactors and three-dimensional stem cell culture for toxicity testing. *Toxicol In Vitro* **2007**, *21*, 1318–1324. [[CrossRef](#)]
8. Goldman, S.M.; Barabino, G.A. Cultivation of agarose-based microfluidic hydrogel promotes the development of large, full-thickness, tissue-engineered articular cartilage constructs. *J. Tissue Eng. Regen Med.* **2017**, *11*, 572–581. [[CrossRef](#)]
9. Hahn, M.S.; McHale, M.K.; Wang, E.; Schmedlen, R.H.; West, J.L. Physiologic pulsatile flow bioreactor conditioning of poly(ethylene glycol)-based tissue engineered vascular grafts. *Ann. Biomed. Eng.* **2007**, *35*, 190–200. [[CrossRef](#)]
10. Huang, S.B.; Wu, M.H.; Wang, S.S.; Lee, G.B. Microfluidic cell culture chip with multiplexed medium delivery and efficient cell/scaffold loading mechanisms for high-throughput perfusion 3-dimensional cell culture-based assays. *Biomed. Microdevices* **2011**, *13*, 415–430. [[CrossRef](#)]
11. Huang, S.B.; Wang, S.S.; Hsieh, C.H.; Lin, Y.C.; Lai, C.S.; Wu, M.H. An integrated microfluidic cell culture system for high-throughput perfusion three-dimensional cell culture-based assays: Effect of cell culture model on the results of chemosensitivity assays. *Lab. Chip* **2013**, *13*, 1133–1143. [[CrossRef](#)] [[PubMed](#)]
12. Jaeger, A.A.; Das, C.K.; Morgan, N.Y.; Pursley, R.H.; McQueen, P.G.; Hall, M.D.; Pohida, T.J.; Gottesman, M.M. Microfabricated polymeric vessel mimetics for 3-D cancer cell culture. *Biomaterials* **2013**, *34*, 8301–8313. [[CrossRef](#)] [[PubMed](#)]
13. Li, Z.; Kreiner, M.; Edrada-Ebel, R.; Cui, Z.; van der Walle, C.F.; Mardon, H.J. Perfusion culture enhanced human endometrial stromal cell growth in alginate-multivalent integrin alpha5beta1 ligand scaffolds. *J. Biomed. Mater. Res. A* **2011**, *99*, 211–220. [[CrossRef](#)] [[PubMed](#)]
14. Lichtenberg, A.; Dumlu, G.; Walles, T.; Maringka, M.; Ringes-Lichtenberg, S.; Ruhparwar, A.; Mertsching, H.; Haverich, A. A multifunctional bioreactor for three-dimensional cell (co)-culture. *Biomaterials* **2005**, *26*, 555–562. [[CrossRef](#)]
15. Ling, Y.; Rubin, J.; Deng, Y.; Huang, C.; Demirci, U.; Karp, J.M.; Khademhosseini, A. A cell-laden microfluidic hydrogel. *Lab. Chip* **2007**, *7*, 756–762. [[CrossRef](#)]
16. Moretti, M.; Freed, L.E.; Padera, R.F.; Lagana, K.; Boschetti, F.; Raimondi, M.T. An integrated experimental-computational approach for the study of engineered cartilage constructs subjected to combined regimens of hydrostatic pressure and interstitial perfusion. *Biomed. Mater. Eng.* **2008**, *18*, 273–278. [[CrossRef](#)]
17. Pagano, G.; Ventre, M.; Iannone, M.; Greco, F.; Maffettone, P.L.; Netti, P.A. Optimizing design and fabrication of microfluidic devices for cell cultures: An effective approach to control cell microenvironment in three dimensions. *Biomicrofluidics* **2014**, *8*, 046503. [[CrossRef](#)]
18. Santoro, R.; Olivares, A.L.; Brans, G.; Wirz, D.; Longinotti, C.; Lacroix, D.; Martin, I.; Wendt, D. Bioreactor based engineering of large-scale human cartilage grafts for joint resurfacing. *Biomaterials* **2010**, *31*, 8946–8952. [[CrossRef](#)]
19. Sriram, R.; Van Criekinge, M.; Hansen, A.; Wang, Z.J.; Vigneron, D.B.; Wilson, D.M.; Keshari, K.R.; Kurhanewicz, J. Real-time measurement of hyperpolarized lactate production and efflux as a biomarker of tumor aggressiveness in an MR compatible 3D cell culture bioreactor. *NMR Biomed.* **2015**, *28*, 1141–1149. [[CrossRef](#)]
20. Wilkes, R.P.; McNulty, A.K.; Feeley, T.D.; Schmidt, M.A.; Kieswetter, K. Bioreactor for application of subatmospheric pressure to three-dimensional cell culture. *Tissue Eng.* **2007**, *13*, 3003–3010. [[CrossRef](#)]
21. Wu, M.H.; Urban, J.P.; Cui, Z.; Cui, Z.F. Development of PDMS microbioreactor with well-defined and homogenous culture environment for chondrocyte 3-D culture. *Biomed. Microdevices* **2006**, *8*, 331–340. [[CrossRef](#)] [[PubMed](#)]
22. Wu, M.H.; Kuo, C.Y. Application of high throughput perfusion micro 3-D cell culture platform for the precise study of cellular responses to extracellular conditions -effect of serum concentrations on the physiology of articular chondrocytes. *Biomed. Microdevices* **2011**, *13*, 131–141. [[CrossRef](#)] [[PubMed](#)]
23. Wu, M.H.; Huang, S.B.; Cui, Z.; Cui, Z.; Lee, G.B. A high throughput perfusion-based microbioreactor platform integrated with pneumatic micropumps for three-dimensional cell culture. *Biomed. Microdevices* **2008**, *10*, 309–319. [[CrossRef](#)]
24. Hsieh, C.H.; Chen, Y.D.; Huang, S.F.; Wang, H.M.; Wu, M.H. The effect of primary cancer cell culture models on the results of drug chemosensitivity assays: The application of perfusion microbioreactor system as cell culture vessel. *Biomed. Res. Int.* **2015**, *2015*, 470283. [[CrossRef](#)] [[PubMed](#)]

25. Altman, G.H.; Horan, R.L.; Martin, I.; Farhadi, J.; Stark, P.R.; Volloch, V.; Richmond, J.C.; Vunjak-Novakovic, G.; Kaplan, D.L. Cell differentiation by mechanical stress. *FASEB J.* **2002**, *16*, 270–272. [[CrossRef](#)]
26. Cochis, A.; Grad, S.; Stoddart, M.J.; Fare, S.; Altomare, L.; Azzimonti, B.; Alini, M.; Rimondini, L. Bioreactor mechanically guided 3D mesenchymal stem cell chondrogenesis using a biocompatible novel thermo-reversible methylcellulose-based hydrogel. *Sci. Rep.* **2017**, *7*, 45018. [[CrossRef](#)]
27. Correia, C.; Pereira, A.L.; Duarte, A.R.; Frias, A.M.; Pedro, A.J.; Oliveira, J.T.; Sousa, R.A.; Reis, R.L. Dynamic culturing of cartilage tissue: The significance of hydrostatic pressure. *Tissue Eng. Part A* **2012**, *18*, 1979–1991. [[CrossRef](#)]
28. Gharravi, A.M.; Orazizadeh, M.; Ansari-Asl, K.; Banoni, S.; Izadi, S.; Hashemitarabar, M. Design and fabrication of anatomical bioreactor systems containing alginate scaffolds for cartilage tissue engineering. *Avicenna J. Med. Biotechnol.* **2012**, *4*, 65–74.
29. Lee, P.Y.; Liu, Y.C.; Wang, M.X.; Hu, J.J. Fibroblast-seeded collagen gels in response to dynamic equibiaxial mechanical stimuli: A biomechanical study. *J. Biomech.* **2018**, *78*, 134–142. [[CrossRef](#)]
30. Moraes, C.; Wang, G.; Sun, Y.; Simmons, C.A. A microfabricated platform for high-throughput unconfined compression of micropatterned biomaterial arrays. *Biomaterials* **2010**, *31*, 577–584. [[CrossRef](#)]
31. Powell, C.A.; Smiley, B.L.; Mills, J.; Vandenberg, H.H. Mechanical stimulation improves tissue-engineered human skeletal muscle. *Am. J. Physiol. Cell. Physiol.* **2002**, *283*, C1557–C1565. [[CrossRef](#)] [[PubMed](#)]
32. Rodling, L.; Volz, E.M.; Raic, A.; Brandle, K.; Franzreb, M.; Lee-Thedieck, C. Magnetic Macroporous Hydrogels as a Novel Approach for Perfused Stem Cell Culture in 3D Scaffolds via Contactless Motion Control. *Adv. Healthc. Mater.* **2018**, *7*, e1701403. [[CrossRef](#)]
33. Rodday, B.; Hirschhaeuser, F.; Walenta, S.; Mueller-Klieser, W. Semiautomatic growth analysis of multicellular tumor spheroids. *J. Biomol. Screen* **2011**, *16*, 1119–1124. [[CrossRef](#)] [[PubMed](#)]
34. Timmins, N.E.; Scherberich, A.; Fruh, J.A.; Heberer, M.; Martin, I.; Jakob, M. Three-dimensional cell culture and tissue engineering in a T-CUP (tissue culture under perfusion). *Tissue Eng.* **2007**, *13*, 2021–2028. [[CrossRef](#)] [[PubMed](#)]
35. Tostoes, R.M.; Leite, S.B.; Serra, M.; Jensen, J.; Bjorquist, P.; Carrondo, M.J.; Brito, C.; Alves, P.M. Human liver cell spheroids in extended perfusion bioreactor culture for repeated-dose drug testing. *Hepatology* **2012**, *55*, 1227–1236. [[CrossRef](#)]
36. Aucamp, J.; Calitz, C.; Bronkhorst, A.J.; Wrzesinski, K.; Hamman, S.; Gouws, C.; Pretorius, P.J. Cell-free DNA in a three-dimensional spheroid cell culture model: A preliminary study. *Int. J. Biochem. Cell. Biol.* **2017**, *89*, 182–192. [[CrossRef](#)] [[PubMed](#)]
37. Botchwey, E.A.; Pollack, S.R.; Levine, E.M.; Laurencin, C.T. Bone tissue engineering in a rotating bioreactor using a microcarrier matrix system. *J. Biomed. Mater. Res.* **2001**, *55*, 242–253. [[CrossRef](#)]
38. Carterson, A.J.; Honer zu Bentrup, K.; Ott, C.M.; Clarke, M.S.; Pierson, D.L.; Vanderburg, C.R.; Buchanan, K.L.; Nickerson, C.A.; Schurr, M.J. A549 lung epithelial cells grown as three-dimensional aggregates: Alternative tissue culture model for *Pseudomonas aeruginosa* pathogenesis. *Infect. Immun.* **2005**, *73*, 1129–1140. [[CrossRef](#)]
39. Carvalho, H.M.; Teel, L.D.; Goping, G.; O'Brien, A.D. A three-dimensional tissue culture model for the study of attach and efface lesion formation by enteropathogenic and enterohaemorrhagic *Escherichia coli*. *Cell. Microbiol.* **2005**, *7*, 1771–1781. [[CrossRef](#)]
40. Chang, T.T.; Hughes-Fulford, M. Molecular mechanisms underlying the enhanced functions of three-dimensional hepatocyte aggregates. *Biomaterials* **2014**, *35*, 2162–2171. [[CrossRef](#)]
41. Cortiella, J.; Niles, J.; Cantu, A.; Brettler, A.; Pham, A.; Vargas, G.; Winston, S.; Wang, J.; Walls, S.; Nichols, J.E. Influence of acellular natural lung matrix on murine embryonic stem cell differentiation and tissue formation. *Tissue Eng. Part A* **2010**, *16*, 2565–2580. [[CrossRef](#)] [[PubMed](#)]
42. Crabbe, A.; Liu, Y.; Sarker, S.F.; Bonenfant, N.R.; Barrila, J.; Borg, Z.D.; Lee, J.J.; Weiss, D.J.; Nickerson, C.A. Recellularization of decellularized lung scaffolds is enhanced by dynamic suspension culture. *PLoS ONE* **2015**, *10*, e0126846. [[CrossRef](#)] [[PubMed](#)]
43. Crabbe, A.; Liu, Y.; Matthijs, N.; Rigole, P.; De La Fuente-Nunez, C.; Davis, R.; Ledesma, M.A.; Sarker, S.; Van Houdt, R.; Hancock, R.E.; et al. Antimicrobial efficacy against *Pseudomonas aeruginosa* biofilm formation in a three-dimensional lung epithelial model and the influence of fetal bovine serum. *Sci. Rep.* **2017**, *7*, 43321. [[CrossRef](#)] [[PubMed](#)]

44. De Weirdt, R.; Crabbe, A.; Roos, S.; Vollenweider, S.; Lacroix, C.; van Pijkeren, J.P.; Britton, R.A.; Sarker, S.; Van de Wiele, T.; Nickerson, C.A. Glycerol supplementation enhances *L. reuteri*'s protective effect against *S. Typhimurium* colonization in a 3-D model of colonic epithelium. *PLoS ONE* **2012**, *7*, e37116. [CrossRef]
45. Detamore, M.S.; Athanasiou, K.A. Use of a rotating bioreactor toward tissue engineering the temporomandibular joint disc. *Tissue Eng.* **2005**, *11*, 1188–1197. [CrossRef]
46. Drummond, C.G.; Nickerson, C.A.; Coyne, C.B. A Three-Dimensional Cell Culture Model To Study Enterovirus Infection of Polarized Intestinal Epithelial Cells. *mSphere* **2016**, *1*. [CrossRef]
47. Hjelm, B.E.; Berta, A.N.; Nickerson, C.A.; Arntzen, C.J.; Herbst-Kralovetz, M.M. Development and characterization of a three-dimensional organotypic human vaginal epithelial cell model. *Biol. Reprod.* **2010**, *82*, 617–627. [CrossRef]
48. Hoganson, D.M.; Owens, G.E.; Meppelink, A.M.; Bassett, E.K.; Bowley, C.M.; Hinkel, C.J.; Finkelstein, E.B.; Goldman, S.M.; Vacanti, J.P. Decellularized extracellular matrix microparticles as a vehicle for cellular delivery in a model of anastomosis healing. *J. Biomed. Mater. Res. A* **2016**, *104*, 1728–1735. [CrossRef]
49. Honer zu Bentrup, K.; Ramamurthy, R.; Ott, C.M.; Emami, K.; Nelman-Gonzalez, M.; Wilson, J.W.; Richter, E.G.; Goodwin, T.J.; Alexander, J.S.; Pierson, D.L.; et al. Three-dimensional organotypic models of human colonic epithelium to study the early stages of enteric salmonellosis. *Microbes Infect.* **2006**, *8*, 1813–1825. [CrossRef]
50. LaMarca, H.L.; Ott, C.M.; Honer Zu Bentrup, K.; Leblanc, C.L.; Pierson, D.L.; Nelson, A.B.; Scandurro, A.B.; Whitley, G.S.; Nickerson, C.A.; Morris, C.A. Three-dimensional growth of extravillous cytotrophoblasts promotes differentiation and invasion. *Placenta* **2005**, *26*, 709–720. [CrossRef]
51. Laniowski, P.; Gomez, A.; Hire, G.; So, M.; Herbst-Kralovetz, M.M. Human Three-Dimensional Endometrial Epithelial Cell Model To Study Host Interactions with Vaginal Bacteria and *Neisseria gonorrhoeae*. *Infect. Immun.* **2017**, *85*. [CrossRef] [PubMed]
52. Lei, X.H.; Ning, L.N.; Cao, Y.J.; Liu, S.; Zhang, S.B.; Qiu, Z.F.; Hu, H.M.; Zhang, H.S.; Liu, S.; Duan, E.K. NASA-approved rotary bioreactor enhances proliferation of human epidermal stem cells and supports formation of 3D epidermis-like structure. *PLoS ONE* **2011**, *6*, e26603. [CrossRef] [PubMed]
53. Li, W.J.; Tuan, R.S. Fabrication and application of nanofibrous scaffolds in tissue engineering. *Curr. Protoc. Cell. Biol.* **2009**, *42*, 25.2. 1–25.2. 12. [CrossRef] [PubMed]
54. Marlovits, S.; Tichy, B.; Truppe, M.; Gruber, D.; Vecsei, V. Chondrogenesis of aged human articular cartilage in a scaffold-free bioreactor. *Tissue Eng.* **2003**, *9*, 1215–1226. [CrossRef]
55. Montani, C.; Steimberg, N.; Boniotti, J.; Biasiotto, G.; Zanella, I.; Diafera, G.; Biunno, I.; Caimi, L.; Mazzoleni, G.; Di Lorenzo, D. Fibroblasts maintained in 3 dimensions show a better differentiation state and higher sensitivity to estrogens. *Toxicol. Appl. Pharmacol.* **2014**, *280*, 421–433. [CrossRef]
56. Nickerson, C.A.; Goodwin, T.J.; Terlonge, J.; Ott, C.M.; Buchanan, K.L.; Uicker, W.C.; Emami, K.; LeBlanc, C.L.; Ramamurthy, R.; Clarke, M.S.; et al. Three-dimensional tissue assemblies: Novel models for the study of *Salmonella enterica* serovar *Typhimurium* pathogenesis. *Infect. Immun.* **2001**, *69*, 7106–7120. [CrossRef]
57. Papadaki, M.; Bursac, N.; Langer, R.; Merok, J.; Vunjak-Novakovic, G.; Freed, L.E. Tissue engineering of functional cardiac muscle: Molecular, structural, and electrophysiological studies. *Am. J. Physiol. Heart Circ. Physiol.* **2001**, *280*, H168–H178. [CrossRef]
58. Papafragkou, E.; Hewitt, J.; Park, G.W.; Greening, G.; Vinje, J. Challenges of culturing human norovirus in three-dimensional organoid intestinal cell culture models. *PLoS ONE* **2014**, *8*, e63485. [CrossRef]
59. Quail, D.F.; Maciel, T.J.; Rogers, K.; Postovit, L.M. A unique 3D in vitro cellular invasion assay. *J. Biomol. Screen* **2012**, *17*, 1088–1095. [CrossRef]
60. Radtke, A.L.; Wilson, J.W.; Sarker, S.; Nickerson, C.A. Analysis of interactions of *Salmonella* type three secretion mutants with 3-D intestinal epithelial cells. *PLoS ONE* **2010**, *5*, e15750. [CrossRef]
61. Rungarunlert, S.; Klincumhom, N.; Bock, I.; Nemes, C.; Techakumphu, M.; Pirity, M.K.; Dinnyes, A. Enhanced cardiac differentiation of mouse embryonic stem cells by use of the slow-turning, lateral vessel (STLV) bioreactor. *Biotechnol. Lett.* **2011**, *33*, 1565–1573. [CrossRef] [PubMed]
62. Sainz, B., Jr.; TenCate, V.; Uprichard, S.L. Three-dimensional Huh7 cell culture system for the study of Hepatitis C virus infection. *Virol. J.* **2009**, *6*, 103. [CrossRef] [PubMed]
63. Samuelson, L.; Gerber, D.A. Improved function and growth of pancreatic cells in a three-dimensional bioreactor environment. *Tissue Eng. Part. C Methods* **2013**, *19*, 39–47. [CrossRef] [PubMed]

64. Sikavitsas, V.I.; Bancroft, G.N.; Mikos, A.G. Formation of three-dimensional cell/polymer constructs for bone tissue engineering in a spinner flask and a rotating wall vessel bioreactor. *J. Biomed. Mater. Res.* **2002**, *62*, 136–148. [[CrossRef](#)]
65. Skardal, A.; Sarker, S.F.; Crabbe, A.; Nickerson, C.A.; Prestwich, G.D. The generation of 3-D tissue models based on hyaluronan hydrogel-coated microcarriers within a rotating wall vessel bioreactor. *Biomaterials* **2010**, *31*, 8426–8435. [[CrossRef](#)]
66. Smit, T.; Calitz, C.; Willers, C.; Svitina, H.; Hamman, J.; Fey, S.J.; Gouws, C.; Wrzesinski, K. Characterization of an Alginic Encapsulated LS180 Spheroid Model for Anti-colorectal Cancer Compound Screening. *ACS Med. Chem. Lett.* **2020**, *11*, 1014–1021. [[CrossRef](#)]
67. Smith, Y.C.; Grande, K.K.; Rasmussen, S.B.; O'Brien, A.D. Novel three-dimensional organoid model for evaluation of the interaction of uropathogenic Escherichia coli with terminally differentiated human urothelial cells. *Infect. Immun.* **2006**, *74*, 750–757. [[CrossRef](#)]
68. Valmikinathan, C.M.; Hoffman, J.; Yu, X. Impact of Scaffold Micro and Macro Architecture on Schwann Cell Proliferation under Dynamic Conditions in a Rotating Wall Vessel Bioreactor. *Mater. Sci. Eng. C Mater. Biol. Appl.* **2011**, *31*, 22–29. [[CrossRef](#)]
69. Alcantara Warren, C.; Destura, R.V.; Sevilleja, J.E.; Barroso, L.F.; Carvalho, H.; Barrett, L.J.; O'Brien, A.D.; Guerrant, R.L. Detection of epithelial-cell injury, and quantification of infection, in the HCT-8 organoid model of cryptosporidiosis. *J. Infect. Dis.* **2008**, *198*, 143–149. [[CrossRef](#)]
70. Wrzesinski, K.; Fey, S.J. Metabolic Reprogramming and the Recovery of Physiological Functionality in 3D Cultures in Micro-Bioreactors. *Bioengineering* **2018**, *5*, 22. [[CrossRef](#)]
71. Yamashita, T.; Takayama, K.; Sakurai, F.; Mizuguchi, H. Billion-scale production of hepatocyte-like cells from human induced pluripotent stem cells. *Biochem. Biophys. Res. Commun.* **2018**, *496*, 1269–1275. [[CrossRef](#)] [[PubMed](#)]
72. Altman, G.H.; Lu, H.H.; Horan, R.L.; Calabro, T.; Ryder, D.; Kaplan, D.L.; Stark, P.; Martin, I.; Richmond, J.C.; Vunjak-Novakovic, G. Advanced bioreactor with controlled application of multi-dimensional strain for tissue engineering. *J. Biomech. Eng.-T Asme* **2002**, *124*, 742–749. [[CrossRef](#)] [[PubMed](#)]
73. Bancroft, G.N.; Sikavitsas, V.I.; van den Dolder, J.; Sheffield, T.L.; Ambrose, C.G.; Jansen, J.A.; Mikos, A.G. Fluid flow increases mineralized matrix deposition in 3D perfusion culture of marrow stromal osteoblasts in a dose-dependent manner. *Proc. Natl. Acad. Sci. USA* **2002**, *99*, 12600–12605. [[CrossRef](#)] [[PubMed](#)]
74. Bancroft, G.N.; Sikavitsas, V.I.; Mikos, A.G. Design of a flow perfusion bioreactor system for bone tissue-engineering applications. *Tissue Eng.* **2003**, *9*, 549–554. [[CrossRef](#)]
75. Bartnikowski, M.; Klein, T.J.; Melchels, F.P.; Woodruff, M.A. Effects of scaffold architecture on mechanical characteristics and osteoblast response to static and perfusion bioreactor cultures. *Biotechnol. Bioeng.* **2014**, *111*, 1440–1451. [[CrossRef](#)]
76. Baudoin, R.; Griscom, L.; Prot, J.M.; Legallais, C.; Leclerc, E. Behavior of HepG2/C3A cell cultures in a microfluidic bioreactor. *Biochem. Eng. J.* **2011**, *53*, 172–181. [[CrossRef](#)]
77. Baudoin, R.; Alberto, G.; Legendre, A.; Paullier, P.; Naudot, M.; Fleury, M.J.; Jacques, S.; Griscom, L.; Leclerc, E. Investigation of expression and activity levels of primary rat hepatocyte detoxication genes under various flow rates and cell densities in microfluidic biochips. *Biotechnol. Prog.* **2014**, *30*, 401–410. [[CrossRef](#)]
78. Birru, B.; Mekala, N.K.; Parcha, S.R. Improved osteogenic differentiation of umbilical cord blood MSCs using custom made perfusion bioreactor. *Biomed. J.* **2018**, *41*, 290–297. [[CrossRef](#)]
79. Candini, O.; Grisendi, G.; Foppiani, E.M.; Brogli, M.; Aramini, B.; Masciale, V.; Spano, C.; Petrachi, T.; Veronesi, E.; Conte, P.; et al. A Novel 3D In Vitro Platform for Pre-Clinical Investigations in Drug Testing, Gene Therapy, and Immuno-oncology. *Sci. Rep.* **2019**, *9*, 7154. [[CrossRef](#)]
80. Carrier, R.L.; Rupnick, M.; Langer, R.; Schoen, F.J.; Freed, L.E.; Vunjak-Novakovic, G. Perfusion improves tissue architecture of engineered cardiac muscle. *Tissue Eng.* **2002**, *8*, 175–188. [[CrossRef](#)]
81. Christoffersson, J.; Bergstrom, G.; Schwanke, K.; Kempf, H.; Zweigerdt, R.; Mandenius, C.F. A Microfluidic Bioreactor for Toxicity Testing of Stem Cell Derived 3D Cardiac Bodies. *Methods Mol. Biol.* **2016**, *1502*, 159–168. [[CrossRef](#)] [[PubMed](#)]
82. Christoffersson, J.; Mandenius, C.-F. Using a Microfluidic Device for Culture and Drug Toxicity Testing of 3D Cells. In *Cell-Based Assays Using iPSCs for Drug Development and Testing*; Mandenius, C.-F., Ross, J., Eds.; Humana: New York, NY, USA, 2019. [[CrossRef](#)]

83. Cimetta, E.; Sirabella, D.; Yeager, K.; Davidson, K.; Simon, J.; Moon, R.T.; Vunjak-Novakovic, G. Microfluidic bioreactor for dynamic regulation of early mesodermal commitment in human pluripotent stem cells. *Lab. Chip.* **2013**, *13*, 355–364. [CrossRef]
84. Costa, P.F.; Vaquette, C.; Baldwin, J.; Chhaya, M.; Gomes, M.E.; Reis, R.L.; Theodoropoulos, C.; Hutmacher, D.W. Biofabrication of customized bone grafts by combination of additive manufacturing and bioreactor knowhow. *Biofabrication* **2014**, *6*, 035006. [CrossRef] [PubMed]
85. Egger, D.; Spitz, S.; Fischer, M.; Handschuh, S.; Glosmann, M.; Friemert, B.; Egerbacher, M.; Kasper, C. Application of a Parallelizable Perfusion Bioreactor for Physiologic 3D Cell Culture. *Cells Tissues Organs* **2017**, *203*, 316–326. [CrossRef] [PubMed]
86. Figallo, E.; Cannizzaro, C.; Gerecht, S.; Burdick, J.A.; Langer, R.; Elvassore, N.; Vunjak-Novakovic, G. Micro-bioreactor array for controlling cellular microenvironments. *Lab. Chip.* **2007**, *7*, 710–719. [CrossRef] [PubMed]
87. Frohlich, M.; Grayson, W.L.; Marolt, D.; Gimble, J.M.; Kregar-Velikonja, N.; Vunjak-Novakovic, G. Bone grafts engineered from human adipose-derived stem cells in perfusion bioreactor culture. *Tissue Eng. Part. A* **2010**, *16*, 179–189. [CrossRef]
88. Fu, C.Y.; Tseng, S.Y.; Yang, S.M.; Hsu, L.; Liu, C.H.; Chang, H.Y. A microfluidic chip with a U-shaped microstructure array for multicellular spheroid formation, culturing and analysis. *Biofabrication* **2014**, *6*, 015009. [CrossRef]
89. Goldstein, A.S.; Juarez, T.M.; Helmke, C.D.; Gustin, M.C.; Mikos, A.G. Effect of convection on osteoblastic cell growth and function in biodegradable polymer foam scaffolds. *Biomaterials* **2001**, *22*, 1279–1288. [CrossRef]
90. Gomes, M.E.; Sikavitsas, V.I.; Behravesh, E.; Reis, R.L.; Mikos, A.G. Effect of flow perfusion on the osteogenic differentiation of bone marrow stromal cells cultured on starch-based three-dimensional scaffolds. *J. Biomed. Mater. Res. Part. A* **2003**, *67*, 87–95. [CrossRef]
91. Gottwald, E.; Lahni, B.; Thiele, D.; Giselbrecht, S.; Welle, A.; Weibe Zahnh, K.F. Chip-based three-dimensional cell culture in perfused micro-bioreactors. *J. Vis. Exp.* **2008**. [CrossRef]
92. Guzzardi, M.A.; Domenici, C.; Ahluwalia, A. Metabolic control through hepatocyte and adipose tissue cross-talk in a multicompartimental modular bioreactor. *Tissue Eng. Part A* **2011**, *17*, 1635–1642. [CrossRef] [PubMed]
93. Hongo, T.; Kajikawa, M.; Ishida, S.; Ozawa, S.; Ohno, Y.; Sawada, J.; Umezawa, A.; Ishikawa, Y.; Kobayashi, T.; Honda, H. Three-dimensional high-density culture of HepG2 cells in a 5-mL radial-flow bioreactor for construction of artificial liver. *J. Biosci. Bioeng.* **2005**, *99*, 237–244. [CrossRef] [PubMed]
94. Izzo, L.; Tunesi, M.; Boeri, L.; Lagana, M.; Giordano, C.; Raimondi, M.T. Influence of the static magnetic field on cell response in a miniaturized optically accessible bioreactor for 3D cell culture. *Biomod. Microdevices* **2019**, *21*, 29. [CrossRef] [PubMed]
95. Janssen, F.W.; Hofland, I.; van Oorschot, A.; Oostra, J.; Peters, H.; van Blitterswijk, C.A. Online measurement of oxygen consumption by goat bone marrow stromal cells in a combined cell-seeding and proliferation perfusion bioreactor. *J. Biomed. Mater. Res. A* **2006**, *79*, 338–348. [CrossRef]
96. Janssen, F.W.; Oostra, J.; Oorschot, A.; van Blitterswijk, C.A. A perfusion bioreactor system capable of producing clinically relevant volumes of tissue-engineered bone: In vivo bone formation showing proof of concept. *Biomaterials* **2006**, *27*, 315–323. [CrossRef]
97. Kleinhans, C.; Mohan, R.R.; Vacun, G.; Schwarz, T.; Haller, B.; Sun, Y.; Kahlig, A.; Kluger, P.; Finne-Wistrand, A.; Walles, H.; et al. A perfusion bioreactor system efficiently generates cell-loaded bone substitute materials for addressing critical size bone defects. *Biotechnol. J.* **2015**, *10*, 1727–1738. [CrossRef]
98. Kostrzewski, T.; Cornforth, T.; Snow, S.A.; Ouro-Gnao, L.; Rowe, C.; Large, E.M.; Hughes, D.J. Three-dimensional perfused human in vitro model of non-alcoholic fatty liver disease. *World J. Gastroenterol.* **2017**, *23*, 204–215. [CrossRef]
99. Leclerc, E.; Sakai, Y.; Fujii, T. Perfusion culture of fetal human hepatocytes in microfluidic environments. *Biochem. Eng. J.* **2004**, *20*, 143–148. [CrossRef]
100. Lee, S.H.; Hong, S.; Song, J.; Cho, B.; Han, E.J.; Kondapavulur, S.; Kim, D.; Lee, L.P. Microphysiological Analysis Platform of Pancreatic Islet beta-Cell Spheroids. *Adv. Healthc. Mater.* **2018**, *7*, 1701111. [CrossRef]
101. Legendre, A.; Baudoine, R.; Alberto, G.; Paullier, P.; Naudot, M.; Bricks, T.; Brochetton, J.; Jacques, S.; Cotton, J.; Leclerc, E. Metabolic characterization of primary rat hepatocytes cultivated in parallel microfluidic biochips. *J. Pharm. Sci.* **2013**, *102*, 3264–3276. [CrossRef]

102. Linti, C.; Zipfel, A.; Schenk, M.; Dauner, M.; Doser, M.; Viebahn, R.; Becker, H.D.; Planck, H. Cultivation of porcine hepatocytes in polyurethane nonwovens as part of a biohybrid liver support system. *Int. J. Artif. Organs* **2002**, *25*, 994–1000. [CrossRef] [PubMed]
103. Neubauer, A.; Nies, C.; Schepkin, V.D.; Hu, R.; Malzacher, M.; Chacon-Caldera, J.; Thiele, D.; Gottwald, E.; Schad, L.R. Tracking protein function with sodium multi quantum spectroscopy in a 3D-tissue culture based on microcavity arrays. *Sci. Rep.* **2017**, *7*, 3943. [CrossRef] [PubMed]
104. Ong, S.M.; Zhang, C.; Toh, Y.C.; Kim, S.H.; Foo, H.L.; Tan, C.H.; van Noort, D.; Park, S.; Yu, H. A gel-free 3D microfluidic cell culture system. *Biomaterials* **2008**, *29*, 3237–3244. [CrossRef] [PubMed]
105. Ostrovidov, S.; Sakai, Y.; Fujii, T. Integration of a pump and an electrical sensor into a membrane-based PDMS microbioreactor for cell culture and drug testing. *Biomed. Microdevices* **2011**, *13*, 847–864. [CrossRef]
106. Powers, M.J.; Domansky, K.; Kaazempur-Mofrad, M.R.; Kalezi, A.; Capitano, A.; Upadhyaya, A.; Kurzawski, P.; Wack, K.E.; Stoltz, D.B.; Kamm, R.; et al. A microfabricated array bioreactor for perfused 3D liver culture. *Biotechnol. Bioeng.* **2002**, *78*, 257–269. [CrossRef]
107. Powers, M.J.; Janigan, D.M.; Wack, K.E.; Baker, C.S.; Beer Stoltz, D.; Griffith, L.G. Functional behavior of primary rat liver cells in a three-dimensional perfused microarray bioreactor. *Tissue Eng.* **2002**, *8*, 499–513. [CrossRef]
108. Prot, J.M.; Aninat, C.; Griscom, L.; Razan, F.; Brochot, C.; Guillouzo, C.G.; Legallais, C.; Corlu, A.; Leclerc, E. Improvement of HepG2/C3a cell functions in a microfluidic biochip. *Biotechnol. Bioeng.* **2011**, *108*, 1704–1715. [CrossRef]
109. Prot, J.M.; Videau, O.; Brochot, C.; Legallais, C.; Benech, H.; Leclerc, E. A cocktail of metabolic probes demonstrates the relevance of primary human hepatocyte cultures in a microfluidic biochip for pharmaceutical drug screening. *Int. J. Pharm.* **2011**, *408*, 67–75. [CrossRef]
110. Radisic, M.; Euloth, M.; Yang, L.; Langer, R.; Freed, L.E.; Vunjak-Novakovic, G. High-density seeding of myocyte cells for cardiac tissue engineering. *Biotechnol. Bioeng.* **2003**, *82*, 403–414. [CrossRef]
111. Sailon, A.M.; Allori, A.C.; Davidson, E.H.; Reformat, D.D.; Allen, R.J.; Warren, S.M. A novel flow-perfusion bioreactor supports 3D dynamic cell culture. *J. Biomed. Biotechnol.* **2009**, *2009*, 873816. [CrossRef]
112. Santoro, M.; Lamhamdi-Cherradi, S.E.; Menegaz, B.A.; Ludwig, J.A.; Mikos, A.G. Flow perfusion effects on three-dimensional culture and drug sensitivity of Ewing sarcoma. *Proc. Natl. Acad. Sci. USA* **2015**, *112*, 10304–10309. [CrossRef]
113. Sikavitsas, V.I.; Bancroft, G.N.; Holtorf, H.L.; Jansen, J.A.; Mikos, A.G. Mineralized matrix deposition by marrow stromal osteoblasts in 3D perfusion culture increases with increasing fluid shear forces. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 14683–14688. [CrossRef] [PubMed]
114. Sikavitsas, V.I.; Bancroft, G.N.; Lemoine, J.J.; Liebschner, M.A.; Dauner, M.; Mikos, A.G. Flow perfusion enhances the calcified matrix deposition of marrow stromal cells in biodegradable nonwoven fiber mesh scaffolds. *Ann. Biomed. Eng.* **2005**, *33*, 63–70. [CrossRef] [PubMed]
115. Silva, A.R.; Paula, A.C.; Martins, T.M.; Goes, A.M.; Pereria, M.M. Synergistic effect between bioactive glass foam and a perfusion bioreactor on osteogenic differentiation of human adipose stem cells. *J. Biomed. Mater. Res. A* **2014**, *102*, 818–827. [CrossRef] [PubMed]
116. Sivaraman, A.; Leach, J.K.; Townsend, S.; Iida, T.; Hogan, B.J.; Stoltz, D.B.; Fry, R.; Samson, L.D.; Tannenbaum, S.R.; Griffith, L.G. A microscale in vitro physiological model of the liver: Predictive screens for drug metabolism and enzyme induction. *Curr. Drug Metab.* **2005**, *6*, 569–591. [CrossRef] [PubMed]
117. Spitkovsky, D.; Lemke, K.; Forster, T.; Romer, R.; Wiedemeier, S.; Hescheler, J.; Sachinidis, A.; Gastrock, G. Generation of Cardiomyocytes in Pipe-Based Microbioreactor Under Segmented Flow. *Cell. Physiol. Biochem.* **2016**, *38*, 1883–1896. [CrossRef]
118. Sun, T.; Norton, D.; Vickers, N.; Sall, L.M.; Neil, S.M.; Ryan, A.J.; Haycock, J.W. Development of a bioreactor for evaluating novel nerve conduits. *Biotechnol. Bioeng.* **2008**, *99*, 1250–1260. [CrossRef]
119. Tania, M.; Hsu, M.N.; Png, S.N.; Leo, H.L.; Toh, G.W.; Birgersson, E. Perfusion enhanced polydimethylsiloxane based scaffold cell culturing system for multi-well drug screening platform. *Biotechnol. Prog.* **2014**, *30*, 418–428. [CrossRef]
120. Thompson, C.A.; Colon-Hernandez, P.; Pomerantseva, I.; MacNeil, B.D.; Nasseri, B.; Vacanti, J.P.; Oesterle, S.N. A novel pulsatile, laminar flow bioreactor for the development of tissue-engineered vascular structures. *Tissue Eng.* **2002**, *8*, 1083–1088. [CrossRef]

121. Toh, Y.C.; Zhang, C.; Zhang, J.; Khong, Y.M.; Chang, S.; Samper, V.D.; van Noort, D.; Hutmacher, D.W.; Yu, H. A novel 3D mammalian cell perfusion-culture system in microfluidic channels. *Lab. Chip.* **2007**, *7*, 302–309. [[CrossRef](#)]
122. Tunisi, M.; Fusco, F.; Fiordaliso, F.; Corbelli, A.; Biella, G.; Raimondi, M.T. Optimization of a 3D Dynamic Culturing System for In Vitro Modeling of Frontotemporal Neurodegeneration-Relevant Pathologic Features. *Front. Aging Neurosci.* **2016**, *8*, 146. [[CrossRef](#)] [[PubMed](#)]
123. Vetsch, J.R.; Betts, D.C.; Muller, R.; Hofmann, S. Flow velocity-driven differentiation of human mesenchymal stromal cells in silk fibroin scaffolds: A combined experimental and computational approach. *PLoS ONE* **2017**, *12*, e0180781. [[CrossRef](#)] [[PubMed](#)]
124. Weise, F.; Fernekorn, U.; Hampl, J.; Klett, M.; Schober, A. Analysis and comparison of oxygen consumption of HepG2 cells in a monolayer and three-dimensional high density cell culture by use of a matrigrid(R). *Biotechnol. Bioeng.* **2013**, *110*, 2504–2512. [[CrossRef](#)] [[PubMed](#)]
125. Wen, Y.; Zhang, X.; Yang, S.T. Microplate-reader compatible perfusion microbioreactor array for modular tissue culture and cytotoxicity assays. *Biotechnol. Prog.* **2010**, *26*, 1135–1144. [[CrossRef](#)] [[PubMed](#)]
126. Wendt, D.; Marsano, A.; Jakob, M.; Heberer, M.; Martin, I. Oscillating perfusion of cell suspensions through three-dimensional scaffolds enhances cell seeding efficiency and uniformity. *Biotechnol. Bioeng.* **2003**, *84*, 205–214. [[CrossRef](#)] [[PubMed](#)]
127. Xin, X.; Wu, Y.; Zang, R.; Yang, S.T. A fluorescent 3D cell culture assay for high throughput screening of cancer drugs down-regulating survivin. *J. Biotechnol.* **2019**, *289*, 80–87. [[CrossRef](#)]
128. Yu, F.; Deng, R.; Hao Tong, W.; Huan, L.; Chan Way, N.; IslamBadhan, A.; Iliescu, C.; Yu, H. A perfusion incubator liver chip for 3D cell culture with application on chronic hepatotoxicity testing. *Sci. Rep.* **2017**, *7*, 14528. [[CrossRef](#)]
129. Zakharyants, A.A.; Burmistrova, O.A.; Poloznikov, A.A. The Use of Human Liver Cell Model and Cytochrome P450 Substrate-Inhibitor Panel for Studies of Dasatinib and Warfarin Interactions. *Bull. Exp. Biol. Med.* **2017**, *162*, 515–519. [[CrossRef](#)]
130. Zhang, C.; Chia, S.M.; Ong, S.M.; Zhang, S.; Toh, Y.C.; van Noort, D.; Yu, H. The controlled presentation of TGF-beta1 to hepatocytes in a 3D-microfluidic cell culture system. *Biomaterials* **2009**, *30*, 3847–3853. [[CrossRef](#)]
131. Zhang, Y.S.; Aleman, J.; Shin, S.R.; Kilic, T.; Kim, D.; Mousavi Shaegh, S.A.; Massa, S.; Riahi, R.; Chae, S.; Hu, N.; et al. Multisensor-integrated organs-on-chips platform for automated and continual in situ monitoring of organoid behaviors. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, E2293–E2302. [[CrossRef](#)]
132. Zhao, F.; Chella, R.; Ma, T. Effects of shear stress on 3-D human mesenchymal stem cell construct development in a perfusion bioreactor system: Experiments and hydrodynamic modeling. *Biotechnol. Bioeng.* **2007**, *96*, 584–595. [[CrossRef](#)]
133. Darnell, M.; Schreiter, T.; Zeilinger, K.; Urbaniak, T.; Soderdahl, T.; Rossberg, I.; Dillner, B.; Berg, A.L.; Gerlach, J.C.; Andersson, T.B. Cytochrome P450-dependent metabolism in HepaRG cells cultured in a dynamic three-dimensional bioreactor. *Drug Metab. Dispos.* **2011**, *39*, 1131–1138. [[CrossRef](#)] [[PubMed](#)]
134. Darnell, M.; Ulvestad, M.; Ellis, E.; Weidolf, L.; Andersson, T.B. In Vitro Evaluation of Major In Vivo Drug Metabolic Pathways Using Primary Human Hepatocytes and HepaRG Cells in Suspension and a Dynamic Three-Dimensional Bioreactor System. *J. Pharmacol. Exp. Ther.* **2012**, *343*, 134–144. [[CrossRef](#)] [[PubMed](#)]
135. De Bartolo, L.; Salerno, S.; Curcio, E.; Piscioneri, A.; Rende, M.; Morelli, S.; Tasselli, F.; Bader, A.; Drioli, E. Human hepatocyte functions in a crossed hollow fiber membrane bioreactor. *Biomaterials* **2009**, *30*, 2531–2543. [[CrossRef](#)]
136. Hoekstra, R.; Nibourg, G.A.; van der Hoeven, T.V.; Plomer, G.; Seppen, J.; Ackermans, M.T.; Camus, S.; Kulik, W.; van Gulik, T.M.; Elferink, R.P.; et al. Phase 1 and phase 2 drug metabolism and bile acid production of HepaRG cells in a bioartificial liver in absence of dimethyl sulfoxide. *Drug Metab. Dispos.* **2013**, *41*, 562–567. [[CrossRef](#)] [[PubMed](#)]
137. Hoffmann, S.A.; Muller-Vieira, U.; Biemel, K.; Knobeloch, D.; Heydel, S.; Lubberstedt, M.; Nussler, A.K.; Andersson, T.B.; Gerlach, J.C.; Zeilinger, K. Analysis of drug metabolism activities in a miniaturized liver cell bioreactor for use in pharmacological studies. *Biotechnol. Bioeng.* **2012**, *109*, 3172–3181. [[CrossRef](#)] [[PubMed](#)]
138. Lubberstedt, M.; Muller-Vieira, U.; Biemel, K.M.; Darnell, M.; Hoffmann, S.A.; Knospel, F.; Wonne, E.C.; Knobeloch, D.; Nussler, A.K.; Gerlach, J.C.; et al. Serum-free culture of primary human hepatocytes in a

- miniaturized hollow-fibre membrane bioreactor for pharmacological in vitro studies. *J. Tissue Eng. Regen Med.* **2015**, *9*, 1017–1026. [[CrossRef](#)]
139. Pless, G.; Steffen, I.; Zeilinger, K.; Sauer, I.M.; Katzenz, E.; Kehr, D.C.; Roth, S.; Mieder, T.; Schwartlander, R.; Muller, C.; et al. Evaluation of primary human liver cells in bioreactor cultures for extracorporeal liver support on the basis of urea production. *Artif. Organs* **2006**, *30*, 686–694. [[CrossRef](#)]
140. Ring, A.; Gerlach, J.; Peters, G.; Pazin, B.J.; Minervini, C.F.; Turner, M.E.; Thompson, R.L.; Triolo, F.; Gridelli, B.; Miki, T. Hepatic maturation of human fetal hepatocytes in four-compartment three-dimensional perfusion culture. *Tissue Eng. Part C Methods* **2010**, *16*, 835–845. [[CrossRef](#)]
141. Schmelzer, E.; Triolo, F.; Turner, M.E.; Thompson, R.L.; Zeilinger, K.; Reid, L.M.; Gridelli, B.; Gerlach, J.C. Three-dimensional perfusion bioreactor culture supports differentiation of human fetal liver cells. *Tissue Eng. Part A* **2010**, *16*, 2007–2016. [[CrossRef](#)]
142. Tapia, F.; Vogel, T.; Genzel, Y.; Behrendt, I.; Hirschel, M.; Gangemi, J.D.; Reichl, U. Production of high-titer human influenza A virus with adherent and suspension MDCK cells cultured in a single-use hollow fiber bioreactor. *Vaccine* **2014**, *32*, 1003–1011. [[CrossRef](#)] [[PubMed](#)]
143. Ulvestad, M.; Darnell, M.; Molden, E.; Ellis, E.; Asberg, A.; Andersson, T.B. Evaluation of organic anion-transporting polypeptide 1B1 and CYP3A4 activities in primary human hepatocytes and HepaRG cells cultured in a dynamic three-dimensional bioreactor system. *J. Pharmacol. Exp. Ther.* **2012**, *343*, 145–156. [[CrossRef](#)] [[PubMed](#)]
144. Zeilinger, K.; Schreiter, T.; Darnell, M.; Soderdahl, T.; Lubberstedt, M.; Dillner, B.; Knobeloch, D.; Nussler, A.K.; Gerlach, J.C.; Andersson, T.B. Scaling down of a clinical three-dimensional perfusion multicompartment hollow fiber liver bioreactor developed for extracorporeal liver support to an analytical scale device useful for hepatic pharmacological in vitro studies. *Tissue Eng. Part C Methods* **2011**, *17*, 549–556. [[CrossRef](#)] [[PubMed](#)]
145. Egger, D.; Fischer, M.; Clementi, A.; Ribitsch, V.; Hansmann, J.; Kasper, C. Development and Characterization of a Parallelizable Perfusion Bioreactor for 3D Cell Culture. *Bioengineering* **2017**, *4*, 51. [[CrossRef](#)] [[PubMed](#)]
146. Kensah, G.; Gruh, I.; Viering, J.; Schumann, H.; Dahlmann, J.; Meyer, H.; Skvorc, D.; Bar, A.; Akhyari, P.; Heisterkamp, A.; et al. A Novel Miniaturized Multimodal Bioreactor for Continuous In Situ Assessment of Bioartificial Cardiac Tissue During Stimulation and Maturation. *Tissue Eng. Part C-Me* **2011**, *17*, 463–473. [[CrossRef](#)] [[PubMed](#)]
147. Mauney, J.R.; Sjostrom, S.; Blumberg, J.; Horan, R.; O’Leary, J.P.; Vunjak-Novakovic, G.; Volloch, V.; Kaplan, D.L. Mechanical stimulation promotes osteogenic differentiation of human bone marrow stromal cells on 3-D partially demineralized bone scaffolds in vitro. *Calcif. Tissue Int.* **2004**, *74*, 458–468. [[CrossRef](#)] [[PubMed](#)]
148. Mol, A.; Driessens, N.J.; Rutten, M.C.; Hoerstrup, S.P.; Bouting, C.V.; Baaijens, F.P. Tissue engineering of human heart valve leaflets: A novel bioreactor for a strain-based conditioning approach. *Ann. Biomed. Eng.* **2005**, *33*, 1778–1788. [[CrossRef](#)]
149. Saini, S.; Wick, T.M. Concentric cylinder bioreactor for production of tissue engineered cartilage: Effect of seeding density and hydrodynamic loading on construct development. *Biotechnol. Prog.* **2003**, *19*, 510–521. [[CrossRef](#)]
150. Scaglione, S.; Zerega, B.; Badano, R.; Benatti, U.; Fato, M.; Quarto, R. A three-dimensional traction/torsion bioreactor system for tissue engineering. *Int. J. Artif. Organs* **2010**, *33*, 362–369. [[CrossRef](#)]
151. Tsimbouri, P.M.; Childs, P.G.; Pemberton, G.D.; Yang, J.; Jayawarna, V.; Orapiriyakul, W.; Burgess, K.; Gonzalez-Garcia, C.; Blackburn, G.; Thomas, D.; et al. Stimulation of 3D osteogenesis by mesenchymal stem cells using a nanovibrational bioreactor. *Nat. Biomed. Eng.* **2017**, *1*, 758–770. [[CrossRef](#)]
152. Wang, B.; Wang, G.; To, F.; Butler, J.R.; Claude, A.; McLaughlin, R.M.; Williams, L.N.; de Jongh Curry, A.L.; Liao, J. Myocardial scaffold-based cardiac tissue engineering: Application of coordinated mechanical and electrical stimulations. *Langmuir* **2013**, *29*, 11109–11117. [[CrossRef](#)] [[PubMed](#)]
153. Weiss, S.; Henle, P.; Roth, W.; Bock, R.; Boeuf, S.; Richter, W. Design and characterization of a new bioreactor for continuous ultra-slow uniaxial distraction of a three-dimensional scaffold-free stem cell culture. *Biotechnol. Prog.* **2011**, *27*, 86–94. [[CrossRef](#)] [[PubMed](#)]
154. Lancaster, M.A.; Renner, M.; Martin, C.A.; Wenzel, D.; Bicknell, L.S.; Hurles, M.E.; Homfray, T.; Penninger, J.M.; Jackson, A.P.; Knoblich, J.A. Cerebral organoids model human brain development and microcephaly. *Nature* **2013**, *501*, 373–379. [[CrossRef](#)] [[PubMed](#)]

155. Qian, X.; Nguyen, H.N.; Song, M.M.; Hadiono, C.; Ogden, S.C.; Hammack, C.; Yao, B.; Hamersky, G.R.; Jacob, F.; Zhong, C.; et al. Brain-Region-Specific Organoids Using Mini-bioreactors for Modeling ZIKV Exposure. *Cell* **2016**, *165*, 1238–1254. [[CrossRef](#)]
156. Rebelo, S.P.; Costa, R.; Silva, M.M.; Marcelino, P.; Brito, C.; Alves, P.M. Three-dimensional co-culture of human hepatocytes and mesenchymal stem cells: Improved functionality in long-term bioreactor cultures. *J. Tissue Eng. Regen Med.* **2017**, *11*, 2034–2045. [[CrossRef](#)]
157. Barrila, J.; Yang, J.; Crabbe, A.; Sarker, S.F.; Liu, Y.; Ott, C.M.; Nelman-Gonzalez, M.A.; Clemett, S.J.; Nydam, S.D.; Forsyth, R.J.; et al. Three-dimensional organotypic co-culture model of intestinal epithelial cells and macrophages to study *Salmonella enterica* colonization patterns. *NPJ Microgravity* **2017**, *3*, 10. [[CrossRef](#)]
158. Crabbe, A.; Sarker, S.F.; Van Houdt, R.; Ott, C.M.; Leys, N.; Cornelis, P.; Nickerson, C.A. Alveolar epithelium protects macrophages from quorum sensing-induced cytotoxicity in a three-dimensional co-culture model. *Cell Microbiol.* **2011**, *13*, 469–481. [[CrossRef](#)]
159. Devarasetty, M.; Wang, E.; Soker, S.; Skardal, A. Mesenchymal stem cells support growth and organization of host-liver colorectal-tumor organoids and possibly resistance to chemotherapy. *Biofabrication* **2017**, *9*, 021002. [[CrossRef](#)]
160. DiStefano, T.; Chen, H.Y.; Panebianco, C.; Kaya, K.D.; Brooks, M.J.; Gieser, L.; Morgan, N.Y.; Pohida, T.; Swaroop, A. Accelerated and Improved Differentiation of Retinal Organoids from Pluripotent Stem Cells in Rotating-Wall Vessel Bioreactors. *Stem Cell Rep.* **2018**, *10*, 300–313. [[CrossRef](#)]
161. Gerecht-Nir, S.; Cohen, S.; Itskovitz-Eldor, J. Bioreactor cultivation enhances the efficiency of human embryoid body (hEB) formation and differentiation. *Biotechnol. Bioeng.* **2004**, *86*, 493–502. [[CrossRef](#)]
162. Goodwin, T.J.; McCarthy, M.; Cohrs, R.J.; Kaufer, B.B. 3D tissue-like assemblies: A novel approach to investigate virus-cell interactions. *Methods* **2015**, *90*, 76–84. [[CrossRef](#)] [[PubMed](#)]
163. Salerno-Goncalves, R.; Fasano, A.; Sztein, M.B. Development of a Multicellular Three-dimensional Organotypic Model of the Human Intestinal Mucosa Grown Under Microgravity. *J. Vis. Exp.* **2016**, *113*, e54148. [[CrossRef](#)] [[PubMed](#)]
164. Skardal, A.; Devarasetty, M.; Rodman, C.; Atala, A.; Soker, S. Liver-Tumor Hybrid Organoids for Modeling Tumor Growth and Drug Response In Vitro. *Ann. Biomed. Eng.* **2015**, *43*, 2361–2373. [[CrossRef](#)] [[PubMed](#)]
165. Wilkinson, D.C.; Mellody, M.; Meneses, L.K.; Hope, A.C.; Dunn, B.; Gomperts, B.N. Development of a Three-Dimensional Bioengineering Technology to Generate Lung Tissue for Personalized Disease Modeling. *Curr. Protoc. Stem Cell Biol.* **2018**, *46*, e56. [[CrossRef](#)] [[PubMed](#)]
166. Cheng, M.; Moretti, M.; Engelmayr, G.C.; Freed, L.E. Insulin-like growth factor-I and slow, bi-directional perfusion enhance the formation of tissue-engineered cardiac grafts. *Tissue Eng. Part A* **2009**, *15*, 645–653. [[CrossRef](#)] [[PubMed](#)]
167. Chung, S.; Sudo, R.; Mack, P.J.; Wan, C.R.; Vickerman, V.; Kamm, R.D. Cell migration into scaffolds under co-culture conditions in a microfluidic platform. *Lab. Chip* **2009**, *9*, 269–275. [[CrossRef](#)]
168. Daley, E.L.H.; Kuttig, J.; Stegemann, J.P. Development of Modular, Dual-Perfused Osteochondral Constructs for Cartilage Repair. *Tissue Eng. Part C Methods* **2019**, *25*, 127–136. [[CrossRef](#)]
169. Ghiaseddin, A.; Pouri, H.; Soleimani, M.; Vasheghani-Farahani, E.; Ahmadi Tafti, H.; Hashemi-Najafabadi, S. Cell laden hydrogel construct on-a-chip for mimicry of cardiac tissue in-vitro study. *Biochem. Biophys. Res. Commun.* **2017**, *484*, 225–230. [[CrossRef](#)]
170. Goldman, S.M.; Barabino, G.A. Spatial Engineering of Osteochondral Tissue Constructs Through Microfluidically Directed Differentiation of Mesenchymal Stem Cells. *Biores Open Access* **2016**, *5*, 109–117. [[CrossRef](#)]
171. Gottwald, E.; Nies, C.; Wuchter, P.; Saffrich, R.; Truckenmuller, R.; Giselbrecht, S. A Microcavity Array-Based 3D Model System of the Hematopoietic Stem Cell Niche. *Methods Mol. Biol.* **2019**, *2017*, 85–95. [[CrossRef](#)]
172. Grayson, W.L.; Marolt, D.; Bhumiratana, S.; Frohlich, M.; Guo, X.E.; Vunjak-Novakovic, G. Optimizing the medium perfusion rate in bone tissue engineering bioreactors. *Biotechnol. Bioeng.* **2011**, *108*, 1159–1170. [[CrossRef](#)] [[PubMed](#)]
173. Hoerstrup, S.P.; Zund, G.; Sodian, R.; Schnell, A.M.; Grunenfelder, J.; Turina, M.I. Tissue engineering of small caliber vascular grafts. *Eur. J. Cardiothorac. Surg.* **2001**, *20*, 164–169. [[CrossRef](#)]
174. Hwa, A.J.; Fry, R.C.; Sivaraman, A.; So, P.T.; Samson, L.D.; Stolz, D.B.; Griffith, L.G. Rat liver sinusoidal endothelial cells survive without exogenous VEGF in 3D perfused co-cultures with hepatocytes. *FASEB J.* **2007**, *21*, 2564–2579. [[CrossRef](#)] [[PubMed](#)]

175. Kenar, H.; Kose, G.T.; Toner, M.; Kaplan, D.L.; Hasirci, V. A 3D aligned microfibrous myocardial tissue construct cultured under transient perfusion. *Biomaterials* **2011**, *32*, 5320–5329. [CrossRef] [PubMed]
176. Kuiper, N.J.; Wang, Q.G.; Cartmell, S.H. A Perfusion Co-Culture Bioreactor for Osteochondral Tissue Engineered Plugs. *J. Biomater. Tiss Eng.* **2014**, *4*, 162–171. [CrossRef]
177. Lee, P.S.; Eckert, H.; Hess, R.; Gelinsky, M.; Rancourt, D.; Krawetz, R.; Cuniberti, G.; Scharnweber, D. Developing a Customized Perfusion Bioreactor Prototype with Controlled Positional Variability in Oxygen Partial Pressure for Bone and Cartilage Tissue Engineering. *Tissue Eng. Part C Methods* **2017**, *23*, 286–297. [CrossRef]
178. Li, Z.; Sun, H.; Zhang, J.; Zhang, H.; Meng, F.; Cui, Z. Development of in vitro 3D TissueFlex(R) islet model for diabetic drug efficacy testing. *PLoS ONE* **2013**, *8*, e72612. [CrossRef]
179. Liu, X.G.; Jiang, H.K. Preparation of an osteochondral composite with mesenchymal stem cells as the single-cell source in a double-chamber bioreactor. *Biotechnol. Lett.* **2013**, *35*, 1645–1653. [CrossRef]
180. Mahmoudifar, N.; Doran, P.M. Osteogenic differentiation and osteochondral tissue engineering using human adipose-derived stem cells. *Biotechnol. Prog.* **2013**, *29*, 176–185. [CrossRef]
181. Mauleon, G.; Fall, C.P.; Eddington, D.T. Precise spatial and temporal control of oxygen within in vitro brain slices via microfluidic gas channels. *PLoS ONE* **2012**, *7*, e43309. [CrossRef]
182. Radisic, M.; Marsano, A.; Maidhof, R.; Wang, Y.; Vunjak-Novakovic, G. Cardiac tissue engineering using perfusion bioreactor systems. *Nat. Protoc.* **2008**, *3*, 719–738. [CrossRef] [PubMed]
183. Rieke, M.; Gottwald, E.; Weibezaehn, K.F.; Layer, P.G. Tissue reconstruction in 3D-spheroids from rodent retina in a motion-free, bioreactor-based microstructure. *Lab. Chip* **2008**, *8*, 2206–2213. [CrossRef] [PubMed]
184. Schepers, A.; Li, C.; Chhabra, A.; Seney, B.T.; Bhatia, S. Engineering a perfusable 3D human liver platform from iPS cells. *Lab. Chip* **2016**, *16*, 2644–2653. [CrossRef] [PubMed]
185. Song, J.J.; Guyette, J.P.; Gilpin, S.E.; Gonzalez, G.; Vacanti, J.P.; Ott, H.C. Regeneration and experimental orthotopic transplantation of a bioengineered kidney. *Nat. Med.* **2013**, *19*, 646–651. [CrossRef]
186. Song, L.; Zhou, Q.; Duan, P.; Guo, P.; Li, D.; Xu, Y.; Li, S.; Luo, F.; Zhang, Z. Successful development of small diameter tissue-engineering vascular vessels by our novel integrally designed pulsatile perfusion-based bioreactor. *PLoS ONE* **2012**, *7*, e42569. [CrossRef]
187. Trietsch, S.J.; Israels, G.D.; Joore, J.; Hankemeier, T.; Vulto, P. Microfluidic titer plate for stratified 3D cell culture. *Lab. Chip* **2013**, *13*, 3548–3554. [CrossRef]
188. van Midwoud, P.M.; Groothuis, G.M.; Merema, M.T.; Verpoorte, E. Microfluidic biochip for the perfusion of precision-cut rat liver slices for metabolism and toxicology studies. *Biotechnol. Bioeng.* **2010**, *105*, 184–194. [CrossRef]
189. van Midwoud, P.M.; Merema, M.T.; Verpoorte, E.; Groothuis, G.M. Microfluidics enables small-scale tissue-based drug metabolism studies with scarce human tissue. *J. Lab. Autom.* **2011**, *16*, 468–476. [CrossRef]
190. Zhou, W.; Chen, Y.; Roh, T.; Lin, Y.; Ling, S.; Zhao, S.; Lin, J.D.; Khalil, N.; Cairns, D.M.; Manousiouthakis, E.; et al. Multifunctional Bioreactor System for Human Intestine Tissues. *ACS Biomater. Sci. Eng.* **2018**, *4*, 231–239. [CrossRef]
191. Pekor, C.; Gerlach, J.C.; Nettleship, I.; Schmelzer, E. Induction of Hepatic and Endothelial Differentiation by Perfusion in a Three-Dimensional Cell Culture Model of Human Fetal Liver. *Tissue Eng. Part C Methods* **2015**, *21*, 705–715. [CrossRef]
192. Visone, R.; Talo, G.; Occhetta, P.; Cruz-Moreira, D.; Lopa, S.; Pappalardo, O.A.; Redaelli, A.; Moretti, M.; Rasponi, M. A microscale biomimetic platform for generation and electro-mechanical stimulation of 3D cardiac microtissues. *APL Bioeng.* **2018**, *2*, 046102. [CrossRef] [PubMed]
193. Esch, E.W.; Bahinski, A.; Huh, D. Organs-on-chips at the frontiers of drug discovery. *Nat. Rev. Drug Discov.* **2015**, *14*, 248–260. [CrossRef] [PubMed]
194. Huh, D.; Matthews, B.D.; Mammoto, A.; Montoya-Zavala, M.; Hsin, H.Y.; Ingber, D.E. Reconstituting organ-level lung functions on a chip. *Science* **2010**, *328*, 1662–1668. [CrossRef] [PubMed]

195. Bein, A.; Shin, W.; Jalili-Firoozinezhad, S.; Park, M.H.; Sontheimer-Phelps, A.; Tovaglieri, A.; Chalkiadaki, A.; Kim, H.J.; Ingber, D.E. Microfluidic Organ-on-a-Chip Models of Human Intestine. *Cell. Mol. Gastroenterol. Hepatol.* **2018**, *5*, 659–668. [[CrossRef](#)] [[PubMed](#)]
196. Wu, Q.; Liu, J.; Wang, X.; Feng, L.; Wu, J.; Zhu, X.; Wen, W.; Gong, X. Organ-on-a-chip: Recent breakthroughs and future prospects. *Biomed. Eng. Online* **2020**, *19*, 9. [[CrossRef](#)] [[PubMed](#)]

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).