

Transverse mass distribution and charge asymmetry in W boson production to third order in QCD

Xuan Chen^{a,b,c,*}, Thomas Gehrmann^c, Nigel Glover^d, Alexander Huss^e, Tong-Zhi Yang^c, Hua Xing Zhu^f

^a Institute for Theoretical Physics, Karlsruhe Institute of Technology, 76131 Karlsruhe, Germany

^b Institute for Astroparticle Physics, Karlsruhe Institute of Technology, 76344 Eggenstein-Leopoldshafen, Germany

^c Physik-Institut, Universität Zürich, Winterthurerstrasse 190, CH-8057 Zürich, Switzerland

^d Institute for Particle Physics Phenomenology, Physics Department, Durham University, Durham, DH1 3LE, UK

^e Theoretical Physics Department, CERN, 1211 Geneva 23, Switzerland

^f Zhejiang Institute of Modern Physics, Department of Physics, Zhejiang University, Hangzhou, 310027, China

ARTICLE INFO

Article history:

Received 25 January 2023

Received in revised form 22 March 2023

Accepted 24 March 2023

Available online 28 March 2023

Editor: B. Grinstein

ABSTRACT

Charged gauge boson production at hadron colliders is a fundamental benchmark for the extraction of electroweak parameters and the understanding of the proton structure. To enable precision phenomenology for this process, we compute the third-order ($N^3\text{LO}$) QCD corrections to the rapidity distribution and charge asymmetry in W boson production and to the transverse mass distribution of its decay products. Our results display substantial QCD corrections in kinematic regions relevant for Tevatron and LHC measurements. We compare the numerical magnitude of the $N^3\text{LO}$ corrections with uncertainties from electroweak input parameters and illustrate their potential impact on the determination of the W boson mass.

© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

Charged electroweak (EW) gauge bosons W^\pm are produced copiously through the charged-current Drell-Yan process at hadron colliders [1]. Measurements of the inclusive and differential properties of W production play a central role in tests of the Standard Model (SM) of particle physics and in the search for novel physics effects beyond it, allowing precision determinations of electroweak parameters such as the W boson mass and weak mixing angle, and of parton distribution functions (PDFs) in the proton. In the past, measurements have been performed at the Fermilab Tevatron and the Large Hadron Collider (LHC), substantially improving our knowledge about the SM [2–9].

Very recently, using a sample of approximately 4 million W bosons collected at the Tevatron, the CDF Collaboration reported a new measurement of the W boson mass using template fits to the transverse mass distribution in W boson decays and the transverse momentum distribution of the decay leptons [10]. The new

measurement displays significant tension with the SM expectation and has an unprecedentedly small uncertainty of ± 9.4 MeV. The new result calls for a careful assessment of the theoretical predictions for the charged-current Drell-Yan process and their associated uncertainties.

Due to their importance for SM measurements, precision predictions for Drell-Yan production have been among the first applications of perturbative QCD at next-to-leading order (NLO, [11]), next-to-next-to-leading order (NNLO, [12]), and the inclusive corrections were accomplished at third order ($N^3\text{LO}$) accuracy in QCD [13] recently. Fully differential NNLO QCD corrections [14–17], including the kinematics of the decay leptons, have been available for a while and are routinely used in the experimental analysis of Drell-Yan data.

Significant efforts have also gone in the derivation of EW [18–20] and mixed QCD-EW corrections [21–25] for W production and into the combination of fixed-order predictions with resummation of large logarithmic corrections [26–29].

In this letter we present for the first time differential predictions for W production at $N^3\text{LO}$ in QCD, including the gauge boson rapidity distributions and associated charge asymmetry as well as the transverse mass distribution. Compared to NNLO, we find large $N^3\text{LO}$ corrections to the rapidity distributions with non-overlapping scale uncertainty bands. For the normalized transverse

* Corresponding author.

E-mail addresses: xuan.chen@uzh.ch (X. Chen), thomas.gehrmann@uzh.ch (T. Gehrmann), e.w.n.glover@durham.ac.uk (N. Glover), alexander.huss@cern.ch (A. Huss), toyang@physik.uzh.ch (T.-Z. Yang), zhuhx@zju.edu.cn (H.X. Zhu).

mass distribution, which plays an important role in the W mass measurement, we observe a high perturbative stability of the predictions from NNLO to N³LO. We quantify the impact of varying EW input parameters around the peak region of the transverse mass distribution, finding substantial effects that could have an important impact on future precision EW measurements using charged-current Drell-Yan production.

2. Methodology

In this letter, we calculate the differential cross section $d\sigma_{DY}$, of charged-current production using the q_T -subtraction formalism [30]:

$$d\sigma_{DY}^{N^3LO} = \mathcal{F}_{DY}^{N^3LO} \otimes d\sigma_{DY}^{LO}|_{q_T < q_T^{cut}} + d\sigma_{DY+jet}^{NNLO}|_{q_T > q_T^{cut}}, \quad (1)$$

where a slicing parameter q_T^{cut} is introduced to separate unresolved and resolved contributions. Our study is based on an established framework at N³LO [31–33] which integrates the unresolved and resolved contributions in a computationally demanding manner. The independence of the results on the unphysical slicing parameter q_T^{cut} serves as a strong check.

The unresolved contribution $\mathcal{F}_{DY}^{N^3LO} \otimes d\sigma_{DY}^{LO}|_{q_T < q_T^{cut}}$ denotes the fixed-order prediction for producing a W boson with transverse momentum q_T less than q_T^{cut} within Soft-Collinear Effective Theory (SCET) [34–38]. It can be expanded into logarithmic terms $\alpha_s^m \ln^n(q_T^{cut}/M_W)$ and constant terms. All logarithmic terms can be predicted through to N³LO [32,39] using the rapidity renormalization group formalism [40]. The key ingredients to achieve N³LO accuracy for colour-singlet production are the constant terms. They arise from the boundary conditions for the renormalization group equation, namely the rapidity-divergent transverse-momentum-dependent soft function [41] and beam functions [42–44] at three loops, as well as the massless QCD form factor [45–47].

The resolved contribution above q_T^{cut} is computed using the NNLOJET code for charged-current Drell-Yan-plus-jet production at NNLO [48,49]. It is fully differential at NNLO accuracy by employing the antenna subtraction method [50–52]. Sufficient numerical precision is mandatory to enable the cancellation between resolved and unresolved contributions at q_T^{cut} . This is achieved through dedicated optimization [29,33,39] of phase space generation and subtraction terms to enable robust coverage in the unresolved regions for small values of the slicing parameter q_T^{cut} . The cancellation of q_T^{cut} -dependent terms between resolved and unresolved contributions in (1) is accurate only up to power-suppressed terms at $\mathcal{O}(\alpha_s^3(q_T^{cut}/M_W)^2)$ which are unaccounted for in the unresolved piece. These terms are found to be sufficiently suppressed to no longer affect the final result for $q_T^{cut} \sim 1.5$ (0.75) GeV at the LHC (Tevatron), which is validated for each LHC (Tevatron) observable by varying q_T^{cut} by ± 0.5 (0.25) GeV.

The decay of the W boson into a charged lepton and a neutrino is described at leading order with a Breit-Wigner parametrisation of the W propagator using a fixed width. To assess the impact of higher order QCD corrections and EW input parameters, we use the PDG [53] values $M_W = 80.379$ GeV and $\Gamma_W = 2.085$ GeV as the default setup and compare predictions with variations of M_W and Γ_W .

3. Results

Applying the q_T -subtraction method described above, we study charged-current Drell-Yan production at fully differential N³LO accuracy in proton-proton collisions with centre-of-mass energy at 13 TeV. We use the central member of NNPDF3.1 and NNPDF4.0 NNLO PDFs [54,55] with $\alpha_s(M_Z) = 0.118$ throughout the calculation and the scale evolution is performed with LHAPDF [56].




Fig. 1. W boson rapidity distributions from LO to N³LO accuracy at the LHC. The coloured bands represent theory uncertainties from 7-point scale variation. The bottom panels show the ratio with respect to NNLO, for three different values of q_T^{cut} .

The electroweak couplings are determined using the G_μ scheme with: $M_Z = 91.1876$ GeV, $\Gamma_Z = 2.4952$ GeV, $G_F = 1.1663787 \times 10^{-5}$ GeV⁻² [53]. The CKM parameters are taken at their PDG values [53] in all Tevatron predictions, while a diagonal CKM matrix is used for LHC predictions. For absolute cross sections, the CKM effects are negligible for LHC energies (0.2%) but relevant at 2% level at Tevatron energies (largely due to the different partonic composition in proton-antiproton collisions). For normalized distributions without fiducial cuts, the CKM effects are negligible throughout. The central factorisation and renormalisation scales are chosen to be the invariant mass of final state leptons, $\mu_F = \mu_R = m_{l\nu}$. To estimate theoretical uncertainties, we adopt the 7-point scale variation of μ_F and μ_R by a factor of two while enforcing $1/2 \leq \mu_F/\mu_R \leq 2$.

In Fig. 1, we show the rapidity distributions of the Drell-Yan pairs from W⁺ and W⁻ decays, with no fiducial cuts applied. Fixed order contributions with up to N³LO accuracy are included with the bottom panels showing their ratio with respect to the central NNLO result. The coloured bands represent theory uncertainties from the 7-point scale variation and the error bars indicate the numerical integration error. Our state-of-the-art predictions at N³LO accuracy amount to a contribution of about -2.5% with respect to NNLO with relatively flat corrections for all rapidities. While the NLO and NNLO scale variation bands overlap, the N³LO prediction is found to be non-overlapping with the previous order




Fig. 2. W boson charge asymmetry distribution from LO to N³LO at the LHC. The coloured bands represent theory uncertainties from 31 scale variations. The bottom panel is the ratio with respect to NNLO.

within the respective scale uncertainties. This feature at N³LO has already been observed for the total cross sections for neutral current [57,58] and charged-current [13] Drell-Yan production and for the neutral-current Drell-Yan rapidity distribution [33] and fiducial cross sections [59]. The relative size of scale variation remains comparable at NNLO and N³LO at about $\pm 1\%$ for central rapidity and slightly increasing at large rapidity. We use three different q_T^{cut} values (1, 1.5 and 2 GeV) to confirm the q_T^{cut} -independence of the results within integration errors. A strong check on our results is provided by the rapidity-integrated charged current Drell-Yan cross section at N³LO, where our results for $q_T^{\text{cut}} = 1.5$ GeV agree with [13] within our numerical integration error of 1.5 per-mille.

The W boson charge asymmetry A_W at hadron colliders reveals details of the proton structure. It has been measured at the Tevatron [60,61] and the LHC [7,9,62] and is defined as

$$A_W(|y_W|) = \frac{d\sigma/d|y_{W^+}| - d\sigma/d|y_{W^-}|}{d\sigma/d|y_{W^+}| + d\sigma/d|y_{W^-}|}. \quad (2)$$

In Fig. 2, we display the predictions of $A_W(|y_W|)$ at 13 TeV centre of mass energy with up to N³LO corrections. We independently vary the scale choices between the numerator and the denominator of Eq. (2) while requiring $1/2 \leq \mu/\mu' \leq 2$ for any pair of scales, leading to 31 combinations. Their envelope is used to estimate the theoretical uncertainty. We observe positive N³LO corrections of about 2% relative to the NNLO predictions. The N³LO contribution is not flat in rapidity. In contrast to the individual rapidity distributions, the charge asymmetry converges well from NLO to N³LO with scale variation uncertainty reduced to about $\pm 1.5\%$ at N³LO.

Finally, we consider the transverse mass distribution in charged-current Drell-Yan production. The transverse mass is constructed as

$$m_T^{W^\pm} = \sqrt{2E_T^{\ell^\pm} E_T^{\nu}(1 - \cos\Delta\phi)}, \quad (3)$$

with $E_T^{\ell^\pm(\nu)}$ denoting the transverse energies of the final state charged lepton and neutrino and $\Delta\phi$ being their azimuthal angle difference. It is a characteristic observable in measurements of M_W [3–5,10] and Γ_W [63,64] at hadron colliders, since its distribution peaks around M_W and the shape of its tail is sensitive to Γ_W . Precise predictions for the $m_T^{W^\pm}$ distribution are vital for the measurement of W boson mass and width, which are based on fitting theory templates for the normalized distribution to data in the experimental analysis. The most recent measurement of M_W by CDFII collaboration reports ± 9.4 MeV overall uncertainty among





Fig. 3. Normalised W[±] transverse mass distribution from LO to N³LO accuracy at the Tevatron without (upper) and with (lower) CDFII fiducial cuts. The coloured bands represent theory uncertainties from 7-point scale variation. The bottom panel is the ratio with respect to NNLO, with different cutoff q_T^{cut} .

which ± 5.2 MeV arises from theoretical modelling [10]. The different sources of modelling uncertainties have subsequently been revisited [65,66], largely supporting the CDFII approach [10] while however not accounting for the state-of-the-art fixed-order predictions in mixed QCD-EW [25] and fixed-order QCD corrections.

Fig. 3 presents the normalized W[±] boson transverse mass distribution at the Tevatron. With the newly computed N³LO corrections, it establishes a new state-of-the-art in the precise description of this observable. The inclusive distribution without fiducial cuts is displayed in the upper frame, while the fiducial cuts on charged lepton and neutrino of the CDFII analysis [10] are applied in the lower frame: $p_{T,\ell}, E_{T,\nu} \in [30, 55]$ GeV, $|\eta_\ell| < 1$ and $p_{T,W^\pm} < 15$ GeV. For the N³LO coefficient, we compensate the linear q_T^{cut} -dependence due to the fiducial cuts through a recoil prescription [67,68] where the unresolved contribution in Eq. (1) is active if all fiducial requirements are satisfied after boosting Born kinematics to finite q_T below q_T^{cut} . At N³LO, corrections are very uniform in the peak region for both inclusive and fiducial distribution, while displaying some kinematical dependence at low $m_T^{W^\pm}$ below 12 GeV in the inclusive distribution and below 68 GeV in the fiducial distribution. Starting from NNLO, we observe a stabilization of scale uncertainties to the level of $\pm 1\%$.

Our newly derived predictions for the $m_T^{W^\pm}$ distribution allow us to compare different sources of theory uncertainty arising from




Fig. 4. Normalized W^\pm transverse mass distribution at the Tevatron with CDFII fiducial cuts. The NLO to N³LO corrections are in the top panel, with different M_W values from PDG, CDFII and L3 in the middle panel and with different Γ_W with PDG central value and $\pm 1\sigma$ uncertainties in the bottom panel. All distributions are compared to the NNLO result with PDG central values. The coloured bands represent theory uncertainties from 7-point scale variation.

higher perturbative orders and from variations of the input parameters for M_W and Γ_W in Fig. 4. For this purpose, we consider the PDG baseline values $M_W = 80.379$ GeV and $\Gamma_W = 2.085$ GeV, supplemented by the values of M_W according to the measurement from CDFII (80.433 GeV) [10] and L3 (80.27 GeV) [69], as well as a variation of Γ_W within the PDG uncertainty of ± 42 MeV [53].

The top panel of Fig. 4 demonstrates that the normalized $m_T^{W^\pm}$ fiducial distribution changes substantially in shape when going from NLO to NNLO, but remains much more stable especially away from the peak region upon inclusion of the newly derived N³LO corrections. The sensitivity to the input parameters can thus be reliably quantified based on the NNLO predictions. The middle panel compares the normalized $m_T^{W^\pm}$ distributions for fixed $\Gamma_W = 2.085$ GeV and M_W values from CDFII [10] and L3 [69], thus quantifying the magnitude and shape of the resulting variations in the distribution. As expected, a strong sensitivity on M_W around the peak region is observed. Compared to the change in the normalized distribution from NLO to NNLO, we observe that an NLO-based template fit could experience a slight pull towards larger values of M_W to compensate for missing NNLO corrections. These variations are to be contrasted with the lower panel of Fig. 4, where Γ_W is varied by its PDG uncertainty of ± 42 MeV for fixed PDG M_W , which basically affects the distributions only above the peak region, with a magnitude comparable to the effect of the M_W variation. In contrast to the sensitivity on Γ_W of the CDFII measurement recently reported in [65], a more realistic assessment of the uncertainties from Γ_W may thus be warranted.

4. Conclusions

In this letter, we have produced state-of-the-art predictions for differential distributions in the charged-current Drell-Yan production to third order in perturbative QCD. We used the q_T -subtraction method at N³LO, by combining an NNLO calculation for the production of a charged Drell-Yan pair at large q_T and leading-

power factorised predictions from SCET at small q_T . The robust numerical quality of both contributions allowed us to consistently check the cancellation of the q_T^{cut} dependence in our predictions.

We presented differential distributions for the rapidity charge asymmetry at the LHC and for $m_T^{W^\pm}$ at the Tevatron. Our results display modest N³LO corrections in asymmetries and normalized distributions, that are usually within the uncertainties of the NNLO predictions. N³LO perturbative uncertainties estimated by scale variations are found to be about $\pm 1\%$ to $\pm 1.5\%$ throughout. Distortions to the shape of the distributions are minimal at N³LO and only become visible outside the peak region of the $m_T^{W^\pm}$ distribution.

On the CDFII fiducial $m_T^{W^\pm}$ distribution [10], we studied the impact of perturbative corrections and of variations of EW input parameters. We observed only minimal effects in going from NNLO to N³LO. Variations of M_W and Γ_W within their respective experimental uncertainties led on the other hand to characteristic shifts in the shape of the normalized transverse mass distributions at a level between 2% to 6% around the peak region.

With the newly derived N³LO corrections, our results establish a new state-of-the-art for the perturbative description of W boson production at hadron colliders. They yield perturbative QCD uncertainties at the sub-per-cent level, which combine with recent results on QCD resummation, electroweak as well as mixed QCDxEW corrections to enable precision physics studies with upcoming LHC data.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Data on all results presented in this paper is available as supplementary material, see below.

Acknowledgements

The authors would like to thank Claude Duhr, Falko Dulat and Bernhard Mistlberger for discussions and Julien Baglio for providing detailed results for the predictions in [13]. We are grateful to Aude Gehrmann-De Ridder, Tom Morgan and Duncan Walker for their contributions to the V+jet process in the NNLOJET code. This work has received funding from the Swiss National Science Foundation (SNF) under contract 200020-204200 and from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme grant agreement 101019620 (ERC Advanced Grant TOPUP). This work is also supported in part by the UK Science and Technology Facilities Council (STFC) through grant ST/T001011/1 and in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under grant 396021762-TRR 257. H. X. Z. is supported by the Natural Science Foundation of China (NSFC) under contract No. 11975200.

Appendix A. Supplementary material

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.physletb.2023.137876>.

References

- [1] S.D. Drell, T.-M. Yan, Phys. Rev. Lett. 25 (1970) 316, Phys. Rev. Lett. 25 (1970) 902 (Erratum).
- [2] T. Aaltonen, et al., CDF, Phys. Rev. Lett. 108 (2012) 151803, arXiv:1203.0275 [hep-ex].

- [3] V.M. Abazov, et al., D0, Phys. Rev. Lett. 108 (2012) 151804, arXiv:1203.0293 [hep-ex].
- [4] M. Aaboud, et al., ATLAS, Eur. Phys. J. C 78 (2018) 110, Eur. Phys. J. C 78 (2018) 898 (Erratum), arXiv:1701.07240 [hep-ex].
- [5] R. Aaij, et al., LHCb, J. High Energy Phys. 01 (2022) 036, arXiv:2109.01113 [hep-ex].
- [6] T.A. Aaltonen, et al., CDF, D0, Phys. Rev. D 97 (2018) 112007, arXiv:1801.06283 [hep-ex].
- [7] G. Aad, et al., ATLAS, Eur. Phys. J. C 79 (2019) 760, arXiv:1904.05631 [hep-ex].
- [8] G. Aad, et al., ATLAS, Eur. Phys. J. C 80 (2020) 616, arXiv:1912.02844 [hep-ex].
- [9] A.M. Sirunyan, et al., CMS, Phys. Rev. D 102 (2020) 092012, arXiv:2008.04174 [hep-ex].
- [10] T. Aaltonen, et al., CDF, Science 376 (2022) 170.
- [11] G. Altarelli, R.K. Ellis, G. Martinelli, Nucl. Phys. B 157 (1979) 461.
- [12] R. Hamberg, W.L. van Neerven, T. Matsuura, Nucl. Phys. B 359 (1991) 343, Nucl. Phys. B 644 (2002) 403–404 (Erratum).
- [13] C. Duhr, F. Dulat, B. Mistlberger, J. High Energy Phys. 11 (2020) 143, arXiv:2007.13313 [hep-ph].
- [14] C. Anastasiou, L.J. Dixon, K. Melnikov, F. Petriello, Phys. Rev. D 69 (2004) 094008, arXiv:hep-ph/0312266.
- [15] K. Melnikov, F. Petriello, Phys. Rev. D 74 (2006) 114017, arXiv:hep-ph/0609070.
- [16] S. Catani, L. Cieri, G. Ferrera, D. de Florian, M. Grazzini, Phys. Rev. Lett. 103 (2009) 082001, arXiv:0903.2120 [hep-ph].
- [17] S. Catani, G. Ferrera, M. Grazzini, J. High Energy Phys. 05 (2010) 006, arXiv:1002.3115 [hep-ph].
- [18] S. Dittmaier, M. Krämer, Phys. Rev. D 65 (2002) 073007, arXiv:hep-ph/0109062.
- [19] U. Baur, D. Wackerth, Phys. Rev. D 70 (2004) 073015, arXiv:hep-ph/0405191.
- [20] C.M. Carloni Calame, G. Montagna, O. Nicrosini, A. Vicini, J. High Energy Phys. 10 (2007) 109, arXiv:0710.1722 [hep-ph].
- [21] S. Dittmaier, A. Huss, C. Schwinn, Nucl. Phys. B 904 (2016) 216, arXiv:1511.08016 [hep-ph].
- [22] S. Dittmaier, T. Schmidt, J. Schwarz, J. High Energy Phys. 12 (2020) 201, arXiv:2009.02229 [hep-ph].
- [23] A. Behring, F. Buccioni, F. Caola, M. Delfo, M. Jaquier, K. Melnikov, R. Rötsch, Phys. Rev. D 103 (2021) 013008, arXiv:2009.10386 [hep-ph].
- [24] L. Buonocore, M. Grazzini, S. Kallweit, C. Savoini, F. Tramontano, Phys. Rev. D 103 (2021) 114012, arXiv:2102.12539 [hep-ph].
- [25] A. Behring, F. Buccioni, F. Caola, M. Delfo, M. Jaquier, K. Melnikov, R. Rötsch, Phys. Rev. D 103 (2021) 113002, arXiv:2103.02671 [hep-ph].
- [26] C. Balazs, C.P. Yuan, Phys. Rev. D 56 (1997) 5558, arXiv:hep-ph/9704258.
- [27] G. Bozzi, S. Catani, G. Ferrera, D. de Florian, M. Grazzini, Phys. Lett. B 696 (2011) 207, arXiv:1007.2351 [hep-ph].
- [28] T. Becher, M. Neubert, D. Wilhelm, J. High Energy Phys. 02 (2012) 124, arXiv:1109.6027 [hep-ph].
- [29] W. Bizon, A. Gehrmann-De Ridder, T. Gehrmann, N. Glover, A. Huss, P.F. Monni, E. Re, L. Rottoli, D.M. Walker, Eur. Phys. J. C 79 (2019) 868, arXiv:1905.05171 [hep-ph].
- [30] S. Catani, M. Grazzini, Phys. Rev. Lett. 98 (2007) 222002, arXiv:hep-ph/0703012.
- [31] L. Cieri, X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss, J. High Energy Phys. 02 (2019) 096, arXiv:1807.11501 [hep-ph].
- [32] G. Billis, M.A. Ebert, J.K.L. Michel, F.J. Tackmann, Eur. Phys. J. Plus 136 (2021) 214, arXiv:1909.00811 [hep-ph].
- [33] X. Chen, T. Gehrmann, N. Glover, A. Huss, T.-Z. Yang, H.X. Zhu, Phys. Rev. Lett. 128 (2022) 052001, arXiv:2107.09085 [hep-ph].
- [34] C.W. Bauer, S. Fleming, M.E. Luke, Phys. Rev. D 63 (2000) 014006, arXiv:hep-ph/0005275.
- [35] C.W. Bauer, S. Fleming, D. Pirjol, I.W. Stewart, Phys. Rev. D 63 (2001) 114020, arXiv:hep-ph/0011336.
- [36] C.W. Bauer, D. Pirjol, I.W. Stewart, Phys. Rev. D 65 (2002) 054022, arXiv:hep-ph/0109045.
- [37] C.W. Bauer, S. Fleming, D. Pirjol, I.Z. Rothstein, I.W. Stewart, Phys. Rev. D 66 (2002) 014017, arXiv:hep-ph/0202088.
- [38] M. Beneke, A.P. Chapovsky, M. Diehl, T. Feldmann, Nucl. Phys. B 643 (2002) 431, arXiv:hep-ph/0206152.
- [39] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss, Y. Li, D. Neill, M. Schulze, I.W. Stewart, H.X. Zhu, Phys. Lett. B 788 (2019) 425, arXiv:1805.00736 [hep-ph].
- [40] J.-Y. Chiu, A. Jain, D. Neill, I.Z. Rothstein, J. High Energy Phys. 05 (2012) 084, arXiv:1202.0814 [hep-ph].
- [41] Y. Li, H.X. Zhu, Phys. Rev. Lett. 118 (2017) 022004, arXiv:1604.01404 [hep-ph].
- [42] M.-x. Luo, T.-Z. Yang, H.X. Zhu, Y.J. Zhu, Phys. Rev. Lett. 124 (2020) 092001, arXiv:1912.05778 [hep-ph].
- [43] M.A. Ebert, B. Mistlberger, G. Vita, J. High Energy Phys. 09 (2020) 146, arXiv:2006.05329 [hep-ph].
- [44] M.-x. Luo, T.-Z. Yang, H.X. Zhu, Y.J. Zhu, J. High Energy Phys. 06 (2021) 115, arXiv:2012.03256 [hep-ph].
- [45] P.A. Baikov, K.G. Chetyrkin, A.V. Smirnov, V.A. Smirnov, M. Steinhauser, Phys. Rev. Lett. 102 (2009) 212002, arXiv:0902.3519 [hep-ph].
- [46] R.N. Lee, A.V. Smirnov, V.A. Smirnov, J. High Energy Phys. 04 (2010) 020, arXiv:1001.2887 [hep-ph].
- [47] T. Gehrmann, E.W.N. Glover, T. Huber, N. Ikizlerli, C. Studerus, J. High Energy Phys. 06 (2010) 094, arXiv:1004.3653 [hep-ph].
- [48] A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, A. Huss, D.M. Walker, Phys. Rev. Lett. 120 (2018) 122001, arXiv:1712.07543 [hep-ph].
- [49] A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, A. Huss, D.M. Walker, Eur. Phys. J. C 79 (2019) 526, arXiv:1901.11041 [hep-ph].
- [50] A. Gehrmann-De Ridder, T. Gehrmann, E.W.N. Glover, J. High Energy Phys. 09 (2005) 056, arXiv:hep-ph/0505111.
- [51] A. Daleo, T. Gehrmann, D. Maitre, J. High Energy Phys. 04 (2007) 016, arXiv:hep-ph/0612257.
- [52] J. Currie, E.W.N. Glover, S. Wells, J. High Energy Phys. 04 (2013) 066, arXiv:1301.4693 [hep-ph].
- [53] P.A. Zyla, et al., Particle Data Group, PTEP 2020 (2020) 083C01.
- [54] R.D. Ball, et al., NNPDF, Eur. Phys. J. C 77 (2017) 663, arXiv:1706.00428 [hep-ph].
- [55] R.D. Ball, et al., NNPDF, Eur. Phys. J. C 82 (2022) 428, arXiv:2109.02653 [hep-ph].
- [56] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht, M. Schönherr, G. Watt, Eur. Phys. J. C 75 (2015) 132, arXiv:1412.7420 [hep-ph].
- [57] C. Duhr, F. Dulat, B. Mistlberger, Phys. Rev. Lett. 125 (2020) 172001, arXiv:2001.07717 [hep-ph].
- [58] C. Duhr, B. Mistlberger, J. High Energy Phys. 03 (2022) 116, arXiv:2111.10379 [hep-ph].
- [59] X. Chen, T. Gehrmann, E.W.N. Glover, A. Huss, P.F. Monni, E. Re, L. Rottoli, P. Torrielli, Phys. Rev. Lett. 128 (2022) 252001, arXiv:2203.01565 [hep-ph].
- [60] T. Aaltonen, et al., CDF, Phys. Rev. Lett. 102 (2009) 181801, arXiv:0901.2169 [hep-ex].
- [61] V.M. Abazov, et al., D0, Phys. Rev. Lett. 112 (2014) 151803, Phys. Rev. Lett. 114 (2015) 049901 (Erratum), arXiv:1312.2895 [hep-ex].
- [62] G. Aad, et al., ATLAS, Phys. Lett. B 701 (2011) 31, arXiv:1103.2929 [hep-ex].
- [63] T. Aaltonen, et al., CDF, Phys. Rev. Lett. 100 (2008) 071801, arXiv:0710.4112 [hep-ex].
- [64] V.M. Abazov, et al., D0, Phys. Rev. Lett. 103 (2009) 231802, arXiv:0909.4814 [hep-ex].
- [65] J. Isaacson, Y. Fu, C.P. Yuan, arXiv:2205.02788 [hep-ph], 2022.
- [66] J. Gao, D. Liu, K. Xie, Chin. Phys. C 46 (2022) 123110, arXiv:2205.03942 [hep-ph].
- [67] S. Camarda, L. Cieri, G. Ferrera, Eur. Phys. J. C 82 (2022) 575, arXiv:2111.14509 [hep-ph].
- [68] L. Buonocore, S. Kallweit, L. Rottoli, M. Wiesemann, Phys. Lett. B 829 (2022) 137118, arXiv:2111.13661 [hep-ph].
- [69] P. Achard, et al., L3, Eur. Phys. J. C 45 (2006) 569, arXiv:hep-ex/0511049.