

Search for heavy resonances and quantum black holes in $e\mu$, $e\tau$, and $\mu\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A search is reported for heavy resonances and quantum black holes decaying into $e\mu$, $e\tau$, and $\mu\tau$ final states in proton-proton collision data recorded by the CMS experiment at the CERN LHC during 2016–2018 at $\sqrt{s} = 13$ TeV, corresponding to an integrated luminosity of 138 fb^{-1} . The $e\mu$, $e\tau$, and $\mu\tau$ invariant mass spectra are reconstructed, and no evidence is found for physics beyond the standard model. Upper limits are set at 95% confidence level on the product of the cross section and branching fraction for lepton flavor violating signals. Three benchmark signals are studied: resonant τ sneutrino production in R parity violating supersymmetric models, heavy Z' gauge bosons with lepton flavor violating decays, and nonresonant quantum black hole production in models with extra spatial dimensions. Resonant τ sneutrinos are excluded for masses up to 4.2 TeV in the $e\mu$ channel, 3.7 TeV in the $e\tau$ channel, and 3.6 TeV in the $\mu\tau$ channel. A Z' boson with lepton flavor violating couplings is excluded up to a mass of 5.0 TeV in the $e\mu$ channel, up to 4.3 TeV in the $e\tau$ channel, and up to 4.1 TeV in the $\mu\tau$ channel. Quantum black holes in the benchmark model are excluded up to the threshold mass of 5.6 TeV in the $e\mu$ channel, 5.2 TeV in the $e\tau$ channel, and 5.0 TeV in the $\mu\tau$ channel. In addition, model-independent limits are extracted to allow comparisons with other models for the same final states and similar event selection requirements. The results of these searches provide the most stringent limits available from collider experiments for heavy particles that undergo lepton flavor violating decays.

KEYWORDS: Beyond Standard Model, Hadron-Hadron Scattering

ARXIV EPRINT: [2205.06709](https://arxiv.org/abs/2205.06709)

Contents

1	Introduction	1
2	The CMS detector	2
3	Collision data and simulated events	4
4	Event reconstruction and selection	4
4.1	Selection of events for the $e\mu$ final state	5
4.2	Selection of events for the $e\tau$ and $\mu\tau$ final states	6
5	Background estimation	8
6	Systematic uncertainties	9
7	Results and their interpretations	10
8	Summary	18
	The CMS collaboration	24

1 Introduction

Extensions of the standard model (SM) can accommodate heavy particles that undergo lepton flavor violating (LFV) decays, motivating thereby searches for deviations from the SM in the $e\mu$, $e\tau$, and $\mu\tau$ final states. This paper reports a search for such phenomena in these final states in proton-proton (pp) collisions at the CERN LHC, and is designed to be as model-independent as possible. Additionally, the results are interpreted in terms of the characteristics of the following possible states: a τ sneutrino ($\tilde{\nu}_\tau$), which can be the lightest SUSY particle (LSP) [1, 2] in R parity violating (RPV) supersymmetric (SUSY) models [3], a heavy Z' gauge boson in LFV models [4], and quantum black holes (QBHs) [5–8]. This is the first analysis searching for high-mass lepton flavor violating signals using the full Run 2 data set, recorded at $\sqrt{s} = 13$ TeV.

In RPV SUSY models, lepton flavor and lepton number are violated at lowest (Born) level in interactions between fermions and their superpartners. For resonant $\tilde{\nu}_\tau$ signals, the trilinear RPV part of the superpotential can be expressed as

$$W_{\text{RPV}} = \frac{1}{2}\lambda_{ijk}L_iL_j\bar{E}_k + \lambda'_{ijk}L_iQ_j\bar{D}_k, \tag{1.1}$$

where i, j , and k are generation indices, L and Q are the $SU(2)_L$ doublet superfields of the leptons and quarks, and \bar{E} and \bar{D} are the respective $SU(2)_L$ singlet superfields of the

charged leptons and down-like quarks. For simplicity, we assume that all RPV couplings vanish, except for those that are connected to the production and decay of the $\tilde{\nu}_\tau$, and we consider a SUSY mass hierarchy with $\tilde{\nu}_\tau$ as the LSP. In this model, the $\tilde{\nu}_\tau$ can be produced resonantly in pp collisions via the λ' coupling, and can decay either into dilepton final states via the λ couplings, or into quarks via λ' couplings. We consider only the final states with two charged leptons. Also, this analysis considers $\tilde{\nu}_\tau$ that decay promptly and are not long-lived [9], which we define as having a transverse displacement less than 1 mm from the production vertex. As long as the λ coupling is larger than 10^{-7} , a $\tilde{\nu}_\tau$ of mass 1 TeV will not be considered to be long-lived [3].

An extension of the SM through the addition of an extra U(1) gauge symmetry provides a massive Z' vector boson [4]. In our search, we assume that the Z' boson has identical couplings to SM particles as the SM Z boson, but that the Z' boson can also decay to the LFV $e\mu$, $e\tau$, and $\mu\tau$ final states, each with an assumed branching fraction of 10%. This value is defined in a benchmark scenario commonly used in these searches [10, 11].

Theories that invoke extra spatial dimensions can lower the fundamental Planck scale to the TeV region. Such theories also provide the possibility of producing microscopic QBHs [5, 6] at the LHC. In contrast to semiclassical thermal black holes that can decay to high-multiplicity final states, QBHs are nonthermal objects, expected to decay predominantly to pairs of particles. We consider the production of spin-0, colorless, neutral QBHs in a model with LFV [12], in which the cross section for QBH production depends on the threshold mass m_{th} in n additional spatial dimensions. The $n = 1$ possibility corresponds to the Randall–Sundrum (RS) brane-world model [13], and $n > 1$ corresponds to the Arkani-Hamed–Dimopoulos–Dvali (ADD) model [14]. While the resonant $\tilde{\nu}_\tau$ and Z' signals are expected to generate narrow peaks in the invariant mass spectrum of the lepton pair, the distribution of the QBH signal is characterized by a sharp edge at the threshold of QBH production, followed by a monotonic decrease at larger masses. Feynman diagrams for all these three models are shown in figure 1.

Similar searches in LFV dilepton mass spectra have been carried out by the CDF [15] and D0 [16] experiments at the Fermilab Tevatron in proton-antiproton collisions at $\sqrt{s} = 1.96$ TeV and by the ATLAS and CMS experiments at the LHC in pp collisions at $\sqrt{s} = 8$ TeV [17, 18] and 13 TeV [10, 11, 19].

This paper is structured as follows. The CMS detector is briefly described in section 2. A description of the collision data and the simulated event samples used in the analysis is given in section 3. The event reconstruction and selection are described in section 4 and the estimation of SM backgrounds is discussed in section 5. Systematic uncertainties are described in section 6, followed by the results and their statistical interpretation in section 7. The paper is summarized in section 8.

Tabulated results are provided in the HEPData record for this analysis [20].

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass




Figure 1. Leading order Feynman diagrams considered in our search. Left: Resonant production of a τ sneutrino in an RPV SUSY model that includes the subsequent decay into two leptons of different flavors. The $\tilde{\nu}_\tau$ is produced from the annihilation of two down quarks via the λ'_{311} coupling, and then decays via the λ couplings. Middle: Resonant production of a Z' boson with subsequent decay into two leptons of different flavors. Right: Production of quantum black holes in a model with extra dimensions that involves subsequent decay into two leptons of different flavors.

and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. Muons are measured in the pseudorapidity range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. Isolated particles of transverse momentum (p_T) of 100 GeV emitted within the pseudorapidity range $|\eta| < 1.4$ have track resolutions of 2.8% in p_T and 10 (30) μm in the transverse (longitudinal) impact parameter [21].

Events of interest are selected using a two-tiered trigger system. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [22]. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [23].

The single-muon trigger efficiency exceeds 90% over the full η acceptance, and the efficiency to reconstruct and identify muons is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative p_T resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps. The p_T resolution in the barrel is better than 7% for muons with p_T up to 1 TeV [24].

The single isolated electron trigger efficiency exceeds 80% over the full η acceptance of $|\eta| < 2.5$, and the efficiency to reconstruct electrons is greater than 95% for electrons with transverse momentum larger than 20 GeV [25]. The electron momentum is estimated by combining the energy measurement in the ECAL with the momentum measurement in the tracker. The momentum resolution for electrons with $p_T \approx 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.7 to 4.5%. It is generally better in the barrel region than in the endcaps, and also depends on the bremsstrahlung energy emitted by the electron as it traverses the material in front of the ECAL [26].

A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in ref. [27].

3 Collision data and simulated events

The data sample used for this analysis was collected during 2016–2018 pp operation at a center-of-mass energy of 13 TeV. After applying data quality requirements, such as requiring minimum fractions of active detector channels, the total integrated luminosity is 138 fb^{-1} .

Simulated samples of signal and background events are produced with several event generators. The RPV SUSY $\tilde{\nu}_\tau$, Z' , and QBH signal events are generated at leading order (LO) precision, using the CALCHEP 3.6 [28], PYTHIA 8.203 [29], and QBH 3.0 [12] Monte Carlo (MC) generators, respectively. The width of the Z' signal is taken as 3% of its mass, similar to that of the Z boson. The RPV $\tilde{\nu}_\tau$ and QBH signals are generated using the CTEQ6L [30] parton distribution functions (PDF), while the Z' boson signals are simulated using the NNPDF 3.0 PDF set in 2016 and the 3.1 PDF set [31] in 2017–2018, respectively. The LO RPV SUSY $\tilde{\nu}_\tau$ signal event yield is normalized to a next-to-leading order (NLO) calculation of the production cross section [32]; in this calculation the factorization and renormalization scales are set to the mass of the $\tilde{\nu}_\tau$. The POWHEG v2.0 event generator [33–36] is used to simulate both top quark pair ($t\bar{t}$) production, the dominant background over most of the dilepton mass region, and single top quark production. Diboson production, a significant background in the high mass region, is simulated at LO using the PYTHIA generator.

The MADGRAPH5_aMC@NLO v2.2.2 generator [37] is used to simulate Drell–Yan+jets production simulated at NLO with the FxFx jet matching and merging [38]. The cross sections used to normalize the contribution of these backgrounds are calculated at next-to-next-to-leading order for WW, ZZ, single top quark, and $t\bar{t}$ processes, and at NLO precision for WZ and Drell–Yan events. The POWHEG and MADGRAPH5_aMC@NLO generators are interfaced with PYTHIA for parton showering, fragmentation, and decays. The PYTHIA parameters for the underlying event description are set to the CUETP8M1 (CP5) tune [39] in 2016 (2017–2018) simulated samples.

The generated events are processed through a full simulation of the CMS detector, based on GEANT4 [40–42]. The simulated events incorporate additional pp interactions within the same or nearby bunch crossings, termed pileup, that are weighted to match the measured distribution of the number of interactions per bunch crossing in data. The simulated event samples are normalized to the integrated luminosity of the data. The products of the total acceptance and efficiency for the three signal models in this analysis are determined through MC simulation. The trigger and object reconstruction efficiencies are corrected to the values measured in data.

4 Event reconstruction and selection

The global event reconstruction is performed using a particle-flow algorithm [43], which aims to reconstruct and identify each individual particle with an optimized combination of all subdetector information. In this process, the identification of the particle type (photon,

electron, muon, and charged or neutral hadron) plays an important role in the determination of the particle’s direction and energy.

The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [44].

To reconstruct an electron candidate, energy deposits in the ECAL are first combined into clusters, assuming that each cluster represents a single particle. The clusters are then combined in a way consistent with bremsstrahlung emission, to produce a single “supercluster”, which represents an electron or photon. These superclusters are used to seed tracking algorithms, and if a resulting track is found, it is assigned to the supercluster to form an electron candidate.

To reconstruct a muon candidate, hits are first fitted separately to trajectories in the inner tracker detector and in the outer-muon system. The two trajectories are then combined in a global-muon track hypothesis.

Hadronic τ lepton decays (τ_h) are reconstructed from jets, using the hadrons-plus-strips algorithm [45], which combines 1 or 3 tracks with energy deposits in the calorimeters, to identify the τ lepton decay modes. Neutral pions are reconstructed as strips with dynamic size in η - ϕ , where ϕ is the azimuthal angle in radians, primarily from reconstructed photons, but also from reconstructed electrons which originate due to conversion of photons. The strip size varies as a function of the p_T of the electron or photon candidate.

To distinguish genuine τ_h decays from jets originating from the hadronization of quarks or gluons, and from electrons or muons, the DEEPTAU algorithm is used [46]. Information from all individual reconstructed particles near the τ_h axis is combined with properties of the τ_h candidate in the event to yield separate discriminants for jet, electron, and muon backgrounds. The probability that a jet is misidentified as τ_h by the DEEPTAU algorithm depends on the p_T and quark flavor of the jet. In simulated events of W boson production in association with jets, this probability has been estimated to be 0.43% for a τ_h identification efficiency of 70%. The probability that an electron (muon) is misidentified as a τ is 2.60 (0.03)% for a τ_h identification efficiency of 80 (>99)%.

4.1 Selection of events for the $e\mu$ final state

For the $e\mu$ selection, at least one prompt, isolated electron and at least one prompt, isolated muon are required in the event. This minimal selection facilitates a reinterpretation of the results in terms of models with more complex signal topologies than a single $e\mu$ pair.

Events that satisfy single electromagnetic-cluster or single-muon triggers, with respective p_T thresholds of 175 and 50 GeV for photons and muons, are selected. In 2018, the p_T threshold of the photon trigger was raised to 200 GeV. Electromagnetic energy deposited by an electron in the calorimeter activates the photon trigger, and the photon trigger is therefore as efficient as the corresponding electron trigger, while its weaker isolation requirements yield an event sample that can also be used in sideband analyses to estimate the background to the signal. Moreover, using only the muon trigger yields about 90% signal efficiency in masses above 1 TeV. Using the photon trigger together with the muon trigger ensures that the signal efficiency is very close to 100% in the high mass region.

The electron candidate must pass the dedicated selection criteria developed for high-energy electrons [25], which require the energy deposition in the ECAL to be consistent with that of an electron. The electron candidates are required to have $p_T > 35$ GeV and $|\eta| < 2.5$. The energy sum in the HCAL within a cone defined by $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.15$ centered around the electron candidate must be $< 5\%$ of its energy after it is corrected for jet activity unrelated to the primary interaction. The electron candidate must have a prompt track in the η - ϕ plane that has at most one missing hit in the pixel tracker and that is well-matched between the tracker and ECAL. The high-energy electron selection also requires electrons to be isolated. The scalar- p_T sum of tracks within a cone of radius $\Delta R = 0.3$ around the candidate's direction, excluding the candidate track, is required to be < 5 GeV. Moreover, the p_T sum of energy depositions in the calorimeters within the same cone, excluding the ECAL supercluster, is required to be $< 3\%$ of the p_T of the candidate [25]. The efficiency of the electron identification requirements, measured in data, is approximately 90% for the highly energetic electrons that are relevant for this search.

Muon candidates are required to have $p_T > 53$ GeV and $|\eta| < 2.4$. The muon candidate must pass the high- p_T muon identification criteria, which require that the transverse and longitudinal impact parameters of muon candidates relative to the primary vertex must be < 0.2 and < 0.5 cm, respectively. The track of the muon candidate must have at least one hit in the pixel detector and hits in at least six silicon-strip layers, and must contain matched segments in at least two muon detector planes. To suppress backgrounds arising from muons within jets, the scalar- p_T sum of all other tracks in the tracker within a cone of $\Delta R = 0.3$ around the muon candidate track is required to be $< 10\%$ of the p_T of the muon candidate. The relative uncertainty in the p_T of the muon track is required to be $< 30\%$. The efficiency of the muon identification and isolation requirements, measured in data, is about 90% for muons with $p_T > 50$ GeV.

To prevent a loss in signal efficiency resulting from the misidentification of the sign of the electron or muon charge at large p_T , the electron and muon are not required to have opposite charges. Since highly energetic muons can produce bremsstrahlung in the ECAL along the direction of the inner-muon trajectory, such muons can be misidentified as electrons. An electron candidate is therefore rejected if there is a muon candidate track with $p_T > 5$ GeV within $\Delta R < 0.1$ of the electron candidate track. Only one $e\mu$ pair is considered per event. When there is more than one $e\mu$ candidate, the pair with the highest invariant mass is selected for the analysis. The statistical interpretation is conducted by comparing the shape of the observed invariant $e\mu$ mass distribution with those expected for the signal and background.

4.2 Selection of events for the $e\tau$ and $\mu\tau$ final states

Events are required to have at least one prompt, isolated light lepton (electron or muon) and at least one prompt, isolated τ_h .

Events that satisfy single-electron triggers with thresholds of 27, 35, and 32 GeV in 2016, 2017, and 2018, respectively, are selected for the $e\tau$ channel. To recover efficiency in the high mass region, events that satisfy single electromagnetic cluster triggers are also

selected for this channel. The electron candidate must pass the same high-energy electron selection used for the $e\mu$ channel, but with an increased p_T threshold of 50 GeV.

The single-muon triggers used in the $e\mu$ channel are also used to collect the data samples in the $\mu\tau$ channel. The muon candidate must pass the same high- p_T muon identification criteria used for the $e\mu$ channel.

The τ_h candidate having the largest transverse momentum must have $p_T > 50$ GeV and $|\eta| < 2.3$. To reduce the rate of jets, electrons, and muons misidentified as τ_h , the τ_h candidate is required to satisfy the tight, loose, and tight working points of the respective DEEPTAU discriminator [46].

The low transverse mass (m_T) region is dominated by misidentified τ_h events, which have no genuine neutrinos, so the missing transverse momentum is small. Thus, a requirement of $m_T > 120$ GeV is applied which helps to reject misidentified τ_h background, where m_T is defined as:

$$m_T = \sqrt{2p_T^\ell p_T^{\text{miss}} [1 - \cos \Delta\phi(\vec{p}_T^\ell, \vec{p}_T^{\text{miss}})]}, \quad (4.1)$$

where ℓ denotes the light lepton, i.e., the electron in the $e\tau$ final state or the muon in the $\mu\tau$ final state, \vec{p}_T^{miss} is the missing transverse momentum vector in the event, and $\Delta\phi$ is the difference in the azimuthal angle between \vec{p}_T^ℓ and \vec{p}_T^{miss} .

A muon veto is applied in the $e\tau$ final state to remove overlap with the $\mu\tau$ and $e\mu$ final states. This veto rejects events if they contain any isolated muon with $p_T > 35$ GeV, $|\eta| < 2.4$, passing the high- p_T muon identification criteria and having tracker-based isolation < 0.15 . Events with a well-separated electron pair are also rejected. A well-separated electron pair is defined as two electrons with $\Delta R(e, e) > 0.5$, each of which has $p_T > 10$ GeV, $|\eta| < 2.5$, and passes a very loose working point ($> 95\%$ efficiency) of the electron identification criteria.

In the $\mu\tau$ channel, in order to avoid overlap with the $e\tau$ and $e\mu$ channels, events are vetoed if they contain an electron with $p_T > 35$ GeV that passes the high-energy electron selection criteria. Events with a well-separated muon pair are also rejected. A well-separated muon pair is defined as two muons passing the high- p_T identification criteria, with $p_T > 10$ GeV, $|\eta| < 2.4$, tracker-based isolation < 0.15 , and with $\Delta R(\mu, \mu) > 0.2$.

If there is more than one $e\tau$ or $\mu\tau$ pair in an event, then the pair with highest invariant mass is chosen. The leptons forming the pair are not required to have opposite electric charges. The statistical interpretations are conducted by comparing the shapes of the observed collinear mass distributions with those expected for the signals and backgrounds. For an $e\tau$ pair or a $\mu\tau$ pair, we use the collinear mass, which provides an estimate of the mass of the new resonance or QBH based on their observed decay products. It is justified by the observation that, since the mass scale of the signal is orders of magnitude higher than that of the τ lepton, the τ lepton decay products are highly Lorentz boosted in the direction of the τ candidate. The neutrino momentum can be approximated to have the same direction as the other, visible decay products of the τ lepton, so the component of \vec{p}_T^{miss} in the direction of the visible τ lepton decay products is used to estimate the neutrino p_T . The variable x_τ^{vis} , which is the fraction of energy carried by the visible decay products of the τ , is defined as $x_\tau^{\text{vis}} = p_T^{\text{vis}} / (p_T^{\text{vis}} + p_{T, \text{coll}}^{\text{miss}})$, where $p_{T, \text{coll}}^{\text{miss}}$ is the part of the p_T^{miss} that

is collinear with the τ_h p_T . The collinear mass m_{coll} can then be derived from the visible mass m_{vis} of the $e\tau$ or $\mu\tau$ system to be $m_{\text{coll}} = m_{\text{vis}}/\sqrt{x_{\tau}^{\text{vis}}}$, where m_{vis} is the invariant mass of the visible τ decay products.

5 Background estimation

The SM background in the LFV dilepton search includes several processes that produce a final state with two different-flavor charged leptons. For all channels, the dominant background contributions originate from $t\bar{t}$ production. Other less significant backgrounds originate from diboson (WW, WZ, and ZZ), $Z \rightarrow \ell^+\ell^-$ ($\ell = e, \mu, \tau$), $W\gamma$, and single top quark production processes. All these backgrounds are estimated from MC simulation. For the $Z \rightarrow \tau\tau$ process, all decay modes of τ lepton have been considered. Multijet and W+jets processes also contribute due to the misidentification of jets as leptons. These backgrounds are estimated from data.

For the $e\mu$ channel, to determine the contributions from W+jets and multijet processes to the $m_{e\mu}$ distribution, a control sample in data is defined using jet-to-electron misidentification rate (F_e). This rate is obtained from data, by selecting electron candidates with relaxed selection requirements, using events collected with a single electromagnetic-cluster trigger. Data sidebands obtained by inverting the selection requirements on either the electron isolation or shower shape variables are used to evaluate the contribution of jets passing the full electron selection in the control sample [47]. The jet-to-electron misidentification rate is then defined as the number of jets passing the full electron selection divided by the number of jet candidates in the sample. The rate is quantified in bins of misidentified electron p_T and pseudorapidity. The measured jet-to-electron rate is then used to estimate the W+jets and multijet contributions to $e\mu$ using data containing muons that pass the single-muon trigger and the full muon selection, and electron candidates satisfying relaxed selection requirements, but failing the full electron selection. Each event is weighted by the factor $F_e/(1 - F_e)$ to determine the overall contribution from the jet backgrounds. A correction is applied to account for the fraction of events in the control sample that have genuine leptons, estimated from simulation. The background originating from jets misidentified as electrons is about 10% of the total background.

Background from jets mimicking muons is estimated in a similar way, where identification criteria are loosened for muons in order to create a control sample enriched with jet candidates that are misidentified as muons. Then the jet-to-muon misidentification rate F_{μ} is defined as the number of jets passing the full muon selection divided by the number of jet candidates in the sample. The rate is quantified in bins of misidentified muon p_T and η as well. Events in data that have one well-identified electron and one candidate satisfying loose requirements of muon identification are selected, with the weight factor $F_{\mu}/(1 - F_{\mu})$ applied to estimate the jet-to-muon misidentification contribution in the signal region. The background originating from jets misidentified as muons is about 2% of the total background. Events with both electron and muon misidentified from jets are not considered since their contribution is expected to be small.

In the $\mu\tau$ and $e\tau$ channels, the most significant background after the $t\bar{t}$ and WW processes comes from W +jets and multijet processes, where jets are misidentified as τ_h candidates. This background is estimated using collision data, in a control sample with the same selection as the signal region, except that the m_T requirement is inverted to $m_T < 120$ GeV. From this low- m_T control sample, two subsamples are constructed to compute the probability for an accompanying jet to be misidentified as a τ_h :

- Subsample A requires τ_h candidates to fail the tight antijet discriminator working point but pass the loose working point.
- Subsample B requires τ_h to pass the tight working point.

A factor F_τ is calculated from these samples, defined as the ratio of the number of events in subsamples B and A. This factor is calculated as a function of the p_T of the τ_h candidate, the ratio of the p_T of the τ_h candidate to the p_T of its parent jet, and the pseudorapidity of the τ_h . The number of events expected from misidentified τ_h in the signal region is estimated from a control sample fulfilling the same criteria as events in the signal region, except that the τ_h candidates pass the looser working point but fail the tight working point of the antijet discriminator. Each event is weighted by the factor F_τ to determine its effective contribution. A correction is applied to account for the fraction of events in the control sample that have genuine τ_h candidates, estimated from simulation.

6 Systematic uncertainties

Various effects impact the shape and the normalization of the invariant or collinear mass distributions and lead to systematic uncertainties in the signal and background estimates. The uncertainty in the modeling of the invariant or collinear mass distributions reflects three types of systematic effects.

The first type includes those that affect both the shape and normalization of the mass distributions. For all channels, the dominant uncertainties arise from the leading $t\bar{t}$ and subleading WW backgrounds. They result in a 30–50% uncertainty in the number of $t\bar{t}$ and WW events at a dilepton mass scale of 2 TeV. The uncertainty in the WW background is estimated from the envelope of the resummed next-to-next-to-leading logarithmic calculation of the soft-gluon contributions to the cross section at NLO, as presented in ref. [48], using changes in the renormalization and factorization scales by factors of 2 and 0.5. Similarly, the uncertainty in the $t\bar{t}$ background is estimated by considering the variations in PDF and factorization scales, as discussed in ref. [49]. Other contributions include the uncertainty in the muon momentum scale, which is approximately 1–2% for 1 TeV muons and depends on their η and ϕ [50, 51]. The uncertainty in the muon efficiency (0.5% for 1 TeV muons) is considered in the $e\mu$ and $\mu\tau$ channels [51], and the uncertainty in the electron efficiency (varies between 1–5%, depending on p_T and η of the electrons) is considered in the $e\mu$ and $e\tau$ channels [25]. In the τ channels, the uncertainties in the τ_h identification (5% for 1 TeV τ leptons) and τ_h energy scale (1.5–4.0% for $p_T(\tau_h) > 100$ GeV, depending on the decay mode) are considered [45]. Uncertainties in the electron p_T scale and resolution, the muon p_T scale and resolution, and the pileup rate are also taken into account, but

they have negligible impact on the total background. The uncertainties in the determination of the trigger efficiencies have a very small impact on the expected event yields. The uncertainty associated with the choice of the PDF in the simulation is evaluated according to the PDF4LHC prescription [52].

The jet energy scale is determined with an uncertainty amounting to a few percent, depending on the jet p_T and η , using the p_T -balance method. This correction is applied to $Z/\gamma^* \rightarrow ee$, $Z/\gamma^* \rightarrow \mu\mu$, γ +jets, dijet, and multijet events [53]. The resulting effect on signal and background expectations is evaluated by varying the energies of jets in simulated events within their uncertainties, recalculating all kinematic observables, such as \vec{p}_T^{miss} , and reapplying the event selection criteria. The effects of uncertainties in the energy scale of the unclustered particles and jet energy resolution are evaluated in a similar way. These systematic uncertainties affect the shapes as well as the normalizations of the collinear mass distributions.

Uncertainties of the second type directly influence only the normalization of the mass distribution. A systematic uncertainty of 1.6% in the integrated luminosity [54–56] is taken for the backgrounds and signals. Among the uncertainties in the cross sections used for the normalization of various simulated backgrounds, the 5% uncertainty in the $t\bar{t}$ background is the most important. A systematic uncertainty of 50% is applied to the estimate of the misidentified jet background derived from data in all three channels. For τ_h final states, this uncertainty is obtained by using the misidentification probability F_τ derived from an independent control sample of $Z (\rightarrow \mu\mu)$ +jets events for the estimation of misidentified jet background. It is found, especially at high masses (≥ 1.5 TeV), that the collinear mass distributions obtained in the signal region using the F_τ calculated from the two independent control samples agree within 50% and are consistent within the statistical uncertainties. Therefore, an overall 50% systematic uncertainty is assigned to the estimation of this background.

Uncertainties of the third type are associated with limited sizes of event samples in the MC simulation of background processes [57]. In contrast to other uncertainties, they are not correlated between bins of the invariant mass distribution.

Taking all systematic uncertainties into account, the resulting relative uncertainty in the background increases with mass. This relative increase does not significantly affect the sensitivity at large mass values, where the expected number of events from SM processes becomes negligible. All the relevant uncertainties are also taken into account in the extraction of various signals.

All uncertainties are considered to be correlated across the different data-taking years, with the exception of the τ_h object-related uncertainties and the uncertainty in the unclustered energy, which are derived from statistically independent sources. No correlation between the different final states is considered, and the analysis results are presented independently for each of the final states.

7 Results and their interpretations

The mass distributions in the $e\mu$, $e\tau$, and $\mu\tau$ channels, shown in figure 2, do not exhibit significant deviations from the expected SM background. The last bins with data events




Figure 2. Invariant mass distributions for the $e\mu$ channel (upper), and collinear mass distributions for the $e\tau$ (lower left) and $\mu\tau$ (lower right) channels. In addition to the observed data (black points) and the SM prediction (filled histograms), the expected signal distributions for three models are shown: the RPV SUSY model with $\lambda = \lambda' = 0.01$ and τ sneutrino mass of 1.6 TeV, LFV Z' ($\mathcal{B} = 0.1$) boson with a mass of 1.6 TeV, and the QBH signal expectation for $n = 4$ and a threshold mass of 1.6 TeV. The bottom panel of each plot shows the ratio of data and SM prediction. The bin width gradually increases with mass.

show minor statistical fluctuations in the $e\tau$ and $\mu\tau$ channels, which are consistent with the SM expectations within 2 standard deviations, as shown later in the limit plots. The expected mass distributions of signal and backgrounds are taken from simulation, with the exception of backgrounds with jets misidentified as leptons, which are estimated using collision data, as discussed in previous sections. Upper limits on the products of the production cross section σ and branching fraction \mathcal{B} are determined using a Bayesian binned-likelihood method [58, 59] with a uniform positive prior probability density for the signal cross section. The nuisance parameters associated with the systematic uncertainties are modeled via log-normal distributions for uncertainties in the normalization. Uncertainties in the shape of the distributions are modeled via “template morphing” techniques [60]. A Markov Chain

Channel	RPV SUSY $\tilde{\nu}_\tau$ (TeV)		LFV Z' (TeV)	QBH m_{th} (TeV)
	$\lambda = \lambda' = 0.01$	$\lambda = \lambda' = 0.1$	$\mathcal{B} = 0.1$	$n = 4$
$e\mu$	2.2 (2.2)	4.2 (4.2)	5.0 (4.9)	5.6 (5.6)
$e\tau$	1.6 (1.6)	3.7 (3.7)	4.3 (4.3)	5.2 (5.2)
$\mu\tau$	1.6 (1.6)	3.6 (3.7)	4.1 (4.2)	5.0 (5.0)

Table 1. The observed and expected (in parentheses) 95% CL lower mass limits on the RPV SUSY, Z' , and QBH signals for the $e\mu$, $e\tau$, and $\mu\tau$ channels.

MC method [61] is used for the integration. All limits presented here are at 95% confidence level (CL).

Model-specific limits for the product of the cross section and the branching fraction were obtained for the RPV SUSY, Z' , and QBH signals and are shown in figures 3, 4, and 5, respectively. These are full Run 2 results based on 2016–2018 data sets. The observed and expected lower mass limits obtained in all three channels are summarized in table 1. A $\tilde{\nu}_\tau$ in RPV SUSY is excluded up to a mass of 4.2 TeV in the $e\mu$ channel, up to 3.7 TeV in the $e\tau$ channel, and up to 3.6 TeV for the $\mu\tau$ channel, in each case for the coupling hypothesis $\lambda = \lambda' = 0.1$. If $\lambda = \lambda' = 0.01$ is assumed, the observed limits drop to 2.2 TeV in the $e\mu$ channel, 1.6 TeV in the $e\tau$ channel, and 1.6 TeV in the $\mu\tau$ channel. These two sets of couplings are used for the results shown in figures 3, 4, and 5, as they were for the results shown in refs. [10, 18]. To provide results for a wider range of couplings, in figure 6 we show the exclusion limits as functions of λ' , for five values of λ .

A Z' ($\mathcal{B} = 0.1$) boson with LFV couplings is excluded up to a mass of 5.0 TeV in the $e\mu$ channel, up to 4.3 TeV in the $e\tau$ channel, and up to 4.1 TeV in the $\mu\tau$ channel. With increasing Z' boson mass, the phase space for the on-shell Z' production in pp collisions at 13 TeV decreases because of decreasing parton-parton luminosity, leading to an increasing fraction of off-shell production at lower masses. This effect leads to weaker Z' boson cross section limits at higher mass values, as shown in figure 4. Quantum black holes derived from an ADD model with $n = 4$ extra dimensions are excluded up to the threshold mass of 5.6 TeV in the $e\mu$ channel, 5.2 TeV in the $e\tau$ channel, and 5.0 TeV in the $\mu\tau$ channel. In all channels, the results of this search significantly improve on the previous mass limits obtained by the ATLAS and CMS Collaborations.

In the narrow-width approximation, the value of $\sigma\mathcal{B}$ scales with the RPV couplings, in all three channels. For example, in the $e\mu$ channel, the following approximation holds:

$$\sigma\mathcal{B} \approx (\lambda'_{311})^2 [(\lambda_{132})^2 + (\lambda_{231})^2] / \{3(\lambda'_{311})^2 + [(\lambda_{132})^2 + (\lambda_{231})^2]\}, \quad (7.1)$$

Using the narrow-width approximation formula for the RPV signal cross section, the cross section limit is translated into exclusion bounds in the plane of mass and coupling of the parameter space of the RPV SUSY model for fixed values of the λ couplings responsible for the decay of the $\tilde{\nu}_\tau$. Limit contours in the plane of mass and coupling for several fixed values of the coupling are shown in figure 6.

Model-independent cross section limits are also obtained. The model-specific shape analysis assumes a certain signal shape in invariant ($e\mu$ channel) or collinear mass (τ




Figure 3. Expected (black dashed line) and observed (black solid line) 95% CL upper limits on the product of the cross section and the branching fraction as a function of the τ sneutrino mass in an RPV SUSY model for the $e\mu$ (upper), $e\tau$ (lower left), and $\mu\tau$ (lower right) channels. The shaded bands represent 68% and 95% uncertainties in the expected limits. The red and blue solid lines show the predicted product of the cross section and the branching fraction as a function of the tau sneutrino mass for two different values of the couplings.

channels). However, alternative new physics processes yielding the LFV final states could cause an excess of a different shape. A model-independent cross section limit is determined using a single bin with a lower threshold on invariant (collinear) mass, and no upper threshold. No assumptions on the shape of the signal distribution are made other than that of a flat product of acceptance times efficiency, $A\varepsilon$, as a function of the mass. The excluded cross section model-independent limit is shown in figure 7. In order to determine the limit for a specific model from the model-independent limit described here, the model-dependent part of the efficiency and acceptance needs to be applied. The experimental efficiencies for the signal are already taken into account.

A factor f_m that reflects the effect of the threshold mass m^{\min} on the signal is determined by counting the events with masses $m > m^{\min}$ and dividing the result by the number of MC-generated events. The reconstruction efficiency is nearly constant over the entire mass range probed here, therefore f_m can be evaluated at the generator level. A




Figure 4. Expected (black dashed line) and observed (black solid line) 95% CL upper limits on the product of the cross section and the branching fraction for a Z' boson with LFV decays, in the $e\mu$ (upper), $e\tau$ (lower left), and $\mu\tau$ (lower right) channels. The shaded bands represent 68% and 95% uncertainties in the expected limits. The red solid lines show the predicted product of the cross section and the branching fraction as a function of the Z' mass assuming $\mathcal{B} = 0.1$.

limit on the product of the cross section and branching fraction $(\sigma\mathcal{B}A\varepsilon)_{\text{excl}}$ can be obtained by dividing the excluded cross section of the model-independent limit $(\sigma\mathcal{B}A\varepsilon)_{\text{MI}}$ given in figure 7 by the calculated fraction $f_m(m^{\text{min}})$:

$$(\sigma\mathcal{B}A\varepsilon)_{\text{excl}}(\text{total}) = \frac{(\sigma\mathcal{B}A\varepsilon)_{\text{MI}}(m^{\text{min}})}{f_m(m^{\text{min}})}. \tag{7.2}$$

Here, \mathcal{B} is the branching fraction of the new particle decaying to the relevant LFV final state. Models with a theoretical cross section $(\sigma\mathcal{B})_{\text{theo}}$ larger than $(\sigma\mathcal{B})_{\text{excl}}$ can be excluded. The fraction of events $f_m(m^{\text{min}})$ must be determined for the particular model under consideration.




Figure 5. Expected (black dashed line) and observed (black solid line) 95% CL upper limits on the product of the cross section and the branching fraction for quantum black hole production in an ADD model with $n = 4$ extra dimensions, in the $e\mu$ (upper), $e\tau$ (lower left), and $\mu\tau$ (lower right) channels. The shaded bands represent 68% and 95% uncertainties in the expected limits. The red solid lines show the predicted product of the cross section and the branching fraction as a function of the QBH threshold mass.




Figure 6. Exclusion limits at 95% CL on the RPV SUSY model in the plane of τ sneutrino mass and λ' coupling, for four values of λ couplings. The regions to the left of and above the curves are excluded. The upper plot corresponds to the $e\mu$ channel, while the lower left and right plots show the $e\tau$ and $\mu\tau$ channels, respectively. The lack of a smooth behavior of the exclusion limits for high λ' values and at high masses where there are no events is an artifact caused by the limited number of discrete mass values of the generated signal samples, in this region.




Figure 7. Model-independent upper limits at 95% CL on the product of the cross section, the branching fraction, acceptance, and efficiency are shown. Observed (expected) limits are shown in black solid (dashed) lines for the $e\mu$ (upper), $e\tau$ (lower left), and $\mu\tau$ (lower right) channels. The shaded bands represent 68% and 95% uncertainties in the expected limits.

8 Summary

A search has been conducted for heavy particles that undergo lepton flavor violating decays into $e\mu$, $e\tau$, and $\mu\tau$ final states. The search is based on proton-proton collision data at $\sqrt{s} = 13$ TeV recorded during 2016–2018 in the CMS detector at the CERN LHC, corresponding to an integrated luminosity of 138 fb^{-1} . The data are consistent with expectations from the standard model. Lower limits at 95% confidence level are set on the mass of supersymmetric τ sneutrinos at 4.2 TeV in $e\mu$, 3.7 TeV in $e\tau$, and 3.6 TeV in $\mu\tau$ channels. A Z' vector boson with lepton flavor violating couplings is excluded for masses below 5.0, 4.3, and 4.1 TeV in the $e\mu$, $e\tau$, and $\mu\tau$ channels, respectively, assuming a branching fraction of 10%. In the context of the Arkani-Hamed–Dimopoulos–Dvali model with four extra dimensions, values of the threshold mass for quantum black hole production less than 5.6, 5.2, and 5.0 TeV are excluded in the $e\mu$, $e\tau$, and $\mu\tau$ channels, respectively. In addition, model-independent limits are provided allowing the results to be interpreted in other models with the same final states and similar kinematic distributions. Limits in the $e\tau$ and $\mu\tau$ final states, as well as model-independent limits, are reported for the first time in the context of a high-mass lepton flavor violation search. These are the first results of a high-mass lepton flavor violation search using the full Run 2 data set, and they are currently the most stringent limits from any collider experiment.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 884104, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the "Excellence of Science — EOS" — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010; the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy — EXC 2121 "Quantum Universe" — 390833306, and under project number 400140256 — GRK2497; the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF "a way of making Europe", and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B05F650021 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] G.R. Farrar and P. Fayet, *Phenomenology of the production, decay, and detection of new hadronic states associated with supersymmetry*, *Phys. Lett. B* **76** (1978) 575 [[INSPIRE](#)].
- [2] F. de Campos et al., *CERN LHC signals for neutrino mass model in bilinear R-parity violating mAMSB*, *Phys. Rev. D* **77** (2008) 115025 [[arXiv:0803.4405](#)] [[INSPIRE](#)].
- [3] R. Barbier et al., *R-parity violating supersymmetry*, *Phys. Rept.* **420** (2005) 1 [[hep-ph/0406039](#)] [[INSPIRE](#)].

- [4] P. Langacker, *The physics of heavy Z' gauge bosons*, *Rev. Mod. Phys.* **81** (2009) 1199 [[arXiv:0801.1345](#)] [[INSPIRE](#)].
- [5] X. Calmet, W. Gong and S.D.H. Hsu, *Colorful quantum black holes at the LHC*, *Phys. Lett. B* **668** (2008) 20 [[arXiv:0806.4605](#)] [[INSPIRE](#)].
- [6] P. Meade and L. Randall, *Black holes and quantum gravity at the LHC*, *JHEP* **05** (2008) 003 [[arXiv:0708.3017](#)] [[INSPIRE](#)].
- [7] S. Dimopoulos and G.L. Landsberg, *Black holes at the LHC*, *Phys. Rev. Lett.* **87** (2001) 161602 [[hep-ph/0106295](#)] [[INSPIRE](#)].
- [8] S.B. Giddings and S.D. Thomas, *High-energy colliders as black hole factories: the end of short distance physics*, *Phys. Rev. D* **65** (2002) 056010 [[hep-ph/0106219](#)] [[INSPIRE](#)].
- [9] P.W. Graham, D.E. Kaplan, S. Rajendran and P. Saraswat, *Displaced supersymmetry*, *JHEP* **07** (2012) 149 [[arXiv:1204.6038](#)] [[INSPIRE](#)].
- [10] CMS collaboration, *Search for lepton-flavor violating decays of heavy resonances and quantum black holes to $e\mu$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **04** (2018) 073 [[arXiv:1802.01122](#)] [[INSPIRE](#)].
- [11] ATLAS collaboration, *Search for lepton-flavor violation in different-flavor, high-mass final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **98** (2018) 092008 [[arXiv:1807.06573](#)] [[INSPIRE](#)].
- [12] D.M. Gingrich, *Quantum black holes with charge, colour, and spin at the LHC*, *J. Phys. G* **37** (2010) 105008 [[arXiv:0912.0826](#)] [[INSPIRE](#)].
- [13] L. Randall and R. Sundrum, *A large mass hierarchy from a small extra dimension*, *Phys. Rev. Lett.* **83** (1999) 3370 [[hep-ph/9905221](#)] [[INSPIRE](#)].
- [14] N. Arkani-Hamed, S. Dimopoulos and G.R. Dvali, *The hierarchy problem and new dimensions at a millimeter*, *Phys. Lett. B* **429** (1998) 263 [[hep-ph/9803315](#)] [[INSPIRE](#)].
- [15] CDF collaboration, *Search for R -parity Violating Decays of τ Sneutrinos to $e\mu$, $\mu\tau$, and $e\tau$ Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. Lett.* **105** (2010) 191801 [[arXiv:1004.3042](#)] [[INSPIRE](#)].
- [16] D0 collaboration, *Search for sneutrino Production in $e\mu$ Final States in 5.3 fb^{-1} of $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. Lett.* **105** (2010) 191802 [[arXiv:1007.4835](#)] [[INSPIRE](#)].
- [17] ATLAS collaboration, *Search for a Heavy Neutral Particle Decaying to $e\mu$, $e\tau$, or $\mu\tau$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector*, *Phys. Rev. Lett.* **115** (2015) 031801 [[arXiv:1503.04430](#)] [[INSPIRE](#)].
- [18] CMS collaboration, *Search for lepton flavour violating decays of heavy resonances and quantum black holes to an $e\mu$ pair in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *Eur. Phys. J. C* **76** (2016) 317 [[arXiv:1604.05239](#)] [[INSPIRE](#)].
- [19] ATLAS collaboration, *Search for new phenomena in different-flavour high-mass dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 541 [[arXiv:1607.08079](#)] [[INSPIRE](#)].
- [20] CMS collaboration, *Search for heavy resonances and quantum black holes in $e\mu$, $e\tau$, and $\mu\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV*, <https://www.hepdata.net/record/ins2081834> [[DOI:10.17182/HEPDATA.127302](#)].

- [21] CMS collaboration, *Description and performance of track and primary-vertex reconstruction with the CMS tracker*, *2014 JINST* **9** P10009 [[arXiv:1405.6569](#)] [[INSPIRE](#)].
- [22] CMS collaboration, *Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *2020 JINST* **15** P10017 [[arXiv:2006.10165](#)] [[INSPIRE](#)].
- [23] CMS collaboration, *The CMS trigger system*, *2017 JINST* **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [24] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, *2018 JINST* **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [25] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, *2021 JINST* **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [26] CMS collaboration, *Performance of electron reconstruction and selection with the CMS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *2015 JINST* **10** P06005 [[arXiv:1502.02701](#)] [[INSPIRE](#)].
- [27] CMS collaboration, *The CMS experiment at the CERN LHC*, *2008 JINST* **3** S08004 [[INSPIRE](#)].
- [28] A. Belyaev, N.D. Christensen and A. Pukhov, *CalcHEP 3.4 for collider physics within and beyond the Standard Model*, *Comput. Phys. Commun.* **184** (2013) 1729 [[arXiv:1207.6082](#)] [[INSPIRE](#)].
- [29] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [30] J. Pumplin et al., *New generation of parton distributions with uncertainties from global QCD analysis*, *JHEP* **07** (2002) 012 [[hep-ph/0201195](#)] [[INSPIRE](#)].
- [31] NNPDF collaboration, *Parton distributions for the LHC Run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [32] H.K. Dreiner, S. Grab, M. Kramer and M.K. Trenkel, *Supersymmetric NLO QCD corrections to resonant slepton production and signals at the Tevatron and the CERN LHC*, *Phys. Rev. D* **75** (2007) 035003 [[hep-ph/0611195](#)] [[INSPIRE](#)].
- [33] P. Nason, *A New method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](#)] [[INSPIRE](#)].
- [34] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [35] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [36] S. Frixione, P. Nason and G. Ridolfi, *A Positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, *JHEP* **09** (2007) 126 [[arXiv:0707.3088](#)] [[INSPIRE](#)].
- [37] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [38] R. Frederix and S. Frixione, *Merging meets matching in MC@NLO*, *JHEP* **12** (2012) 061 [[arXiv:1209.6215](#)] [[INSPIRE](#)].

- [39] CMS collaboration, *Event generator tunes obtained from underlying event and multiparton scattering measurements*, *Eur. Phys. J. C* **76** (2016) 155 [[arXiv:1512.00815](#)] [[INSPIRE](#)].
- [40] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [41] J. Allison et al., *GEANT4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [[INSPIRE](#)].
- [42] J. Allison et al., *Recent developments in GEANT4*, *Nucl. Instrum. Meth. A* **835** (2016) 186 [[INSPIRE](#)].
- [43] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, *2017 JINST* **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].
- [44] D. Contardo et al., *Technical proposal for the Phase-II Upgrade of the CMS detector*, *CERN-LHCC-2015-010* (2015) [[DOI:10.17181/CERN.VU8I.D59J](#)].
- [45] CMS collaboration, *Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV*, *2018 JINST* **13** P10005 [[arXiv:1809.02816](#)] [[INSPIRE](#)].
- [46] CMS collaboration, *Identification of hadronic tau lepton decays using a deep neural network*, *2022 JINST* **17** P07023 [[arXiv:2201.08458](#)] [[INSPIRE](#)].
- [47] CMS collaboration, *Search for physics beyond the standard model in dilepton mass spectra in proton-proton collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **04** (2015) 025 [[arXiv:1412.6302](#)] [[INSPIRE](#)].
- [48] B.D. Pecjak, D.J. Scott, X. Wang and L.L. Yang, *Resummed differential cross sections for top-quark pairs at the LHC*, *Phys. Rev. Lett.* **116** (2016) 202001 [[arXiv:1601.07020](#)] [[INSPIRE](#)].
- [49] M. Czakon et al., *Top-pair production at the LHC through NNLO QCD and NLO EW*, *JHEP* **10** (2017) 186 [[arXiv:1705.04105](#)] [[INSPIRE](#)].
- [50] CMS collaboration, *Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2021) 208 [[arXiv:2103.02708](#)] [[INSPIRE](#)].
- [51] CMS collaboration, *Performance of the reconstruction and identification of high-momentum muons in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *2020 JINST* **15** P02027 [[arXiv:1912.03516](#)] [[INSPIRE](#)].
- [52] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, *J. Phys. G* **43** (2016) 023001 [[arXiv:1510.03865](#)] [[INSPIRE](#)].
- [53] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, *2017 JINST* **12** P02014 [[arXiv:1607.03663](#)] [[INSPIRE](#)].
- [54] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [[arXiv:2104.01927](#)] [[INSPIRE](#)].
- [55] CMS collaboration, *CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-17-004*, CERN, Geneva (2018).
- [56] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-18-002*, CERN, Geneva (2019).
- [57] R.J. Barlow and C. Beeston, *Fitting using finite Monte Carlo samples*, *Comput. Phys. Commun.* **77** (1993) 219 [[INSPIRE](#)].



















- [58] G. Cowan, *Statistics*, in PARTICLE DATA GROUP collaboration, *Review of Particle Physics*, *Chin. Phys. C* **40** (2016) 100001 [[INSPIRE](#)].
- [59] THEATLAS et al. collaborations, *Procedure for the LHC Higgs boson search combination in Summer 2011*, [CMS-NOTE-2011-005](#), CERN, Geneva (2011).
- [60] M. Baak, S. Gadatsch, R. Harrington and W. Verkerke, *Interpolation between multi-dimensional histograms using a new non-linear moment morphing method*, *Nucl. Instrum. Meth. A* **771** (2015) 39 [[arXiv:1410.7388](#)] [[INSPIRE](#)].
- [61] B.C. Allanach and C.G. Lester, *Sampling using a ‘bank’ of clues*, *Comput. Phys. Commun.* **179** (2008) 256 [[arXiv:0705.0486](#)] [[INSPIRE](#)].

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Tumasyan¹ 













Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , A. Escalante Del Valle , R. Frühwirth² , M. Jeitler² , N. Krammer , L. Lechner , D. Liko , I. Mikulec , P. Paulitsch, F.M. Pitters, J. Schieck² , R. Schöfbeck , D. Schwarz , S. Tempel , W. Waltenberger , C.-E. Wulz² 

















Universiteit Antwerpen, Antwerpen, Belgium

M.R. Darwish³ , E.A. De Wolf, T. Janssen , T. Kello⁴, A. Lelek , H. Rejeb Sfar, P. Van Mechelen , S. Van Putte , N. Van Remortel 










Vrije Universiteit Brussel, Brussel, Belgium

F. Blekman , E.S. Bols , J. D'Hondt , M. Delcourt , H. El Faham , S. Lowette , S. Moortgat , A. Morton , D. Müller , A.R. Sahasransu , S. Tavernier , W. Van Doninck, D. Vannerom 


















Université Libre de Bruxelles, Bruxelles, Belgium

D. Beghin, B. Bilin , B. Clerbaux , G. De Lentdecker , L. Favart , A.K. Kalsi⁵ , K. Lee , M. Mahdavihorrani , I. Makarenko , L. Moureaux , S. Paredes , L. Pétré , A. Popov , N. Postiau, E. Starling , L. Thomas , M. Vanden Bemden, C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

T. Cornelis , D. Dobur , J. Knolle , L. Lambrecht , G. Mestdach, M. Niedziela , C. Rendón, C. Roskas , A. Samalan, K. Skovpen , M. Tytgat , B. Vermassen, L. Wezenbeek 





















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , F. Bury , C. Caputo , P. David , C. De-laere , I.S. Donertas , A. Giammanco , K. Jaffel , Sa. Jain , V. Lemaitre, K. Mondal , J. Prisciandaro, A. Taliercio , M. Teklishyn , T.T. Tran , P. Vischia , S. Wertz 




Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , C. Hensel , A. Moraes , P. Rebello Teles 

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Alves Gallo Pereira , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁶, E.M. Da Costa , G.G. Da Silveira⁷ , D. De Jesus Damiao , V. Dos Santos Sousa , S. Fonseca De Souza , C. Mora Herrera , K. Mota Amarilo , L. Mundim , H. Nogima , A. Santoro , S.M. Silva Do Amaral , A. Sznajder , M. Thiel , F. Torres Da Silva De Araujo⁸ , A. Vilela Pereira 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁷ , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , D. S. Lemos , P.G. Mercadante , S.F. Novaes , Sandra S. Padula 




Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Rodozov, M. Shopova , G. Sultanov 





University of Sofia, Sofia, Bulgaria

A. Dimitrov , T. Ivanov , L. Litov , B. Pavlov , P. Petkov , A. Petrov

















Beihang University, Beijing, China

T. Cheng , T. Javaid⁹, M. Mittal , L. Yuan 


Department of Physics, Tsinghua University, Beijing, China

M. Ahmad , G. Bauer, C. Dozen , Z. Hu , J. Martins¹⁰ , Y. Wang, K. Yi^{11,12}

Institute of High Energy Physics, Beijing, China

E. Chapon , G.M. Chen⁹ , H.S. Chen⁹ , M. Chen , F. Iemmi , A. Kapoor , D. Leggat, H. Liao , Z.-A. Liu¹³ , V. Milosevic , F. Monti , R. Sharma , J. Tao , J. Thomas-Wilsker , J. Wang , H. Zhang , J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. An , Y. Ban , C. Chen, A. Levin , Q. Li , X. Lyu, Y. Mao, S.J. Qian , D. Wang , J. Xiao , H. Yang

Sun Yat-Sen University, Guangzhou, China

M. Lu , Z. You 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

X. Gao⁴ , H. Okawa , Y. Zhang 

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , M. Xiao 




Universidad de Los Andes, Bogota, Colombia
















































C. Avila , A. Cabrera , C. Florez , J. Fraga 

Universidad de Antioquia, Medellin, Colombia

J. Mejia Guisao , F. Ramirez , J.D. Ruiz Alvarez , C.A. Salazar González 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic , N. Godinovic , D. Lelas , I. Puljak 

University of Split, Faculty of Science, Split, CroatiaZ. Antunovic, M. Kovac , T. Sculac **Institute Rudjer Boskovic, Zagreb, Croatia**V. Brigljevic , D. Ferencek , D. Majumder , M. Roguljic , A. Starodumov¹⁴ , T. Susa **University of Cyprus, Nicosia, Cyprus**A. Attikis , K. Christoforou , A. Ioannou, G. Kole , M. Kolosova , S. Konstantinou , J. Mousa , C. Nicolaou, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka **Charles University, Prague, Czech Republic**M. Finger¹⁴ , M. Finger Jr.¹⁴ , A. Kveton **Escuela Politecnica Nacional, Quito, Ecuador**E. Ayala **Universidad San Francisco de Quito, Quito, Ecuador**E. Carrera Jarrin **Academy of Scientific Research and Technology of the Arab Republic of Egypt,
Egyptian Network of High Energy Physics, Cairo, Egypt**S. Elgammal¹⁵, A. Ellithi Kamel¹⁶**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum,
Egypt**M.A. Mahmoud , Y. Mohammed **National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**S. Bhowmik , R.K. Dewanjee , K. Ehataht , M. Kadastik, S. Nandan , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken **Department of Physics, University of Helsinki, Helsinki, Finland**P. Eerola , H. Kirschenmann , K. Osterberg , M. Voutilainen **Helsinki Institute of Physics, Helsinki, Finland**S. Bharthuar , E. Brücken , F. Garcia , J. Havukainen , M.S. Kim , R. Kinnunen, T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Lotti, L. Martikainen , M. Myllymäki , J. Ott , H. Siikonen , E. Tuominen , J. Tuominiemi **Lappeenranta-Lahti University of Technology, Lappeenranta, Finland**P. Luukka , H. Petrow , T. Tuuva**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**C. Amendola , M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , P. Jarry , B. Lenzi , E. Locci , J. Malcles , J. Rander, A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro¹⁷ , M. Titov , G.B. Yu 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

S. Ahuja^{ID}, F. Beaudette^{ID}, M. Bonanomi^{ID}, A. Buchot Perraguin^{ID}, P. Busson^{ID}, A. Cappati^{ID}, C. Charlot^{ID}, O. Davignon^{ID}, B. Diab^{ID}, G. Falmagne^{ID}, S. Ghosh^{ID}, R. Granier de Cassagnac^{ID}, A. Hakimi^{ID}, I. Kucher^{ID}, J. Motta^{ID}, M. Nguyen^{ID}, C. Ochando^{ID}, P. Paganini^{ID}, J. Rembser^{ID}, R. Salerno^{ID}, U. Sarkar^{ID}, J.B. Sauvan^{ID}, Y. Sirois^{ID}, A. Tarabini^{ID}, A. Zabi^{ID}, A. Zghiche^{ID}

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram¹⁸^{ID}, J. Andrea, D. Apparu^{ID}, D. Bloch^{ID}, G. Bourgatte, J.-M. Brom^{ID}, E.C. Chabert^{ID}, C. Collard^{ID}, D. Darej, J.-C. Fontaine¹⁸, U. Goerlach^{ID}, C. Grimault, A.-C. Le Bihan^{ID}, E. Nibigira^{ID}, P. Van Hove^{ID}

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

E. Asilar^{ID}, S. Beauceron^{ID}, C. Bernet^{ID}, G. Boudoul^{ID}, C. Camen, A. Carle, N. Chanon^{ID}, D. Contardo^{ID}, P. Depasse^{ID}, H. El Mamouni, J. Fay^{ID}, S. Gascon^{ID}, M. Gouzevitch^{ID}, B. Ille^{ID}, I.B. Laktineh, H. Lattaud^{ID}, A. Lesauvage^{ID}, M. Lethuillier^{ID}, L. Mirabito, S. Perries, K. Shchablo, V. Sordini^{ID}, L. Torterotot^{ID}, G. Touquet, M. Vander Donckt^{ID}, S. Viret

Georgian Technical University, Tbilisi, Georgia

I. Bagaturia¹⁹^{ID}, I. Lomidze^{ID}, Z. Tsamalaidze¹⁴^{ID}

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta^{ID}, L. Feld^{ID}, K. Klein^{ID}, M. Lipinski^{ID}, D. Meuser^{ID}, A. Pauls^{ID}, N. Röwert^{ID}, J. Schulz, M. Teroerde^{ID}

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany


























A. Dodonova^{ID}, D. Eliseev^{ID}, M. Erdmann^{ID}, P. Fackeldey^{ID}, B. Fischer^{ID}, S. Ghosh^{ID}, T. Hebbeker^{ID}, K. Hoepfner^{ID}, F. Ivone^{ID}, L. Mastrolorenzo, M. Merschmeyer^{ID}, A. Meyer^{ID}, G. Mocellin^{ID}, S. Mondal^{ID}, S. Mukherjee^{ID}, D. Noll^{ID}, A. Novak^{ID}, T. Pook^{ID}, A. Pozdnyakov^{ID}, Y. Rath, H. Reithler^{ID}, J. Roemer, A. Schmidt^{ID}, S.C. Schuler, A. Sharma^{ID}, L. Vigilante, S. Wiedenbeck^{ID}, S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany




























C. Dziwok^{ID}, G. Flügge^{ID}, W. Haj Ahmad²⁰^{ID}, O. Hlushchenko, T. Kress^{ID}, A. Nowack^{ID}, O. Pooth^{ID}, D. Roy^{ID}, A. Stahl²¹^{ID}, T. Ziemons^{ID}, A. Zotz^{ID}

Deutsches Elektronen-Synchrotron, Hamburg, Germany






























H. Aarup Petersen, M. Aldaya Martin^{ID}, P. Asmuss, S. Baxter^{ID}, M. Bayatmakou^{ID}, O. Behnke, A. Bermúdez Martínez^{ID}, S. Bhattacharya^{ID}, A.A. Bin Anuar^{ID}, K. Borras²²^{ID}, D. Brunner^{ID}, A. Campbell^{ID}, A. Cardini^{ID}, C. Cheng, F. Colombina, S. Consuegra Rodríguez^{ID}, G. Correia Silva^{ID}, V. Danilov, M. De Silva^{ID}, L. Didukh^{ID}, G. Eckerlin, D. Eckstein, L.I. Estevez Banos^{ID}, O. Filatov^{ID}, E. Gallo²³^{ID}, A. Geiser^{ID}, A. Giraldi^{ID}, A. Grohsjean^{ID}, M. Guthoff^{ID}, A. Jafari²⁴^{ID}, N.Z. Jomhari^{ID}, A. Kasem²²^{ID}, M. Kasemann^{ID}, H. Kaveh^{ID}, C. Kleinwort^{ID}, R. Kogler^{ID}, D. Krücker^{ID}, W. Lange, J. Lidrych^{ID}, K. Lipka^{ID}, W. Lohmann²⁵^{ID}, R. Mankel^{ID}, I.-A. Melzer-Pellmann^{ID},

M. Mendizabal Morentin , J. Metwally, A.B. Meyer , M. Meyer , J. Mnich , A. Mussgiller , Y. Otariid, D. Pérez Adán , D. Pitzl, A. Raspereza, B. Ribeiro Lopes , J. Rübenach, A. Saggio , A. Saibel , M. Savitskyi , M. Scham²⁶ , V. Scheurer, S. Schnake , P. Schütze , C. Schwanenberger²³ , M. Shchedrolosiev , R.E. Sosa Ricardo , D. Stafford, N. Tonon , M. Van De Klundert , R. Walsh , D. Walter , Q. Wang , Y. Wen , K. Wichmann, L. Wiens , C. Wissing , S. Wuchterl 

University of Hamburg, Hamburg, Germany

R. Aggleton, S. Albrecht , S. Bein , L. Benato , P. Connor , K. De Leo , M. Eich, F. Feindt, A. Fröhlich, C. Garbers , E. Garutti , P. Gunnellini, M. Hajheidari, J. Haller , A. Hinzmann , G. Kasieczka , R. Klanner , T. Kramer , V. Kutzner , J. Lange , T. Lange , A. Lobanov , A. Malara , A. Nigamova , K.J. Pena Rodriguez , M. Rieger , O. Rieger, P. Schleper , M. Schröder , J. Schwandt , J. Sonnneveld , H. Stadie , G. Steinbrück , A. Tews, I. Zoi 










Karlsruher Institut fuer Technologie, Karlsruhe, Germany

J. Bechtel , S. Brommer , M. Burkart, E. Butz , R. Caspart , T. Chwalek , W. De Boer[†], A. Dierlamm , A. Droll, K. El Morabit , N. Faltermann , M. Giffels , J.O. Gosewisch, A. Gottmann , F. Hartmann²¹ , C. Heidecker, U. Husemann , P. Keicher, R. Koppenhöfer , S. Maier , M. Metzler, S. Mitra , Th. Müller , M. Neukum, A. Nürnberg , G. Quast , K. Rabbertz , J. Rauser, D. Savoiu , M. Schnepf, D. Seith, I. Shvetsov, H.J. Simonis , R. Ulrich , J. Van Der Linden , R.F. Von Cube , M. Wassmer , M. Weber , S. Wieland , R. Wolf , S. Wozniewski , S. Wunsch

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis, A. Stakia 







National and Kapodistrian University of Athens, Athens, Greece

M. Diamantopoulou, D. Karasavvas, P. Kontaxakis , C.K. Koraka , A. Manousakis-Katsikakis , A. Panagiotou, I. Papavergou , N. Saoulidou , K. Theofilatos , E. Tziiferi , K. Vellidis , E. Vourliotis 












National Technical University of Athens, Athens, Greece

G. Bakas , K. Kousouris , I. Papakrivopoulos , G. Tsipolitis, A. Zacharopoulou






University of Ioánnina, Ioánnina, Greece





K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, P. Gianneios , P. Katsoulis, P. Kokkas , N. Manthos , I. Papadopoulos , J. Strologas 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary




























































M. Csanád , K. Farkas , M.M.A. Gadallah²⁷ , S. Lökös²⁸ , P. Major , K. Mandal , A. Mehta , G. Pásztor , A.J. Rádl , O. Surányi , G.I. Veres 

Wigner Research Centre for Physics, Budapest, Hungary

M. Bartók²⁹ , G. Bencze, C. Hajdu , D. Horvath^{30,31} , F. Sikler , V. Veszpremi 

Institute of Nuclear Research ATOMKI, Debrecen, HungaryS. Czellar, D. Fasanella , F. Fienga , J. Karancsi²⁹ , J. Molnar, Z. Szillasi, D. Teyssier **Institute of Physics, University of Debrecen, Debrecen, Hungary**P. Raics, Z.L. Trocsanyi³² , B. Ujvari **Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary**T. Csorgo³³ , F. Nemes³³ , T. Novak **Indian Institute of Science (IISc), Bangalore, India**

S. Choudhury

Panjab University, Chandigarh, IndiaS. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra⁵ , R. Gupta, A. Kaur , H. Kaur , M. Kaur , P. Kumari , M. Meena , K. Sandeep , J.B. Singh , A. K. Viridi **University of Delhi, Delhi, India**A. Ahmed , A. Bhardwaj , B.C. Choudhary , M. Gola, S. Keshri , A. Kumar , M. Naimuddin , P. Priyanka , K. Ranjan , A. Shah **Saha Institute of Nuclear Physics, HBNI, Kolkata, India**M. Bharti³⁴, R. Bhattacharya , S. Bhattacharya , D. Bhowmik, S. Dutta , S. Dutta, B. Gomber³⁵ , M. Maity³⁶, P. Palit , P.K. Rout , G. Saha , B. Sahu , S. Sarkar, M. Sharan, S.Thakur³⁴ **Indian Institute of Technology Madras, Madras, India**P.K. Behera , S.C. Behera , P. Kalbhor , J.R. Komaragiri³⁷ , D. Kumar³⁷ , A. Muhammad , L. Panwar³⁷ , R. Pradhan , P.R. Pujahari , A. Sharma , A.K. Sikdar , P.C. Tiwari³⁷ **Bhabha Atomic Research Centre, Mumbai, India**D. Dutta , V. Jha, V. Kumar , D.K. Mishra, K. Naskar³⁸ , P.K. Netrakanti, L.M. Pant, P. Shukla **Tata Institute of Fundamental Research-A, Mumbai, India**T. Aziz, S. Dugad, M. Kumar , G.B. Mohanty **Tata Institute of Fundamental Research-B, Mumbai, India**S. Banerjee , R. Chudasama , M. Guchait , S. Karmakar , S. Kumar , G. Majumder , K. Mazumdar , S. Mukherjee **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**S. Bahinipati³⁹ , C. Kar , P. Mal , T. Mishra , V.K. Muraleedharan Nair Bindhu⁴⁰ , A. Nayak⁴⁰ , P. Saha , N. Sur , S.K. Swain, D. Vats⁴⁰ **Indian Institute of Science Education and Research (IISER), Pune, India**A. Alpana , S. Dube , B. Kansal , A. Laha , S. Pandey , A. Rastogi , S. Sharma 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi⁴¹ , E. Khazaie , M. Zeinali⁴² 


































Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Chenarani⁴³ , S.M. Etesami , M. Khakzad , M. Mohammadi Najafabadi 





























University College Dublin, Dublin, Ireland

M. Grunewald 







INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia^{a,b} , R. Aly^{a,c,44} , C. Aruta^{a,b} , A. Colaleo^a , D. Creanza^{a,c} ,
N. De Filippis^{a,c} , M. De Palma^{a,b} , A. Di Florio^{a,b} , A. Di Pilato^{a,b} ,
W. Elmetenawee^{a,b} , L. Fiore^a , A. Gelmi^{a,b} , M. Gul^a , G. Iaselli^{a,c} , M. Ince^{a,b} ,
S. Lezki^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^{a,b} , V. Mastrapasqua^{a,b} ,
S. My^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} ,
D. Ramos^a , A. Ranieri^a , G. Selvaggi^{a,b} , L. Silvestris^a , F.M. Simone^{a,b} ,
Ü. Sözbilir^a , R. Venditti^a , P. Verwilligen^a 
















INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , L. Borgonovi^a , R. Campanini^{a,b} ,
P. Capiluppi^{a,b} , A. Castro^{a,b} , F.R. Cavallo^a , C. Ciocca^a , M. Cuffiani^{a,b} ,
G.M. Dallavalle^a , T. Diotallevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , P. Giacomelli^a ,
L. Giommi^{a,b} , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,45} , L. Lunerti^{a,b} ,
S. Marcellini^a , G. Masetti^a , F.L. Navarria^{a,b} , A. Perrotta^a , F. Primavera^{a,b} ,
A.M. Rossi^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b} 

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Albergo^{a,b,46} , S. Costa^{a,b,46} , A. Di Mattia^a , R. Potenza^{a,b} , A. Tricomi^{a,b,46} ,
C. Tuve^{a,b} 

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

G. Barbagli^a , A. Cassese^a , R. Ceccarelli^{a,b} , V. Ciulli^{a,b} , C. Civinini^a ,
R. D'Alessandro^{a,b} , E. Focardi^{a,b} , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^{a,b} ,
M. Meschini^a , S. Paoletti^a , R. Seidita^{a,b} , G. Sguazzoni^a , L. Viliani^a 


























INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , D. Piccolo 











INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Bozzo^{a,b} , F. Ferro^a , R. Mulargia^{a,b} , E. Robutti^a , S. Tosi^{a,b} 


























INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , G. Boldrini^a , F. Brivio^{a,b} , F. Cettorelli^{a,b} , F. De Guio^{a,b} ,
M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^{a,b} ,
M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a ,
L. Moroni^a , M. Paganoni^{a,b} , D. Pedrini^a , B.S. Pinolini^a , S. Ragazzi^{a,b} ,
N. Redaelli^a , T. Tabarelli de Fatis^{a,b} , D. Valsecchi^{a,b,21} , D. Zuolo^{a,b} 











INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Università G. Marconi^d, Roma, Italy

S. Buontempo^a , F. Carnevali^{a,b}, N. Cavallo^{a,c} , A. De Iorio^{a,b} , F. Fabozzi^{a,c} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,47} , S. Meola^{a,d,21} , P. Paolucci^{a,21} , B. Rossi^a , C. Sciacca^{a,b} 















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

P. Azzi^a , N. Bacchetta^a , D. Bisello^{a,b} , P. Bortignon^a , A. Bragagnolo^{a,b} , R. Carlin^{a,b} , P. Checchia^a , T. Dorigo^a , U. Dosselli^a , F. Gasparini^{a,b} , U. Gasparini^{a,b} , G. Grosso^a, S.Y. Hoh^{a,b} , L. Layer^{a,48}, E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , J. Pazzini^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , G. Strong^a , M. Tosi^{a,b} , H. Yarar^{a,b}, M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b} , G. Zumerle^{a,b} 

































INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

C. Aimè^{a,b} , A. Braghieri^a , S. Calzaferri^{a,b} , D. Fiorina^{a,b} , P. Montagna^{a,b} , S.P. Ratti^{a,b}, V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^a , P. Vitulo^{a,b} 







INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

P. Asenov^{a,49} , G.M. Bilei^a , D. Ciangottini^{a,b} , L. Fanò^{a,b} , M. Magherini^{a,b} , G. Mantovani^{a,b}, V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,49} , A. Piccinelli^{a,b} , M. Presilla^{a,b} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 
















INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

P. Azzurri^a , G. Bagliesi^a , V. Bertacchi^{a,c} , L. Bianchini^a , T. Boccali^a , E. Bossini^{a,b} , R. Castaldi^a , M.A. Ciocci^{a,b} , V. D'Amante^{a,d} , R. Dell'Orso^a , M.R. Di Domenico^{a,d} , S. Donato^a , A. Giassi^a , F. Ligabue^{a,c} , E. Manca^{a,c} , G. Mandorli^{a,c} , D. Matos Figueiredo^a , A. Messineo^{a,b} , F. Palla^a , S. Parolia^{a,b} , G. Ramirez-Sanchez^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,c} , A. Scribano^a , N. Shafiei^{a,b} , P. Spagnolo^a , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , A. Venturi^a , P.G. Verdini^a 

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Barria^a , M. Campana^{a,b} , F. Cavallari^a , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a , E. Longo^{a,b} , P. Meridiani^a , G. Organtini^{a,b} , F. Pandolfi^a , R. Paramatti^{a,b} , C. Quaranta^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , R. Tramontano^{a,b} 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^a , R. Bellan^{a,b} , A. Bellora^{a,b} , J. Berenguer Antequera^{a,b} , C. Biino^a , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , B. Kiani^{a,b} , F. Legger^a 

C. Mariotti^a, S. Maselli^a, E. Migliore^{a,b}, E. Monteil^{a,b}, M. Monteno^a, M.M. Obertino^{a,b}, G. Ortona^a, L. Pacher^{a,b}, N. Pastrone^a, M. Pelliccioni^a, M. Ruspa^{a,c}, K. Shchelina^a, F. Siviero^{a,b}, V. Sola^a, A. Solano^{a,b}, D. Soldi^{a,b}, A. Staiano^a, M. Tornago^{a,b}, D. Trocino^a, A. Vagnerini^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

S. Belforte^a, V. Candelise^{a,b}, M. Casarsa^a, F. Cossutti^a, A. Da Rold^{a,b}, G. Della Ricca^{a,b}, G. Sorrentino^{a,b}, F. Vazzoler^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra, C. Huh, B. Kim, D.H. Kim, G.N. Kim, J. Kim, J. Lee, S.W. Lee, C.S. Moon, Y.D. Oh, S.I. Pak, S. Sekmen, Y.C. Yang

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

H. Kim, D.H. Moon

Hanyang University, Seoul, Korea

B. Francois, T.J. Kim, J. Park

Korea University, Seoul, Korea

S. Cho, S. Choi, B. Hong, K. Lee, K.S. Lee, J. Lim, J. Park, S.K. Park, J. Yoo

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh, A. Gurtu

Sejong University, Seoul, Korea

H. S. Kim, Y. Kim

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi, S. Jeon, J. Kim, J.S. Kim, S. Ko, H. Kwon, H. Lee, S. Lee, B.H. Oh, M. Oh, S.B. Oh, H. Seo, U.K. Yang, I. Yoon

University of Seoul, Seoul, Korea

W. Jang, D.Y. Kang, Y. Kang, S. Kim, B. Ko, J.S.H. Lee, Y. Lee, J.A. Merlin, I.C. Park, Y. Roh, M.S. Ryu, D. Song, I.J. Watson, S. Yang

Yonsei University, Department of Physics, Seoul, Korea

S. Ha, H.D. Yoo

Sungkyunkwan University, Suwon, Korea

M. Choi, H. Lee, Y. Lee, I. Yu

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy, Y. Maghrbi

Riga Technical University, Riga, Latvia

K. Dreimanis, V. Veckalns

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Carvalho Antunes De Oliveira , A. Juodagalvis , A. Rinkevicius ,
G. Tamulaitis 








National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

N. Bin Norjoharuddeen , W.A.T. Wan Abdullah, M.N. Yusli, Z. Zolkapli

Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , M. León Coello , J.A. Murillo Quijada ,
A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , E. De La Cruz-Burelo , I. Heredia-De La Cruz ⁵⁰ ,
R. Lopez-Fernandez , C.A. Mondragon Herrera, D.A. Perez Navarro ,
A. Sánchez Hernández 


Universidad Iberoamericana, Mexico City, Mexico

S. Carrillo Moreno, C. Oropeza Barrera , F. Vazquez Valencia 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Pedraza , H.A. Salazar Ibarquen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

J. Mijuskovic ⁵¹, N. Raicevic 







University of Auckland, Auckland, New Zealand

D. Krofcheck 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan ,
M.A. Shah, M. Shoaib , M. Waqas 

AGH University of Science and Technology Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

V. Avati, L. Grzanka , M. Malawski 














National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , B. Boimska , M. Górski , M. Kazana , M. Szeper ,
P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski 































Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , P. Bargassa , D. Bastos , A. Boletti , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , T. Niknejad , M. Pisano , J. Seixas , O. Toldaiev , J. Varela 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

P. Adzic⁵² , M. Dordevic , P. Milenovic , J. Milosevic 















Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Aguilar-Benitez , J. Alcaraz Maestre , A. Álvarez Fernández , I. Bachiller , M. Barrio Luna , Cristina F. Bedoya , C.A. Carrillo Montoya , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. León Holgado , D. Moran , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , L. Romero , S. Sánchez Navas , L. Urda Gómez , C. Willmott 




















Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz , R. Reyes-Almanza 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , C. Erice , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , E. Palencia Cortezon , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , N. Trevisani , C. Vico Villalba 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , C. Fernandez Madrazo , P.J. Fernández Manteca , A. García Alonso , G. Gomez , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , J. Piedra Gomez , C. Prieels , A. Ruiz-Jimeno , L. Scodellaro , I. Vila , J.M. Vizan Garcia 


















University of Colombo, Colombo, Sri Lanka









M.K. Jayananda , B. Kailasapathy⁵³ , D.U.J. Sonnadara , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna , K. Liyanage , N. Perera , N. Wickramage 

CERN, European Organization for Nuclear Research, Geneva, Switzerland


































T.K. Aarrestad , D. Abbaneo , J. Alimena , E. Auffray , G. Auzinger , J. Baechler , P. Baillon[†] , D. Barney , J. Bendavid , M. Bianco , A. Bocci , C. Caillol , T. Camporesi , M. Capeans Garrido , G. Cerminara , N. Chernyavskaya , S.S. Chhibra 

M. Cipriani , L. Cristella , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , M. Dünser , N. Dupont, A. Elliott-Peisert, N. Emriskova, F. Fallavollita⁵⁴, A. Florent , L. Forthomme , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill, F. Glege , L. Gouskos , M. Haranko , J. Hege-man , V. Innocente , T. James , P. Janot , J. Kaspar , J. Kieseler , M. Komm , N. Kratochwil , C. Lange , S. Laurila , P. Lecoq , A. Lintuluoto , K. Long , C. Lourenço , B. Maier , L. Malgeri , S. Mallios, M. Mannelli , A.C. Marini , F. Meijers , S. Mersi , E. Meschi , F. Moortgat , M. Mulders , S. Orfanelli, L. Orsini, F. Pantaleo , E. Perez, M. Peruzzi , A. Petrilli , G. Petrucciani , A. Pfeiffer , M. Pierini , D. Piparo , M. Pitt , H. Qu , T. Quast, D. Rabady , A. Racz, G. Reales Gutiérrez, M. Rovere , H. Sakulin , J. Salfeld-Nebgen , S. Scarfi, C. Schäfer, M. Selvaggi , A. Sharma , P. Silva , W. Snoeys , P. Sphicas⁵⁵ , S. Summers , K. Tatar , V.R. Tavolaro , D. Treille , P. Tropea , A. Tsirou, G.P. Van Onsem , J. Wanczyk⁵⁶ , K.A. Wozniak , W.D. Zeuner
























Paul Scherrer Institut, Villigen, Switzerland

L. Caminada⁵⁷ , A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , M. Missiroli⁵⁷ , L. Noehte⁵⁷ , T. Rohe 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

K. Androsov⁵⁶ , M. Backhaus , P. Berger, A. Calandri , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , C. Dorfer , F. Eble , K. Gedia , F. Glessgen , T.A. Gómez Espinosa , C. Grab , D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , L. Marchese , C. Martin Perez , M.T. Meinhard , F. Nessi-Tedaldi , J. Niedziela , F. Pauss , V. Perovic , S. Pigazzini , M.G. Ratti , M. Reichmann , C. Reissel , T. Reitenspiess , B. Ristic , D. Ruini, D.A. Sanz Becerra , V. Stampf, J. Steggemann⁵⁶ , R. Wallny , D.H. Zhu 











Universität Zürich, Zurich, Switzerland

C. Amsler⁵⁸ , P. Bärttschi , C. Botta , D. Brzhechko, M.F. Canelli , K. Cormier , A. De Wit , R. Del Burgo, J.K. Heikkilä , M. Huwiler , W. Jin , A. Jofre-Hei , B. Kilminster , S. Leontsinis , S.P. Liehti , A. Macchiolo , P. Meiring , V.M. Mikuni , U. Molinatti , I. Neutelings , A. Reimers , P. Robmann, S. Sanchez Cruz , K. Schweiger , M. Senger , Y. Takahashi 




National Central University, Chung-Li, Taiwan

C. Adloff⁵⁹, C.M. Kuo, W. Lin, A. Roy , T. Sarkar³⁶ , S.S. Yu 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, Y. Chao , K.F. Chen , P.H. Chen , P.s. Chen, H. Cheng , W.-S. Hou , Y.y. Li , R.-S. Lu , E. Paganis , A. Psallidas, A. Steen , H.y. Wu, E. Yazgan , P.r. Yu

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

B. Asavapibhop , C. Asawatangtrakuldee , N. Srimanobhas 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

F. Boran⁶⁰, S. Damarseckin⁶⁰, Z.S. Demiroglu, F. Dolek, I. Dumanoglu⁶¹, E. Eskut, Y. Guler⁶², E. Gurpinar Guler⁶², C. Isik, O. Kara, A. Kayis Topaksu, U. Kiminsu, G. Onengut, K. Ozdemir⁶³, A. Polatoz, A.E. Simsek, B. Tali⁶⁴, U.G. Tok, S. Turkcapar, I.S. Zorbakir

Middle East Technical University, Physics Department, Ankara, Turkey

G. Karapinar, K. Ocalan⁶⁵, M. Yalvac⁶⁶

Bogazici University, Istanbul, Turkey

B. Akgun, I.O. Atakisi, E. Gülmez, M. Kaya⁶⁷, O. Kaya⁶⁸, Ö. Özçelik, S. Tekten⁶⁹, E.A. Yetkin⁷⁰

Istanbul Technical University, Istanbul, Turkey

A. Cakir, K. Cankocak⁶¹, Y. Komurcu, S. Sen⁷¹

Istanbul University, Istanbul, Turkey

S. Cerci⁶⁴, I. Hos⁷², B. Isildak⁷³, B. Kaynak, S. Ozkorucuklu, H. Sert, D. Sunar Cerci⁶⁴, C. Zorbilmez

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

B. Grynyov

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk

University of Bristol, Bristol, United Kingdom







D. Anthony, E. Bhal, S. Bologna, J.J. Brooke, A. Bundock, E. Clement, D. Cussans, H. Flacher, J. Goldstein, G.P. Heath, H.F. Heath, L. Kreczko, B. Krikler, S. Paramesvaran, S. Seif El Nasr-Storey, V.J. Smith, N. Stylianou⁷⁴, K. Walkingshaw Pass, R. White

Rutherford Appleton Laboratory, Didcot, United Kingdom

K.W. Bell, A. Belyaev⁷⁵, C. Brew, R.M. Brown, D.J.A. Cockerill, C. Cooke, K.V. Ellis, K. Harder, S. Harper, M.-L. Holmberg, J. Linacre, K. Manolopoulos, D.M. Newbold, E. Olaiya, D. Petyt, T. Reis, T. Schuh, C.H. Shepherd-Themistocleous, I.R. Tomalin, T. Williams

Imperial College, London, United Kingdom

R. Bainbridge, P. Bloch, S. Bonomally, J. Borg, S. Breeze, O. Buchmuller, V. Cepaitis, G.S. Chahal⁷⁶, D. Colling, P. Dauncey, G. Davies, M. Della Negra, S. Fayer, G. Fedi, G. Hall, M.H. Hassanshahi, G. Iles, J. Langford, L. Lyons, A.-M. Magnan, S. Malik, A. Martelli, D.G. Monk, J. Nash⁷⁷, M. Pesaresi, B.C. Radburn-Smith, D.M. Raymond, A. Richards, A. Rose, E. Scott, C. Seez

A. Shtipliyski, A. Tapper , K. Uchida , T. Virdee²¹ , M. Vojinovic , N. Wardle , S.N. Webb , D. Winterbottom

Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid , L. Teodorescu, S. Zahid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , B. Caraway , J. Dittmann , K. Hatakeyama , A.R. Kanuganti , B. McMaster , N. Pastika , M. Saunders , S. Sawant , C. Sultantawibul , J. Wilson 
















Catholic University of America, Washington, DC, USA

R. Bartek , A. Dominguez , R. Uniyal , A.M. Vargas Hernandez 

















The University of Alabama, Tuscaloosa, Alabama, USA

A. Buccilli , S.I. Cooper , D. Di Croce , S.V. Gleyzer , C. Henderson , C.U. Perez , P. Rumerio⁷⁸ , C. West 


















Boston University, Boston, Massachusetts, USA

A. Akpınar , A. Albert , D. Arcaro , C. Cosby , Z. Demiragli , E. Fontanesi , D. Gastler , S. May , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , S. Yuan , D. Zou














Brown University, Providence, Rhode Island, USA

G. Benelli , B. Burkle , X. Coubez²², D. Cutts , M. Hadley , U. Heintz , J.M. Hogan⁷⁹ , T. KWON , G. Landsberg , K.T. Lau , D. Li, M. Lukasik, J. Luo , M. Narain, N. Pervan , S. Sagir⁸⁰ , F. Simpson , E. Usai , W.Y. Wong, X. Yan , D. Yu , W. Zhang

University of California, Davis, Davis, California, USA

J. Bonilla , C. Brainerd , R. Breedon , M. Calderon De La Barca Sanchez , M. Chertok , J. Conway , P.T. Cox , R. Erbacher , G. Haza , F. Jensen , O. Kukral , R. Lander, M. Mulhearn , D. Pellett , B. Regnery , D. Taylor , Y. Yao , F. Zhang 






















University of California, Los Angeles, California, USA

M. Bachtis , R. Cousins , A. Datta , D. Hamilton , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , W.A. Nash , S. Regnard , D. Saltzberg , B. Stone , V. Valuev 

















University of California, Riverside, Riverside, California, USA

K. Burt, Y. Chen, R. Clare , J.W. Gary , M. Gordon, G. Hanson , G. Karapostoli , O.R. Long , N. Manganeli , M. Olmedo Negrete, W. Si , S. Wimpenny, Y. Zhang














University of California, San Diego, La Jolla, California, USA

J.G. Branson, P. Chang , S. Cittolin, S. Cooperstein , N. Deelen , D. Diaz , J. Duarte , R. Gerosa , L. Giannini , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , M. Pieri , B.V. Sathia Narayanan , V. Sharma , M. Tadel , F. Würthwein , Y. Xiang , A. Yagil 











University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

N. Amin, C. Campagnari , M. Citron , A. Dorsett , V. Dutta , J. Incandela , M. Kilpatrick , J. Kim , B. Marsh, H. Mei , M. Oshiro , M. Quinnan , J. Richman , U. Sarica , F. Setti , J. Sheplock , P. Siddireddy, D. Stuart , S. Wang 

California Institute of Technology, Pasadena, California, USA

A. Bornheim , O. Cerri, I. Dutta , J.M. Lawhorn , N. Lu , J. Mao , H.B. Newman , T. Q. Nguyen , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , Z. Zhang , R.Y. Zhu 

















Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , M.B. Andrews , P. Bryant , T. Ferguson , A. Harilal , C. Liu , T. Mudholkar , M. Paulini , A. Sanchez , W. Terrill 

































































University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hassani , G. Karathanasis , E. MacDonald, R. Patel, A. Perloff , C. Savard , K. Stenson , K.A. Ulmer , S.R. Wagner 
























Cornell University, Ithaca, New York, USA

J. Alexander , S. Bright-Thonney , X. Chen , Y. Cheng , D.J. Cranshaw , S. Hogan , J. Monroy , J.R. Patterson , D. Quach , J. Reichert , M. Reid , A. Ryd , W. Sun , J. Thom , P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , G. Apollinari , A. Apresyan , A. Apyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , K.F. Di Petrillo , V.D. Elvira , Y. Feng , J. Freeman , Z. Gecse , L. Gray , D. Green, S. Grünendahl , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma , B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid , K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , V. O'Dell, V. Papadimitriou , K. Pedro , C. Pena⁸¹ , O. Prokofyev, F. Ravera , A. Reinsvold Hall⁸² , L. Ristori , E. Sexton-Kennedy , N. Smith , A. Soha , L. Spiegel , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , H.A. Weber 

University of Florida, Gainesville, Florida, USA

P. Avery , D. Bourilkov , L. Cadamuro , V. Cherepanov , F. Errico , R.D. Field, D. Guerrero , B.M. Joshi , M. Kim, E. Koenig , J. Konigsberg , A. Korytov , K.H. Lo, K. Matchev , N. Menendez , G. Mitselmakher , A. Muthirakalayil Madhu , N. Rawal , D. Rosenzweig , S. Rosenzweig , J. Rotter , K. Shi , J. Wang , Z. Wu , E. Yigitbasi , X. Zuo 

Florida State University, Tallahassee, Florida, USA

T. Adams^{[ID](#)}, A. Askew^{[ID](#)}, R. Habibullah^{[ID](#)}, V. Hagopian^{[ID](#)}, K.F. Johnson, R. Khurana, T. Kolberg^{[ID](#)}, G. Martinez, H. Prosper^{[ID](#)}, C. Schiber, O. Viazlo^{[ID](#)}, R. Yohay^{[ID](#)}, J. Zhang

Florida Institute of Technology, Melbourne, Florida, USA

M.M. Baarmand^{[ID](#)}, S. Butalla^{[ID](#)}, T. Elkafrawy^{[ID](#)}⁸³, M. Hohlmann^{[ID](#)}, R. Kumar Verma^{[ID](#)}, D. Noonan^{[ID](#)}, M. Rahmani, F. Yumiceva^{[ID](#)}

University of Illinois at Chicago (UIC), Chicago, Illinois, USA

M.R. Adams^{[ID](#)}, H. Becerril Gonzalez^{[ID](#)}, R. Cavanaugh^{[ID](#)}, S. Dittmer^{[ID](#)}, O. Evdokimov^{[ID](#)}, C.E. Gerber^{[ID](#)}, D.A. Hangal^{[ID](#)}, D.J. Hofman^{[ID](#)}, A.H. Merrit^{[ID](#)}, C. Mills^{[ID](#)}, G. Oh^{[ID](#)}, T. Roy^{[ID](#)}, S. Rudrabhatla^{[ID](#)}, M.B. Tonjes^{[ID](#)}, N. Varelas^{[ID](#)}, J. Viinikainen^{[ID](#)}, X. Wang^{[ID](#)}, Z. Ye^{[ID](#)}

The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini^{[ID](#)}, K. Dilsiz^{[ID](#)}⁸⁴, L. Emediato^{[ID](#)}, R.P. Gandrajula^{[ID](#)}, O.K. Köseyan^{[ID](#)}, J.-P. Merlo, A. Mestvirishvili^{[ID](#)}⁸⁵, J. Nachtman^{[ID](#)}, H. Ogul^{[ID](#)}⁸⁶, Y. Onel^{[ID](#)}, A. Penzo^{[ID](#)}, C. Snyder, E. Tiras^{[ID](#)}⁸⁷

Johns Hopkins University, Baltimore, Maryland, USA

O. Amram^{[ID](#)}, B. Blumenfeld^{[ID](#)}, L. Corcodilos^{[ID](#)}, J. Davis^{[ID](#)}, M. Eminizer^{[ID](#)}, A.V. Gritsan^{[ID](#)}, S. Kyriacou^{[ID](#)}, P. Maksimovic^{[ID](#)}, J. Roskes^{[ID](#)}, M. Swartz^{[ID](#)}, T.Á. Vámi^{[ID](#)}

The University of Kansas, Lawrence, Kansas, USA

A. Abreu^{[ID](#)}, J. Anguiano^{[ID](#)}, C. Baldenegro Barrera^{[ID](#)}, P. Baringer^{[ID](#)}, A. Bean^{[ID](#)}, A. Bylinkin^{[ID](#)}, Z. Flowers^{[ID](#)}, T. Isidori^{[ID](#)}, S. Khalil^{[ID](#)}, J. King^{[ID](#)}, G. Krintiras^{[ID](#)}, A. Kropivnitskaya^{[ID](#)}, M. Lazarovits^{[ID](#)}, C. Le Mahieu^{[ID](#)}, C. Lindsey, J. Marquez^{[ID](#)}, N. Minafra^{[ID](#)}, M. Murray^{[ID](#)}, M. Nickel^{[ID](#)}, C. Rogan^{[ID](#)}, C. Royon^{[ID](#)}, R. Salvatico^{[ID](#)}, S. Sanders^{[ID](#)}, E. Schmitz^{[ID](#)}, C. Smith^{[ID](#)}, J.D. Tapia Takaki^{[ID](#)}, Q. Wang^{[ID](#)}, Z. Warner, J. Williams^{[ID](#)}, G. Wilson^{[ID](#)}

Kansas State University, Manhattan, Kansas, USA

S. Duric, A. Ivanov^{[ID](#)}, K. Kaadze^{[ID](#)}, D. Kim, Y. Maravin^{[ID](#)}, T. Mitchell, A. Modak, K. Nam

Lawrence Livermore National Laboratory, Livermore, California, USA

F. Rebassoo^{[ID](#)}, D. Wright^{[ID](#)}













University of Maryland, College Park, Maryland, USA

E. Adams^{[ID](#)}, A. Baden^{[ID](#)}, O. Baron, A. Belloni^{[ID](#)}, S.C. Eno^{[ID](#)}, N.J. Hadley^{[ID](#)}, S. Jabeen^{[ID](#)}, R.G. Kellogg^{[ID](#)}, T. Koeth^{[ID](#)}, Y. Lai^{[ID](#)}, S. Lascio^{[ID](#)}, A.C. Mignerey^{[ID](#)}, S. Nabili^{[ID](#)}, C. Palmer^{[ID](#)}, M. Seidel^{[ID](#)}, A. Skuja^{[ID](#)}, L. Wang^{[ID](#)}, K. Wong^{[ID](#)}
















Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

D. Abercrombie, G. Andreassi, R. Bi, W. Busza^{[ID](#)}, I.A. Cali^{[ID](#)}, Y. Chen^{[ID](#)}, M. D'Alfonso^{[ID](#)}, J. Eysermans^{[ID](#)}, C. Freer^{[ID](#)}, G. Gomez-Ceballos^{[ID](#)}, M. Goncharov, P. Harris, M. Hu^{[ID](#)}, M. Klute^{[ID](#)}, D. Kovalskyi^{[ID](#)}, J. Krupa^{[ID](#)}, Y.-J. Lee^{[ID](#)}, C. Mironov^{[ID](#)}, C. Paus^{[ID](#)}, D. Rankin^{[ID](#)}, C. Roland^{[ID](#)}, G. Roland^{[ID](#)}, Z. Shi^{[ID](#)}, G.S.F. Stephans^{[ID](#)}, J. Wang, Z. Wang^{[ID](#)}, B. Wyslouch^{[ID](#)}

University of Minnesota, Minneapolis, Minnesota, USA

R.M. Chatterjee, A. Evans , J. Hiltbrand , Sh. Jain , M. Krohn , Y. Kubota , J. Mans , M. Revering , R. Rusack , R. Saradhy , N. Schroeder , N. Strobbe , M.A. Wadud 




















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , M. Bryson, S. Chauhan , D.R. Claes , C. Fangmeier , L. Finco , F. Golf , C. Joo , I. Kravchenko , M. Musich , I. Reed , J.E. Siado , G.R. Snow[†], W. Tabb , A. Wightman , F. Yan , A.G. Zecchinelli 

State University of New York at Buffalo, Buffalo, New York, USA

G. Agarwal , H. Bandyopadhyay , L. Hay , I. Iashvili , A. Kharchilava , C. McLean , D. Nguyen , J. Pekkanen , S. Rappoccio , A. Williams 






















Northeastern University, Boston, Massachusetts, USA

G. Alverson , E. Barberis , Y. Haddad , Y. Han , A. Hortiangtham , A. Krishna , J. Li , G. Madigan , B. Marzocchi , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari , A. Tishelman-Charny , T. Wamorkar , B. Wang , A. Wise-carver , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Bhattacharya , J. Bueghly, Z. Chen , A. Gilbert , T. Gunter , K.A. Hahn , Y. Liu , N. Odell , M.H. Schmitt , M. Velasco




















University of Notre Dame, Notre Dame, Indiana, USA

R. Band , R. Bucci, M. Cremonesi, A. Das , N. Dev , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , C. Jessop , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , K. Mohrman , C. Moore , Y. Musienko¹⁴ , R. Ruchti , A. Townsend , M. Wayne , M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, USA

B. Bylsma, L.S. Durkin , B. Francis , C. Hill , M. Nunez Ornelas , K. Wei, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, USA

F.M. Addesa , B. Bonham , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg , N. Haubrich , S. Higginbotham , A. Kalogeropoulos , G. Kopp , S. Kwan , D. Lange , D. Marlow , K. Mei , I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , S. Norberg

Purdue University, West Lafayette, Indiana, USA














A.S. Bakshi , V.E. Barnes , R. Chawla , S. Das , L. Gutay, M. Jones , A.W. Jung , D. Kondratyev , A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki 

S. Piperov , A. Purohit , J.F. Schulte , M. Stojanovic , J. Thieman , F. Wang ,
R. Xiao , W. Xie 











Purdue University Northwest, Hammond, Indiana, USA

J. Dolen , N. Parashar 
















Rice University, Houston, Texas, USA

D. Acosta , A. Baty , T. Carnahan , M. Decaro, S. Dildick , K.M. Ecklund ,
S. Freed, P. Gardner, F.J.M. Geurts , A. Kumar , W. Li , B.P. Padley , R. Redjimi,
W. Shi , A.G. Stahl Leiton , S. Yang , L. Zhang⁸⁸, Y. Zhang 

University of Rochester, Rochester, New York, USA

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , C. Fallon, T. Ferbel ,
M. Galanti, A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , E. Ranken ,
R. Taus 






Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , A. Gandrakota , Y. Gershtein , E. Halkiadakis , A. Hart ,
M. Heindl , O. Karacheban²⁵ , I. Laflotte , A. Lath , R. Montalvo, K. Nash, M. Os-
herson , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas,
H. Wang 










University of Tennessee, Knoxville, Tennessee, USA

H. Acharya, A.G. Delannoy , S. Fiorendi , S. Spanier 



Texas A&M University, College Station, Texas, USA

O. Bouhali⁸⁹ , M. Dalchenko , A. Delgado , R. Eusebi , J. Gilmore , T. Huang ,
T. Kamon⁹⁰ , H. Kim , S. Luo , S. Malhotra, R. Mueller , D. Overton , D. Rath-
jens , A. Safonov 














Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , V. Hegde , S. Kunori, K. Lamichhane , S.W. Lee ,
T. Mengke, S. Muthumuni , T. Peltola , I. Volobouev , Z. Wang, A. Whitbeck 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , S. Greene, A. Gurrola , W. Johns , A. Melo , H. Ni, K. Padeken ,
F. Romeo , P. Sheldon , S. Tuo , J. Velkovska 












University of Virginia, Charlottesville, Virginia, USA

M.W. Arenton , B. Cardwell , B. Cox , G. Cummings , J. Hakala , R. Hirosky ,
M. Joyce , A. Ledovsky , A. Li , C. Neu , C.E. Perez Lara , B. Tannenwald ,
S. White 

Wayne State University, Detroit, Michigan, USA

N. Poudyal 

University of Wisconsin - Madison, Madison, Wisconsin, USA

S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts , C. Galloni,
H. He , M. Herndon , A. Hervé , U. Hussain, A. Lanaro, A. Loeliger , R. Loveless 

J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, S. Trembath-Reichert, W. Vetens 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev, V. Andreev , Yu. Andreev , T. Aushev , M. Azarkin , A. Babaev , A. Belyaev , V. Blinov⁹¹, E. Boos , V. Borshch , D. Budkouski , V. Bunichev , O. Bychkova, V. Chekhovsky, R. Chistov⁹¹ , M. Danilov⁹¹ , A. Dermenev , T. Dimova⁹¹ , I. Dremin , M. Dubinin⁸¹ , L. Dudko , V. Epshteyn⁹² , A. Ershov , G. Gavrillov , V. Gavrillov , S. Gninenko , V. Golovtcov , N. Golubev , I. Golutvin, I. Gorbunov , A. Gribushin , V. Ivanchenko , Y. Ivanov , V. Kachanov , L. Kardapoltsev⁹¹ , V. Karjavine , A. Karneyeu , V. Kim⁹¹ , M. Kirakosyan, D. Kirpichnikov , M. Kirsanov , V. Klyukhin , O. Kodolova⁹³ , D. Konstantinov , V. Koronkov , A. Kozyrev⁹¹ , N. Krasnikov , E. Kuznetsova⁹⁴, A. Lanev , A. Litomin, N. Lychkovskaya , V. Makarenko⁹¹ , A. Malakhov , V. Matveev⁹¹ , V. Murzin , A. Nikitenko⁹⁵ , S. Obraztsov , V. Okhotnikov , V. Oreshkin , A. Oskin, I. Ovtin⁹¹ , V. Palichik , P. Parygin⁹⁶ , A. Pashenkov, V. Perelygin , S. Petrushanko , G. Pivovarov , S. Polikarpov⁹¹ , V. Popov, O. Radchenko⁹¹ , M. Savina , V. Savrin , D. Seitova, V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen⁹¹ , S. Slabospitskii , I. Smirnov, V. Smirnov , D. Sosnov , A. Stepenov , V. Sulimov , E. Tcherniaev , A. Terkulov , O. Teryaev , M. Toms⁹⁷ , A. Toropin , L. Uvarov , A. Uzunian , E. Vlasov⁹⁸ , S. Volkov, A. Vorobyev, N. Voytishin , B.S. Yuldashev⁹⁹, A. Zarubin , I. Zhizhin , A. Zhokin 

[†] Deceased

¹ Also at Yerevan State University, Yerevan, Armenia

² Also at TU Wien, Vienna, Austria

³ Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt

⁴ Also at Université Libre de Bruxelles, Bruxelles, Belgium

⁵ Also at Punjab Agricultural University, Ludhiana, India

⁶ Also at Universidade Estadual de Campinas, Campinas, Brazil

⁷ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁸ Also at The University of the State of Amazonas, Manaus, Brazil

⁹ Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰ Also at UFMS, Nova Andradina, Brazil

¹¹ Also at Nanjing Normal University Department of Physics, Nanjing, China

¹² Now at The University of Iowa, Iowa City, Iowa, USA

¹³ Also at University of Chinese Academy of Sciences, Beijing, China

¹⁴ Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹⁵ Now at British University in Egypt, Cairo, Egypt

¹⁶ Now at Cairo University, Cairo, Egypt

¹⁷ Also at Purdue University, West Lafayette, Indiana, USA

¹⁸ Also at Université de Haute Alsace, Mulhouse, France

¹⁹ Also at Ilia State University, Tbilisi, Georgia

- 20 Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- 21 Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- 22 Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
- 23 Also at University of Hamburg, Hamburg, Germany
- 24 Also at Isfahan University of Technology, Isfahan, Iran
- 25 Also at Brandenburg University of Technology, Cottbus, Germany
- 26 Also at Forschungszentrum Jülich, Juelich, Germany
- 27 Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- 28 Also at Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary
- 29 Also at Institute of Physics, University of Debrecen, Debrecen, Hungary
- 30 Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary
- 31 Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- 32 Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- 33 Also at Wigner Research Centre for Physics, Budapest, Hungary
- 34 Also at Shoolini University, Solan, India
- 35 Also at University of Hyderabad, Hyderabad, India
- 36 Also at University of Visva-Bharati, Santiniketan, India
- 37 Also at Indian Institute of Science (IISc), Bangalore, India
- 38 Also at Indian Institute of Technology (IIT), Mumbai, India
- 39 Also at IIT Bhubaneswar, Bhubaneswar, India
- 40 Also at Institute of Physics, Bhubaneswar, India
- 41 Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- 42 Also at Sharif University of Technology, Tehran, Iran
- 43 Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- 44 Also at Helwan University, Cairo, Egypt
- 45 Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- 46 Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- 47 Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- 48 Also at Università di Napoli 'Federico II', Napoli, Italy
- 49 Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- 50 Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- 51 Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France
- 52 Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia
- 53 Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- 54 Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy
- 55 Also at National and Kapodistrian University of Athens, Athens, Greece
- 56 Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- 57 Also at Universität Zürich, Zurich, Switzerland
- 58 Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- 59 Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- 60 Also at Şirnak University, Sirmak, Turkey
- 61 Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- 62 Also at Konya Technical University, Konya, Turkey

- 63 Also at Izmir Bakircay University, Izmir, Turkey
64 Also at Adiyaman University, Adiyaman, Turkey
65 Also at Necmettin Erbakan University, Konya, Turkey
66 Also at Bozok Universititesi Rektörlüğü, Yozgat, Turkey
67 Also at Marmara University, Istanbul, Turkey
68 Also at Milli Savunma University, Istanbul, Turkey
69 Also at Kafkas University, Kars, Turkey
70 Also at Istanbul Bilgi University, Istanbul, Turkey
71 Also at Hacettepe University, Ankara, Turkey
72 Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
73 Also at Ozyegin University, Istanbul, Turkey
74 Also at Vrije Universiteit Brussel, Brussel, Belgium
75 Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
76 Also at IPPP Durham University, Durham, United Kingdom
77 Also at Monash University, Faculty of Science, Clayton, Australia
78 Also at Università di Torino, Torino, Italy
79 Also at Bethel University, St. Paul, Minnesota, USA
80 Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
81 Also at California Institute of Technology, Pasadena, California, USA
82 Also at United States Naval Academy, Annapolis, Maryland, USA
83 Also at Ain Shams University, Cairo, Egypt
84 Also at Bingol University, Bingol, Turkey
85 Also at Georgian Technical University, Tbilisi, Georgia
86 Also at Sinop University, Sinop, Turkey
87 Also at Erciyes University, Kayseri, Turkey
88 Also at Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China
89 Also at Texas A&M University at Qatar, Doha, Qatar
90 Also at Kyungpook National University, Daegu, Korea
91 Also at another institute or international laboratory covered by a cooperation agreement with CERN
92 Now at Istanbul University, Istanbul, Turkey
93 Also at Yerevan Physics Institute, Yerevan, Armenia
94 Now at University of Florida, Gainesville, Florida, USA
95 Also at Imperial College, London, United Kingdom
96 Now at University of Rochester, Rochester, New York, USA
97 Now at Baylor University, Waco, Texas, USA
98 Now at INFN Sezione di Torino, Università di Torino, Torino, Italy; Università del Piemonte Orientale, Novara, Italy
99 Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan