

Absence of embedded eigenvalues of Pauli and Dirac operators

Dirk Hundertmark, Hynek Kovařík

CRC Preprint 2023/26, December 2023

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Wave
phenomena

Participating universities



Funded by



ABSENCE OF EMBEDDED EIGENVALUES OF PAULI AND DIRAC OPERATORS

DIRK HUNDERTMARK AND HYNEK KOVAŘÍK

ABSTRACT. We consider eigenvalues of the Pauli operator in \mathbb{R}^3 embedded in the continuous spectrum. In our main result we prove the absence of such eigenvalues above a threshold which depends on the asymptotic behavior of the magnetic and electric field at infinity. We show moreover that the decay conditions on the magnetic and electric field are sharp. Analogous results are obtained for purely magnetic Dirac operators.

Keywords: Pauli operator, Dirac operator, embedded eigenvalues.

MSC 2020: 35Q40, 35P05

1. Introduction and outline of the paper

In this paper we study the point spectrum of the Pauli operator in $L^2(\mathbb{R}^3, \mathbb{C}^2)$ formally given by

$$H_{A,V} = (\sigma \cdot (P - A))^2 + V. \quad (1.1)$$

Here $P = -i\nabla$ denotes the momentum operator, $A \in L^2_{\text{loc}}(\mathbb{R}^3, \mathbb{R}^3)$ is a magnetic vector potential, $\sigma = (\sigma_1, \sigma_2, \sigma_3)$ is the set of Pauli matrices, see equation (2.1) below, and V is a potential function which associates to each $x \in \mathbb{R}^3$ a two by two hermitian matrix $V(x)$. We refer to equations (4.4) and (4.5) for a more precise definition of $H_{A,V}$. The free Pauli operator $H_{A,0}$ represents a quantum Hamiltonian of a particle with spin $\frac{1}{2}$ interacting with a magnetic field $B = \text{curl } A$, see e.g. [23] for further reading and references.

Our aim is to find sharp conditions on B and V under which the operator $H_{A,V}$ has no eigenvalues above certain critical energy.

The absence of discrete eigenvalues of $H_{A,V}$, also in dimensions higher than three, can be deduced from the results of [10], where the authors show, via the method of multipliers, that if B and V satisfy certain smallness assumptions, then $H_{A,V}$ has no eigenvalues at all.

The absence of eigenvalues at the threshold of the essential spectrum, typically zero, is also well understood, at least in the case $V = 0$. A sharp criterion for zero to be an eigenvalue of $H_{A,0}$ was recently established in [12, 13], see also [6, 7]. In particular, it is proved in [13] that $H_{A,0}$ can have a zero energy eigenfunction only if $\|A\|_{L^3(\mathbb{R}^3)}$ exceeds certain explicit value. Examples of magnetic fields which produce zero energy eigenfunctions of $H_{A,0}$, and which show that the criterion of [13] is sharp, can be found in [1, 11, 19, 21]. We will give more comments on this question in Remark 6.6.

What is not well understood so far is the question of absence of eigenvalues embedded in the essential spectrum, which is of fundamental importance e.g. for the validity of a limiting absorption principle, for the scattering theory, as well as for dispersive estimates. One could of course apply the result of [10], since the conditions stated there guarantee not only the absence of discrete eigenvalues, but also the absence of *all eigenvalues*, [10, Thm. 3.5]. However, this automatically implies that such conditions are way too strong if

©2023 by the authors. Faithful reproduction of this article, in its entirety, by any means is permitted for non-commercial purposes.

one is interested only in embedded eigenvalues, since creating discrete eigenvalues is usually much “easier” than creating eigenvalues embedded in the essential spectrum. Indeed, consider the generic case in which $\sigma_{\text{es}}(H_{A,0}) = [0, \infty)$. Then any negative and sufficiently strong potential V will create negative eigenvalues, but it should typically not create positive eigenvalues, at least when B and V decay fast enough at infinity. Hence in order to exclude all eigenvalues, one has to impose global smallness assumptions on B and V , see [10, Thm. 3.5]. On the other hand, embedded eigenvalues belong to the essential spectrum and therefore their absence or existence should depend only on the behavior of B and V at infinity.

In this paper we prove that the operator $H_{A,V}$ cannot have eigenvalues above an energy level $\Lambda \geq 0$ allowing, at the same time, $H_{A,V}$ to have discrete and/or threshold eigenvalues, see Theorem 6.5. We provide an explicit expression for Λ which shows, in agreement with the above heuristics, that Λ depends only on the behavior of B and V at infinity. In particular, no global bounds on B and V are needed.

Let us describe the main result of this paper more in detail. In Theorem 6.5 it is proved, under rather mild regularity and decay conditions on B and V , that $H_{A,V}$ has no eigenvalues larger than

$$\Lambda = \Lambda(B, V) := \frac{1}{4} \left(\beta + \omega_1 + \sqrt{(\beta + \omega_1)^2 + 2\omega_2} \right)^2, \quad (1.2)$$

where β, ω_1 and ω_2 are non-negative constants which depend, in a weak sense, on the spacial asymptotics of B and V . We refer to Assumption 3.9 and equation (3.10) for a full definition of β and ω_j . If B and V are regular enough at infinity, then the values of β and ω_j are determined from their pointwise asymptotics. Indeed, splitting the potential into a sum of its short-range and long-range component; $V = V^s + V^\ell$, we find

$$\beta \leq \limsup_{|x| \rightarrow \infty} |\tilde{B}(x)|, \quad \omega_1 \leq \limsup_{|x| \rightarrow \infty} |x V^s(x)|_{\mathbb{C}^2}, \quad \text{and} \quad \omega_2 \leq \limsup_{|x| \rightarrow \infty} |(x \cdot \nabla V^\ell(x))_+|_{\mathbb{C}^2}.$$

see Lemma A.1.

Remark 1.1. It is illustrative to compare Theorem 6.5 with classical results on the absence of positive eigenvalues of non-magnetic Schrödinger operators [2, 17, 22]. If $B = 0$, then by choosing $V^s = V$ and $V^\ell = 0$ we obtain $\Lambda = \omega_1^2$ which generalizes the result of Kato [17]. On the other hand, by choosing V^s such that $V^s(x) = o(|x|^{-1})$, and setting $V^\ell = V - V^s$ we get $\Lambda = \omega_2/2$, and recover thus the results of Agmon [2] and Simon [22].

To prove Theorem 6.5 we adapt a version of the quadratic form method of [4], which in turn is inspired by the approach invented by Froese and Herbst for non-magnetic Schrödinger operators in [14, 15]. However, due to the spinor structure of the operator $H_{A,V}$ and of its wave-functions, the technique of [4, 14, 15] cannot be applied directly. The problem is that the operator-valued matrix $H_{A,V}$ is, contrary to the two-dimensional case, non-diagonal. Consequently, a direct application of the above mentioned technique, developed for scalar magnetic operators, is not feasible. It is therefore necessary to implement the fundamental ingredients of [4] in such a way that the spinor structure of $H_{A,V}$ be taken into account. To do so we make use of multiplication and commutation relations for the Pauli matrices and of their convenient interplay with the Poincaré gauge for the vector potential A . This is yet another example of the importance of choosing a gauge which suits best the problem in question. In our case the choice of the Poincaré gauge, together with the properties of the Pauli matrices, allows us to prove a matrix-valued versions of the virial-type identities for the weighted commutator between $H_{A,V}$ and the generator of dilations, see equations (5.10) and (5.15). With the help of these identities we then show that any eigenfunction of $H_{A,V}$ with eigenvalue larger than Λ must identically vanish. We would like to point out that although the identities (5.10) and (5.15) are identical to their scalar counterparts obtained in [4], due to the spinor structure of the problem under consideration their derivation is essentially different.

The paper is organized in the following way. In the first two sections we collect necessary prerequisites and state our hypotheses. In Sections 4 and 5 we prove some preliminary results concerning dilations and commutator properties of $H_{A,V}$. The main result is stated and proved in Section 6. In Section 7 we construct an example which shows that the critical energy $\Lambda(B, V)$ given by (1.2) is sharp. As a consequence of Theorem 6.5 we also establish sufficient conditions for the absence of embedded eigenvalues of the magnetic Dirac operator, see Theorem 8.1 and Corollary 8.2. In Appendix A we show that all the hypothesis stated in Section 3 are satisfied under some mild pointwise conditions on B and V .

2. Prerequisites

2.1. Basic setup. We identify the magnetic field with the vector-field $B : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with components (B_1, B_2, B_3) . A vector potential is a vector field $A : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ which generates a magnetic fields via $B = \text{curl}A$, in the distributional sense. We recall the well-known Pauli matrices $\sigma_j : \mathbb{C}^2 \rightarrow \mathbb{C}^2$;

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (2.1)$$

In what follows we use the shorthand

$$z \cdot \sigma = \sum_{j=1}^3 z_j \sigma_j \quad z \in \mathbb{C}^3. \quad (2.2)$$

The Pauli matrices satisfy the following multiplication and commutation relations,

$$\sigma_j \sigma_k = \delta_{jk} \mathbf{1} + i \sum_{m=1}^3 \varepsilon_{jkm} \sigma_m \quad (2.3)$$

$$[\sigma_j, \sigma_k] = 2i \sum_{m=1}^3 \varepsilon_{jkm} \sigma_m. \quad (2.4)$$

Here $\mathbf{1}$ is the unite 2×2 matrix, and ε_{jkm} denotes the Levi-Civita permutation symbol. In particular, $\sigma_j^2 = \mathbf{1}$ for $j = 1, 2, 3$.

Given a magnetic field B and a point $w \in \mathbb{R}^3$ let $\tilde{B}_w(x) := B(x+w)[x]$. More precisely, \tilde{B}_w is a vector-field on \mathbb{R}^3 defined by

$$\tilde{B}_w(x+w) = B(x+w) \wedge x. \quad (2.5)$$

Making use of translations, we will often assume $w = 0$, in which case we will simply write \tilde{B} .

2.2. Notation. If $A \in L^2_{\text{loc}}(\mathbb{R}^3, \mathbb{R}^3)$ is a magnetic vector potential, the the magnetic Sobolev space is defined by

$$\mathcal{H}^1(\mathbb{R}^3, \mathbb{C}^2) := \mathcal{D}(P - A) = \left\{ \varphi \in L^2(\mathbb{R}^3, \mathbb{C}^2) : (P - A) \varphi \in L^2(\mathbb{R}^3, \mathbb{C}^2) \right\}, \quad (2.6)$$

equipped with the graph norm

$$\|u\|_{\mathcal{H}^1} = \left(\|(P - A)u\|_{L^2(\mathbb{R}^3, \mathbb{C}^2)}^2 + \|u\|_{L^2(\mathbb{R}^3, \mathbb{C}^2)}^2 \right)^{1/2}. \quad (2.7)$$

The corresponding scalar Sobolev space will be denoted by

$$\mathcal{H}^1(\mathbb{R}^3) = \{u \in L^2(\mathbb{R}^3) : (P - A)u \in L^2(\mathbb{R}^3)\}.$$

Given a set M and two functions $f_1, f_2 : M \rightarrow \mathbb{R}$, we write $f_1(x) \lesssim f_2(x)$ if there exists a numerical constant c such that $f_1(x) \leq c f_2(x)$ for all $x \in M$. The symbol $f_1(x) \gtrsim f_2(x)$ is defined analogously. Moreover, we use the notation

$$f_1(x) \sim f_2(x) \quad \Leftrightarrow \quad f_1(x) \lesssim f_2(x) \wedge f_2(x) \lesssim f_1(x),$$

and

$$\limsup_{|x| \rightarrow \infty} f(x) = L \quad \Leftrightarrow \quad \lim_{r \rightarrow \infty} \operatorname{ess\,sup}_{|x| \geq r} f(x) = L, \quad (2.8)$$

and similarly for $\liminf_{|x| \rightarrow \infty} f(x)$. We will use $\partial_j = \frac{\partial}{\partial x_j}$ for the usual partial derivatives in the weak sense, i.e., as distributions.

The scalar product on a Hilbert space \mathcal{H} will be denoted by $\langle \cdot, \cdot \rangle_{\mathcal{H}}$. If $\mathcal{H} = L^2(\mathbb{R}^3, \mathbb{C}^2)$, we omit the subscript and write

$$\langle \varphi, \psi \rangle_{L^2(\mathbb{R}^3, \mathbb{C}^2)} = \langle \varphi, \psi \rangle, \quad \varphi, \psi \in L^2(\mathbb{R}^3, \mathbb{C}^2).$$

Accordingly, for any $\varphi \in L^r(\mathbb{R}^3, \mathbb{C}^2)$ with $1 \leq r \leq \infty$ we will use the shorthand

$$\|\varphi\|_r := \|\varphi\|_{L^r(\mathbb{R}^3, \mathbb{C}^2)}$$

for the L^r -norm of φ . By the symbol

$$\mathcal{U}_R(x) = \{y \in \mathbb{R}^3 : |x - y| < R\}$$

we denote the ball of radius R centered at a point $x \in \mathbb{R}^3$. If $x = 0$, we abbreviate $\mathcal{U}_R = \mathcal{U}_R(0)$.

Given a Hermitian matrix valued function $\mathbb{R}^3 \ni x \mapsto M(x) : \mathbb{C}^2 \rightarrow \mathbb{C}^2$, we denote by $\lambda(x)$ and $\mu(x)$ its eigenvalues. The norm of M is then equal to

$$|M(x)|_{\mathbb{C}^2} = \max \{ |\lambda(x)|, |\mu(x)| \}.$$

Accordingly we define

$$|M(x)_+|_{\mathbb{C}^2} = \max \{ \lambda(x)_+, \mu(x)_+ \}. \quad (2.9)$$

Convention: In the sequel we will use latin letters for functions with values in \mathbb{C} , and greek letters for functions with values in \mathbb{C}^2 . In particular, we will often identify a spinor φ with two scalar fields as follows;

$$\varphi = \begin{pmatrix} u \\ v \end{pmatrix}. \quad (2.10)$$

Throughout the paper we will often make use of the polarisation identity which, for the reader's convenience, we now briefly recall; given a sesquilinear form s on a Hilbert space \mathcal{H} , and any $\varphi, \psi \in \mathcal{H}$, we have

$$s(\varphi, \psi) = \frac{1}{4} \left[s(\varphi + \psi, \varphi + \psi) - s(\varphi - \psi, \varphi - \psi) + is(\varphi - i\psi, \varphi - i\psi) - is(\varphi + i\psi, \varphi + i\psi) \right]. \quad (2.11)$$

2.3. The Poincaré gauge. For a given magnetic field B and a point $w \in \mathbb{R}^3$ we define the vector field \tilde{B}_w by equation (2.5), and put

$$A_w(x) := \int_0^1 \tilde{B}_w(tx) dt, \quad (2.12)$$

which is the vector potential in the Poincaré gauge. Using translations, it is no loss of generality to assume $w = 0$, in which case we will simply write A for the vector potential given by (2.12). Note that when $w = 0$, then A given by (2.12) satisfies

$$x \cdot A(x) = 0 \quad \forall x \in \mathbb{R}^3. \quad (2.13)$$

It is easy to see that for A given by (2.12) one has $A \in L^2_{\text{loc}}(\mathbb{R}^3, \mathbb{R}^3)$ for bounded magnetic fields B and this extends to a large class of singular magnetic fields, see [4, Lem. 2.9]. Except otherwise stated, we will always use the Poincaré gauge.

3. Hypotheses

In this section we formulate general sufficient conditions on B and V under which our main result, Theorem 6.5, holds true. In Appendix A we will show that all these conditions are satisfied under rather mild assumptions on B and V , see in particular Lemma A.1, Proposition A.2 and Proposition A.4.

Assumption 3.1. The matrix valued function $V : \mathbb{R}^3 \rightarrow M(2, \mathbb{C})$ is Hermitian. i.e.

$$(V(x))_{jk} = \overline{(V(x))_{kj}} \quad \forall x \in \mathbb{R}^3, \quad \forall j, k = 1, 2. \quad (3.1)$$

If the potential is split as $V = V^s + V^\ell$, then V^s and V^ℓ also satisfy (3.1).

Remark 3.2. Similarly as in the scalar non-magnetic case, see in particular [15, Thm. 2.1], our results could be extended to all matrix valued V , possibly non-Hermitian, for which the associated Pauli operator $H_{A,V}$ has real spectrum. For the sake of brevity, we will stick to Assumption 3.1 throughout the paper.

Assumption 3.3. The magnetic field B is such that for some $w \in \mathbb{R}^3$

$$\mathbb{R}^3 \ni x \mapsto |x - w|^{-1} \log_+^2 \left(\frac{R}{|x - w|} \right) |\tilde{B}_w(x)|^2 \in L_{\text{loc}}^1(\mathbb{R}^3) \quad (3.2)$$

for all $R > 0$.

We have already pointed out that without loss of generality we may assume $w = 0$. In view of [4, Lem. 2.9] condition (3.2) assures that the corresponding vector potential in the Poincaré gauge is locally square integrable. The latter property is essential in order to define the Pauli operator through the associated quadratic form.

3.1. Global relative bounds.

Assumption 3.4. The scalar fields $|B|^2$ and $|\tilde{B}|^2$ are relatively form bounded w.r.t. $(P - A)^2$, where A is the Poincaré gauge vector potential corresponding to B . That is,

$$\| |B|\varphi \|_2^2 + \| |\tilde{B}|\varphi \|_2^2 \lesssim \| (P - A)\varphi \|_2^2 + \|\varphi\|_2^2 \quad \forall \varphi \in \mathcal{D}(P - A). \quad (3.3)$$

Here we abuse the notation and use the same symbol $P - A$ for the operator in $L^2(\mathbb{R}^3)$ as well as for the operator in $L^2(\mathbb{R}^3, \mathbb{C}^2)$ acting as $\mathbf{1}(P - A)$.

Assumption 3.5. The potential V is relatively form small w.r.t. $(P - A)^2$, that is, there exist constants $\alpha_0 < 1$ and C such that

$$|\langle \varphi, V\varphi \rangle| \leq \alpha_0 \| (P - A)\varphi \|_2^2 + C \|\varphi\|_2^2 \quad \forall \varphi \in \mathcal{D}(P - A). \quad (3.4)$$

In order to control the virial $x \cdot \nabla V$, we decompose the potential as $V = V^s + V^\ell$. The splitting $V = V^s + V^\ell$ is arbitrary, as long as the conditions below are satisfied.

3.2. Behaviour at infinity. Below we quantify the notions of boundedness and vanishing at infinity w.r.t. $(P - A)^2$.

Definition 3.6 (Boundedness at infinity). A potential W is bounded from above at infinity with respect to $(P - A)^2$ if for some $R_0 > 0$ its quadratic form domain contains all $\varphi \in \mathcal{D}(P - A)$ with $\text{supp}(\varphi) \in \mathcal{U}_{R_0}^c$ and for $R \geq R_0$ there exist positive α_R, γ_R with $\lim_{R \rightarrow \infty} \alpha_R = 0$ and $\liminf_{R \rightarrow \infty} \gamma_R < \infty$ such that

$$\langle \varphi, W\varphi \rangle \leq \alpha_R \| (P - A)\varphi \|_2^2 + \gamma_R \|\varphi\|_2^2 \quad \text{for all } \varphi \in \mathcal{D}(P - A) \text{ with } \text{supp}(\varphi) \subset \mathcal{U}_R^c \quad (3.5)$$

By monotonicity we may assume, without loss of generality, that α_R and γ_R are decreasing in $R \geq R_0$.

Assumption 3.7. The positive part of the potential V vanishes at infinity w.r.t. $(P - A)^2$ in the following sense: there exist positive α_R, γ_R with $\alpha_R, \gamma_R \rightarrow 0$ as $R \rightarrow \infty$ such that

$$\langle \varphi, V\varphi \rangle_+ \leq \alpha_R \|(P - A)\varphi\|_2^2 + \gamma_R \|\varphi\|_2^2 \quad \text{for all } \varphi \in \mathcal{D}(P - A) \text{ with } \text{supp}(\varphi) \subset \mathcal{U}_R^c. \quad (3.6)$$

Moreover, if we split $V = V^s + V^\ell$, then also the positive parts of V^s and V^ℓ vanish at infinity in the sense defined above. By monotonicity we may assume, without loss of generality, that α_R and γ_R are decreasing in $R \geq R_0$.

Assumption 3.8. The potential V is bounded at infinity w.r.t. $(P - A)^2$ in the sense of Definition 3.6. Moreover, if we split $V = V^s + V^\ell$, then also V^s is bounded at infinity w.r.t. $(P - A)^2$ in the sense of Definition 3.6.

Assumption 3.9. There exist positive sequences $(\varepsilon_j)_j, (\beta_j)_j$ and $(R_j)_j$ with $\varepsilon_j \rightarrow 0$ and $R_j \rightarrow \infty$ as $j \rightarrow \infty$, such that for all $\varphi \in \mathcal{D}(P - A)$ with $\text{supp}(\varphi) \subset \mathcal{U}_j^c = \{x \in \mathbb{R}^3 : |x| \geq R_j\}$

$$\|\tilde{B}|\varphi\|_2^2 \leq \varepsilon_j \|(P - A)\varphi\|_2^2 + \beta_j^2 \|\varphi\|_2^2 \quad (3.7)$$

For the decomposition $V = V^s + V^\ell$ of the potential, we also assume that there exist positive sequences $(\omega_{1,j})_j$ and $(\omega_{2,j})_j$ such that for all $\varphi \in \mathcal{D}(P - A)$ with $\text{supp}(\varphi) \subset \mathcal{U}_j^c$

$$\|xV^s\varphi\|_2^2 \leq \varepsilon_j \|(P - A)\varphi\|_2^2 + \omega_{1,j}^2 \|\varphi\|_2^2 \quad (3.8)$$

$$\langle \varphi, x \cdot \nabla V^\ell \varphi \rangle \leq \varepsilon_j \|(P - A)\varphi\|_2^2 + \omega_{2,j} \|\varphi\|_2^2 \quad (3.9)$$

By monotonicity we may assume, without loss of generality, that the sequences $\beta_j, \omega_{1,j}$, and $\omega_{2,j}$ in Assumptions 3.9 are decreasing. We define

$$\beta := \lim_{j \rightarrow \infty} \beta_j, \quad \omega_k := \lim_{j \rightarrow \infty} \omega_{k,j}, \quad k = 1, 2. \quad (3.10)$$

3.3. Unique continuation at infinity. For a unique continuation type argument at infinity, we also need a quantitative version of relative form boundedness.

Assumption 3.10. If $V = V^s + V^\ell$, then we assume that for all $\varphi \in \mathcal{D}(P - A)$

$$\|\tilde{B}|\varphi\|_2^2 + \|xV^s\varphi\|_2^2 \leq \frac{\alpha_1^2}{4} \|(P - A)\varphi\|_2^2 + C\|\varphi\|_2^2, \quad (3.11)$$

$$\langle \varphi, x \cdot \nabla V^\ell \varphi \rangle \leq \alpha_2 \|(P - A)\varphi\|_2^2 + C\|\varphi\|_2^2, \quad (3.12)$$

$$|\langle \varphi, V^s \varphi \rangle| \leq \alpha_3 \|(P - A)\varphi\|_2^2 + C\|\varphi\|_2^2 \quad (3.13)$$

for some $C > 0$ and α_j such that

$$\alpha_1 + \alpha_2 + 3\alpha_3 < 1. \quad (3.14)$$

Remark 3.11. By the diamagnetic inequality

$$|P|\varphi| \leq |(P - A)\varphi| \quad \text{a.e.} \quad \text{for all } \varphi \in \mathcal{D}(P - A), \quad (3.15)$$

see e.g. [18], it suffices to verify the conditions of Assumptions 3.4-3.10 with $(P - A)$ replaced by P .

4. Preliminary results

In this section we collect several technical results which will be needed later.

Lemma 4.1. *Let $A \in L^2_{\text{loc}}(\mathbb{R}^3)$ and let $B = \text{curl } A$. Suppose moreover that B satisfy Assumption 3.4. Then*

$$\langle \sigma \cdot (P - A)\varphi, \sigma \cdot (P - A)\varphi \rangle = \|(P - A)\varphi\|_2^2 + \langle \varphi, \sigma \cdot B\varphi \rangle \quad \forall \varphi \in \mathcal{H}^1(\mathbb{R}^3, \mathbb{C}^2).$$

Proof. The claim follows by a direct calculation from (2.3) and (2.4). ■

Lemma 4.2. *Let B satisfy Assumption 3.4. For any $\eta > 0$ there exists $C_\eta \in \mathbb{R}$ such that*

$$\|(P - A)\varphi\|_2^2 \leq (1 + \eta)\|\sigma \cdot (P - A)\varphi\|_2^2 + C_\eta\|\varphi\|_2^2 \quad (4.1)$$

holds for all $\varphi \in \mathcal{D}(P - A)$.

Proof. Let $\varphi \in \mathcal{D}(P - A)$. A short calculation shows that

$$\|\sigma \cdot w\varphi\|_2 = \| |w| \varphi \|_2 \quad \forall w \in \mathbb{C}^3. \quad (4.2)$$

Hence by (3.3), Lemma 4.1 and Cauchy-Schwarz inequality,

$$\begin{aligned} \|(P - A)\varphi\|_2^2 &= \|\sigma \cdot (P - A)\varphi\|_2^2 - \langle \varphi, \sigma \cdot B\varphi \rangle \leq \|\sigma \cdot (P - A)\varphi\|_2^2 + \| |B| \varphi \|_2 \|\varphi\|_2 \\ &\leq \|\sigma \cdot (P - A)\varphi\|_2^2 + \varepsilon \|(P - A)\varphi\|_2^2 + C_\varepsilon \|\varphi\|_2^2 \end{aligned}$$

for any $0 < \varepsilon < 1$ and some C_ε , independent of φ . Inequality (4.1) now follows upon setting $\frac{1}{1-\varepsilon} = 1 + \eta$. ■

An immediate consequence of Lemma 4.2 is the following

Corollary 4.3. *Let B and V satisfy Assumptions 3.4, 3.8 and 3.10. Then for all $\varphi \in \mathcal{D}(P - A)$,*

$$\begin{aligned} |\langle \varphi, V\varphi \rangle| &\leq \alpha_0 \|(P - A)\varphi\|_2^2 + C_0\|\varphi\|_2^2 \\ \|\tilde{B}\varphi\|_2^2 + \|xV^s\varphi\|_2^2 &\leq \frac{\alpha_1^2}{4} \|(P - A)\varphi\|_2^2 + C_1\|\varphi\|_2^2, \\ \langle \varphi, x \cdot \nabla V^\ell \varphi \rangle &\leq \alpha_2 \|(P - A)\varphi\|_2^2 + C_2\|\varphi\|_2^2, \\ |\langle \varphi, V^s \varphi \rangle| &\leq \alpha_3 \|(P - A)\varphi\|_2^2 + C_3\|\varphi\|_2^2 \end{aligned}$$

where $\alpha_0 < 1$, and $\alpha_j, j = 1, 2, 3$ satisfy (3.14).

Another consequence of Lemma 4.2 is the identity

$$\mathcal{D}(P - A) = \mathcal{D}(\sigma \cdot (P - A)), \quad (4.3)$$

which holds whenever Assumption 3.4 is satisfied. This allows us to define the sesquilinear form

$$Q_{A,0}(\varphi, \psi) = \langle \sigma \cdot (P - A)\varphi, \sigma \cdot (P - A)\psi \rangle, \quad \varphi, \psi \in \mathcal{D}(P - A). \quad (4.4)$$

By standard arguments one verifies that the quadratic form $Q_{A,0}(\varphi, \varphi)$ is closed. In view of Lemma 4.2 and Assumption 3.5 the quadratic form associated to

$$Q_{A,V}(\varphi, \psi) = Q_{A,0}(\varphi, \psi) + \langle \varphi, V\psi \rangle, \quad \varphi, \psi \in \mathcal{D}(P - A) \quad (4.5)$$

is then closed as well. Now we can define the Hamiltonians $H_{A,0}$ and $H_{A,V}$ as the unique self-adjoint operators associated to $Q_{A,0}$ and $Q_{A,V}$ respectively.

For the next result we need to introduce some additional notation. Given a vector field $v : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ we define the operator

$$D_v = \frac{1}{2}(v \cdot P + P \cdot v), \quad D := D_x \quad \text{if } v = x. \quad (4.6)$$

Lemma 4.4. *Let B satisfy Assumption 3.3. Let $g, F \in C^1(\mathbb{R}^3)$ $g, F \in C^1(\mathbb{R}^3; \mathbb{R})$ be radial functions such that $\nabla F = xg$, and such that $x \cdot \nabla g$ and $|x|g$ are bounded. Put $v = \nabla F$. Then*

$$\mathcal{D}(P - A) \subset \mathcal{D}(D_v) = \mathcal{D}(gD) \quad (4.7)$$

Proof. In the sense of distributions,

$$2D_v = gx \cdot P + P \cdot (gx) = gx \cdot P + gP \cdot x - ix \cdot \nabla g = 2gD - ix \cdot \nabla g.$$

So if $\varphi \in \mathcal{D}(gD)$ and $x \cdot \nabla g$ is bounded, then

$$D_v \varphi = gD\varphi - ix \cdot \nabla g \varphi \in L^2(\mathbb{R}^3, \mathbb{C}^2).$$

Hence $\varphi \in \mathcal{D}(D_v)$. Conversely, if $\varphi \in \mathcal{D}(D_v)$ and $x \cdot \nabla g$ is bounded, then $\varphi \in \mathcal{D}(gD)$. This proves the equality $\mathcal{D}(D_v) = \mathcal{D}(gD)$. Moreover, since $x \cdot A(x) = 0$,

$$2gD = g(x \cdot P + P \cdot x) = 2gx \cdot P - 3i = 2gx \cdot (P - A) - 3i.$$

So if $\varphi \in \mathcal{D}(P - A)$, and $|x|g$ is bounded, then $gD\varphi \in L^2(\mathbb{R}^3, \mathbb{C}^2)$. Hence $\mathcal{D}(P - A) \subset \mathcal{D}(gD) = \mathcal{D}(D_v)$. ■

Lemma 4.5. *Under the assumptions of Lemma 4.4,*

$$\operatorname{Re} \langle (\sigma \cdot v) \varphi, \sigma \cdot (P - A) \varphi \rangle = \langle \varphi, D_v \mathbf{1} \varphi \rangle \quad \forall \varphi \in \mathcal{D}(P - A). \quad (4.8)$$

Remark 4.6. We note that thanks to Lemma 4.4, the right hand side of (4.8) is well defined for all $\varphi \in \mathcal{D}(P - A)$. Lemma 4.4 is also used implicitly in Lemma 5.1 and in Proposition 5.3.

Proof of Lemma 4.5. Let $\varphi \in C_0^\infty(\mathbb{R}^3, \mathbb{C}^2)$. Then

$$\operatorname{Re} \langle (\sigma \cdot v) \varphi, \sigma \cdot (P - A) \varphi \rangle = \operatorname{Re} \langle \varphi, (\sigma \cdot v) \sigma \cdot (P - A) \varphi \rangle = \operatorname{Re} \langle \varphi, (\sigma \cdot v) \sigma \cdot P \varphi \rangle - \operatorname{Re} \langle \varphi, (\sigma \cdot v) \sigma \cdot A \varphi \rangle.$$

In view of (2.3),

$$(\sigma \cdot v)(\sigma \cdot A) = \sum_{j,k=1}^3 v_j A_k \sigma_j \sigma_k = (v \cdot A) \mathbf{1} + i \sum_{m=1}^3 \left(\sum_{j,k=1}^3 \varepsilon_{mjk} v_j A_k \right) \sigma_m = (v \cdot A) \mathbf{1} + i(v \wedge A) \cdot \sigma.$$

Hence

$$\operatorname{Re} \langle \varphi, (\sigma \cdot v) \sigma \cdot A \varphi \rangle = \frac{1}{2} \langle \varphi, [(\sigma \cdot v)(\sigma \cdot A) + (\sigma \cdot A)(\sigma \cdot v)] \varphi \rangle = \langle \varphi, (v \cdot A) \mathbf{1} \varphi \rangle,$$

because $v \wedge A + A \wedge v = 0$. But $v(x) = g(|x|)x$ by assumption, and A is in the Poincaré gauge. So $v \cdot A = 0$, see (2.13). We thus have

$$\operatorname{Re} \langle (\sigma \cdot v) \varphi, \sigma \cdot (P - A) \varphi \rangle = \operatorname{Re} \langle \varphi, (\sigma \cdot v)(\sigma \cdot P) \varphi \rangle = \frac{1}{2} \langle \varphi, [(\sigma \cdot v)(\sigma \cdot P) + (\sigma \cdot P)(\sigma \cdot v)] \varphi \rangle. \quad (4.9)$$

Now, using (2.3) we get

$$\begin{aligned}
 (\sigma \cdot v)(\sigma \cdot P) + (\sigma \cdot P)(\sigma \cdot v) &= \sum_{j,k=1}^3 (v_j P_k + P_j v_k) \sigma_j \sigma_k = \sum_{j,k=1}^3 (v_j P_k + P_j v_k) \left(\delta_{jk} \mathbf{1} + i \sum_{m=1}^3 \varepsilon_{jkm} \sigma_m \right) \\
 &= (v \cdot P + P \cdot v) \mathbf{1} + i \sum_{j,k,m=1}^3 (v_j P_k + P_j v_k) \varepsilon_{jkm} \sigma_m \\
 &= (v \cdot P + P \cdot v) \mathbf{1} + i \sum_{j,k,m=1}^3 (v_j P_k - P_k v_j) \varepsilon_{jkm} \sigma_m \\
 &= (v \cdot P + P \cdot v) \mathbf{1} + \sum_{j,k,m=1}^3 (\partial_k v_j) \varepsilon_{jkm} \sigma_m = (v \cdot P + P \cdot v) \mathbf{1} = 2D_v \mathbf{1},
 \end{aligned}$$

where we have used the identity

$$\sum_{j,k,m=1}^3 (\partial_k v_j) \varepsilon_{jkm} \sigma_m = (\operatorname{curl} v) \cdot \sigma = (\operatorname{curl} \nabla F) \cdot \sigma = 0.$$

Summing up, we have

$$\operatorname{Re} \langle (\sigma \cdot v) \varphi, \sigma \cdot (P - A) \varphi \rangle = \langle \varphi, D_v \mathbf{1} \varphi \rangle \quad \forall \varphi \in C_0^\infty(\mathbb{R}^3, \mathbb{C}^2). \quad (4.10)$$

Since $v = \nabla F$ is bounded, this identity extends by density to all $\varphi \in \mathcal{D}(P - A)$. \blacksquare

5. Dilations and the commutator

In this section we will define the commutator $[H_{A,V}, D]$ in the sense of quadratic form and derive a matrix-valued version of the weighted virial identities. The latter are our main technical tools in the proof of absence of positive eigenvalues. In some places we make use of technical results obtained in [4].

5.1. Dilations. For $t \in \mathbb{R}$ define the unitary dilation operator $U_t : L^2(\mathbb{R}^3, \mathbb{C}^2) \rightarrow L^2(\mathbb{R}^3, \mathbb{C}^2)$ by

$$(U_t f)(x) = e^{\frac{3t}{2}} f(e^t x) \quad x \in \mathbb{R}^3. \quad (5.1)$$

Then $U_t = e^{itD}$ on $L^2(\mathbb{R}^3, \mathbb{C}^2)$. Let

$$iG_t = \frac{U_t - U_{-t}}{2t} \quad t \in \mathbb{R}. \quad (5.2)$$

It is easily seen that G_t is bounded and symmetric on $L^2(\mathbb{R}^3, \mathbb{C}^2)$. We will use it to approximate the operator D in the limit $t \rightarrow 0$.

As in [4] we define the commutator of H and D by

$$\langle \varphi, i[H_{A,V}, D] \varphi \rangle := \lim_{t \rightarrow 0} \langle \varphi, [H_{A,V}, iG_t] \varphi \rangle := 2 \lim_{t \rightarrow 0} \operatorname{Re} Q_{A,V}(\varphi, iG_t \varphi), \quad (5.3)$$

provided the limit on the right hand side exists. Recall that $\mathcal{D}(Q_{A,V})$ is invariant under dilations, see [4, Prop. 3.3], hence $Q_{A,V}(\varphi, iG_t \varphi)$ is well defined for any $t \neq 0$.

Lemma 5.1. *Let B satisfy Assumption 3.3. Then*

$$\langle \varphi, i[H_{A,0}, D] \varphi \rangle = 2 \lim_{t \rightarrow 0} \operatorname{Re} Q_{A,0}(\varphi, iG_t \varphi) = 2 \|\sigma \cdot (P - A) \varphi\|_2^2 + 2 \operatorname{Re} \langle \sigma \cdot \tilde{B} \varphi, \sigma \cdot (P - A) \varphi \rangle \quad (5.4)$$

for all $\varphi \in \mathcal{D}(P - A)$.

Proof. Let φ be given by (2.10) with $u, v \in \mathcal{H}^1(\mathbb{R}^3)$. A short calculation shows that

$$(P_j - A_j)U_t u = e^t U_t (P_j - A_j) u + X_t^j u \quad \text{with} \quad X_t^j = U_t (e^t A_j - A_j (e^{-t} \cdot))$$

for any $j = 1, 2, 3$. Hence for any $w \in L^2(\mathbb{R}^3)$,

$$\langle w, (P_j - A_j)(U_t - U_{-t})u \rangle = \langle w, (e^t U_t - e^{-t} U_{-t})(P_j - A_j)u \rangle + \langle w, (X_t^j - X_{-t}^j)u \rangle. \quad (5.5)$$

Since

$$\lim_{t \rightarrow 0} t^{-1} X_{\pm t}^j u = \pm \tilde{B}_j u \quad \text{in} \quad L^2(\mathbb{R}^3),$$

see [4, Prop. 3.6], we deduce from (5.2) that

$$2 \operatorname{Re} \lim_{t \rightarrow 0} \operatorname{Re} \langle w, (P_j - A_j) i G_t u \rangle_{L^2(\mathbb{R}^3)} = 2 \operatorname{Re} \langle w, (P_j - A_j) u \rangle + 2 \operatorname{Re} \langle w, \tilde{B}_j u \rangle_{L^2(\mathbb{R}^3)}.$$

After an elementary, but lengthy calculation we then obtain

$$\begin{aligned} \langle \varphi, i [H_{A,0}, D] \varphi \rangle &= 2 \lim_{t \rightarrow 0} \operatorname{Re} \langle \sigma \cdot (P - A) \varphi, \sigma \cdot (P - A) i G_t \varphi \rangle \\ &= 2 \|\sigma \cdot (P - A) \varphi\|_2^2 + 2 \operatorname{Re} \langle \sigma \cdot \tilde{B} \varphi, \sigma \cdot (P - A) \varphi \rangle, \end{aligned}$$

as claimed. ■

Lemma 5.2. *Let B satisfy Assumption 3.3 and let $W : \mathbb{R}^3 \rightarrow \mathbb{R}$ be a potential with form domain containing $\mathcal{D}(P - A)$, such that the distribution $x \cdot \nabla W$ extends to a quadratic form which is form bounded with respect to $(P - A)^2$. Then*

$$2 \lim_{t \rightarrow 0} \langle u, W i G_t v \rangle_{L^2(\mathbb{R}^3)} = -\langle u, x \cdot \nabla W v \rangle_{L^2(\mathbb{R}^3)} \quad (5.6)$$

for all $u, v \in \mathcal{H}^1(\mathbb{R}^3)$.

Proof. By [4, Lemma 3.7, Eq. (3.32)] we have

$$2 \lim_{t \rightarrow 0} \langle u, W i G_t u \rangle_{L^2(\mathbb{R}^3)} = -\langle u, x \cdot \nabla W u \rangle_{L^2(\mathbb{R}^3)}$$

The claim thus follows again from the polarisation identity (2.11). ■

5.2. The commutator. The following result provides a matrix-operator version of a magnetic virial theorem.

Proposition 5.3. *Let B and V satisfy Assumptions 3.3-3.5. Suppose moreover that $x \cdot \nabla V$ is form bounded with respect to $(P - A)^2$. Then for all $\varphi \in \mathcal{D}(\sigma \cdot (P - A))$, the limit $\lim_{t \rightarrow 0} \operatorname{Re} (Q_{A,V}(\varphi, i G_t \varphi))$ in (5.3) exists. Moreover,*

$$\langle \varphi, [H_{A,V}, iD] \varphi \rangle = 2 \|\sigma \cdot (P - A) \varphi\|_2^2 + 2 \operatorname{Re} \langle \sigma \cdot \tilde{B} \varphi, \sigma \cdot (P - A) \varphi \rangle - \langle \varphi, x \cdot \nabla V \varphi \rangle. \quad (5.7)$$

Proof. Let $\varphi \in \mathcal{D}(\sigma \cdot (P - A))$ be given by (2.10). In view of Lemma 5.1 it suffices to show that

$$\langle \varphi, [V, iD] \varphi \rangle = 2 \lim_{t \rightarrow 0} \operatorname{Re} \langle \varphi, V i G_t \varphi \rangle = -\langle \varphi, x \cdot \nabla V \varphi \rangle. \quad (5.8)$$

Let V_{jk} denote the matrix elements of V . By hypothesis of the proposition we have

$$\begin{aligned} & \left| \langle u, x \cdot \nabla V_{11} u \rangle_{L^2(\mathbb{R}^3)} + \langle v, x \cdot \nabla V_{22} v \rangle_{L^2(\mathbb{R}^3)} + \langle u, x \cdot \nabla V_{12} v \rangle_{L^2(\mathbb{R}^3)} + \langle v, x \cdot \nabla V_{12} u \rangle_{L^2(\mathbb{R}^3)} \right| \\ & \lesssim \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + \|(P - A)v\|_{L^2(\mathbb{R}^3)}^2 + \|u\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{L^2(\mathbb{R}^3)}^2, \end{aligned} \quad (5.9)$$

for all $u, v \in \mathcal{H}^1(\mathbb{R}^3)$. Applying the above inequality first with $v = 0$ and then with $u = 0$ shows that $x \cdot \nabla V_{11}$ and $x \cdot \nabla V_{22}$ are relatively form bounded with respect to $(P - A)^2$ in $L^2(\mathbb{R}^3)$. Hence if we return to (5.9) and put $u = v$, then using the triangle inequality we deduce that also the quadratic form

$$\langle u, x \cdot \nabla V_{12} u \rangle_{L^2(\mathbb{R}^3)} + \langle u, x \cdot \nabla V_{21} u \rangle_{L^2(\mathbb{R}^3)} = 2 \operatorname{Re} \langle u, x \cdot \nabla V_{12} u \rangle_{L^2(\mathbb{R}^3)}$$

is relatively bounded with respect to $(P - A)^2$ in $L^2(\mathbb{R}^3)$. Equation (5.8), and hence the claim, thus follows from Lemma 5.2 and (2.11). \blacksquare

Remark 5.4. For rigorous results on virial identities, which have a long history in mathematics and physics, we refer e.g. to [24] and [3].

5.3. Exponentially weighted commutator. The crucial ingredient for the proof of our main result, see Theorem 6.5 below, is finding two different expressions for the weighted commutator $\langle e^F \psi, [H_{A,V}, D] e^F \psi \rangle$, when F is a weight function and ψ is a weak eigenfunction of $H_{A,V}$. This is provided by the following Lemma and by the subsequent equation (5.15).

Lemma 5.5. *Let B and V satisfy Assumptions 3.3-3.5. Assume that $x \cdot \nabla V$ is form bounded with respect to $(P - A)^2$. Let $F \in C^2(\mathbb{R}^3; \mathbb{R})$ be a bounded radial function, such that $\nabla F = xg$, and assume that $g \geq 0$ and that the functions $\nabla(|\nabla F|^2)$, $(1 + |\cdot|^2)g$, $x \cdot \nabla g$ and $(x \cdot \nabla)^2 g$ are bounded. Let $\psi \in \mathcal{D}(P - A)$ be a weak eigenfunction of $H_{A,V}$, i.e., $E\langle \varphi, \psi \rangle = Q_{A,V}(\varphi, \psi)$ for some $E \in \mathbb{R}$ and all $\varphi \in \mathcal{D}(P - A)$. Then*

$$\langle \psi_F, i[H_{A,V}, D] \psi_F \rangle = -4 \|\sqrt{g} D \psi_F\|_2^2 + \langle \psi_F, ((x \cdot \nabla)^2 g - x \cdot \nabla |\nabla F|^2) \psi_F \rangle, \quad (5.10)$$

where $\psi_F = e^F \psi$.

Proof. Note that in the sense of quadratic forms

$$e^F H_{A,V} e^{-F} = H_{A,V} + i[(\sigma \cdot \nabla F) \sigma \cdot (P - A) + (\sigma \cdot (P - A)) \sigma \cdot \nabla F] - |\nabla F|^2. \quad (5.11)$$

Hence

$$\begin{aligned} \langle e^F \psi, [H_{A,V}, iD] e^F \psi \rangle &= 2 \operatorname{Re} \langle H_{A,V} e^F \psi, iD e^F \psi \rangle = 2 \operatorname{Re} \langle e^F H_{A,V} e^{-F} e^F \psi, iD e^F \psi \rangle \\ &\quad - 2 \operatorname{Re} \langle ((\sigma \cdot \nabla F) \sigma \cdot (P - A) + (\sigma \cdot (P - A)) \sigma \cdot \nabla F) \psi_F, D \psi_F \rangle \\ &\quad + 2 \operatorname{Re} \langle |\nabla F|^2 \psi_F, iD \psi_F \rangle \\ &= 2 \operatorname{Re} \langle ((\sigma \cdot \nabla F) \sigma \cdot (P - A) + (\sigma \cdot (P - A)) \sigma \cdot \nabla F) \psi_F, D \psi_F \rangle \\ &\quad - \langle \psi_F, x \cdot \nabla |\nabla F|^2 \psi_F \rangle, \end{aligned} \quad (5.12)$$

where we have used the fact that

$$\operatorname{Re} \langle e^F H_{A,V} e^{-F} e^F \psi, iD e^F \psi \rangle = E \operatorname{Re} \langle \psi_F, iD \psi_F \rangle = 0.$$

Now since F is radial and A is in the Poincaré gauge (2.13), it follows from (2.3) that

$$(\sigma \cdot \nabla F)(\sigma \cdot A) + (\sigma \cdot A)(\sigma \cdot \nabla F) = 2(\nabla F \cdot A) \mathbb{1} = 0.$$

On the other hand, still using (2.3) we obtain

$$\begin{aligned} (\sigma \cdot \nabla F)(\sigma \cdot P) + (\sigma \cdot P)(\sigma \cdot \nabla F) &= (\nabla F \cdot P + P \cdot \nabla F) \mathbb{1} + i \sum_{j,k,m=1}^3 (\partial_j F P_k + P_j \partial_k F) \varepsilon_{jkm} \sigma_m \\ &= (gx \cdot P + P \cdot xg) \mathbb{1} + i \sum_{j,k,m=1}^3 (\partial_j F P_k - P_k \partial_j F) \varepsilon_{jkm} \sigma_m \\ &= (gD - ix \cdot \nabla g) \mathbb{1} + \sum_{m=1}^3 \left(\sum_{j,k=1}^3 \partial_k \partial_j F \varepsilon_{jkm} \right) \sigma_m. \end{aligned}$$

Since $\sum_{j,k=1}^3 \partial_k \partial_j F \varepsilon_{jkm} = 0$ for all $m = 1, 2, 3$, the last equation in combination with (5.12) gives

$$\begin{aligned} \langle \psi_F, [H_{A,V}, iD] \psi_F \rangle &= -\operatorname{Re} \langle (gD - ix \cdot \nabla g) \psi_F, D \psi_F \rangle - \langle \psi_F, x \cdot \nabla |\nabla F|^2 \psi_F \rangle \\ &= -4 \|\sqrt{g} D \psi_F\|_2^2 + \langle \psi_F, ((x \cdot \nabla)^2 g - x \cdot \nabla |\nabla F|^2) \psi_F \rangle. \end{aligned}$$

■

In view of the fact that $D_{\nabla F}$ is symmetric equation (5.11) and Lemma 4.5 imply

$$Q_{A,V}(\varphi, \varphi) = Q_{A,V}(e^{-F}\varphi, e^F\varphi) + \langle \nabla F \varphi, \nabla F \varphi \rangle \quad (5.13)$$

for all $\varphi \in \mathcal{D}(P-A)$. By inserting $\varphi = \psi_F$ in the above equation, which is allowed because $\psi_F \in \mathcal{D}(P-A)$, we get

$$Q_{A,V}(\psi_F, \psi_F) = \|\sigma \cdot (P-A)\psi_F\|_2^2 + \langle \psi_F, V\psi_F \rangle = \langle \psi_F, (E + |\nabla F|^2)\psi_F \rangle. \quad (5.14)$$

A combination with (5.7) thus gives

$$\begin{aligned} \langle \psi_F, i[H_{A,V}, D]\psi_F \rangle &= \langle \psi_F, (E + |\nabla F|^2)\psi_F \rangle + \|\sigma \cdot (P-A)\psi_F\|_2^2 + 2\operatorname{Re} \langle \sigma \cdot \tilde{B}\psi_F, \sigma \cdot (P-A)\psi_F \rangle \\ &\quad - \langle \psi_F, (V + x \cdot \nabla V)\psi_F \rangle. \end{aligned} \quad (5.15)$$

Lemma 5.6. *Let B and V satisfy Assumptions 3.3, 3.5, and 3.10. Assume that ψ and F satisfy conditions of Lemma 5.5. Then there exist constants $\kappa > 0$ and $c_\kappa > 0$ such that*

$$\langle \psi_F, [H_{A,V}, iD]\psi_F \rangle \geq \kappa \langle \psi_F, |\nabla F|^2 \psi_F \rangle - c_\kappa \|\psi_F\|_2^2. \quad (5.16)$$

Proof. Below we denote by c a generic constant whose value might change from line to line. By Proposition 5.3, the Cauchy-Schwarz inequality and Corollary 4.3,

$$\begin{aligned} \langle \psi_F, [H_{A,V}, iD]\psi_F \rangle &\geq \|\sigma \cdot (P-A)\psi_F\|_2^2 - 2\|\sigma \cdot (P-A)\psi_F\|_2 (\|\tilde{B}\psi_F\|_2 + \|xV^s\psi_F\|_2) \\ &\quad - (\alpha_2 + 3\alpha_3)\|\sigma \cdot (P-A)\psi_F\|_2^2 - c\|\psi_F\|_2^2. \end{aligned}$$

Now let $\kappa > 0$ and split

$$\|\sigma \cdot (P-A)\psi_F\|_2^2 = (1-\kappa)\|\sigma \cdot (P-A)\psi_F\|_2^2 + \kappa\|\sigma \cdot (P-A)\psi_F\|_2^2.$$

Using equation (5.15) together with Corollary 4.3 we find

$$\begin{aligned} \langle \psi_F, [H_{A,V}, iD]\psi_F \rangle &\geq (2-\kappa)\|\sigma \cdot (P-A)\psi_F\|_2^2 + \kappa \langle \psi_F, |\nabla F|^2 \psi_F \rangle - (\alpha_2 + d\alpha_3 + \kappa\alpha_0)\|\sigma \cdot (P-A)\psi_F\|_2^2 \\ &\quad - 2\|\sigma \cdot (P-A)\psi_F\|_2 (\|\tilde{B}\psi_F\|_2 + \|xV^s\psi_F\|_2) - c\|\psi_F\|_2^2, \end{aligned}$$

and

$$\begin{aligned} 2\|\sigma \cdot (P-A)\psi_F\|_2 (\|\tilde{B}\psi_F\|_2 + \|xV^s\psi_F\|_2) &\leq \alpha_1\|\sigma \cdot (P-A)\psi_F\|_2^2 + 2C_1\|(P-A)\psi_F\|_2 \|\psi_F\|_2 \\ &\leq (\alpha_1 + \kappa)\|\sigma \cdot (P-A)\psi_F\|_2^2 + \frac{C_1}{\kappa}\|\psi_F\|_2^2. \end{aligned}$$

Hence

$$\langle \psi_F, [H_{A,V}, iD]\psi_F \rangle \geq (1-2\kappa - \kappa\alpha_0 - \alpha_1 - \alpha_2 - 3\alpha_3)\|\sigma \cdot (P-A)\psi_F\|_2^2 + \kappa \langle \psi_F, |\nabla F|^2 \psi_F \rangle - c_\kappa \|\psi_F\|_2^2.$$

If we now set $\kappa = (2 + \alpha_0)^{-1}(1 - \alpha_1 - \alpha_2 - 3\alpha_3)$, then $\kappa > 0$, see (3.14), and the claim follows. ■

5.4. The virial. Below we provide a matrix version of the Kato form of the virial.

Lemma 5.7. *Let B and V satisfy Assumptions 3.3-3.5. Suppose that V and $|x|^2V^2$ are relatively form bounded with respect to $(P-A)^2$. Then*

$$\langle \varphi, x \cdot \nabla V \varphi \rangle = 2\operatorname{Im} \langle xV\varphi, (P-A)\varphi \rangle - 3\langle \varphi, V\varphi \rangle \quad (5.17)$$

for all $\varphi \in \mathcal{D}(P-A)$.

Proof. Let V_{jk} be the matrix elements of V and let $W : \mathbb{R}^3 \rightarrow \mathbb{R}$. By [4, Lemma 3.12],

$$\langle u, x \cdot \nabla W u \rangle_{L^2(\mathbb{R}^3)} = 2 \operatorname{Im} \langle u, x W (P - A)u \rangle_{L^2(\mathbb{R}^3)} - 3 \langle u, W u \rangle_{L^2(\mathbb{R}^3)} \quad (5.18)$$

holds for all $u \in \mathcal{H}^1(\mathbb{R}^3)$ provided W and $|x|^2 W^2$ are relatively form bounded with respect to $(P - A)^2$ in $L^2(\mathbb{R}^3)$. To prove the statement of the lemma we have to verify that equation (5.18) can be applied with $W = V_{11}$, $W = V_{22}$ and $W = V_{12}$, cf. (3.1). Reasoning in the same way as in the proof of Proposition 5.3 we verify that V_{11} , V_{22} and V_{12} are relatively form bounded with respect to $(P - A)^2$ in $L^2(\mathbb{R}^3)$. In order to verify the relative form boundedness of $|x|^2 V_{11}^2$, $|x|^2 V_{22}^2$ and $|x|^2 |V_{12}|^2$ we note that since

$$V^2 = \begin{pmatrix} V_{11}^2 + |V_{12}|^2 & V_{12}(V_{11} + V_{22}) \\ V_{21}(V_{11} + V_{22}) & V_{22}^2 + |V_{12}|^2 \end{pmatrix},$$

the assumptions of the lemma imply that

$$\begin{aligned} & \langle u, |x|^2 (V_{11}^2 + |V_{12}|^2) u \rangle_{L^2(\mathbb{R}^3)} + \langle v, |x|^2 (V_{22}^2 + |V_{12}|^2) v \rangle_{L^2(\mathbb{R}^3)} + \langle u, |x|^2 V_{12} (V_{11} + V_{22}) v \rangle_{L^2(\mathbb{R}^3)} \\ & + \langle v, |x|^2 V_{21} (V_{11} + V_{22}) u \rangle_{L^2(\mathbb{R}^3)} \lesssim \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + \|(P - A)v\|_{L^2(\mathbb{R}^3)}^2 + \|u\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{L^2(\mathbb{R}^3)}^2, \end{aligned} \quad (5.19)$$

for all $u, v \in \mathcal{H}^1(\mathbb{R}^3)$. As in the proof of Proposition 5.3 we apply (5.19) with $v = 0$ and $u = 0$ respectively, and deduce that $|x|^2 (V_{11}^2 + |V_{12}|^2)$ and $|x|^2 (V_{22}^2 + |V_{12}|^2)$ are relatively form bounded with respect to $(P - A)^2$ in $L^2(\mathbb{R}^3)$. Hence (5.18) holds for $W = V_{jk}$ with any $j, k = 1, 2$. In view of (2.11), this proves equation (5.17). \blacksquare

Corollary 5.8. *Let B and V satisfy Assumptions 3.3-3.5. Assume moreover, that the potential V splits as $V = V^s + V^\ell$ where V^s and $|x|^2 (V^s)^2$ are relatively form bounded with respect to $(P - A)^2$ and the distribution $x \cdot \nabla V^\ell$ extend to a quadratic form which is form bounded with respect to $(P - A)^2$. Then*

$$\langle \varphi, x \cdot \nabla V \varphi \rangle = 2 \operatorname{Im} \langle x V^s \varphi, (P - A) \varphi \rangle - 3 \langle \varphi, V^s \varphi \rangle + \langle \varphi, x \cdot \nabla V^\ell \varphi \rangle \quad (5.20)$$

for all $\varphi \in \mathcal{D}(P - A)$.

Proof. The claim follows from Lemmas 5.2 and 5.7. \blacksquare

6. The main result

Once we have established the virial identities (5.10) and (5.15), we can follow the strategy of [14, 15, 4]. This is done in two steps. First we show that if eigenfunctions corresponding to energies larger than Λ exist, then they decay faster than exponentially. Second, we prove that such eigenfunctions have to vanish identically.

6.1. Super-exponential decay. Given $x \in \mathbb{R}^3$, $\lambda > 0$, we set

$$\langle x \rangle_\lambda := \sqrt{\lambda + |x|^2}.$$

If $\lambda = 1$, we omit the subscript and write $\langle x \rangle_1 = \langle x \rangle$.

We have

Proposition 6.1. *Assume that B and V satisfy Assumptions 3.1-3.9 and that the magnetic field A corresponding to B is in the Poincaré gauge. Furthermore, assume that ψ is a weak eigenfunction of the magnetic Schrödinger operator $H_{A,V}$ corresponding to the energy $E \in \mathbb{R}$, and that there exist $\bar{\mu} \geq 0$ and $\lambda > 0$ such that $x \mapsto e^{\bar{\mu} \langle x \rangle_\lambda} \psi(x) \in L^2(\mathbb{R}^3, \mathbb{C}^2)$. If $E + \bar{\mu}^2 > \Lambda$ with Λ given by (1.2), then*

$$x \mapsto e^{\mu \langle x \rangle_\lambda} \psi(x) \in L^2(\mathbb{R}^3, \mathbb{C}^2) \quad \forall \mu > 0, \quad \forall \lambda > 0. \quad (6.1)$$

The proof of Proposition 6.1 requires some preliminaries. Obviously it suffices to prove the statement for $\lambda = 1$. First we consider the case $\bar{\mu} = 0$ and choose

$$F_{\mu,\varepsilon}(x) = \frac{\mu}{\varepsilon} \left(1 - e^{-\varepsilon\langle x \rangle}\right), \quad (6.2)$$

for some $\mu \geq 0$ and $\varepsilon > 0$. We have $F_{\mu,\varepsilon}(x) \rightarrow \mu\langle x \rangle$ as $\varepsilon \rightarrow 0$. Moreover, the identity

$$\nabla F_{\mu,\varepsilon} = \mu\langle x \rangle^{-1} e^{-\varepsilon\langle x \rangle} x \quad (6.3)$$

implies

$$g_{\mu,\varepsilon}(x) = \mu\langle x \rangle^{-1} e^{-\varepsilon\langle x \rangle}. \quad (6.4)$$

Let

$$\mu_* = \sup \left\{ \mu \geq 0 : e^{\mu\langle x \rangle} \psi \in L^2(\mathbb{R}^3, \mathbb{C}^2) \right\},$$

be the maximal exponential decay rate of ψ . To prove (6.1) we have to show that $\mu_* = \infty$. We will argue by contradiction.

Lemma 6.2. *Suppose that $0 \leq \mu_* < \infty$. Then there exist decreasing sequences μ_n and ε_n such that $\mu_n \rightarrow \mu_*$ and $\varepsilon_n \rightarrow 0$, as $n \rightarrow \infty$, and such that, writing $F_n := F_{\mu_n, \varepsilon_n}$, we have*

$$a_n := \|e^{F_n} \psi\|_2 \rightarrow \infty \quad \text{as } n \rightarrow \infty, \quad (6.5)$$

Proof. For a fixed x and μ we have

$$\partial_\varepsilon F_{\mu,\varepsilon}(x) = -\frac{\mu}{\varepsilon^2} \left(1 - (1 + \varepsilon\langle x \rangle) e^{-\varepsilon\langle x \rangle}\right). \quad (6.6)$$

On the other hand, a short calculation shows that the function $t \mapsto (1+t)e^{-t}$ is strictly decreasing on $(0, \infty)$. It follows that

$$\partial_\varepsilon F_{\mu,\varepsilon}(x) < 0 \quad \forall \varepsilon > 0, \quad \mu > 0, \quad x \in \mathbb{R}^3.$$

Thus $F_{\mu,\varepsilon}(x)$ is strictly decreasing in ε for any $\mu > 0$ and

$$F_{\mu,\varepsilon}(x) \nearrow \mu\langle x \rangle \quad \text{as } \varepsilon \searrow 0.$$

By setting $\mu_n = \mu_* + \frac{1}{n}$, we then have

$$\lim_{\varepsilon \searrow 0} \|e^{F_{\mu_n, \varepsilon}} \psi\|_2 = \|e^{\mu_n\langle x \rangle} \psi\|_2 = +\infty \quad \forall n \geq 1, \quad (6.7)$$

by monotone convergence. Now we construct the sequence ε_n as follows. Take ε_1 such that $\|e^{F_{\mu_1, \varepsilon_1}} \psi\|_2 > 1$, and for each $n \geq 2$ we choose $\varepsilon_n < \varepsilon_{n-1}$ so that

$$\|e^{F_{\mu_n, \varepsilon_n}} \psi\|_2 > n,$$

which is possible in view of (6.7). This proves the claim. ■

Now let $g_n(x) := g_{\mu_n, \varepsilon_n}$, and define

$$\varphi_n = \frac{e^{F_n} \psi}{\|e^{F_n} \psi\|_2}. \quad (6.8)$$

Since $\mu_n \rightarrow \mu_*$, and since

$$F_n(x) \leq \mu_n\langle x \rangle, \quad (6.9)$$

for any compact set $\omega \subset \mathbb{R}^3$ it holds

$$\langle \varphi_n, \chi_\omega \varphi_n \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (6.10)$$

where χ denotes the characteristic functions. Hence if W is bounded and $W(x) \rightarrow 0$ as $x \rightarrow \infty$, then

$$\langle \varphi_n, W \varphi_n \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (6.11)$$

We will also need the following auxiliary Lemmas.

Lemma 6.3. *Let F_n , g_n , ψ , and φ_n be given as above. If $0 < \mu_* < \infty$, then*

$$\lim_{n \rightarrow \infty} \langle e^{F_n} \psi, \varepsilon_n \langle x \rangle e^{F_n} \psi \rangle = 0. \quad (6.12)$$

Moreover, if $0 \leq \mu_* < \infty$, then

$$\lim_{n \rightarrow \infty} \langle \nabla F_n \varphi_n, \nabla F_n \varphi_n \rangle = \mu_*^2 \quad (6.13)$$

and

$$\lim_{n \rightarrow \infty} \langle \varphi_n, ((x \cdot \nabla)^2 g_n - x \cdot \nabla |\nabla F_n|^2) \varphi_n \rangle = 0 \quad (6.14)$$

Proof. Let $\delta > 0$. Since $\|\varphi_n\|_2 = 1$, it follows that

$$\langle \varphi_n, \varepsilon_n \langle x \rangle \varphi_n \rangle \leq \delta + \langle \varphi_n, \mathbf{1}_{\{\varepsilon_n \langle x \rangle > \delta\}} \varepsilon_n \langle x \rangle \varphi_n \rangle. \quad (6.15)$$

Next we note that the mapping $t \mapsto \frac{1-e^{-t}}{t}$ is decreasing on $(0, \infty)$. Hence

$$\bar{\gamma}_\delta := \sup_{t \geq \delta} \frac{1 - e^{-t}}{t} < 1. \quad (6.16)$$

This shows that for any x such that $\varepsilon_n \langle x \rangle \geq \delta$ we have

$$F_n = \frac{\mu_n \langle x \rangle}{\varepsilon_n \langle x \rangle} (1 - e^{-\varepsilon_n \langle x \rangle}) \leq \mu_n \bar{\gamma}_\delta \langle x \rangle.$$

Let κ be such that $\bar{\gamma}_\delta < \kappa < 1$. If $0 < \mu_* < \infty$ then, by the definition of μ_* , ψ decays exponentially with any rate μ satisfying $\kappa \mu_* < \mu < \mu_*$. Since $\mu_n \bar{\gamma}_\delta \rightarrow \bar{\gamma}_\delta \mu_* < \kappa \mu_*$ as $n \rightarrow \infty$, this implies

$$\limsup_{n \rightarrow \infty} \langle e^{F_n} \psi, \mathbf{1}_{\{\varepsilon_n \langle x \rangle > \delta\}} \langle x \rangle e^{F_n} \psi \rangle \leq \limsup_{n \rightarrow \infty} \langle e^{\mu_n \bar{\gamma}_\delta \langle x \rangle} \psi, \langle x \rangle e^{\mu_n \bar{\gamma}_\delta \langle x \rangle} \psi \rangle \leq \langle e^{\kappa \mu_* \langle x \rangle} \psi, \langle x \rangle e^{\kappa \mu_* \langle x \rangle} \psi \rangle < \infty.$$

Equation (6.12) thus follows from (6.5) and (6.15).

To prove the remaining claims of the Lemma we need the identity

$$|\nabla F_n|^2 = \mu_n^2 (1 - \langle x \rangle^{-2}) e^{-2\varepsilon_n \langle x \rangle}, \quad (6.17)$$

which follows by a direct calculation from equation (6.3). Hence

$$\mu_n^2 - \langle \nabla F_n \varphi_n, \nabla F_n \varphi_n \rangle = \mu_n^2 \left(\langle \varphi_n, (1 - e^{-2\varepsilon_n \langle x \rangle}) \varphi_n \rangle + \langle \varphi_n, \langle x \rangle^{-2} e^{-2\varepsilon_n \langle x \rangle} \varphi_n \rangle \right), \quad (6.18)$$

where we have used again the fact that $\|\varphi_n\|_2 = 1$. If $\mu_* = \lim_{n \rightarrow \infty} \mu_n = 0$, then (6.18) shows

$$|\mu_n^2 - \langle \nabla F_n \varphi_n, \nabla F_n \varphi_n \rangle| \leq 2\mu_n^2 \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

and hence (6.13) with $\mu_* = 0$. If $\mu_* > 0$, then we insert the bound $0 \leq 1 - e^{-2\varepsilon_n \langle x \rangle} \leq 2\varepsilon_n \langle x \rangle$ into (6.18) and get

$$|\mu_n^2 - \langle \nabla F_n \varphi_n, \nabla F_n \varphi_n \rangle| \leq \mu_n^2 (2\langle \varphi_n, \varepsilon_n \langle x \rangle \varphi_n \rangle + \langle \varphi_n, \langle x \rangle^{-2} \varphi_n \rangle) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

In view of (6.12) and (6.11) this proves (6.13) in the case $\mu_* > 0$.

It remains to prove (6.14). From the definitions of F_n and g_n we deduce, after a short calculation, that

$$|(x \cdot \nabla)^2 g_n - x \cdot \nabla |\nabla F|^2| \lesssim \mu_n (\mu_n + 1) [\langle x \rangle^{-2} + \langle x \rangle^{-1} + \varepsilon_n \langle x \rangle + \varepsilon_n^2 \langle x \rangle] e^{-\varepsilon_n \langle x \rangle} \quad (6.19)$$

This and that boundedness of mapping $t \mapsto te^{-t}$ on $[0, +\infty)$ implies that if $\mu_* = 0$, then

$$|\langle \varphi_n, ((x \cdot \nabla)^2 g_n - x \cdot \nabla |\nabla F|^2) \varphi_n \rangle| \lesssim \mu_n (\mu_n + 1) \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

If $0 < \mu_* < \infty$, then we use (6.19) to estimate

$$|\langle \varphi_n, ((x \cdot \nabla)^2 g_n - x \cdot \nabla |\nabla F|^2) \varphi_n \rangle| \lesssim \langle \varphi_n, (\langle x \rangle^{-2} + \langle x \rangle^{-1}) \varphi_n \rangle + \langle \varphi_n, \varepsilon_n \langle x \rangle \varphi_n \rangle \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Here we have used again equations (6.12) and (6.11). This completes the proof of (6.14) and hence of the Lemma. \blacksquare

Lemma 6.4. *Let $0 \leq \mu_* < \infty$ and F_n , g_n , and φ_n be given as above. If V satisfies Assumptions 3.5 and 3.7, then*

$$\limsup_{n \rightarrow \infty} \langle \varphi_n, V \varphi_n \rangle =: \nu \leq 0 \quad (6.20)$$

$$\liminf_{n \rightarrow \infty} \langle \sigma \cdot (P - A) \varphi_n, \sigma \cdot (P - A) \varphi_n \rangle \geq E + \mu_*^2 - \nu. \quad (6.21)$$

Moreover, if the magnetic field B satisfy Assumptions 3.4 and 3.9, then

$$\limsup_{n \rightarrow \infty} |\langle \sigma \cdot \tilde{B} \varphi_n, \sigma \cdot (P - A) \varphi_n \rangle| \leq \beta(E + \mu_*^2 - \nu)^{1/2}. \quad (6.22)$$

Finally, if one splits $V = V^s + V^\ell$, with V^s and V^ℓ satisfying Assumptions 3.9 and 3.10, then

$$\limsup_{n \rightarrow \infty} \langle \varphi_n, x \cdot \nabla V \varphi_n \rangle \leq 2\omega_1(E + \mu_*^2 - \nu)^{1/2} + \omega_2. \quad (6.23)$$

Proof. First we prove that

$$\limsup_{n \rightarrow \infty} |\langle \varphi_n, V \varphi_n \rangle| < \infty. \quad (6.24)$$

Indeed, by Lemma 4.2 and equation (3.4)

$$|\langle \varphi_n, V \varphi_n \rangle| \leq \alpha_0(1 + \eta) \|\sigma \cdot (P - A) \varphi_n\|_2^2 + C$$

for any $\eta > 0$ and some $C > 0$ independent of n . Using (5.14) with $F = F_n$ we then further get

$$|\langle \varphi_n, V \varphi_n \rangle| \leq \alpha_0(1 + \eta) \langle \varphi_n, (E + |\nabla F_n|^2) \varphi_n \rangle + \alpha_0(1 + \eta) |\langle \varphi_n, V \varphi_n \rangle| + C,$$

and (6.24) follow by choosing η small enough so that $\alpha_0(1 + \eta) < 1$ and letting $n \rightarrow \infty$, see (6.13).

To prove (6.20) we let $j_m : [0, \infty) \rightarrow \mathbb{R}_+$, $m = 1, 2$, be infinitely often differentiable on $(0, \infty)$ with $j_1(r) = 1$ for $0 \leq r \leq 1$, $j_1(r) > 0$ for $r \leq 3/2$, $j_1(r) = 0$ for $r \geq 7/4$, and $j_2(r) = 0$ for $r \leq 5/4$, $j_2(r) > 0$ for $r \geq 3/2$, $j_2(r) = 1$ for $r \geq 2$. Then $\inf_{r \geq 0} (j_1^2(r) + j_2^2(r)) > 0$ and thus

$$\xi_1 := \frac{j_1}{\sqrt{j_1^2 + j_2^2}}, \quad \xi_2 := \frac{j_2}{\sqrt{j_1^2 + j_2^2}}$$

are infinitely often differentiable with bounded derivatives and $\xi_1^2 + \xi_2^2 = 1$. Given $R \geq 1$ we set

$$\xi_{R-}(x) := \xi_1(|x|/R), \quad \xi_{R+}(x) := \xi_2(|x|/R).$$

Note that $\xi_{R+}, \xi_{R-} \in C^\infty(\mathbb{R}^3)$ that the all partial derivatives of ξ_{R+} and ξ_{R-} . Moreover, ξ_{R-} has compact support, and $\text{supp}(\xi_{R+}) \subset \mathcal{U}_R^c = \{x \in \mathbb{R}^3 : |x| \geq R\}$. By construction,

$$\langle \varphi_n, V \varphi_n \rangle_+ = \langle \xi_{R-}^2 \varphi_n, V \varphi_n \rangle_+ + \langle \xi_{R+}^2 \varphi_n, V \varphi_n \rangle_+.$$

From [4, Lemma 4.6] it follows that

$$\sup_{R \geq 1} \sup_{n \in \mathbb{N}} \|(P - A) \xi_{R+} \varphi_n\| < \infty, \quad \text{and} \quad \forall R \geq 1 : \limsup_{n \rightarrow \infty} \|(P - A) \xi_{R-} \varphi_n\| = 0.$$

Hence a combination of Lemma 4.1, Assumption 3.4 and equation (6.11) applied with $W = \xi_{R-}$ gives

$$\sup_{R \geq 1} \sup_{n \in \mathbb{N}} \|\sigma \cdot (P - A) \xi_{R+} \varphi_n\| < \infty, \quad \text{and} \quad \forall R \geq 1 : \limsup_{n \rightarrow \infty} \|\sigma \cdot (P - A) \xi_{R-} \varphi_n\| = 0. \quad (6.25)$$

Let us now treat the terms containing V . Since V is form bounded with respect to $(P - A)^2$, it follows from Lemma 4.2 and equations (6.11), (6.25) that for a fixed $R \geq 1$ we have

$$\langle \xi_{R-}^2 \varphi_n, V \varphi \rangle_+ = \langle \xi_{R-} \varphi_n, V \xi_{R-} \varphi_n \rangle_+ \lesssim \|\sigma \cdot (P - A) \xi_{R-} \varphi_n\|_2^2 + \|\xi_{R-} \varphi_n\|_2^2 \rightarrow 0, \text{ as } n \rightarrow \infty$$

Moreover, since V_+ vanishes at infinity w.r.t. $(P - A)^2$, there exist sequences α_R, γ_R with $\alpha_R, \gamma_R \rightarrow 0$ as $R \rightarrow \infty$ such that

$$\langle \xi_{R+}^2 \varphi_n, V \varphi_n \rangle_+ = \langle \xi_{R+} \varphi_n, V \xi_{R+} \varphi_n \rangle_+ \leq \alpha_R \|\sigma \cdot (P - A) \xi_{R+} \varphi_n\|_2^2 + \gamma_R \|\xi_{R+} \varphi_n\|_2^2.$$

Equation (6.25) then shows that

$$\limsup_{n \rightarrow \infty} \langle \xi_{R+}^2 \varphi_n, V \varphi_n \rangle_+ \lesssim \alpha_R + \gamma_R \rightarrow 0, \text{ as } R \rightarrow \infty,$$

which proves (6.20). Next, from (5.14), (6.13) and (6.20) we obtain

$$\liminf_{n \rightarrow \infty} \langle \sigma \cdot (P - A) \varphi_n, \sigma \cdot (P - A) \varphi_n \rangle = \liminf_{n \rightarrow \infty} (E + \langle \nabla F_n \varphi_n, \nabla F_n \varphi_n \rangle - \langle \varphi_n, V \varphi_n \rangle) \geq E + \mu_*^2 - \nu.$$

Hence (6.21). To treat the term with $|\tilde{B}|$ we argue in the same way as for V and conclude that for any fixed R ,

$$\limsup_{n \rightarrow \infty} \langle \varphi_n, |\tilde{B}|^2 \varphi_n \rangle \leq \limsup_{n \rightarrow \infty} \langle \xi_{R+}^2 \varphi_n |\tilde{B}|^2, \varphi_n \rangle \lesssim \varepsilon_R + \beta_R^2,$$

where we used Assumption 3.9 and equation (6.25). Since $\varepsilon_R \rightarrow 0$ and $\beta_R \rightarrow \beta$, as $R \rightarrow \infty$, with the help of (4.2) we get

$$\limsup_{n \rightarrow \infty} \|\sigma \cdot \tilde{B} \varphi_n\|_2 \leq \beta.$$

Moreover, equations (6.13) and (5.14) imply

$$\limsup_{n \rightarrow \infty} \|\sigma \cdot (P - A) \varphi_n\|_2 \leq \sqrt{E + \mu_*^2 - \nu}. \quad (6.26)$$

Hence

$$|\langle \sigma \cdot \tilde{B} \varphi_n, \sigma \cdot (P - A) \varphi_n \rangle| \leq \|\sigma \cdot \tilde{B} \varphi_n\|_2 \|\sigma \cdot (P - A) \varphi_n\|_2 \leq \beta \sqrt{E + \mu_*^2 - \nu},$$

which proves (6.22). If the potential splits as $V = V^s + V^\ell$ with V^s, V^ℓ satisfying Assumptions 3.10 and 3.9, then one can argue exactly as above to conclude with

$$\limsup_{n \rightarrow \infty} |\langle x V^s \varphi_n, \sigma \cdot (P - A) \varphi_n \rangle| \leq \omega_1 \sqrt{E + \mu_*^2 - \nu} \quad \text{and} \quad \limsup_{n \rightarrow \infty} |\langle \varphi_n, x \cdot \nabla V^\ell \varphi_n \rangle| \leq \omega_2.$$

Moreover, if V^s and V^ℓ satisfying Assumptions 3.10 and 3.9, and $\varphi \in \mathcal{D}(P - A)$ with $\text{supp}(\varphi) \subset \{|x| \geq R\}$, then using Lemma 4.2 we get

$$|\langle \varphi, V^s \varphi \rangle| = |\langle |x|^{-1} \varphi, |x| V^s \varphi \rangle| \leq \| |x|^{-1} \varphi \|_2 \| |x| V^s \varphi \|_2 \lesssim R^{-1} \|\varphi\|_2 (\|\sigma \cdot (P - A) \varphi\|_2^2 + \|\varphi\|_2^2)^{1/2}.$$

Thus $\lim_{n \rightarrow \infty} \langle \varphi_n, V^s \varphi_n \rangle = 0$, and Corollary 5.8 gives

$$\limsup_{n \rightarrow \infty} \langle \varphi, x \cdot \nabla V \varphi \rangle \leq 2\omega_1 (E + \mu_*^2 - \nu)^{1/2} + \omega_2. \quad \blacksquare$$

Proof of Proposition 6.1. Assume that $0 \leq \mu_*^2 < \infty$. One easily verifies that F_n and g_n satisfy the assumptions of Lemma 5.5. The latter in combination with equation (6.14) shows that

$$\limsup_{n \rightarrow \infty} \langle \varphi_n, [H, iD] \varphi_n \rangle \leq 0. \quad (6.27)$$

On the other hand, equation (5.7) and Lemma 6.4 imply the lower bound

$$\begin{aligned} \liminf_{n \rightarrow \infty} \langle \varphi_n, [H, iD] \varphi_n \rangle &\geq 2(E + \mu_*^2 - \nu) - 2(\beta + \omega_1)(E + \mu_*^2 - \nu)^{1/2} - \omega_2 \\ &= 2 \left[\left(\sqrt{E + \mu_*^2 - \nu} - \frac{\beta + \omega_1}{2} \right)^2 - \left(\frac{\beta + \omega_1}{2} \right)^2 - \frac{\omega_2}{2} \right]. \end{aligned}$$

Hence if

$$\sqrt{E + \mu_*^2 - \nu} > \frac{1}{2}(\beta + \omega_1 + \sqrt{(\beta + \omega_1)^2 + 2\omega_2}) = \sqrt{\Lambda},$$

then

$$\liminf_{n \rightarrow \infty} \langle \varphi_n, i[H, D] \varphi_n \rangle > 0,$$

which contradicts (6.27). Thus $\mu_* = \infty$ and (6.1) follows. \blacksquare

6.2. Absence of embedded eigenvalues. We are now in position to prove our main result.

Theorem 6.5. *Let B and V satisfy Assumptions 3.1-3.10. Then the Pauli operator $H_{A,V}$ has no eigenvalues in the interval (Λ, ∞) , where Λ is given by (1.2).*

Proof. Assume that $E\langle \varphi, \psi \rangle = Q_{A,V}(\varphi, \psi)$ holds for all $\varphi \in \mathcal{D}(Q_{A,V}) = \mathcal{D}(P - A)$, and that $E > \Lambda$. From Proposition 6.1 we then deduce that

$$x \mapsto e^{\mu\langle x \rangle_\lambda} \psi(x) \in L^2(\mathbb{R}^3, \mathbb{C}^2) \quad \forall \mu > 0, \quad \forall \lambda > 0,$$

where $\langle x \rangle_\lambda = (\lambda + |x|^2)^{1/2}$. Let $\mu > 0, \varepsilon > 0, \lambda > 0$, and define

$$F(x) = F_{\mu,\varepsilon,\lambda}(x) = \frac{\mu}{\varepsilon} \left(1 - e^{-\varepsilon\langle x \rangle_\lambda} \right).$$

Then

$$\nabla F_{\mu,\varepsilon,\lambda}(x) = x g_{\mu,\varepsilon,\lambda}(x), \quad \text{with} \quad g_{\mu,\varepsilon,\lambda}(x) = \frac{\mu e^{-\varepsilon\langle x \rangle_\lambda}}{\sqrt{\lambda + |x|^2}}.$$

Let $\psi_{\mu,\varepsilon,\lambda} = e^{F_{\mu,\varepsilon,\lambda}} \psi$. Lemma 5.6 and equation (5.10) give

$$\kappa \langle \psi_{\mu,\varepsilon,\lambda}, |\nabla F_{\mu,\varepsilon,\lambda}|^2 \psi_{\mu,\varepsilon,\lambda} \rangle \leq \langle \psi_{\mu,\varepsilon,\lambda}, ((x \cdot \nabla)^2 g_{\mu,\varepsilon,\lambda} - x \cdot \nabla |\nabla F_{\mu,\varepsilon,\lambda}|^2) \psi_{\mu,\varepsilon,\lambda} \rangle + C \|\psi_{\mu,\varepsilon,\lambda}\|_2^2 \quad (6.28)$$

for all $\mu, \varepsilon, \lambda > 0$ and some constant C independent of μ, λ and ε . Now a direct calculation shows that

$$\lim_{\varepsilon \rightarrow 0} x \cdot \nabla |\nabla F_{\mu,\varepsilon,\lambda}(x)|^2 = 2\lambda\mu^2 \langle x \rangle_\lambda^{-1} (1 - \langle x \rangle_\lambda^{-2}) > 0, \quad (6.29)$$

and

$$\lim_{\varepsilon \rightarrow 0} (x \cdot \nabla)^2 g_{\mu,\varepsilon,\lambda}(x) = -2\lambda\mu \langle x \rangle_\lambda^{-3} |x|^2 < 0. \quad (6.30)$$

Since

$$\lim_{\varepsilon \rightarrow 0} F_{\mu,\varepsilon,\lambda}(x) := F_{\mu,\lambda}(x) = \mu \langle x \rangle_\lambda,$$

in view of Proposition 6.1 we can pass to limit $\varepsilon \rightarrow 0$ in (6.28) to obtain

$$\kappa \mu^2 \langle \psi_{\mu,\lambda}, \frac{|x|^2}{\lambda + |x|^2} \psi_{\mu,\lambda} \rangle \leq C \|\psi_{\mu,\lambda}\|_2^2 \quad \forall \mu, \lambda > 0, \quad (6.31)$$

where

$$\psi_{\mu,\lambda}(x) := e^{\mu\langle x \rangle_\lambda} \psi(x) \in L^2(\mathbb{R}^3, \mathbb{C}^2).$$

Using once again Proposition 6.1 together with the monotone convergence theorem we arrive, by letting $\lambda \rightarrow 0$, at the inequality

$$\kappa \mu^2 \|\psi_\mu\|_2^2 \leq C \|\psi_\mu\|_2^2 \quad \forall \mu > 0, \quad (6.32)$$

where $\psi_\mu(x) = e^{\mu|x|} \psi(x)$. This is of course impossible for μ large enough. Hence $\psi_\mu = 0$ and the claim follows. \blacksquare

Remark 6.6. The statement of Theorem 6.5 cannot be extended to the interval $[\Lambda, \infty)$. Indeed, the result of Loss and Yau [19] shows that if

$$B(x) = \frac{12}{(1 + |x|^2)^3} (2x_1x_3 - 2x_2, 2x_2x_3 + 2x_1, 1 - x_1^2 - x_2^2 + x_3^2), \quad (6.33)$$

then zero is an eigenvalue of $H_{A,0}$. More precisely, Loss and Yau proved that there exists $A : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with $\text{curl}A = B$ such that

$$\sigma \cdot (P - A) \varphi = 0, \quad \varphi = \frac{1 + i\sigma \cdot x}{(1 + |x|^2)^{3/2}} \varphi_0, \quad (6.34)$$

where φ_0 is an arbitrary normalized spinor. In this case we have $\Lambda = 0$, see equation (6.33) and Lemma A.1. Hence Theorem 6.5 guarantees the absence of eigenvalues in the interval $(0, \infty)$. The fact that our technique cannot be applied to exclude zero eigenvalue is reflected also by the power-like decay of φ at infinity, see (6.34), which is in stark contrast to the super-exponential decay of eigenfunctions with positive eigenvalues, cf. Proposition 6.1.

For more examples of magnetic fields supporting a zero eigenvalue we refer to [1, 11, 21]. It should be pointed out, however, that the existence of a zero eigenvalue of $H_{A,0}$ is an exceptional event. Indeed, it was proven in [6] that those magnetic fields for which zero is not an eigenvalue of $H_{A,0}$ form a dense set in $L^{3/2}(\mathbb{R}^3; \mathbb{R}^3)$.

As a simple consequence of Theorem 6.5 we obtain sufficient conditions for absence of positive eigenvalues of $H_{A,V}$.

Corollary 6.7. *Let B, V satisfy assumptions of Proposition A.2 and suppose moreover that $B(x) = o(|x|^{-1})$ and $\|V(x)\| = o(|x|^{-1})$ as $|x| \rightarrow \infty$. Then the operator $H_{A,V}$ has no positive eigenvalues.*

Proof. We use the splitting $V^s = V$, $V^\ell = 0$. From the assumptions of the corollary and from Proposition A.2 we get $\beta = \omega_1 = \omega_2 = 0$. The claim thus follows from Theorem 6.5. \blacksquare

7. Example

In this section we construct an example which indicates the sharpness of the critical energy Λ . Consider the radial magnetic field

$$B(x) = (0, 0, b(r)), \quad b(r) = \frac{b_0}{\sqrt{1 + r^2}}, \quad r = \sqrt{x_1^2 + x_2^2} \quad (7.1)$$

The vector potential associated to B in the Poincaré gauge is then given by

$$A(x) = \frac{(-x_2, x_1, 0)}{r} \int_0^r b(s) s ds =: (a_1(x_1, x_2), a_2(x_1, x_2), 0).$$

Let $v : \mathbb{R} \rightarrow (-\infty, 0]$ be a bounded compactly supported function such that $\int_{\mathbb{R}} v < 0$, and let

$$V(x) = \begin{pmatrix} 0 & 0 \\ 0 & v(x_3) \end{pmatrix}.$$

Then

$$H_{A,\varepsilon V} = \begin{pmatrix} h_+ \oplus P_3^2 & 0 \\ 0 & h_- \oplus (P_3^2 + \varepsilon v) \end{pmatrix}, \quad (7.2)$$

where h_\pm are the operators in $L^2(\mathbb{R}^2)$ acting on their domains as

$$h_\pm = (P_1 - a_1)^2 + (P_2 - a_2)^2 \pm b.$$

Obviously, the operators h_{\pm} are non-negative being the components of the associated two-dimensional Pauli operator. In addition, since $b(r) \rightarrow 0$ as $r \rightarrow \infty$, the structure of the spectra of h_{\pm} is the same as that of the two-dimensional magnetic Schrödinger operator $(P_1 - a_1)^2 + (P_2 - a_2)^2$. In particular, from the well-known example of Miller-Simon [20], with a numerical error corrected in [4, Sec. 6.1], it follows that the spectrum of h_{\pm} is *dense pure point* in $[0, b_0^2)$ and *absolutely continuous* in $[b_0^2, \infty)$.

Hence if $\varepsilon > 0$ is small enough such that the operator $P_3^2 + \varepsilon v$ in $L^2(\mathbb{R})$ has exactly one discrete negative eigenvalue $-\lambda(\varepsilon)$, then by (7.2),

$$\sigma_{\text{es}}(H_{A,\varepsilon V}) = [-\lambda(\varepsilon), \infty),$$

and the spectrum of $H_{A,\varepsilon V}$ is *dense pure point* in $[-\lambda(\varepsilon), b_0^2 - \lambda(\varepsilon))$ and *absolutely continuous* in $[b_0^2 - \lambda(\varepsilon), \infty)$.

On the other hand, it is easily verified that B and V satisfy assumptions of Theorem 6.5, see Proposition A.2. Moreover, putting $V^s = 0$ and $V^{\ell} = V$ gives $\omega_1 = \omega_2 = 0$. Therefore, for any $\varepsilon > 0$ we have $\Lambda(B, \varepsilon V) = \beta^2 = b_0^2$. Since $\lambda(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$, the above example shows that the threshold energy Λ cannot be improved.

Remark 7.1. Examples of magnetic fields which produce embedded eigenvalues of $H_{A,V}$ above any fixed energy were found in [5, Thm. 5.1] and [9, Thm. 3.1].

8. The Dirac operator

The magnetic Dirac operator in $L^2(\mathbb{R}^3, \mathbb{C}^4)$ is given by

$$\mathbb{D} = \begin{pmatrix} m\mathbf{1} & \sigma \cdot (P - A) \\ \sigma \cdot (P - A) & -m\mathbf{1} \end{pmatrix}, \quad (8.1)$$

where $m \geq 0$ is a constant. From Assumption 3.4 and equation (4.3) it follows that $\mathcal{D}(\mathbb{D}) = \mathcal{D}(P - A)$. Recall also that

$$\sigma(\mathbb{D}) = \sigma_{\text{es}}(\mathbb{D}) = (-\infty, -m] \cup [m, \infty).$$

We have

Theorem 8.1. *Let B satisfy the Assumptions 3.3, 3.4, 3.9 and 3.10 (with $V = 0$). Suppose that $A \in L^2_{\text{loc}}(\mathbb{R}^3; \mathbb{R}^3)$ is such that $\text{curl } A = B$. Then the Dirac operator \mathbb{D} has no eigenvalues in*

$$(-\infty, -\sqrt{\beta^2 + m^2}) \cup (\sqrt{\beta^2 + m^2}, \infty),$$

where β is given by (3.10).

Proof. Since the spectrum of \mathbb{D} is gauge invariant, we may suppose without loss of generality that A is given by (2.12). Note that

$$\mathbb{D}^2 = \begin{pmatrix} H_{A,0} + m^2\mathbf{1} & 0 \\ 0 & H_{A,0} + m^2\mathbf{1} \end{pmatrix}, \quad (8.2)$$

in the sense of quadratic forms on $\mathcal{D}(P - A)$. This means that if $\mathbb{D}\psi = E\psi$ for some $\psi \in \mathcal{D}(P - A)$, then ψ is a weak eigenfunction of $H_{A,0}$ relative to eigenvalue $E^2 - m^2$. Since $\Lambda = \beta^2$, in view of Theorem 6.5 we must have $E^2 - m^2 \leq \beta^2$. ■

Corollary 8.2. *Let B be such that $|\tilde{B}| \in L^p_{\text{loc}}(\mathbb{R}^3)$ for some $p > 3$, and suppose that $B(x) = o(|x|^{-1})$ as $|x| \rightarrow \infty$. Then the operator \mathbb{D} has no eigenvalues in $(-\infty, -m) \cup (m, \infty)$.*

Proof. This is a combination of Proposition A.2 and Theorem 8.1. ■

Remark 8.3. As in the case of Pauli operator we note that the claim of Corollary 8.2 cannot be extended to the set $(-\infty, -m] \cup [m, \infty)$. Indeed, the magnetic field given by (6.33) satisfies assumptions of Corollary 8.2, but the associated Dirac operator \mathbb{D} has eigenvalues m and $-m$. To see this, consider the spinor $\varphi \in L^2(\mathbb{R}^3, \mathbb{C}^2)$ given by (6.34). Then, with a slight abuse of notation,

$$\mathbb{D} \begin{pmatrix} \varphi \\ 0 \end{pmatrix} = m \begin{pmatrix} \varphi \\ 0 \end{pmatrix} \quad \text{and} \quad \mathbb{D} \begin{pmatrix} 0 \\ \varphi \end{pmatrix} = -m \begin{pmatrix} 0 \\ \varphi \end{pmatrix}.$$

One should mention that sufficient conditions for the absence of all eigenvalues of \mathbb{D} were established in [10]. Indeed, it was proved there that when $A \in W_{\text{loc}}^{1,3}(\mathbb{R}^3)$, then the operator \mathbb{D} has no eigenvalues in $(-\infty, -m] \cup [m, \infty)$, and therefore no eigenvalues at all, if the functional inequality

$$\int_{\mathbb{R}^3} |x|^2 |B|^2 |u|^2 \leq c^2 \int_{\mathbb{R}^3} |(P - A)u|^2 \tag{8.3}$$

holds for all $u \in C_0^\infty(\mathbb{R}^3)$ and with a constant c which satisfies

$$c \left(11 + \frac{3^{3/2}}{2} \sqrt{c} \right) < 1,$$

see [10, Thm. 3.6].

Remark 8.4. Non existence of eigenvalues of the perturbed Dirac operator $\mathbb{D} + \mathbf{1}q$ was studied already by Kalf [16]. He proved that if

$$|x|(|q(x)| + |B(x)|) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty, \tag{8.4}$$

then the operator $\mathbb{D} + \mathbf{1}q$ has no eigenvalues in $\mathbb{R} \setminus [-m, m]$. Note that (8.4) implies $\beta = 0$. The result of Kalf was later extended to matrix valued potentials in [8].

APPENDIX A. Pointwise and local L^p conditions

Here we formulate sufficient conditions which guarantee the validity of Assumptions 3.7, 3.8, 3.9.

A.1. Pointwise conditions.

Lemma A.1. *Given a magnetic field B and potential $V = V^s + V^\ell$ assume that $|\tilde{B}|, |V|_{\mathbb{C}^2}, |xV^s|_{\mathbb{C}^2}$, and $|x \cdot \nabla V^\ell|_{\mathbb{C}^2}$ are bounded outside of a compact set, and that*

$$\lim_{|x| \rightarrow \infty} |V_+(x)|_{\mathbb{C}^2} = 0.$$

Then Assumptions 3.7, 3.8 and 3.9 are satisfied and

$$\beta \leq \limsup_{|x| \rightarrow \infty} |\tilde{B}(x)|, \quad \omega_1 \leq \limsup_{|x| \rightarrow \infty} |xV^s(x)|_{\mathbb{C}^2}, \quad \text{and} \quad \omega_2 \leq \limsup_{|x| \rightarrow \infty} |(x \cdot \nabla V^\ell(x))_+|_{\mathbb{C}^2}. \tag{A.1}$$

Proof. This is a straightforward consequence of the definitions of β, ω_1 and ω_2 . ■

A.2. Local L^p conditions. The conditions of Lemma A.1 can be relaxed by considering potentials which are not necessarily bounded at infinity, but which belong to $L^p_{\text{loc}}(\mathbb{R}^3)$ for a suitable p .

Proposition A.2. *Let B, V satisfy conditions of Lemma A.1, and let V satisfy Assumption 3.1. Suppose moreover that $|\tilde{B}| \in L^p_{\text{loc}}(\mathbb{R}^3)$ and $|V^{s,\ell}(\cdot)|_{\mathbb{C}^2} \in L^p_{\text{loc}}(\mathbb{R}^3)$ for some $p > 3$. Then all the hypotheses of Section 3 are satisfied with $\alpha_j = 0$ for $j = 0, 1, 2, 3$, and the constants $\beta, \omega_1, \omega_2$ satisfy (A.1).*

Proof. If $|\tilde{B}| \in L^p_{\text{loc}}(\mathbb{R}^3)$ with $p > 3$, then it is easily seen that Assumption 3.3 holds. In view of Lemma A.1 it thus remains to prove Assumptions 3.4, 3.5 and 3.10. Given a matrix valued function M on \mathbb{R}^3 and a test function (2.10) with $u, v \in \mathcal{D}(P - A)$, we have

$$\|M\varphi\|_2^2 \leq \| \|M(x)\|_{\mathbb{C}^2} \varphi \|_2^2 = \int_{\mathbb{R}^3} \|M(x)\|_{\mathbb{C}^2}^2 (|u(x)|^2 + |v(x)|^2) dx, \quad (\text{A.2})$$

$$|\langle \varphi, M\varphi \rangle| \leq \langle \varphi, \|M(x)\|_{\mathbb{C}^2} \varphi \rangle = \int_{\mathbb{R}^3} \|M(x)\|_{\mathbb{C}^2} (|u(x)|^2 + |v(x)|^2) dx. \quad (\text{A.3})$$

So let $u \in \mathcal{D}(P - A)$, and let $W \in L^p_{\text{loc}}(\mathbb{R}^3)$, $p > 3$, be bounded outside a compact set $K \subset \mathbb{R}^3$. The compactness of the Sobolev embedding $H^1(K) \hookrightarrow L^s(K)$, $2 \leq s < 6$ and the diamagnetic inequality imply that for any $\varepsilon > 0$ there exists C_ε such that

$$\left(\int_K |u|^s dx \right)^{\frac{2}{s}} \leq \varepsilon \|\nabla|u|\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2 \leq \varepsilon \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2 \quad \forall s \in [2, 6). \quad (\text{A.4})$$

Equation (A.4) and the Hölder inequality give

$$\|Wu\|_{L^2(\mathbb{R}^3)}^2 \leq \|W\|_{L^\infty(K^c)}^2 \|u\|_{L^2(\mathbb{R}^3)}^2 + \|W\|_{L^p(K)}^2 \left(\int_K |u|^{p'} dx \right)^{\frac{2}{p'}} \leq \varepsilon \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2, \quad (\text{A.5})$$

where $p' \in (2, 6)$ satisfies $\frac{2}{p} + \frac{2}{p'} = 1$. By the hypotheses of the proposition we can apply the above estimate with W replaced by $|B|, |\tilde{B}|$ and $\|xV^s\|_{\mathbb{C}^2}$ respectively. This in combination with (A.2) implies Assumption 3.4 and the upper bound (3.11) of Assumption 3.10 with $\alpha_1 = 0$. In the same way we get

$$\int_K |W||u|^2 dx \leq \|W\|_{L^p(K)} (\varepsilon \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2).$$

Inserting $W = \|V\|_{\mathbb{C}^2}$ in the above estimate and using (A.3) we obtain Assumption 3.5 with $\alpha_0 = 0$, and estimate (3.13) with $\alpha_3 = 0$.

To prove (3.12) consider W as above and take R large enough such that $K \subset \mathcal{U}_R$. Integration by parts yields

$$\int_{\mathcal{U}_R} x \cdot \nabla W |u|^2 dx = R \int_{\partial\mathcal{U}_R} W |u|^2 dS - 2 \int_{\mathcal{U}_R} W (|u|^2 + \text{Re}(\bar{u}x \cdot \nabla u)) dx. \quad (\text{A.6})$$

Now let $\varepsilon > 0$. Since the trace embedding $H^1(\mathcal{U}_R) \hookrightarrow L^2(\partial\mathcal{U}_R)$ is compact and $W \in L^\infty(\partial\mathcal{U}_R)$, there exists C_ε such that

$$R \int_{\partial\mathcal{U}_R} W |u|^2 dS \leq \varepsilon \int_{\mathcal{U}_R} |\nabla|u|\|^2 dx + C_\varepsilon \int_{\mathcal{U}_R} |u|^2 dx \leq \varepsilon \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2, \quad (\text{A.7})$$

where we have used also the diamagnetic inequality. As for the second term in (A.6), we note that $x \cdot \nabla u = x \cdot (P - A)u$, see (2.13), and hence

$$\begin{aligned} \left| \int_{\mathcal{U}_R} W \operatorname{Re}(\bar{u} x \cdot \nabla u) dx \right| &= \left| \int_{\mathcal{U}_R} W \operatorname{Im}(\bar{u} x \cdot (P - A)u) dx \right| \leq R \int_{\mathcal{U}_R} |W| |u| |(P - A)u| dx \\ &\leq R \|Wu\|_{L^2(\mathcal{U}_R)} \|(P - A)u\|_{L^2(\mathbb{R}^3)} \leq \frac{R^2}{\sqrt{\varepsilon}} \|Wu\|_{L^2(\mathcal{U}_R)}^2 + \sqrt{\varepsilon} \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 \\ &\leq \sqrt{\varepsilon} \left(R^2 \|W\|_{L^p(\mathcal{U}_R)}^2 + 1 \right) \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2. \end{aligned}$$

where we have used the estimate

$$\int_{\mathcal{U}_R} W^2 |u|^2 dx \leq \|W\|_{L^p(\mathcal{U}_R)}^2 \left(\varepsilon \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2 \right),$$

see (A.5). Putting the above estimates together and using $W \in L^\infty(K^c)$ we find that

$$|\langle u, x \cdot \nabla W u \rangle_{L^2(\mathbb{R}^3)}| \leq \varepsilon \|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + C_\varepsilon \|u\|_{L^2(\mathbb{R}^3)}^2.$$

The polarisation identity (2.11) now gives

$$|\langle v, x \cdot \nabla W u \rangle_{L^2(\mathbb{R}^3)}| \leq \varepsilon (\|(P - A)u\|_{L^2(\mathbb{R}^3)}^2 + \|(P - A)v\|_{L^2(\mathbb{R}^3)}^2) + C_\varepsilon (\|u\|_{L^2(\mathbb{R}^3)}^2 + \|v\|_{L^2(\mathbb{R}^3)}^2).$$

Applying the above estimate with W replaced by the matrix elements of V^ℓ yields inequality (3.12) with $\alpha_2 = 0$. \blacksquare

A.3. Uniformly local L^p conditions. In order to include potentials with stronger singularities than those allowed by Proposition A.2, we introduce the class

$$L_{\text{loc,unif}}^p = \left\{ f : \sup_{x \in \mathbb{R}^3} \int_{\mathcal{U}_1(x)} |f(y)|^p dy < \infty \right\}, \quad p > \frac{3}{2} \quad (\text{A.8})$$

equipped with the norm

$$\|f\|_{L_{\text{loc,unif}}^p} = \sup_{x \in \mathbb{R}^3} \left(\int_{\mathcal{U}_1(x)} |f(y)|^p dy \right)^{1/p}. \quad (\text{A.9})$$

Definition A.3. Let $f \in L_{\text{loc,unif}}^p$. We say that g is equivalent to f at infinity, and write $g \sim f$ if $g \in L^\infty(\mathbb{R}^3)$ and if

$$\limsup_{R \rightarrow \infty} \left\| \mathbb{1}_{\mathcal{U}_R^c} (f - g) \right\|_{L_{\text{loc,unif}}^p} = 0. \quad (\text{A.10})$$

Given $f \in L_{\text{loc,unif}}^p$, we define

$$\gamma(f) = \inf \left(\|g\|_\infty : g \sim f \right).$$

We then have

Proposition A.4. *Let B, V satisfy conditions of Lemma A.1, and let V satisfy Assumption 3.1. Suppose moreover that $|\tilde{B}| \in L_{\text{loc,unif}}^p(\mathbb{R}^3)$ and $|V^{s,\ell}(\cdot)|_{\mathbb{C}^2} \in L_{\text{loc,unif}}^p(\mathbb{R}^3)$ for some $p > 3/2$. Then all the hypotheses of Section 3 are satisfied with $\alpha_j = 0$ for $j = 0, 1, 2, 3$, and the constants $\beta, \omega_1, \omega_2$ satisfy*

$$\beta \leq \limsup_{|x| \rightarrow \infty} \gamma(|\tilde{B}(x)|), \quad \omega_1 \leq \gamma(|xV^s(x)|_{\mathbb{C}^2}), \quad \omega_2 \leq \gamma(|x \cdot \nabla V^\ell(x)|_{\mathbb{C}^2}) \quad (\text{A.11})$$

Proposition A.4 is a matrix valued version of the results established in [4, Sec. A.1]. We therefore omit the proof and refer to [4].

Acknowledgments: Hynek Kovařík has been partially supported by Gruppo Nazionale per Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). Dirk Hundertmark acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 258734477 – SFB 1173.

REFERENCES

- [1] C. Adam, B. Muratori, and C. Nash: *Zero modes of the Dirac operator in three dimensions*. Phys. Rev.D (3), 60(12):125001, 8, 1999.
- [2] S. Agmon, *Lower bounds for solutions of Schrödinger equations*. J. Anal. Math., **23** (1970) 1–25.
- [3] S. Albeverio, *On Bound States in the Continuum of N-body Systems and the Virial Theorem* Ann. Physics **71** (1972) 167–276.
- [4] S. Avramska-Lukarska, D. Hundertmark, H. Kovařík: *Absence of positive eigenvalues for magnetic Schrödinger operators*. Calc. Var. **62**, 63 (2023).
- [5] J. Avron, I. Herbst, B. Simon: *Schrödinger operators with magnetic fields. I* Duke Math. J., **45** (1978) 847–884.
- [6] A. Balinsky, W.D. Evans: *On the zero models of Pauli Operators*. J. Func. Anal. **179** (2001) 120–135.
- [7] R. Benguria, H. Van Den Bosch: *A criterion for the existence of zero modes for the Pauli operator with fastly decaying fields*. J. Math. Phys. **56** (2015) pp. 052104.
- [8] A. Berthier, V. Gerogescu: *On the Point Spectrum of Dirac Operators*. J. Func. Anal. **71** (1987) 309–338.
- [9] J.-F. Bony, V. Bruneau, G. D. Raikov: *Resonances and spectral shift function singularities for magnetic quantum Hamiltonians*. RIMS Kokyuroku Bessatsu **B45** (2014) 77–100.
- [10] L. Cossetti, L. Fanelli, D. Krejčířík: *Absence of eigenvalues of Dirac and Pauli Hamiltonians via the method of multipliers*. Comm. Math. Phys. **379** (2020) 633–691.
- [11] D.M. Elton: *The local structure of zero mode producing magnetic potentials*. Comm. Math. Phys. **229** (2002) 121–139.
- [12] R.L. Frank, M. Loss: *Which magnetic fields support a zero mode?* J. Reine Angew. Math. **788** (2022) 1–36.
- [13] R.L. Frank, M. Loss: *A sharp criterion for zero modes of the Dirac equation*. J. Eur. Math. Soc., to appear. arXiv:2201.03610.
- [14] R. Froese, I. Herbst: *Exponential Bounds and Absence of Positive Eigenvalues for Many-Body Schrödinger Operators*. Comm. Math. Phys., **87** (1982), 429–447.
- [15] R. Froese, I. Herbst, M. Hoffmann-Ostenhof, T. Hoffmann-Ostenhof: *On the absence of positive eigenvalues for one-body Schrödinger operators*. J. Anal. Math. **41** (1982), 272–284. Reine Angew. Math., **571**, (2004) 107–130.
- [16] H. Kalf: *Nonexistence of eigenvalues of Dirac operators*. Proc. Roy. Soc. Edinburgh Sect. A **89** (1981) 309–317.
- [17] T. Kato, *Growth Properties of Solutions of the Reduced Wave Equation With a Variable Coefficient*. Commun. Pure Appl. Math., **12** (1959), 403–425.
- [18] T. Kato, *Schrödinger operators with singular potentials*. Israel J. Math. **13** (1972) 135–148.
- [19] M. Loss and H.-T. Yau: *Stability of Coulomb systems with magnetic fields. III. Zero energy bound states of the Pauli operator*. Comm. Math. Phys. **104** (1986) 282–290.
- [20] K. Miller, B. Simon: *Quantum Magnetic Hamiltonians with Remarkable Spectral Properties*. Phys. Rev. Lett. **44** (1980), 1706–1707.
- [21] C Ross and B.J. Schroers: *Magnetic zero-modes, vortices and Cartan geometry*. Lett. Math.Phys., 108(4):949–983, 2018.
- [22] B. Simon, *On positive eigenvalues of one body Schrödinger operators*. Commun. Pure Appl. Math. **22** (1967), 531–538.
- [23] B. Thaller: *The Dirac Equation*. Texts and Monographs in Physics, Springer 1992.
- [24] J. Weidmann, *The virial theorem and its application to the spectral theory of Schrödinger operators*. Bull. Amer. Math. Soc. **73** (1967), 452–456.

DIRK HUNDERTMARK, DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ANALYSIS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76128 KARLSRUHE, GERMANY, AND DEPARTMENT OF MATHEMATICS UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN 1409 W. GREEN STREET URBANA, ILLINOIS 61801-2975

Email address: `dirk.hundertmark@kit.edu`

HYNEK KOVAŘÍK, DICATAM, SEZIONE DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI BRESCIA, VIA BRANZE 38 - 25123, BRESCIA, ITALY

Email address: `hynek.kovarik@unibs.it`