

Corrigendum to “Tumorigenicity of IL-1 α - and IL-1 β -Deficient Fibrosarcoma Cells” [Neoplasia, Volume 10, Issue 6, June 2008, Pages 549–562]



Irina Nazarenko ^{*,1}, Rachid Marhaba ^{*,1}, Eli Reich [†], Elena Voronov [†], Mario Vitacolonna ^{*}, Dagmar Hildebrand ^{*}, Elena Elter ^{*}, Mohini Rajasagi ^{*}, Ron N. Apte [†], Margot Zöller ^{*,†}

^{*} Department of Tumor Progression and Tumor Defense, German Cancer Research Center Heidelberg, Germany

[†] The Shraga Segal Department of Microbiology and Immunology and the Cancer Center, Ben Gurion University Beer-Sheva, Israel

[‡] Department of Applied Genetics, University of Karlsruhe. Karlsruhe, Germany


The authors regret an accidental duplication of the migration assays showed on the Fig. 3D for *wt2* and *IL-1 α* conditions that was brought to our attention in the originally published manuscript. This is an accidental error, and the data now removed from the Fig. 3D shown below. Furthermore, the accompanying figure legend has been corrected to

reflect this change, indicated in bolded text.

This error does not affect the interpretation of the data. No changes on the main text are required. The authors remain committed to the highest standards of accuracy in the published work and greatly apologize to any inconvenience caused.

DOI of original article: <https://doi.org/10.1593/neo.08286>.

¹ These authors equally contributed to this study.



(caption on next page)

Fig. 3. Tumor-associated IL-1, matrix adhesion, migration, and angiogenesis induction. (A–G) Three wt, IL-1 α –/–, IL-1 β –/–, and IL-1 $\alpha\beta$ –/– fibrosarcoma lines were tested for the following: (A) Adhesion molecule expression by flow cytometry. The percentage of stained cells and the mean intensity of staining are shown. (B) Adhesion to components of the extracellular matrix. Adhesion was measured after a 2-hour incubation by crystal violet staining. (C) Transwell migration, evaluated after a 4-hour incubation by crystal violet staining. (D) Migration into a wounded monolayer 48 hours after wounding. **A representative example of one wt, IL-1 β –/–, and IL-1 $\alpha\beta$ –/– fibrosarcoma line is shown.** (E) Cable formation on matrigel evaluated after 24 hours. (F) uPAR expression was evaluated by flow cytometry. The percentage of stained cells and the mean intensity of staining are shown. (G) MMP2 and MMP9 expression in 3 wt, IL-1 α –/–, IL-1 β –/–, and IL-1 $\alpha\beta$ –/– fibrosarcoma lines as revealed by zymography. (A–C and F) Mean \pm SD of three distinct lines are presented, and significant differences between wt, IL-1 α –/–, IL-1 β –/–, and IL-1 $\alpha\beta$ –/– lines are indicated by an asterisk. All experiments were repeated at least three times and revealed comparable results.
