

RECEIVED: October 5, 2024

REVISED: July 22, 2025

ACCEPTED: September 28, 2025

PUBLISHED: October 27, 2025

Study of same-sign W boson scattering and anomalous couplings in events with one tau lepton from pp collisions at $\sqrt{s} = 13$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A first study is presented of the cross section for the scattering of same-sign W boson pairs via the detection of a τ lepton. The data from proton-proton collisions at the center-of-mass energy of 13 TeV were collected by the CMS detector at the LHC, and correspond to an integrated luminosity of 138 fb^{-1} . Events were selected that contain two jets with large pseudorapidity and large invariant mass, one τ lepton, one light lepton (e or μ), and significant missing transverse momentum. The measured cross section for electroweak same-sign WW scattering is $1.44^{+0.63}_{-0.56}$ times the standard model prediction. In addition, a search is presented for the indirect effects of processes beyond the standard model via the effective field theory framework, in terms of dimension-6 and dimension-8 operators.

KEYWORDS: Hadron-Hadron Scattering, Vector Boson Production

ARXIV EPRINT: [2410.04210](https://arxiv.org/abs/2410.04210)

Contents

1	Introduction	1
2	The CMS detector	3
3	Simulated samples	4
4	Event reconstruction	5
5	Analysis strategy and background estimation	7
6	Systematic uncertainties	13
7	Measurement of the SM SSWW VBS processes	14
8	Effective field theory interpretation	16
9	Summary	20
	The CMS collaboration	28

1 Introduction

The discovery of the Higgs boson [1] provided evidence that fermions and bosons acquire their masses through the Brout-Englert-Higgs mechanism of electroweak (EW) spontaneous symmetry breaking [2–4]. In this framework, vector boson scattering (VBS) processes play a special role. This occurs because in the standard model (SM) the unitarity of the scattering amplitude depends on a cancellation of diagrams involving the mediation of the Higgs boson between longitudinally polarized vector bosons. As a result, the contribution of the longitudinal polarization component to vector boson scattering is very small. Therefore, for even small deviations from the SM couplings of the Higgs boson to the vector bosons, the VBS cross section would diverge from the SM expectations with increasing center-of-mass energy. The measurement of VBS processes thus provides an indirect probe of physics beyond the SM (BSM), even for scenarios in which new resonances are not energetically accessible at the LHC [5]. The theoretical calculation of the effects of different sources of deviations from the SM is sensitive to the method chosen to “unitarize” the process at higher energies [6, 7].

At tree level, the topology commonly identified as VBS appears through purely EW diagrams of order α_{EW}^6 , where $\alpha_{\text{EW}} = g_W^2/4\pi$ is the electroweak SU(2) coupling. However, additional non-VBS electroweak diagrams in the same order also contribute, and the ensemble of all indistinguishable contributions constitutes the electroweak WW production process. For simplicity, we refer to this entire production mechanism at order α_{EW}^6 as “EW VBS” in the rest of this article. Because it arises from higher-order electroweak interactions, EW VBS generally leads to a small cross section. Along with these purely EW terms, experimental




Figure 1. Representative tree-level Feynman diagrams contributing to the process $qq' \rightarrow \tau^\pm \nu_\tau \ell^\pm \nu_\ell jj$, $\ell = e, \mu$, leading to cross sections of order α_{EW}^6 (left) and $\alpha_S^2 \alpha_{EW}^4$ (right).

VBS signatures also include irreducible contributions that enter at order $\alpha_S^2 \alpha_{EW}^4$, where α_S is the strong coupling. These contributions are referred to as quantum chromodynamics (QCD) irreducible contributions. Typical Feynman diagrams for these processes are shown in figure 1.

The ATLAS collaboration provided the first evidence of same-sign W pair (SSWW) production via VBS in 2014 by studying final states with electrons and muons [8]. The first observation, in the same final states, was presented by CMS in 2017 [9]. Among the EW-mediated processes, SSWW EW VBS has the largest cross section ratio between the EW and QCD production modes [10].

The study presented here of $pp \rightarrow W^\pm W^\pm + \text{jets}$ is based on data from proton-proton collisions at $\sqrt{s} = 13 \text{ TeV}$ collected by the CMS experiment at the CERN LHC from 2016 to 2018 and corresponding to an integrated luminosity of 138 fb^{-1} . We investigate a heretofore unexplored final state characterized by the decay of one of the scattered W bosons into a τ lepton that subsequently decays into hadrons (hadronic τ candidate, τ_h). The final state thus consists of a charged light lepton $\ell = e, \mu$, the corresponding neutrino ν_ℓ , one τ_h candidate, the corresponding ν_τ , and two jets produced by the quarks recoiling from the production of the W boson pair. Contributions to this final state could arise from BSM scenarios that favor the τ over light leptons. Examples of these scenarios are provided by the $SU(2)_l \times SU(2)_h \times U(1)$ family models, denoted in the following as G(221) models, where the first two families of quarks and leptons transform according to the $SU(2)_l$ group and the third family according to the $SU(2)_h$ group [11–13]. In these frameworks, a moderate mixing between the W and W' gauge fields suppresses the W' -quark coupling while retaining a sizable HWW' coupling. In the context of the latest direct searches for τ resonances performed by CMS [14] and ATLAS [15], this setup is not excluded for $m_{W'} = 3.0 \text{ TeV}$ and can still yield an enhancement of the $pp \rightarrow \ell \nu_\ell \tau \nu_\tau + \text{jets}$ cross section.

The sensitivity to indirect BSM effects can be probed within the standard model effective field theory framework [16, 17]. This theory is referred to in the literature as “SMEFT”, and we adopt the formulation of ref. [17]. Assuming that new physics with energy scale $\Lambda_{BSM} \gg \Lambda_{SM}$ (where Λ_{SM} is a characteristic energy value, such as the Higgs field vacuum expectation value) induces only perturbative effects in EW VBS processes, the theory is implemented by introducing the following effective Lagrangian:

$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{SM} + \sum_{D_\alpha > 4} \sum_{\alpha} \frac{c_\alpha^{(D_\alpha)}}{\Lambda_{BSM}^{D_\alpha - 4}} \mathcal{O}_\alpha^{(D)}, \quad (1.1)$$

where the operators $\mathcal{O}_\alpha^{(D)}$ are constructed with SM fields at some dimension D_α , and $c_\alpha^{(D_\alpha)}$ are the Wilson coefficients. In this way, the contribution $\mathcal{A}_{\text{BSM}}$ to the total scattering amplitude due to the EFT operators is given by:

$$|\mathcal{A}_{\text{BSM}}|^2 = \sum_i^{D_i > 4} \left[\frac{c_i^{(D_i)}}{\Lambda_{\text{BSM}}^{D_i-4}} 2 \text{Re} |\mathcal{A}_{\text{SM}}^* \mathcal{A}_{\mathcal{O}_i^{(D_i)}}| + \frac{c_i^{(D_i)^2}}{\Lambda_{\text{BSM}}^{2(D_i-4)}} |\mathcal{A}_{\mathcal{O}_i^{(D_i)}}|^2 \right] \\ + \sum_{j \neq k}^{D_j, D_k > 4} \frac{c_j^{(D_j)} c_k^{(D_k)}}{\Lambda_{\text{BSM}}^{D_j+D_k-8}} \text{Re} |\mathcal{A}_{\mathcal{O}_j^{(D_j)}}^* \mathcal{A}_{\mathcal{O}_k^{(D_k)}}|, \quad (1.2)$$

where the first summation runs over the interference terms between SM and one $\mathcal{O}_i^{(D)}$ operator, the second one over the quadratic contributions of the operators, and the last one over interference between two different operators. The $\mathcal{O}_\alpha^{(D)}$, as well as the corresponding $c_\alpha^{(D_\alpha)}$, are classified according to their dimension D_α to provide a first categorization of their physics effects, as explained below.

A deviation from SM expectations in the observed data yields would result in non-zero Wilson coefficient values for operators sensitive to such deviations. This effect would thus provide guidance for characterizing BSM effects. Effective field theory interpretations of search outcomes have previously been presented by the ATLAS and CMS Collaborations [18–28], along with phenomenological reinterpretations of the electroweak VBS differential cross section measurements [29]. Assuming that the new operators still respect lepton-number conservation, those with odd dimensions are ruled out. As a consequence, in this study, we investigate operators of dimension-6 (dim-6) and dimension-8 (dim-8), which induce anomalous triple and quartic gauge couplings, modify boson-fermion couplings, and introduce fermion contact interactions [30–32]. We refer to any contribution of dimension greater than four as an EFT contribution. Specifically, the dim-6 operators are implemented in the standard SMEFT framework via the Warsaw basis [17], while the dim-8 operators are introduced using the Eboli basis [33].

In this paper, we introduce a machine-learning approach to identifying the single- τ_h final state in the SSWW VBS process and measuring the signal strength, defined as the ratio of the observed signal yield to that predicted by the model, for both the EW contribution (with the QCD contribution fixed) and for the combined EW+QCD combination of these processes (with the ratio between the EW and QCD contributions fixed). The same approach is implemented to develop models capable of discriminating possible EFT contributions from SM processes.

In the remainder of this paper, section 2 describes the CMS detector, section 3 presents the methods for simulating events, and section 4 describes the particle reconstruction. The selection of signal and control samples is described in section 5, along with the background estimation, and in section 6 we discuss systematic uncertainties. The results of the measurement are given in section 7, and of the EFT interpretation in section 8. We conclude with a summary in section 9.

2 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within this magnetic field volume are a silicon

pixel and silicon strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass-and-scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the coverage in pseudorapidity η provided by the barrel and endcap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in ref. [34].

The silicon tracker used in 2016 measured charged particles within the range $|\eta| < 2.5$. For nonisolated particles of transverse momentum p_T in the range $1 < p_T < 10$ GeV and $|\eta| < 1.4$, the track resolutions were typically 1.5% in p_T and 25–90 (45–150) μm in the transverse (longitudinal) impact parameter. For isolated particles with $p_T = 100$ GeV emitted at $|\eta| < 1.4$, the resolutions are approximately 2.8% in p_T , and in impact parameter 10 μm (transverse) and 30 μm (longitudinal) [35]. At the start of 2017, a new pixel detector was installed [36]; the upgraded tracker measured particles up to $|\eta| < 3.0$ with typical resolutions of 1.5% in p_T and 20–75 μm in the transverse impact parameter for nonisolated particles of $1 < p_T < 10$ GeV [37].

Events of interest are selected using a two-tiered trigger system [38]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz. The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software, optimized for fast processing that reduces the event rate to around 1 kHz before data storage.

3 Simulated samples

Monte Carlo (MC) simulation is used in the analysis for the design of the event selection, evaluation of signal efficiencies, and estimation of some backgrounds. The EW VBS signal samples are simulated at leading order (LO) with six EW and zero QCD vertices with the MADGRAPH5_aMC@NLO v2.6.5 generator [39], requiring a final state with $\ell\nu_\ell, \tau\nu_\tau$ pairs from the decays of the two W bosons (figure 1, left). The MADGRAPH5_aMC@NLO generator is also used to simulate the QCD-mediated SSWW process, which is generated at LO with up to three additional partons in the matrix element calculations that have two QCD vertices at the tree level (figure 1, right). The interference between the SSWW EW and QCD diagrams, including the terms of order $\alpha_S\alpha_{EW}^5$, contributes less than 4% to the inclusive cross section for the EW signal over the phase space region of interest of the analysis and is therefore neglected.

A complete set of next-to-LO (NLO) QCD and EW corrections for the SSWW VBS processes, described above in the leptonic decay channel for each W boson, have been computed as a function of the invariant mass of the VBS jet system [40–42]. They reduce the LO cross section of the EW SSWW process by 10–15%, with the correction increasing in magnitude with increasing dilepton and dijet invariant masses. This type of correction is applied only to the EW VBS process, and is not applied to the EFT modifications of the EW VBS processes.

The effects of the EFT dim-6 and dim-8 operators are simulated with MADGRAPH5_aMC@NLO at LO for the EW VBS processes. We do not include them in the background

processes, as they would represent irrelevant contributions. For the dim-6 class, we introduce five bosonic operators acting on the scattering of the W bosons ($\mathcal{Q}_W$, $\mathcal{Q}_{HW}$, $\mathcal{Q}_{HWB}$, $\mathcal{Q}_{H\Box}$, $\mathcal{Q}_{HD}$, where the second subscript in $\mathcal{Q}_{H\Box}$ refers to the d'Alembertian that appears in the operator), two fermionic operators acting on contact interactions between fermions ($\mathcal{Q}_{ll}^{(1)}$, $\mathcal{Q}_{qq}^{(1)}$), and four mixed operators acting on interactions between massive bosons and fermions ($\mathcal{Q}_{Hl}^{(1)}$, $\mathcal{Q}_{Hq}^{(1)}$, $\mathcal{Q}_{Hl}^{(3)}$), defined in the Warsaw basis [17] via the SMEFTSIM [43, 44] package. Some of the background processes of our study could be sensitive to the EFT dim-6 operators under investigation. However, these EFT background effects are better constrained in phase spaces orthogonal to the one covered by this analysis, and do not represent significant non-SM contributions. For the dim-8 category, we introduce nine operators modifying the interaction between two scattering W bosons ($\mathcal{O}_{S0}$, $\mathcal{O}_{S1}$, $\mathcal{O}_{M0}$, $\mathcal{O}_{M1}$, $\mathcal{O}_{M7}$, $\mathcal{O}_{T0}$, $\mathcal{O}_{T1}$, $\mathcal{O}_{T2}$ defined in the Eboli basis [30, 33]. In both cases, the effects of the new EFT operators are evaluated and stored using the MADGRAPH5_aMC@NLO reweighting technique [45], and no LO-to-NLO correction is imposed on their contribution.

The POWHEG v2 generator [46–48] is used to simulate, at NLO accuracy in QCD, the following processes: production of $t\bar{t}$ pairs in which each t quark decays to a b quark and a lepton pair (dileptonic $t\bar{t}$); tW ; Higgs boson production mediated by gluon-gluon and vector-boson fusion; and diboson production.

Production of $t\bar{t}W$, $t\bar{t}Z$, $t\bar{t}\gamma$, triple vector boson, vector boson associated with a Higgs boson, and Drell-Yan background events are simulated at NLO accuracy in QCD using the MADGRAPH5_aMC@NLO generator. The tZq process is simulated at NLO in the four-flavor scheme using MADGRAPH5_aMC@NLO. We generally refer to ZZ , $Z\gamma$, $W\gamma$, WZ , tribosons, associated production of a quark-antiquark top pair with a γ , Z , or W boson, and tZq processes as “others”. Higgs boson production via gluon-gluon fusion and vector-boson fusion, as well as the production of a vector boson in association with a Higgs boson, are collectively referred to as “opposite sign” (OS) since they, alongside DY, contribute to the final event selection when one of the leptons has a misreconstructed charge. OS and DY contributions are treated coherently in the statistical fits described in sections 7 and 8.

The NNPDF3.1 next-to-NLO [49] parton distribution functions (PDFs) are used in the simulation of the background and signal samples. The generators used for signal and background processes are interfaced with the PYTHIA 8.306 [50] program, with the CP5 tune [51], to model parton showering and hadronization.

Additional collisions in the same or adjacent bunch crossings (pileup) are included by superimposing simulated minimum bias interactions onto the hard-scattering process, with a multiplicity distribution matching the one that is observed in the data. Simulated events are propagated through the full GEANT4-based simulation [52] of the CMS detector.

4 Event reconstruction

Events are selected for the signal measurement and the estimation of most of the backgrounds that have passed a trigger requiring the presence of either: one muon with $p_T > 24$ GeV in 2016 and 2018 or $p_T > 27$ GeV in 2017; one electron with $p_T > 27$ GeV in 2016 or $p_T > 32$ GeV in 2017 and 2018. Data are selected for the additional control samples used for background estimation that were recorded with triggers requiring the presence of either

a jet with $p_T > 40$ GeV or $H_T > 350$ GeV, where H_T is the scalar sum of the jet transverse momenta in an event.

Particle candidates are processed with an optimized combination of all subdetector information using the CMS particle-flow (PF) algorithm [53] that reconstructs and identifies each individual particle in the event. The missing transverse momentum vector $\vec{p}_T^{\text{miss}}$ is defined as the projection onto the plane perpendicular to the beam axis of the negative vector sum of the momenta of all reconstructed PF objects in an event. Its magnitude is referred to as p_T^{miss} .

Jets are reconstructed by clustering PF candidates using the anti- k_T jet finding algorithm [54, 55] with a distance parameter of 0.4. Jets are calibrated in the simulation, and separately in data, accounting for energy deposits of neutral particles from pileup and any nonlinear detector response. The effect of pileup is mitigated through a charged-hadron subtraction technique [56] that removes the energy of charged hadrons not originating from the primary vertex (PV) of the event. Corrections to jet energies to account for the detector response are propagated to p_T^{miss} . Jets are required to have $p_T > 30$ GeV, $|\eta| < 5$, and to meet jet quality criteria with measured efficiencies that are almost 100%, for both data and simulated samples [57, 58].

The PV is taken to be the reconstructed vertex with the largest value of summed physics-object p_T^2 , as described in section 9.4.1 of ref. [59].

A deep neural network-based tagger, DEEPJET [60–62], is used to identify jets stemming from the hadronization of b quarks, utilizing information from the tracks, neutral particles, and the secondary vertices within the jet. The efficiency and purity of the resulting “b jets” are classified in terms of various working points. The analysis uses a medium working point that correctly identifies b jets with an efficiency of about 70%, and a loose working point with an efficiency of about 85%. The misidentification rates for gluon or light-flavor quark jets for these two working points are 1.0% and 10%, respectively.

Electrons (muons) are reconstructed by associating a track reconstructed in the tracking detectors with a cluster of energy in the ECAL (track in the muon system). The candidates are required to originate from the PV, pass quality selection criteria, and be isolated from other activity in the event. For this purpose we define a lepton relative isolation variable I_{rel} based on the energy deposited on a cone $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.3$ (0.4) around the electron (muon). Specifically, $I_{\text{rel}} = (E_{\text{ch}} + E_{\text{nh}} + E_{\text{ph}} - 0.5 E_{\text{ch}}^{\text{PU}})/p_T$, where E_{ch} is the transverse energy deposited by charged hadrons from the PV, E_{nh} and E_{ph} are the transverse energies of the neutral hadrons and photons, respectively, and p_T is the electron or muon transverse momentum. The term $0.5 E_{\text{ch}}^{\text{PU}}$ accounts for the contribution of neutral particles from pileup vertices, taken as half the energy of the charged particles from pileup vertices.

The quality criteria for the light lepton selection are based on the isolation from other particles in the event, the impact parameter of the candidate with respect to the PV, and the electromagnetic shower shape in the electron case. These are implemented by their inclusion in multivariate discriminators. In this analysis we make use of both “loose” and “tight” working points of these discriminators. The efficiency for loose (tight) electrons is 90 (80)% [63]. The corresponding efficiencies for muons are 99 (95)% [64].

For a loose electron (muon) we require $p_T > 15$ GeV and $|\eta| < 2.5$ (2.4) and $I_{\text{rel}}^{\text{electron}} < 0.20$ ($I_{\text{rel}}^{\text{muon}} < 0.40$). For a tight electron (muon), the criteria are $p_T > 30$ GeV (35 GeV for electrons in 2017 and 2018), and $I_{\text{rel}}^{\text{electron}} < 0.08$ ($I_{\text{rel}}^{\text{muon}} < 0.15$).

Hadronically decaying τ leptons τ_h are reconstructed from jets using the hadrons-plus-strips algorithm [65], which combines 1 or 3 tracks with energy deposits in the calorimeters to identify the τ decay modes. Neutral pions are reconstructed as strips with dynamic size in the (η, ϕ) plane from reconstructed electrons and photons, where the strip size varies as a function of the p_T of the electron or photon candidate. To further distinguish genuine τ_h decays from jets originating from the hadronization of quarks or gluons, and from electrons or muons, we make use of the DEEPTAU algorithm [66]. Information from all individual reconstructed particles near the τ_h axis is combined with properties of the τ_h candidate and the event to provide separate discriminators against hadronic jets (D_j), electrons (D_e), and muons (D_μ). Similarly to the selection of electrons and muons, we employ a loose set of criteria to select τ_h with $p_T > 30$ GeV, $|\eta| < 2.3$, and satisfying D_j, D_e, D_μ working points for which the genuine τ_h identification efficiencies are 70, 98, and 99.5%, respectively. We also make use of a working point with a tighter D_j threshold, for which the identification efficiency for genuine τ_h is 50%.

5 Analysis strategy and background estimation

The analysis targets the EW VBS production of same-sign WW, with one of the W bosons decaying to a τ lepton and the other into a μ or an e, in association with two jets originating from the scattered incoming partons. Events are first selected by requiring one electron or muon, one τ_h candidate, in each case satisfying the tight identification criteria, no additional loose leptons (e, μ , or τ_h candidates), and at least two jets with a pseudorapidity separation $|\Delta\eta| > 2.5$. Among all the possible jet pairs that satisfy the latter requirement, the pair with the highest invariant mass m_{jj} is chosen. We further define the signal region (SR) and several disjoint control regions (CRs) to estimate and validate the background predictions, as specified in the following paragraphs.

The SR is designed to enhance the yield of the EW VBS signal while minimizing that of the background. Events with a same-sign $\ell\tau_h$ pair, $p_T^{\text{miss}} > 50$ GeV, and $m_{jj} > 500$ GeV are selected. In this region, almost 95% of the background events are contain nonprompt lepton candidates, which arise from jets misreconstructed as e, μ , or τ_h , including genuine leptons from the decays of hadrons within jets. About 2% of background events arise from $Z/\gamma^* + \text{jets}$ and 1% from dileptonic $t\bar{t}$ production. Including both the $e\tau_h$ and $\mu\tau_h$ channels, the SR comprises a total of 1047 observed events, with an expected 30 EW SSWW VBS events, 3 QCD-mediated SSWW VBS events, and 998 background events.

Nonprompt leptons are produced mainly by QCD-mediated multijet, associated W + jets, and hadronic and semileptonic $t\bar{t}$ production. They are estimated from data CRs by the “pass-fail” method described in detail in ref. [67]. This method estimates the probability that a nonprompt lepton passes the tight selection criteria by using a region depleted of prompt leptons to determine transfer factors, functions of their p_T and η . These are then used to calculate the nonprompt contributions in the main SRs and CRs, applying them to auxiliary regions defined as the SRs and main CRs, but requiring that at least one of the light lepton and τ_h pass the loose selection, while failing the tight one.

For this background source, we define a “QCD-enriched” CR and a “nonprompt” validation region (VR). To define both the CR and the VR, the transverse mass $m_T(\ell, p_T^{\text{miss}})$ of the




Figure 2. Distributions in the invariant mass of the dijet system for the data and the pre-fit background prediction for the (left) $e\tau_h$ and (right) $\mu\tau_h$ nonprompt VRs. The stacked filled histograms show the background components, with the overflow count included in the last bin. The expectation for the EW SSWW signal is shown by the dashed red line. The hatched error band shows the bin-by-bin statistical uncertainty. The lower panels show the ratio of data to the total background prediction, with statistical uncertainties indicated by error bars and shading, respectively. In all the panels, the vertical bars represent the statistical uncertainty assigned to the observed number of events.

system comprising the light lepton and p_T^{miss} is introduced as follows:

$$m_T(\ell, p_T^{\text{miss}}) = \sqrt{2p_T^\ell p_T^{\text{miss}} [1 - \cos \Delta\phi]}, \quad (5.1)$$

where $\Delta\phi$ is the azimuthal separation between the lepton momentum vector and $\vec{p}_T^{\text{miss}}$. The QCD-enriched CR is used to perform the first step of the nonprompt-lepton background estimations; it contains events with only one loosely identified lepton (e, μ, τ_h), $p_T^{\text{miss}} < 50$ GeV, and $m_T(\ell, p_T^{\text{miss}}) < 50$ GeV selected from data collected with a jet-based trigger. The nonprompt VR serves to validate the yield estimate from the pass-fail method; it contains events with an SS $\ell\tau_h$ pair and $p_T^{\text{miss}} < 50$ GeV passing the same lepton-based triggers mentioned in section 4. Lepton candidates in this VR arise mainly from $W + \text{jets}$ and QCD multijet production. The orthogonality of the QCD-enriched CR and nonprompt VR is ensured by the fact that the triggers used to define them do not overlap. The data and estimated background m_{jj} distributions are compared in the nonprompt VRs for the electron and muon final states in figure 2. The background yields are evaluated before (“pre-fit”) the maximum likelihood (ML) fit introduced at the end of this section. The plots show that the data generally agree with the prediction within uncertainties.

Finally, we define $t\bar{t}$ and OS/DY CRs to constrain the MC simulations of these background sources. Events with an OS $\ell\tau_h$ pair and no loose b-tagged jets are selected for the OS/DY CR; events with an OS $\ell\tau_h$ pair, at least one “medium” b-tagged jet, and $p_T^{\text{miss}} > 50$ GeV are selected for the $t\bar{t}$ CR. Both the $t\bar{t}$ and OS/DY CR are included in the simultaneous ML fit along with the SR. The normalization of the dileptonic $t\bar{t}$ is allowed to freely float. For the OS+DY background, we apply a normalization uncertainty correlated between the OS/DY CR and the SR, as described in section 6. For the $t\bar{t}$ normalization, the post-fit

Region	1 T ℓ , 1 T τ_h , no L ℓ/τ_h	≥ 2 jets with $ \Delta\eta > 2.5$	SS ℓ, τ_h	$p_T^{\text{miss}} > 50$ GeV	Additional requirements
SR	✓	✓	✓	✓	$m_{jj} > 500$ GeV
nonprompt VR	✓	✓	✓	×	
$t\bar{t}$ CR	✓	✓	×	✓	b-tagged jet (“medium”)
OS/DY CR	✓	✓	×	✓	b-tagged jet veto (“loose”)

Table 1. Definitions of the SR, CRs, and VR defined for this analysis. The ✓ symbol indicates that the requirement described in the column heading is applied in that region, whereas the × symbol means that the opposite selection is applied. The symbols T and L refer to the tight and loose selection rules, respectively. The SR and three CRs (nonprompt, $t\bar{t}$, OS) are selected from an inclusive lepton trigger.

results agree with the pre-fit ones, as the observed value for the $t\bar{t}$ normalization parameter is 1.01 ± 0.04 . A summary of the analysis phase space with the definitions of the SR and CRs is given in table 1, with the exclusion of the QCD-enriched CR.

Because of the large background and complex signal topology, sets of significant features to separate signals and backgrounds are combined in three machine-learning discriminators, each targeting a different signal. The discriminators are the outputs of feed-forward deep neural networks (DNNs). The three DNN models are devised to separate the SM EW VBS (SM DNN), EFT dim-6 (dim-6 DNN), and EFT dim-8 (dim-8 DNN) from the SM background processes. In particular, in order to optimize the range of sensitivity of each of the EFT DNN models, we use a balanced mixture (i.e., an equal combination) of contributions from different EFT operators, which include both linear and quadratic terms, as the signal sample. This allows us to avoid the complexities associated with the use of several single-operator models, while preserving discrimination power across the various operators to which this analysis is mostly sensitive. For the first EFT DNN model the signal is represented by a mixture of EFT linear and quadratic contributions weighted with unity values for c_{HW} and c_W . For the second model a mixture of linear and quadratic contributions from all the dim-8 EFT operators included in the study is considered, with the corresponding Wilson coefficient values approximately equal to the 95% confidence level (CL) limits obtained performing a statistical fit with m_{o1} (defined in eq. (5.3) below) with the strategy outlined in section 8. In both cases, the SM EW VBS contribution is considered part of the background sample. The optimized models consist of 1, 2, and 4 hidden layers. Adam Optimizer [68], early stopping, dropout, and L2 regularization [69] techniques are utilized to avoid the overfitting effect, consisting of excessive adaptation to training data of the DNN model.

The sets of input variables for the DNNs, detailed in table 2, are each constructed with a dedicated optimization that includes some quantities introduced to exploit the particular kinematical properties of the EW VBS SSWW reaction [70]. The agreement of the input variables used to train the DNN models has been thoroughly checked in the CRs introduced at the beginning of this section. First, there are the transverse masses

$$m_{1T}^2 = \left(\sqrt{m_{\tau\ell}^2 + p_T^{\tau\ell^2}} + p_T^{\text{miss}} \right)^2 - |\vec{p}_T^{\tau\ell} + \vec{p}_T^{\text{miss}}|^2, \text{ and} \quad (5.2)$$

$$m_{o1}^2 = \left(p_T^{\tau} + p_T^{\ell} + p_T^{\text{miss}} \right)^2 - |\vec{p}_T^{\tau} + \vec{p}_T^{\ell} + \vec{p}_T^{\text{miss}}|^2. \quad (5.3)$$

The variable m_{1T} is the transverse mass of the $\tau\ell$ system with p_T^{miss} . For the second quantity, m_{o1} , the τ , ℓ momenta, and p_T^{miss} are treated as if coming from a system with a null invariant mass when calculating the transverse mass of the three objects. These two variables are a proxy for the energy of the scattering W boson pair, as well as for the angular distribution of the decay objects coming from that pair. With these quantities it is possible to access direct information on the process of interest for this study, more complete than the invariant mass of the VBS jet pair and the p_T of the leptons. In this way, it is possible to enhance the discrimination of the SSWW VBS signal processes against the background, especially when investigating the sensitivity to EFT contributions, as shown for m_{o1} in figure 3. The three transverse masses introduced in eqs. (5.1), (5.2), and (5.3) are among the most important features for all of the DNNs.

Furthermore, the transverse masses $m_T(\tau_h, \vec{p}_T^{\text{miss}})$ and $m_T(\ell + \tau_h, \vec{p}_T^{\text{miss}})$, defined similarly to $m_T(\ell, \vec{p}_T^{\text{miss}})$ in eq. (5.1), are introduced. Next, we define the event Zeppenfeld variable [71]:

$$z_{\text{event}} = \frac{1}{2} \frac{\eta_\ell - \left| \frac{\eta_{j1} + \eta_{j2}}{2} \right|}{\eta_{j1} - \eta_{j2}} + \frac{1}{2} \frac{\eta_{\tau_h} - \left| \frac{\eta_{j1} + \eta_{j2}}{2} \right|}{\eta_{j1} - \eta_{j2}}, \quad (5.4)$$

introduced to exploit the centrality of the leptons in the EW VBS processes with respect to the scattered VBS jets j_1, j_2 . Finally we add the component $p_{T,lj}^{\text{rel}}$ of the ℓ or τ_h momentum $\vec{p}_l$ perpendicular to the momentum $\vec{p}_j$ of VBS jet j :

$$p_{T,lj}^{\text{rel}} = \frac{|\vec{p}_l \times \vec{p}_j|}{|\vec{p}_j|}. \quad (5.5)$$

This variable evaluates how close a lepton is to the flight direction of a jet. For EW VBS processes, this quantity is expected to be distributed towards values larger than for the other processes, since the light lepton and the τ_h produced by the scattered W bosons are far from the VBS jets. The pre-fit m_{o1} distributions in the SRs for the electron and muon final states are shown in figure 3.

For the measurement of the SM VBS processes under investigation, the statistical analysis is implemented with an ML fit to extract the signal strength taking the asymptotic limit of Wilks's theorem [72] to infer the test statistic's dependency on the signal strength. The ML fit is implemented with the CMS statistical analysis tool COMBINE [73], which is based on the ROOFIT [74] and ROOSTATS [75] frameworks. To validate the results obtained by relying on the asymptotic limit, we perform the same measurement by generating pseudo-experiments for the signal and the background, taking into account their statistical fluctuations. The outcomes from the pseudo-experiments are consistent with the ones returned by applying Wilks's theorem, and in the following, the latter are presented. Data yields in both SRs and CRs are incorporated in the likelihood via Poisson probability density functions. The inputs to the fit are the distributions in the DNN output of the data, the signal, and the backgrounds estimated as described above. The distributions in the SRs and CRs are affected by common sources of systematic uncertainty, described in the next section, and thus their expectations are treated as correlated in the fit.

The statistical analysis for the investigation of the EFT contributions in the EW VBS processes of interest is also based on an ML fit, considering that the expected number of events




Figure 3. Distributions in m_{τ_1} transverse mass for the data and the pre-fit background prediction for the (left) $e\tau_h$ and (right) $\mu\tau_h$ SRs. The stacked filled histograms show the background components, and the overflow count is included in the last bin. The expectations for the EW SSWW signal, the $\mathcal{O}_W$ dim-6 operator with $c_W = 1 \text{ TeV}^{-2}$, and the $\mathcal{Q}_{T1}$ dim-8 operator with $f_{T1} = 1 \text{ TeV}^{-4}$ are shown by the dashed red, blue, and green lines, respectively. For the latter two, the interference with SM and pure EFT contributions are summed together with the SM contribution. The hatched error band shows the bin-by-bin statistical uncertainty. The lower panels show the ratio of data to the total background prediction, with statistical uncertainties indicated by error bars and shading, respectively. In all the panels, the vertical bars represent the statistical uncertainty assigned to the observed number of events.

N_{exp} inherits the quadratic dependence on the EFT Wilson coefficients from the scattering amplitude $|\mathcal{A}_{\text{BSM}}|$ reported in eq. (1.2). When the contribution of a single EFT dim-6 (with c_i Wilson coefficient) or dim-8 (with f_α Wilson coefficient) operator is considered, and all the others are set to null values, the expected number of events can be written as follows:

$$N_{\text{exp}}^i = N_{\text{SM}} + \frac{c_i}{\Lambda^2} N_{\text{Lin}}^i + \frac{c_i^2}{\Lambda^4} N_{\text{Quad}}^i, \quad (5.6)$$

$$N_{\text{exp}}^\alpha = N_{\text{SM}} + \frac{f_\alpha}{\Lambda^4} N_{\text{Lin}}^\alpha + \frac{f_\alpha^2}{\Lambda^8} N_{\text{Quad}}^\alpha, \quad (5.7)$$

where N_{SM} stands for the contribution from the SM processes, N_{Lin} for the one from the interference of the considered EFT operator with the SM EW VBS processes, and N_{Quad} for the pure term produced by the specific operator. In the following, this fit setup will be referred to as a 1D dim-6 or dim-8 EFT study, respectively. When two EFT dim-6 operators with Wilson coefficients c_i, c_j are considered active, and all the other ones are set as negligible, the same quantity reads:

$$N_{\text{exp}}^{i,j} = N_{\text{SM}} + \sum_{k=i,j} \left(\frac{c_k}{\Lambda^2} N_{\text{Lin}}^k + \frac{c_k^2}{\Lambda^4} N_{\text{Quad}}^k \right) + \frac{c_i c_j}{\Lambda^4} N_{\text{Cross}}^{ij}, \quad (5.8)$$

where N_{Cross} represents the interference between the two EFT operators under study. This fit setup will be called a two-dimensional (2D) same-dimension EFT study in the rest of this paper. Finally, when considering one EFT dim-6 and one dim-8 operator with Wilson

Input variable	SM DNN	dim-6 DNN	dim-8 DNN
$\tau_h p_T$	✓	✓	✓
ℓp_T	✓	✓	✓
$\tau_h \eta$		✓	
$\ell \eta$		✓	
leading VBS jet p_T	✓	✓	✓
subleading VBS jet p_T	✓	✓	✓
leading VBS jet mass		✓	✓
subleading VBS jet mass		✓	✓
VBS jet pair $\Delta\phi$		✓	
m_{jj}	✓	✓	
m_{1T}	✓	✓	✓
m_{o1}	✓	✓	✓
$m_T(\tau_h, \vec{p}_T^{\text{miss}})$			✓
$m_T(\ell, \vec{p}_T^{\text{miss}})$	✓	✓	✓
$m_T(\ell + \tau_h, \vec{p}_T^{\text{miss}})$			✓
$p_T^{\text{rel}}(\ell, j_1)$		✓	
$p_T^{\text{rel}}(\ell, j_2)$		✓	
$p_T^{\text{rel}}(\tau_h, j_1)$		✓	
$p_T^{\text{rel}}(\tau_h, j_2)$		✓	
$\Delta\phi(\ell, j_1)$		✓	
$\Delta\phi(\ell, j_2)$		✓	
$\Delta\phi(\tau_h, j_1)$		✓	
$\Delta\phi(\tau_h, j_2)$		✓	
$p_T, \text{leading } \tau_h \text{ track} / p_T, \tau_h$	✓	✓	
z_{event}		✓	

Table 2. List of the input variables for the three DNN models developed in this study. The check mark indicates that the variable is included in the DNN model identified in the column header.

coefficients c_i, f_α to be active, the expected number of events becomes:

$$N_{\text{exp}}^{i,\alpha} = N_{\text{SM}} + \frac{c_i}{\Lambda^2} N_{\text{Lin}}^i + \frac{c_i^2}{\Lambda^4} N_{\text{Quad}}^i + \frac{f_\alpha}{\Lambda^4} N_{\text{Lin}}^\alpha + \frac{f_\alpha^2}{\Lambda^8} N_{\text{Quad}}^\alpha. \quad (5.9)$$

This fit setup will be referred to as a 2D different-dimension EFT study. It neglects the contributions due to the possible interference between the dim-6 and the dim-8 operator, for which there are no clear theoretical predictions. The absence of this type of EFT contribution precludes consideration of any linear correlation between the two different classes of operators. Consequently, within this theoretical framework, our physics interpretation is limited to the independent effects arising from the interplay between two operators of different dimensions. For every EFT statistical study, the signal strength of the SM EW SSWW VBS contribution is fixed to 1.

6 Systematic uncertainties

Systematic uncertainties in the signal and background yields are introduced as log-normal or Gaussian nuisance parameters in the ML fit, both for the measurement of the EW VBS SSWW processes and for the EFT investigations.

The uncertainties determined by the CMS luminosity monitoring are partially correlated among the data sets [76–78], resulting in overall uncertainties of 1.2, 2.3, and 2.5% for the 2016, 2017, and 2018 integrated luminosities, respectively. This uncertainty affects only the integrated yields, not the shapes of the distributions.

Uncertainties at the matrix-element level, which impact both the normalization and shape of the background and signal processes, are evaluated through separate variations of the renormalization and factorization scales. Specifically, the renormalization and factorization scales are varied independently by a factor of 2 and 0.5. The resulting uncertainties are then combined by taking the envelope of these variations relative to the nominal expectation, without considering the combinations where one scale is varied by a factor of 2.0 and the other one by a factor of 0.5 [79, 80]. These uncertainties are considered uncorrelated across different process categories but are correlated across the data-taking years.

The MC samples are generated using a default PDF set, as mentioned in section 3, and event weights corresponding to the 100 PDF alternative set members are also stored, evaluated with the MADGRAPH5_amc@NLO reweighting technique introduced in section 3. These are used to evaluate the PDF systematic uncertainties according to the procedure recommended by the PDF4LHC group [81], which is based on the same strategy explained in the previous paragraph. They are correlated among the data-taking years and processes.

Among the possible theoretical uncertainties affecting MC simulations are those related to the QCD-induced parton-shower modeling. These uncertainties are divided into initial-state radiation (ISR) and final-state radiation (FSR) and are estimated by separately varying each of them by the same up and down factor. The resulting shape distortions induced by the alternative scenarios are used to evaluate the magnitude of the associated uncertainties. They are considered correlated among the data-taking years and processes.

Uncertainties in b tagging and mistagging data-to-simulation scale factors (SFs) are applied to reproduce the corresponding efficiencies measured in the data, and implemented in the ML fit as correlated among the data-taking years.

Systematic uncertainties related to the pileup modeling are introduced as a $\pm 4.6\%$ variation in the total inelastic cross section of 69.2 mb that is used to estimate the data pileup distributions [82]. They are correlated among the data-taking years.

The impact of the uncertainty in the trigger efficiency measurement is estimated by varying the SFs within their uncertainties separately for each data-taking year and final state. This uncertainty is treated as uncorrelated among data-taking years.

In 2016 and 2017, a portion of trigger primitives in the ECAL was associated with the wrong bunch crossing, leading to a trigger mistiming effect and a nonnegligible decrease in the trigger efficiency that is not modeled in the simulated samples [83]. Events have been corrected for this effect with a per-event weight, and the corresponding uncertainties have been propagated throughout the analysis chain. They are correlated among the data-taking years. In the following, we will refer to the effect described in this paragraph as trigger mistiming.

In simulated events, reconstructed four-momenta of all of the jets are simultaneously varied according to the η - and p_T -dependent uncertainties in the jet energy scale; they are correlated among data-taking years. These variations are then propagated to the $\vec{p}_T^{\text{miss}}$. Moreover, to properly evaluate the systematic effect coming from differences in the jet energy resolution between data and simulations, smearing is also applied to the latter by varying the jet resolutions according to their uncertainties; they are uncorrelated among the data-taking years [84]. Because of the high efficiency of the jet quality requirements, no SFs or associated uncertainties for those are applied.

Systematics related to uncertainties affecting unclustered energy in the calorimeters are included in the fit and correlated among data-taking years.

Systematic uncertainties due to SFs used to match the efficiencies in light lepton reconstruction, identification, isolation, and energy scale and resolution as measured in the MC samples with those observed in data, are evaluated by varying the corresponding event weights by the SF uncertainty. They are uncorrelated over the data-taking years.

For the uncertainty arising from charge sign misreconstruction in the $e\tau_h$ channel we assign a uniform uncertainty of 15% to the distributions of the background processes, consistent with the data-background agreement observed in the OS/DY CR.

Statistical uncertainties related to the τ lepton identification SFs and the corrections to their energy scale and resolution [65] induce a systematic effect on the expected signal and background distributions. Their impact is evaluated following a procedure similar to the one applied for light-lepton systematic uncertainties. They are uncorrelated among the data-taking years.

The LO to NLO corrections to the VBS signals come with statistical uncertainties that are propagated to the fit [40–42].

For the signal and background processes estimated from simulation, the precision of the modeling is limited by the event count in the MC samples. The corresponding statistical uncertainties are therefore taken as systematic uncertainties applied to each bin of the corresponding distribution, according to the lite Barlow-Beeston method [85].

The estimate of the nonprompt background is affected by the statistical uncertainties of the auxiliary regions used to measure the transfer factors. This uncertainty is propagated by the lite Barlow-Beeston method [85]. We assign a further 30% normalization uncertainty based on a closure test performed in the nonprompt VR. This uncertainty is treated as uncorrelated among the data-taking years.

In the following, we collectively refer to the statistical uncertainties assigned to the backgrounds, extracted from data CRs or from simulation, as background statistical. It represents the dominant source of overall uncertainty. The impacts of the systematic uncertainties in the signal strength, as extracted from the ML fit, are summarized in table 3.

7 Measurement of the SM SSW VBS processes

We extract values of the signal strength and the statistical significance from the fit of the SM SSW VBS signal, with two separate interpretations. As the primary interpretation, we measure the purely EW signal strength keeping the QCD SSW production contribution fixed to the SM prediction. In the second interpretation, we measure the signal strength

Uncertainty source	$+\Delta\mu$	$-\Delta\mu$
Theory (PDF, scales, ISR, FSR)	+0.21	-0.12
Nonprompt background estimation	+0.062	-0.070
Jet energy scale, resolution, and identification	+0.077	-0.059
Trigger mistiming	+0.071	-0.040
$t\bar{t}$ normalization	+0.048	-0.037
Pileup	+0.042	-0.010
Luminosity	+0.025	-0.026
LO-to-NLO VBS corrections	+0.037	-0.024
τ_h energy scale and identification	+0.040	-0.010
b tagging and mistagging	+0.031	-0.008
Unclustered energy	+0.007	-0.005
Charge misidentification	+0.009	-0.009
Lepton reconstruction, identification, and isolation	+0.002	-0.003
MC sample size (bin-by-bin uncertainty)	+0.28	-0.28
Total systematic	+0.36	-0.32
Data sample size	+0.50	-0.46
Total	+0.62	-0.56

Table 3. The impact of each systematic uncertainty, together with the impact of the data statistical uncertainty, on the signal strength μ , as extracted from the fit to measure the SM SSW VBS signal with the DNN output distributions. Upper and lower uncertainties are given for the various sources.

treating as signal the combined EW and QCD SSWW processes, fixing the ratio between the two contributions to the SM value. For the primary result of the EW signal strength measurement, the post-fit distributions of the output score of the SM DNN (SM DNN output) of the SRs are shown in figure 4 for both the electron and muon flavors.

In figure 4, the data are compared with the background estimated before (pre-fit) and after (post-fit) the simultaneous fit of the SRs and CRs. The pulls shown in the lower panels are defined, for each bin, as

$$\text{Pull} = \frac{n_{\text{data}} - n_{\text{post-fit}}}{\sqrt{\sigma_{\text{data}}^2 - \sigma_{\text{post-fit}}^2}}, \quad (7.1)$$

where n_{data} and $n_{\text{post-fit}}$ are respectively measured event numbers and post-fit background predictions, and σ_{data} and $\sigma_{\text{post-fit}}$ are their corresponding uncertainties; when $\sigma_{\text{data}} < \sigma_{\text{post-fit}}$, the pull is set to 0. The quadratic difference of the uncertainties appearing in the denominator is taken to account for the correlation between the data and the post-fit prediction. The observed (expected) EW signal strength is $1.44_{-0.56}^{+0.63}$ ($1.00_{-0.53}^{+0.60}$), corresponding to a signal significance of 2.7 (1.9) standard deviations. The simultaneous measurement of the EW and QCD-associated SSWW production results in an observed (expected) signal strength equal to $1.43_{-0.54}^{+0.60}$ ($1.00_{-0.51}^{+0.57}$), with a significance of 2.9 (2.0) standard deviations. The largest contribution to the overall uncertainty is the statistical uncertainty of the data, as reported in table 3.

Tabulated results are provided in the HEPData record for this analysis [86].




Figure 4. Distribution of the DNN output for the (left) $e\tau_h$ and (right) $\mu\tau_h$ SR. The data points are overlaid on the post-fit background plus signal (stacked histograms). The overflow is included in the last bin. The middle panels show ratios of the data to the pre-fit background prediction and post-fit background yield in yellow and green, respectively. The corresponding colored bands indicate the systematic component of the uncertainty. The lower panels show the distributions of the pulls, defined in the text. The vertical bars represent the statistical uncertainty assigned to the observed number of events for the upper panel, and its propagation to the quantities shown in the middle panel.

8 Effective field theory interpretation

The sensitivity of the measurement to the dim-6 and dim-8 EFT operators, considered one at a time, is estimated from a likelihood scan performed by varying the corresponding Wilson coefficients as they appear in the quadratic parametrization of the signal yields given by eqs. (5.6) and (5.7). For this part of the study, we use the distributions of the dim-6 and dim-8 DNN outputs, respectively. No significant deviations from the SM predictions are observed. The 68 and 95% confidence intervals on the Wilson coefficients are extracted from the scan and reported in table 4. Although CMS and ATLAS have published dim-8 VBS analyses (as reported in section 1), and ATLAS has set limits on the dim-6 c_W and c_{HWB} Wilson coefficients via the reinterpretation of differential cross section measurements [28], our study for the first time investigates EFT dim-6 effects directly at the reconstruction level. With this strategy, it is possible to tailor the analysis for maximum sensitivity to EFT-induced deviations in the event kinematics, specifically through DNN models optimized towards EFT effects. In addition, in contrast to reinterpretation approaches, reconstruction-level studies can fully capture correlations among multiple dim-6 operators. As a result, we obtain the first detector-level one- and two-dimensional limits on a broad set of EFT dim-6 contributions in VBS processes.

In addition, a 2D likelihood scan is performed over pairs of Wilson coefficients as they appear in eqs. (5.8) and (5.9), exploiting the distributions of m_{o1} , which we find to be the

Wilson coefficient		68% CL interval(s)		95% CL interval	
		Observed	Expected	Observed	Expected
dim-6	$c_u^{(1)}/\Lambda^2$	$[-11.6, 0.045]$	$[-12.9, -8.03] \cup [-2.95, 1.91]$	$[-13.5, 2.11]$	$[-14.6, 3.53]$
	$c_{qq}^{(1)}/\Lambda^2$	$[-0.341, 0.416]$	$[-0.501, 0.576]$	$[-0.605, 0.681]$	$[-0.742, 0.818]$
	c_W/Λ^2	$[-0.513, 0.481]$	$[-0.681, 0.669]$	$[-0.842, 0.818]$	$[-0.987, 0.974]$
	c_{HW}/Λ^2	$[-5.48, 4.31]$	$[-7.00, 6.09]$	$[-8.68, 7.60]$	$[-9.99, 9.05]$
	c_{HWB}/Λ^2	$[-30.7, 89.2]$	$[-41.7, 69.6]$	$[-49.7, 110]$	$[-66.6, 96.4]$
	$c_{H\Box}/\Lambda^2$	$[-12.0, 14.0]$	$[-16.6, 18.1]$	$[-20.9, 22.7]$	$[-24.7, 26.3]$
	c_{HD}/Λ^2	$[-15.3, 31.5]$	$[-24.6, 34.7]$	$[-31.4, 45.5]$	$[-38.2, 48.8]$
	$c_{Hl}^{(1)}/\Lambda^2$	$[-38.2, 39.5]$	$[-28.8, 29.9]$	$[-69.3, 68.3]$	$[-49.4, 49.7]$
	$c_{Hl}^{(3)}/\Lambda^2$	$[-0.045, 8.58]$	$[-1.43, 2.23] \cup [5.88, 9.54]$	$[-1.59, 9.94]$	$[-2.64, 10.8]$
	$c_{Hq}^{(1)}/\Lambda^2$	$[-3.27, 3.44]$	$[-4.53, 4.42]$	$[-5.55, 5.60]$	$[-6.56, 6.44]$
dim-8	f_{T0}/Λ^4	$[-0.774, 0.842]$	$[-1.02, 1.08]$	$[-1.32, 1.38]$	$[-1.52, 1.58]$
	f_{T1}/Λ^4	$[-0.319, 0.381]$	$[-0.426, 0.480]$	$[-0.552, 0.613]$	$[-0.640, 0.695]$
	f_{T2}/Λ^4	$[-0.851, 1.12]$	$[-1.15, 1.37]$	$[-1.51, 1.76]$	$[-1.75, 1.98]$
	f_{M0}/Λ^4	$[-8.07, 7.70]$	$[-9.89, 9.74]$	$[-13.1, 12.8]$	$[-14.6, 14.5]$
	f_{M1}/Λ^4	$[-9.54, 11.15]$	$[-12.5, 13.3]$	$[-16.4, 17.7]$	$[-18.7, 19.6]$
	f_{M7}/Λ^4	$[-17.6, 15.3]$	$[-20.3, 19.2]$	$[-27.6, 25.8]$	$[-29.9, 28.8]$
	f_{S0}/Λ^4	$[-9.60, 9.82]$	$[-11.6, 12.0]$	$[-15.9, 16.1]$	$[-17.4, 17.9]$
	f_{S1}/Λ^4	$[-40.9, 41.3]$	$[-37.4, 38.8]$	$[-60.9, 61.8]$	$[-57.2, 58.6]$
	f_{S2}/Λ^4	$[-40.9, 41.3]$	$[-37.4, 38.8]$	$[-60.9, 61.8]$	$[-57.2, 58.6]$

Table 4. Observed and expected 68 and 95% 1D confidence level (CL) intervals on the Wilson coefficients associated with the EFT dim-6 and dim-8 operators considered. The results reported here are obtained by fixing the Wilson coefficients other than the one of interest to their SM values in the fit procedure. These limits are reported in units of TeV^{-2} (TeV^{-4}) for the dim-6 (dim-8) Wilson coefficients.

most sensitive variable to EFT effects among the kinematic quantities considered. This choice ensures the stability of the 2D fits and at the same time enhances the physical interpretability of the results in the broader context of EFT reinterpretation. For the same-dimension fits, we consider pairs of EFT operators that both modify either the $WW \rightarrow WW$ amplitude or the W pairing with fermions. For the different-dimension fits, we consider pairs that modify the $WW \rightarrow WW$ amplitude. The operators in these pairs typically impact the scattering amplitude with similar magnitudes, when the dim-6 (dim-8) coefficients are set to TeV^{-2} (TeV^{-4}). The results are reported in figures 5 and 6.

For the same-dimension pairs shown in figure 5, combining the effects of two operators leads to a broadening of the 68 and 95% CL intervals extracted with the corresponding one-dimensional fit to the m_{o1} distributions, an effect that is slightly more pronounced for the operators to which this analysis is less sensitive. On average, this effect results in a 10% difference for the observed limits and a 13% difference for the expected limits. In general, the contours exhibit an elliptical shape, with correlation effects driven by the interference




Figure 5. Observed (black) and expected (red) 68 (solid) and 95% (dashed) CL contours for $-2 \ln \Delta \mathcal{L}$ as functions of the reported dim-6 bosonic (upper two rows) and mixed (lower row) Wilson coefficient pairs. When there are two contours for the same CL value, the constrained set of Wilson coefficient values is represented by the area between the two of them if they are concentric, otherwise it consists of the internal areas of the contours.

between the operators in each pair, leading to either negative or positive linear correlations that tilt the axes of the ellipses. Additionally, for some operators, the interference with the SM contribution is more significant than the quadratic term, resulting in a double-minima structure in the two-dimensional $-2 \ln \Delta \mathcal{L}$ profiles in the direction of the Wilson coefficient




Figure 6. Observed (black) and expected (red) 68 (solid) and 95% (dashed) CL contours for $-2 \ln \Delta\mathcal{L}$ as functions of the reported (dim-6, dim-8) Wilson coefficient pairs.

associated with the operator introducing such effect. For the observed contours, these effects are further enhanced by data fluctuations that the EFT operators can accommodate. These effects, either individually or in combination, are visible in the upper right, middle right, and lower panels of figure 5, where operators $\mathcal{Q}_{HWB}$, $\mathcal{Q}_{H\Box}$, $\mathcal{Q}_{Hl}^{(3)}$, $\mathcal{Q}_{Hq}^{(1)}$ are involved. Notably, they are most pronounced for the $\mathcal{Q}_{HWB}$, $\mathcal{Q}_{H\Box}$ pair (middle right of figure 5), where significant mutual interference and relatively more important linear terms (compared to the quadratic contributions at unity Wilson coefficient values) are observed.

For the different-dimension pairs, the results indicate that adding the linear and quadratic dim-8 contributions to those of a given dim-6 operator leads to a broadening of the one-dimensional 68% and 95% confidence intervals for the dim-6 operators extracted from the same $m_{\phi 1}$ distributions, regardless of the specific dim-8 operator considered. This broadening is quantified by comparing the 1D intervals obtained from individual fits with the intervals derived from the 2D limits for one coefficient when the other is fixed at its null value. On average, this effect on the dim-6 Wilson coefficients results in a 10% difference in the observed limits and a 12% difference in the expected limits. A similar broadening is observed for the dim-8 Wilson coefficients in the pairs displayed in figure 6, with average differences of 10% for the observed limits and 13% for the expected limits. Moreover, by setting $c/\Lambda^2 = 1$ for the dim-6 operator and $f/\Lambda^4 = 1$ for the dim-8 operator, we can directly compare the effects of the dim-6 and dim-8 operators in each pair and relate them to the shape of the corresponding 2D contours. When the contributions of these operators are comparable, the 68% and 95% confidence contours in the 2D fit appear more circular; conversely, if the effect of the dim-8 operator is negligible relative to that of the dim-6 operator, the contours tend to be more rectangular. For example, in the bottom left of figure 6 the 2D fit for operators $\mathcal{Q}_W$ and $\mathcal{O}_{T0}$ shows circular contours when the effects are similar, whereas the plot on top left of figure 6 illustrates that when the dim-8 contribution of the $\mathcal{O}_{S0}$ operator is negligible compared to the dim-6 contribution of the operator $\mathcal{Q}_W$, the contours become rectangular. In addition, the contour does not show evidence of linear correlations, as the possible interference between the dim-6 and the dim-8 operators in the pair is neglected in this study.

These considerations lead to the conclusion that, within the EFT framework, the one-dimensional constraints on a Wilson coefficient associated with a given operator could be biased by neglect of the contributions of other operators arising in the same physical process and at the same or different power of Λ_{BSM} , with the same or different dimension. This bias is not an experimental artifact but stems from the fundamental structure of the SMEFT model. Contributions from dim-6 and dim-8 operators can interfere nontrivially and can be visible at the reconstruction level. This finding aligns with the EFT theory community's recommendation to perform studies that simultaneously investigate dim-6 and dim-8 contributions and to encompass as wide a range as possible of operator classes and processes [87–91].

It is worth noting that the 2D EFT fits with two dim-6 operators represent the first results of this type for an analysis investigating EW VBS processes, and the 2D EFT fits with one dim-6 and one dim-8 operator represent the first results ever for combinations of different-dimension EFT operators in the same physics processes.

9 Summary

Electroweak (EW) production of a same-sign W boson pair in association with two jets, with a hadronically decaying τ lepton in the final state, is investigated for the first time, together with an interpretation of possible deviations from the standard model expectations in terms of effective field theory (EFT) operators of dimension 6 and 8. The analysis is performed with a sample of proton-proton collisions at $\sqrt{s} = 13$ TeV recorded by the CMS experiment at the CERN LHC in 2016–2018, corresponding to an integrated luminosity of

138 fb^{-1} . Events are selected with the requirement of one τ lepton together with one light lepton (e or μ) of the same sign, missing transverse momentum, and two jets with large pseudorapidity separation and large dijet invariant mass. Deep neural network algorithms are employed to discriminate different types of signal events from the main backgrounds, significantly boosting the sensitivity of the search.

The amplitude for same-sign WW production includes terms that account for strong interactions between partons with W boson radiation. A small fraction of these QCD-mediated events falls within the acceptance of the search. The measured cross section for EW same-sign WW scattering, extracted with the QCD-mediated amplitudes fixed to the standard model (SM) expectations, is $1.44^{+0.63}_{-0.56}$ times the SM prediction. The observed (expected) significance of the EW signal is 2.7 (1.9) standard deviations. A measurement of the combined EW and residual QCD-mediated contributions yields an observed (expected) significance of 2.9 (2.0) standard deviations.

Also presented are the limits on dimension-6 EFT operator contributions, including both one operator and two operators active at the same time, extracted with a reconstruction-level strategy for the first time in vector boson scattering. This is the first study of the combined effects of EFT operators with different dimensions, showing that focusing on one dimensionality can lead to an overestimate of the sensitivity to the corresponding EFT operator class, and that the contributions of terms combining operators with different dimensions should not be neglected, in accordance with the latest recommendations from the EFT theory community.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science — EOS” — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany’s Excellence Strategy — EXC 2121 “Quantum Universe” — 390833306, and under project number 400140256 — GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC — National Research Center for High Performance Computing, Big Data and Quantum Computing and FAIR — Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

Data Availability Statement. Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the [CMS data preservation, re-use and open access policy](#)

Code Availability Statement. The CMS core software is publicly available on [GitHub](#).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](#)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS and CMS collaborations, *Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **08** (2016) 045 [[arXiv:1606.02266](#)] [[INSPIRE](#)].
- [2] P.W. Higgs, *Broken symmetries, massless particles and gauge fields*, *Phys. Lett.* **12** (1964) 132 [[INSPIRE](#)].
- [3] P.W. Higgs, *Broken Symmetries and the Masses of Gauge Bosons*, *Phys. Rev. Lett.* **13** (1964) 508 [[INSPIRE](#)].
- [4] F. Englert and R. Brout, *Broken Symmetry and the Mass of Gauge Vector Mesons*, *Phys. Rev. Lett.* **13** (1964) 321 [[INSPIRE](#)].
- [5] J. Chang, K. Cheung, C.-T. Lu and T.-C. Yuan, *WW scattering in the era of post-Higgs-boson discovery*, *Phys. Rev. D* **87** (2013) 093005 [[arXiv:1303.6335](#)] [[INSPIRE](#)].
- [6] W. Kilian, T. Ohl, J. Reuter and M. Sekulla, *High-Energy Vector Boson Scattering after the Higgs Discovery*, *Phys. Rev. D* **91** (2015) 096007 [[arXiv:1408.6207](#)] [[INSPIRE](#)].
- [7] C. Garcia-Garcia, M. Herrero and R.A. Morales, *Unitarization effects in EFT predictions of WZ scattering at the LHC*, *Phys. Rev. D* **100** (2019) 096003 [[arXiv:1907.06668](#)] [[INSPIRE](#)].
- [8] ATLAS collaboration, *Evidence for Electroweak Production of $W^\pm W^\pm jj$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector*, *Phys. Rev. Lett.* **113** (2014) 141803 [[arXiv:1405.6241](#)] [[INSPIRE](#)].
- [9] CMS collaboration, *Observation of electroweak production of same-sign W boson pairs in the two jet and two same-sign lepton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **120** (2018) 081801 [[arXiv:1709.05822](#)] [[INSPIRE](#)].
- [10] R. Covarelli, M. Pellen and M. Zaro, *Vector-Boson scattering at the LHC: Unraveling the electroweak sector*, *Int. J. Mod. Phys. A* **36** (2021) 2130009 [[arXiv:2102.10991](#)] [[INSPIRE](#)].
- [11] C.-W. Chiang, N.G. Deshpande, X.-G. He and J. Jiang, *The Family $SU(2)_l \times SU(2)_h \times U(1)$ Model*, *Phys. Rev. D* **81** (2010) 015006 [[arXiv:0911.1480](#)] [[INSPIRE](#)].
- [12] A. Hayreter, X.-G. He and G. Valencia, *LHC constraints on W' , Z' that couple mainly to third generation fermions*, *Eur. Phys. J. C* **80** (2020) 912 [[arXiv:1912.06344](#)] [[INSPIRE](#)].
- [13] R. Calabrese et al., *Top-flavor scheme in the context of W' searches at LHC*, *Phys. Rev. D* **104** (2021) 055006 [[arXiv:2104.06720](#)] [[INSPIRE](#)].
- [14] CMS collaboration, *Search for a W' boson decaying to a τ lepton and a neutrino in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **792** (2019) 107 [[arXiv:1807.11421](#)] [[INSPIRE](#)].
- [15] ATLAS collaboration, *Search for high-mass resonances in final states with a τ -lepton and missing transverse momentum with the ATLAS detector*, *Phys. Rev. D* **109** (2024) 112008 [[arXiv:2402.16576](#)] [[INSPIRE](#)].
- [16] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl. Phys. B* **268** (1986) 621 [[INSPIRE](#)].
- [17] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **10** (2010) 085 [[arXiv:1008.4884](#)] [[INSPIRE](#)].
- [18] CMS collaboration, *Study of Exclusive Two-Photon Production of W^+W^- in pp Collisions at $\sqrt{s} = 7$ TeV and Constraints on Anomalous Quartic Gauge Couplings*, *JHEP* **07** (2013) 116 [[arXiv:1305.5596](#)] [[INSPIRE](#)].

- [19] CMS collaboration, *Evidence for exclusive $\gamma\gamma \rightarrow W^+W^-$ production and constraints on anomalous quartic gauge couplings in pp collisions at $\sqrt{s} = 7$ and 8 TeV*, *JHEP* **08** (2016) 119 [[arXiv:1604.04464](#)] [[INSPIRE](#)].
- [20] ATLAS collaboration, *Measurement of $W^\pm W^\pm$ vector-boson scattering and limits on anomalous quartic gauge couplings with the ATLAS detector*, *Phys. Rev. D* **96** (2017) 012007 [[arXiv:1611.02428](#)] [[INSPIRE](#)].
- [21] ATLAS collaboration, *Search for anomalous electroweak production of WW/WZ in association with a high-mass dijet system in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. D* **95** (2017) 032001 [[arXiv:1609.05122](#)] [[INSPIRE](#)].
- [22] ATLAS collaboration, *Measurement of $WW/WZ \rightarrow \ell\nu qq'$ production with the hadronically decaying boson reconstructed as one or two jets in pp collisions at $\sqrt{s} = 8$ TeV with ATLAS, and constraints on anomalous gauge couplings*, *Eur. Phys. J. C* **77** (2017) 563 [[arXiv:1706.01702](#)] [[INSPIRE](#)].
- [23] CMS collaboration, *Measurement of electroweak WZ boson production and search for new physics in $WZ +$ two jets events in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **795** (2019) 281 [[arXiv:1901.04060](#)] [[INSPIRE](#)].
- [24] CMS collaboration, *Search for anomalous electroweak production of vector boson pairs in association with two jets in proton-proton collisions at 13 TeV*, *Phys. Lett. B* **798** (2019) 134985 [[arXiv:1905.07445](#)] [[INSPIRE](#)].
- [25] ATLAS collaboration, *Measurements of $W^+W^- + \geq 1$ jet production cross-sections in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **06** (2021) 003 [[arXiv:2103.10319](#)] [[INSPIRE](#)].
- [26] CMS and TOTEM collaborations, *Search for high-mass exclusive $\gamma\gamma \rightarrow WW$ and $\gamma\gamma \rightarrow ZZ$ production in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2023) 229 [[arXiv:2211.16320](#)] [[INSPIRE](#)].
- [27] ATLAS collaboration, *Measurement and interpretation of same-sign W boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **04** (2024) 026 [[arXiv:2312.00420](#)] [[INSPIRE](#)].
- [28] ATLAS collaboration, *Differential cross-section measurements of the production of four charged leptons in association with two jets using the ATLAS detector*, *JHEP* **01** (2024) 004 [[arXiv:2308.12324](#)] [[INSPIRE](#)].
- [29] J.J. Ethier, R. Gomez-Ambrosio, G. Magni and J. Rojo, *SMEFT analysis of vector boson scattering and diboson data from the LHC Run II*, *Eur. Phys. J. C* **81** (2021) 560 [[arXiv:2101.03180](#)] [[INSPIRE](#)].
- [30] O.J.P. Éboli and M.C. Gonzalez-Garcia, *Classifying the bosonic quartic couplings*, *Phys. Rev. D* **93** (2016) 093013 [[arXiv:1604.03555](#)] [[INSPIRE](#)].
- [31] A. Falkowski et al., *Anomalous Triple Gauge Couplings in the Effective Field Theory Approach at the LHC*, *JHEP* **02** (2017) 115 [[arXiv:1609.06312](#)] [[INSPIRE](#)].
- [32] R. Bellan et al., *A sensitivity study of VBS and diboson WW to dimension-6 EFT operators at the LHC*, *JHEP* **05** (2022) 039 [[arXiv:2108.03199](#)] [[INSPIRE](#)].
- [33] E.S. Almeida, O.J.P. Éboli and M.C. Gonzalez-Garcia, *Unitarity constraints on anomalous quartic couplings*, *Phys. Rev. D* **101** (2020) 113003 [[arXiv:2004.05174](#)] [[INSPIRE](#)].
- [34] CMS collaboration, *The CMS Experiment at the CERN LHC*, 2008 *JINST* **3** S08004 [[INSPIRE](#)].

- [35] CMS collaboration, *Description and Performance of Track and Primary-Vertex Reconstruction with the CMS Tracker*, **2014 JINST** **9** P10009 [[arXiv:1405.6569](#)] [[INSPIRE](#)].
- [36] CMS TRACKER GROUP collaboration, *The CMS Phase-1 Pixel Detector Upgrade*, **2021 JINST** **16** P02027 [[arXiv:2012.14304](#)] [[INSPIRE](#)].
- [37] CMS collaboration, *Track impact parameter resolution for the full pseudo rapidity coverage in the 2017 dataset with the CMS Phase-1 Pixel detector*, **CMS-DP-2020-049** (2020).
- [38] CMS collaboration, *The CMS trigger system*, **2017 JINST** **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [39] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, **JHEP** **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [40] CMS collaboration, *Measurements of production cross sections of WZ and same-sign WW boson pairs in association with two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV*, **Phys. Lett. B** **809** (2020) 135710 [[arXiv:2005.01173](#)] [[INSPIRE](#)].
- [41] B. Biedermann, A. Denner and M. Pellen, *Large electroweak corrections to vector-boson scattering at the Large Hadron Collider*, **Phys. Rev. Lett.** **118** (2017) 261801 [[arXiv:1611.02951](#)] [[INSPIRE](#)].
- [42] B. Biedermann, A. Denner and M. Pellen, *Complete NLO corrections to W^+W^+ scattering and its irreducible background at the LHC*, **JHEP** **10** (2017) 124 [[arXiv:1708.00268](#)] [[INSPIRE](#)].
- [43] I. Brivio, *SMEFTsim 3.0 — a practical guide*, **JHEP** **04** (2021) 073 [[arXiv:2012.11343](#)] [[INSPIRE](#)].
- [44] I. Brivio, Y. Jiang and M. Trott, *The SMEFTsim package, theory and tools*, **JHEP** **12** (2017) 070 [[arXiv:1709.06492](#)] [[INSPIRE](#)].
- [45] O. Mattelaer, *On the maximal use of Monte Carlo samples: re-weighting events at NLO accuracy*, **Eur. Phys. J. C** **76** (2016) 674 [[arXiv:1607.00763](#)] [[INSPIRE](#)].
- [46] P. Nason, *A New method for combining NLO QCD with shower Monte Carlo algorithms*, **JHEP** **11** (2004) 040 [[hep-ph/0409146](#)] [[INSPIRE](#)].
- [47] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with Parton Shower simulations: the POWHEG method*, **JHEP** **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [48] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, **JHEP** **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [49] NNPDF collaboration, *Parton distributions from high-precision collider data*, **Eur. Phys. J. C** **77** (2017) 663 [[arXiv:1706.00428](#)] [[INSPIRE](#)].
- [50] C. Bierlich et al., *A comprehensive guide to the physics and usage of PYTHIA 8.3*, **SciPost Phys. Codeb.** **2022** (2022) 8 [[arXiv:2203.11601](#)] [[INSPIRE](#)].
- [51] CMS collaboration, *Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements*, **Eur. Phys. J. C** **80** (2020) 4 [[arXiv:1903.12179](#)] [[INSPIRE](#)].
- [52] GEANT4 collaboration, *GEANT4 — A Simulation Toolkit*, **Nucl. Instrum. Meth. A** **506** (2003) 250 [[INSPIRE](#)].
- [53] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, **2017 JINST** **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].

- [54] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [55] M. Cacciari, G.P. Salam and G. Soyez, *FastJet User Manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [56] M. Cacciari and G.P. Salam, *Pileup subtraction using jet areas*, *Phys. Lett. B* **659** (2008) 119 [[arXiv:0707.1378](#)] [[INSPIRE](#)].
- [57] CMS collaboration, *Jet Performance in pp Collisions at 7 TeV*, CMS-PAS-JME-10-003, CERN, Geneva (2010).
- [58] CMS collaboration, *Jet algorithms performance in 13 TeV data*, CMS-PAS-JME-16-003, CERN, Geneva (2017).
- [59] D. Contardo et al., *Technical Proposal for the Phase-II Upgrade of the CMS Detector*, CERN-LHCC-2015-010 (2015) [[DOI:10.17181/CERN.VU8I.D59J](#)].
- [60] E. Bols et al., *Jet Flavour Classification Using DeepJet*, 2020 *JINST* **15** P12012 [[arXiv:2008.10519](#)] [[INSPIRE](#)].
- [61] CMS collaboration, *Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV*, 2018 *JINST* **13** P05011 [[arXiv:1712.07158](#)] [[INSPIRE](#)].
- [62] CMS collaboration, *Measurement of the single top quark and antiquark production cross sections in the t channel and their ratio in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **800** (2020) 135042 [[arXiv:1812.10514](#)] [[INSPIRE](#)].
- [63] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, 2021 *JINST* **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [64] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, 2018 *JINST* **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [65] CMS collaboration, *Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV*, 2018 *JINST* **13** P10005 [[arXiv:1809.02816](#)] [[INSPIRE](#)].
- [66] CMS collaboration, *Identification of hadronic tau lepton decays using a deep neural network*, 2022 *JINST* **17** P07023 [[arXiv:2201.08458](#)] [[INSPIRE](#)].
- [67] CMS collaboration, *Measurement of Higgs Boson Production and Properties in the WW Decay Channel with Leptonic Final States*, *JHEP* **01** (2014) 096 [[arXiv:1312.1129](#)] [[INSPIRE](#)].
- [68] D.P. Kingma and J. Ba, *Adam: A Method for Stochastic Optimization*, [arXiv:1412.6980](#) [[INSPIRE](#)].
- [69] I. Goodfellow, Y. Bengio and A. Courville, *Deep learning*, MIT Press (2016), <http://www.deeplearningbook.org>.
- [70] A.J. Barr et al., *Guide to transverse projections and mass-constraining variables*, *Phys. Rev. D* **84** (2011) 095031 [[arXiv:1105.2977](#)] [[INSPIRE](#)].
- [71] D.L. Rainwater, R. Szalapski and D. Zeppenfeld, *Probing color singlet exchange in $Z +$ two jet events at the CERN LHC*, *Phys. Rev. D* **54** (1996) 6680 [[hep-ph/9605444](#)] [[INSPIRE](#)].
- [72] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [Erratum *ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].

- [73] CMS collaboration, *The CMS Statistical Analysis and Combination Tool: Combine*, *Comput. Softw. Big Sci.* **8** (2024) 19 [[arXiv:2404.06614](#)] [[INSPIRE](#)].
- [74] W. Verkerke and D.P. Kirkby, *The RooFit toolkit for data modeling*, *eConf C* **0303241** (2003) MOLT007 [[physics/0306116](#)] [[INSPIRE](#)].
- [75] L. Moneta et al., *The RooStats Project*, *PoS ACAT2010* (2010) 057 [[arXiv:1009.1003](#)] [[INSPIRE](#)].
- [76] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [[arXiv:2104.01927](#)] [[INSPIRE](#)].
- [77] CMS collaboration, *CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-17-004*, CERN, Geneva (2018).
- [78] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-18-002*, CERN, Geneva (2019).
- [79] M. Cacciari et al., *The t anti- t cross-section at 1.8-TeV and 1.96-TeV: A Study of the systematics due to parton densities and scale dependence*, *JHEP* **04** (2004) 068 [[hep-ph/0303085](#)] [[INSPIRE](#)].
- [80] S. Catani, D. de Florian, M. Grazzini and P. Nason, *Soft gluon resummation for Higgs boson production at hadron colliders*, *JHEP* **07** (2003) 028 [[hep-ph/0306211](#)] [[INSPIRE](#)].
- [81] J. Butterworth et al., *PDF4LHC recommendations for LHC Run II*, *J. Phys. G* **43** (2016) 023001 [[arXiv:1510.03865](#)] [[INSPIRE](#)].
- [82] CMS collaboration, *Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2018) 161 [[arXiv:1802.02613](#)] [[INSPIRE](#)].
- [83] CMS collaboration, *Performance of the CMS electromagnetic calorimeter in pp collisions at $\sqrt{s} = 13$ TeV*, *2024 JINST* **19** P09004 [[arXiv:2403.15518](#)] [[INSPIRE](#)].
- [84] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, *2017 JINST* **12** P02014 [[arXiv:1607.03663](#)] [[INSPIRE](#)].
- [85] J.S. Conway, *Incorporating Nuisance Parameters in Likelihoods for Multisource Spectra*, in the proceedings of the *PHYSTAT 2011*, Geneva, Switzerland (2011) [[DOI:10.5170/CERN-2011-006.115](#)] [[arXiv:1103.0354](#)] [[INSPIRE](#)].
- [86] HEPData record for this analysis, HEPData (collection) (2024), <https://doi.org/10.17182/hepdata.154440>.
- [87] C. Hays, A. Martin, V. Sanz and J. Setford, *On the impact of dimension-eight SMEFT operators on Higgs measurements*, *JHEP* **02** (2019) 123 [[arXiv:1808.00442](#)] [[INSPIRE](#)].
- [88] S. Dawson, M. Forsslund and M. Schnubel, *SMEFT matching to Z' models at dimension eight*, *Phys. Rev. D* **110** (2024) 015002 [[arXiv:2404.01375](#)] [[INSPIRE](#)].
- [89] T. Corbett et al., *Impact of dimension-eight SMEFT operators in the electroweak precision observables and triple gauge couplings analysis in universal SMEFT*, *Phys. Rev. D* **107** (2023) 115013 [[arXiv:2304.03305](#)] [[INSPIRE](#)].
- [90] CMS collaboration, *Combined effective field theory interpretation of Higgs boson, electroweak vector boson, top quark, and multi-jet measurements*, [arXiv:2504.02958](#) [[INSPIRE](#)].
- [91] ATLAS collaboration, *Combined effective field theory interpretation of Higgs boson and weak boson production and decay with ATLAS data and electroweak precision observables*, *ATL-PHYS-PUB-2022-037*, CERN, Geneva (2022).

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan ¹

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic ,
P.S. Hussain , M. Jeitler ², N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck ²,
R. Schöfbeck , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz ²

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans , J. D’Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen ,
S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat ³, G.P. Van Onsem ,
S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios ,
J. Jaramillo , A. Khalilzadeh , F.A. Khan , K. Lee , M. Mahdavihorrani , A. Malara ,
S. Paredes , M.A. Shahzad , L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx ,
K. Mota Amarilo , A. Samalan , K. Skovpen , N. Van Den Bossche , J. van der Linden ,
L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere ,
I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre , J. Lidrych ,
P. Mastrapasqua , T.T. Tran , S. Wertz 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Alves Gallo Pereira , E. Coelho , G. Correia Silva , C. Hensel ,
T. Menezes De Oliveira , C. Mora Herrera ⁴, A. Moraes , P. Rebello Teles , M. Soeiro ,
A. Vilela Pereira ⁴

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho ,
J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira ⁶, D. De Jesus Damiao ,
S. Fonseca De Souza , R. Gomes De Souza , M. Macedo , J. Martins ⁷, L. Mundim ,
H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores ,
I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova ,
G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, K. Yi ^{8,9}

Institute of High Energy Physics, Beijing, China

G.M. Chen ¹⁰, H.S. Chen ¹⁰, M. Chen ¹⁰, F. Iemmi , C.H. Jiang, A. Kapoor ¹¹, H. Liao ,
Z.-A. Liu ¹², R. Sharma ¹³, J.N. Song¹², J. Tao , C. Wang¹⁰, J. Wang , Z. Wang¹⁰, H. Zhang ,
J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , S. Deng , B. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao,
S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang, L. Zhang , Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹⁴, B. Li, J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

X. Gao¹⁵, Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin, C. Lu, M. Xiao

Universidad de Los Andes, Bogota, Colombia

C. Avila, D.A. Barbosa Trujillo, A. Cabrera, C. Florez, J. Fraga, J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

F. Ramirez, C. Rendón, M. Rodriguez, A.A. Ruales Barbosa, J.D. Ruiz Alvarez

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic, N. Godinovic, D. Lelas, A. Sculac

University of Split, Faculty of Science, Split, Croatia

M. Kovac, A. Petkovic, T. Sculac

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa, V. Brigljevic, B.K. Chitroda, D. Ferencek, K. Jakovcic, S. Mishra, A. Starodumov¹⁶, T. Susa

University of Cyprus, Nicosia, Cyprus

A. Attikis, K. Christoforou, A. Hadjiagapiou, C. Leonidou, J. Mousa, C. Nicolaou, L. Paizanos, F. Ptochos, P.A. Razis, H. Rykaczewski, H. Saka, A. Stepennov

Charles University, Prague, Czech Republic

M. Finger, M. Finger Jr., A. Kveton

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

A.A. Abdelalim^{17,18}, S. Elgammal¹⁹, A. Ellithi Kamel²⁰

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M. Abdullah Al-Mashad, M.A. Mahmoud

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht, M. Kadastik, T. Lange, S. Nandan, C. Nielsen, J. Pata, M. Raidal, L. Tani, C. Veelken

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann, K. Osterberg, M. Voutilainen

Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew ,
K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , L. Martikainen ,
M. Myllymäki , M.m. Rantanen , H. Siikonen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour ,
P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi ,
L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro ²¹, P. Simkina , M. Titov ,
M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , F. Damas ,
O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert ,
R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen ,
C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza ,
A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram ²², J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert ,
C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet ,
G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo ,
P. Depasse , C. Dozen ²³, H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch ,
C. Greenberg , G. Grenier , B. Ille , E. Jourdhuy, I.B. Laktineh, M. Lethuillier , L. Mirabito,
S. Perries, A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

I. Lomidze , T. Toriashvili ²⁴, Z. Tsamalaidze ²⁵

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser ,
A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann ,
M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung 

M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo ²⁶, S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman ²⁷, K. Borras ²⁸, A. Campbell , A. Cardini , C. Cheng , F. Colombina , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo ²⁷, A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaeck , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁹, W. Lohmann ³⁰, F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otariid, J. Park , E. Ranken , A. Raspereza ²⁸, D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham ^{31,28}, S. Schnake ²⁸, P. Schütze , C. Schwanenberger ²⁷, D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford , F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁸, C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoii , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ³², R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , J.M. Lawhorn , M. Link, A. Lintuluoto , B. Maier , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis , A. Papadopoulos³², A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, Z. Painesis , I. Papavergou , I. Paraskevas , N. Saoulidou ,
K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos ,
E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas ,
P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath ^{33,34}, K. Márton, A.J. Rádl ³⁵, F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkúti ³⁶, M.M.A. Gadallah ³⁷, Á. Kadlecik , P. Major ,
G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo ³⁶, F. Nemes ³⁶, T. Novak 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra ³⁸, A. Kaur ,
A. Kaur , H. Kaur , M. Kaur , S. Kumar , K. Sandeep , T. Sheokand, J.B. Singh ,
A. Singla 

University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar ,
M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman ³⁹, S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor ,
S. Kamble , J.R. Komaragiri ⁴⁰, D. Kumar ⁴⁰, P.R. Pujahari , N.R. Saha , A. Sharma ,
A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , B. Parida , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar ,
G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati , C. Kar , D. Maity , P. Mal , T. Mishra ,
V.K. Muraleedharan Nair Bindhu , K. Naskar , A. Nayak , S. Nayak, K. Pal , P. Sadangi,
S.K. Swain , S. Varghese , D. Vats 

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya , A. Alpana , S. Dube , B. Gomber , P. Hazarika , B. Kansal , A. Laha ,
B. Sahu , S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi , A. Jafari , M. Zeinali 

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie ,
M. Mohammadi Najafabadi , S. Tizchang 

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia ^{a,b}, A. Colaleo ^{a,b}, D. Creanza ^{a,c}, B. D'Anzi ^{a,b}, N. De Filippis ^{a,c},
M. De Palma ^{a,b}, W. Elmetenawee ^{a,b,17}, L. Fiore ^a, G. Iaselli ^{a,c}, L. Longo ^a, M. Louka^{a,b},
G. Maggi ^{a,c}, M. Maggi ^a, I. Margjeka ^a, V. Mastrapasqua ^{a,b}, S. My ^{a,b}, S. Nuzzo ^{a,b},
A. Pellecchia ^{a,b}, A. Pompili ^{a,b}, G. Pugliese ^{a,c}, R. Radogna ^{a,b}, D. Ramos ^a, A. Ranieri ^a,
L. Silvestris ^a, F.M. Simone ^{a,c}, Ü. Sözbilir ^a, A. Stamerra ^{a,b}, D. Troiano ^{a,b}, R. Venditti ^{a,b},
P. Verwilligen ^a, A. Zaza ^{a,b}

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi ^a, C. Battilana ^{a,b}, D. Bonacorsi ^{a,b}, P. Capiluppi ^{a,b}, A. Castro ^{†,a,b},
F.R. Cavallo ^a, M. Cuffiani ^{a,b}, G.M. Dallavalle ^a, T. Diotallevi ^{a,b}, F. Fabbri ^a,
A. Fanfani ^{a,b}, D. Fasanella ^a, P. Giacomelli ^a, L. Giommi ^{a,b}, C. Grandi ^a, L. Guiducci ^{a,b},
S. Lo Meo ^{a,50}, M. Lorusso ^{a,b}, L. Lunerti ^a, S. Marcellini ^a, G. Masetti ^a, F.L. Navarria ^{a,b},
G. Paggi ^{a,b}, A. Perrotta ^a, F. Primavera ^{a,b}, A.M. Rossi ^{a,b}, S. Rossi Tisbeni ^{a,b},
T. Rovelli ^{a,b}, G.P. Siroli ^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa ^{a,b,51}, A. Di Mattia ^a, A. Lapertosa ^a, R. Potenza^{a,b}, A. Tricomi ^{a,b,51}, C. Tuve ^{a,b}

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras^{Ⓘ^a}, G. Barbagli^{Ⓘ^a}, G. Bardelli^{Ⓘ^{a,b}}, B. Camaiani^{Ⓘ^{a,b}}, A. Cassese^{Ⓘ^a}, R. Ceccarelli^{Ⓘ^a}, V. Ciulli^{Ⓘ^{a,b}}, C. Civinini^{Ⓘ^a}, R. D'Alessandro^{Ⓘ^{a,b}}, E. Focardi^{Ⓘ^{a,b}}, T. Kello^{Ⓘ^a}, G. Latino^{Ⓘ^{a,b}}, P. Lenzi^{Ⓘ^{a,b}}, M. Lizzo^{Ⓘ^a}, M. Meschini^{Ⓘ^a}, S. Paoletti^{Ⓘ^a}, A. Papanastassiou^{a,b}, G. Sguazzoni^{Ⓘ^a}, L. Viliani^{Ⓘ^a}

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi[Ⓘ], S. Bianco[Ⓘ], S. Meola^{Ⓘ⁵²}, D. Piccolo[Ⓘ]

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

P. Chatagnon^{Ⓘ^a}, F. Ferro^{Ⓘ^a}, E. Robutti^{Ⓘ^a}, S. Tosi^{Ⓘ^{a,b}}

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^{Ⓘ^a}, F. Brivio^{Ⓘ^a}, F. Cetorelli^{Ⓘ^{a,b}}, F. De Guio^{Ⓘ^{a,b}}, M.E. Dinardo^{Ⓘ^{a,b}}, P. Dini^{Ⓘ^a}, S. Gennai^{Ⓘ^a}, R. Gerosa^{Ⓘ^{a,b}}, A. Ghezzi^{Ⓘ^{a,b}}, P. Govoni^{Ⓘ^{a,b}}, L. Guzzi^{Ⓘ^a}, M.T. Lucchini^{Ⓘ^{a,b}}, M. Malberti^{Ⓘ^a}, S. Malvezzi^{Ⓘ^a}, A. Massironi^{Ⓘ^a}, D. Menasce^{Ⓘ^a}, L. Moroni^{Ⓘ^a}, M. Paganoni^{Ⓘ^{a,b}}, S. Palluotto^{Ⓘ^{a,b}}, D. Pedrini^{Ⓘ^a}, A. Perego^{Ⓘ^{a,b}}, B.S. Pinolini^a, G. Pizzati^{Ⓘ^{a,b}}, S. Ragazzi^{Ⓘ^{a,b}}, T. Tabarelli de Fatis^{Ⓘ^{a,b}}

INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^{Ⓘ^a}, A. Cagnotta^{Ⓘ^{a,b}}, F. Carnevali^{a,b}, N. Cavallo^{Ⓘ^{a,c}}, F. Fabozzi^{Ⓘ^{a,c}}, A.O.M. Iorio^{Ⓘ^{a,b}}, L. Lista^{Ⓘ^{a,b,53}}, P. Paolucci^{Ⓘ^{a,32}}, B. Rossi^{Ⓘ^a}

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino^{Ⓘ^a}, P. Azzi^{Ⓘ^a}, N. Bacchetta^{Ⓘ^{a,54}}, D. Bisello^{Ⓘ^{a,b}}, P. Bortignon^{Ⓘ^a}, G. Bortolato^{a,b}, A. Bragagnolo^{Ⓘ^{a,b}}, A.C.M. Bulla^{Ⓘ^a}, R. Carlin^{Ⓘ^{a,b}}, T. Dorigo^{Ⓘ^a}, F. Gasparini^{Ⓘ^{a,b}}, U. Gasparini^{Ⓘ^{a,b}}, S. Giorgetti^a, E. Lusiani^{Ⓘ^a}, M. Margoni^{Ⓘ^{a,b}}, A.T. Meneguzzo^{Ⓘ^{a,b}}, M. Migliorini^{Ⓘ^{a,b}}, M. Passaseo^{Ⓘ^a}, J. Pazzini^{Ⓘ^{a,b}}, P. Ronchese^{Ⓘ^{a,b}}, R. Rossin^{Ⓘ^{a,b}}, F. Simonetto^{Ⓘ^{a,b}}, M. Tosi^{Ⓘ^{a,b}}, A. Triossi^{Ⓘ^{a,b}}, S. Ventura^{Ⓘ^a}, M. Zanetti^{Ⓘ^{a,b}}, P. Zotto^{Ⓘ^{a,b}}, A. Zucchetta^{Ⓘ^{a,b}}, G. Zumerle^{Ⓘ^{a,b}}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

C. Aimè^{Ⓘ^a}, A. Braghieri^{Ⓘ^a}, S. Calzaferri^{Ⓘ^a}, D. Fiorina^{Ⓘ^a}, P. Montagna^{Ⓘ^{a,b}}, V. Re^{Ⓘ^a}, C. Riccardi^{Ⓘ^{a,b}}, P. Salvini^{Ⓘ^a}, I. Vai^{Ⓘ^{a,b}}, P. Vitulo^{Ⓘ^{a,b}}

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{Ⓘ^{a,b}}, M.E. Ascioti^{a,b}, G.M. Bilei^{Ⓘ^a}, C. Carrivale^{a,b}, D. Ciangottini^{Ⓘ^{a,b}}, L. Fanò^{Ⓘ^{a,b}}, M. Magherini^{Ⓘ^{a,b}}, V. Mariani^{Ⓘ^{a,b}}, M. Menichelli^{Ⓘ^a}, F. Moscatelli^{Ⓘ^{a,55}}, A. Rossi^{Ⓘ^{a,b}}, A. Santocchia^{Ⓘ^{a,b}}, D. Spiga^{Ⓘ^a}, T. Tedeschi^{Ⓘ^{a,b}}

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C.A. Alexe^{Ⓘ^{a,c}}, P. Asenov^{Ⓘ^{a,b}}, P. Azzurri^{Ⓘ^a}, G. Bagliesi^{Ⓘ^a}, R. Bhattacharya^{Ⓘ^a}, L. Bianchini^{Ⓘ^{a,b}}, T. Boccali^{Ⓘ^a}, E. Bossini^{Ⓘ^a}, D. Bruschini^{Ⓘ^{a,c}}, R. Castaldi^{Ⓘ^a}, M.A. Ciocci^{Ⓘ^{a,b}}, M. Cipriani^{Ⓘ^{a,b}},

V. D'Amante ^{a,d}, R. Dell'Orso ^a, S. Donato ^a, A. Giassi ^a, F. Ligabue ^{a,c}, A.C. Marini ^a,
D. Matos Figueiredo ^a, A. Messineo ^{a,b}, M. Musich ^{a,b}, F. Palla ^a, A. Rizzi ^{a,b},
G. Rolandi ^{a,c}, S. Roy Chowdhury ^a, T. Sarkar ^a, A. Scribano ^a, P. Spagnolo ^a,
R. Tenchini ^a, G. Tonelli ^{a,b}, N. Turini ^{a,d}, F. Vaselli ^{a,c}, A. Venturi ^a, P.G. Verдини ^a

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

C. Baldenegro Barrera ^{a,b}, P. Barria ^a, C. Basile ^{a,b}, F. Cavallari ^a, L. Cunqueiro Mendez ^{a,b},
D. Del Re ^{a,b}, E. Di Marco ^{a,b}, M. Diemoz ^a, F. Errico ^{a,b}, E. Longo ^{a,b}, J. Mijuskovic ^{a,b},
G. Organtini ^{a,b}, F. Pandolfi ^a, R. Paramatti ^{a,b}, C. Quaranta ^{a,b}, S. Rahatlou ^{a,b},
C. Rovelli ^a, F. Santanastasio ^{a,b}, L. Soffi ^a

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane ^{a,b}, R. Arcidiacono ^{a,c}, S. Argiro ^{a,b}, M. Arneodo ^{a,c}, N. Bartosik ^a,
R. Bellan ^{a,b}, A. Bellora ^{a,b}, C. Biino ^a, C. Borca ^{a,b}, N. Cartiglia ^a, M. Costa ^{a,b},
R. Covarelli ^{a,b}, N. Demaria ^a, L. Finco ^a, M. Grippo ^{a,b}, B. Kiani ^{a,b}, F. Legger ^a,
F. Luongo ^{a,b}, C. Mariotti ^a, L. Markovic ^{a,b}, S. Maselli ^a, A. Mecca ^{a,b}, L. Menzio ^{a,b},
P. Meridiani ^a, E. Migliore ^{a,b}, M. Monteno ^a, R. Mulargia ^a, M.M. Obertino ^{a,b},
G. Ortona ^a, L. Pacher ^{a,b}, N. Pastrone ^a, M. Pelliccioni ^a, M. Ruspa ^{a,c}, F. Siviero ^{a,b},
V. Sola ^{a,b}, A. Solano ^{a,b}, A. Staiano ^a, C. Tarricone ^{a,b}, D. Trocino ^a, G. Umoret ^{a,b},
R. White ^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar ^{a,b}, S. Belforte ^a, V. Candelise ^{a,b}, M. Casarsa ^a, F. Cossutti ^a, K. De Leo ^a,
G. Della Ricca ^{a,b}

Kyungpook National University, Daegu, Korea

S. Dogra ^a, J. Hong ^a, B. Kim ^a, J. Kim ^a, D. Lee ^a, H. Lee ^a, S.W. Lee ^a, C.S. Moon ^a, Y.D. Oh ^a,
M.S. Ryu ^a, S. Sekmen ^a, B. Tae ^a, Y.C. Yang ^a

Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim ^a

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak ^a, P. Gwak ^a, H. Kim ^a, D.H. Moon ^a

Hanyang University, Seoul, Korea

E. Asilar ^a, J. Choi ^a, D. Kim ^a, T.J. Kim ^a, J.A. Merlin ^a, Y. Ryou ^a

Korea University, Seoul, Korea

S. Choi ^a, S. Han ^a, B. Hong ^a, K. Lee ^a, K.S. Lee ^a, S. Lee ^a, J. Yoo ^a

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh ^a, S. Yang ^a

Sejong University, Seoul, KoreaH. S. Kim , Y. Kim, S. Lee**Seoul National University, Seoul, Korea**J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , S. Ko , H. Kwon , H. Lee ,
J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon **University of Seoul, Seoul, Korea**W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh,
I.J. Watson **Yonsei University, Department of Physics, Seoul, Korea**S. Ha , H.D. Yoo **Sungkyunkwan University, Suwon, Korea**M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu **College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait**T. Beyrouthy , Y. Gharbia **Kuwait University - College of Science - Department of Physics, Safat, Kuwait**F. Alazemi **Riga Technical University, Riga, Latvia**K. Dreimanis , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs, A. Potrebko , M. Seidel ,
D. Sidiropoulos Kontos **University of Latvia (LU), Riga, Latvia**N.R. Strautnieks **Vilnius University, Vilnius, Lithuania**M. Ambrozias , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis **National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**I. Yusuff ⁵⁶, Z. Zolkapli**Universidad de Sonora (UNISON), Hermosillo, Mexico**J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo **Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo ,
I. Heredia-De La Cruz ⁵⁷, R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,
A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, MexicoC. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García **Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada **University of Montenegro, Podgorica, Montenegro**I. Bubanja , N. Raicevic **University of Canterbury, Christchurch, New Zealand**P.H. Butler **National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan **AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland**V. Avati, L. Grzanka , M. Malawski **National Centre for Nuclear Research, Swierk, Poland**H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski **Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad **Warsaw University of Technology, Warsaw, Poland**K. Pozniatek , W. Zabolotny **Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi ,
G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo ,
T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff **Faculty of Physics, University of Belgrade, Belgrade, Serbia**P. Adzic , P. Milenovic **VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**D. Devetak, M. Dordevic , J. Milosevic , V. Rekovic**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero ,
M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris ,
A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz ,
O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , E. Martin Viscasillas 

D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra ,
A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre ,
J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras ,
I. Gonzalez Caballero , J.R. González Fernández , P. Leguina , E. Palencia Cortezon ,
J. Prado Pico , C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote ,
C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros ,
M. Fernandez , G. Gomez , C. Lasasoa García , R. Lopez Ruiz , C. Martinez Rivero ,
P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos ,
J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy ⁵⁸, D.D.C. Wickramaratna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna ⁵⁹, K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney ,
A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta ,
E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski ,
A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni ,
W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber ,
V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban ³⁰, S. Laurila ,
P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman,
A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat ,
M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer ,
M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , M. Rovere , H. Sakulin ,
S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva ,
P. Sphicas ⁶⁰, A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea ,
D. Walter , J. Wanczyk ⁶¹, J. Wang, K.A. Wozniak ⁶², S. Wuchterl , P. Zehetner , P. Zejdl ,
W.D. Zeuner

PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua ⁶³, L. Caminada ⁶³, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram ,
H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶³, L. Noehte ⁶³, T. Rohe 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov ⁶¹, M. Backhaus , G. Bonomelli , A. Calandri , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'archiac ⁶¹, A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , C. Martin Perez , A. Mascellani ⁶¹, F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , B. Ristic , F. Riti , R. Seidita , J. Stegmann ⁶¹, A. Tarabini , D. Valsecchi , R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler ⁶⁴, P. Bäertschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , F. Meng , U. Molinatti , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff⁶⁵, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari ⁴⁰, S.S. Yu 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , P.s. Chen, Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu ⁶⁶, E. Eskut , Y. Guler ⁶⁷, E. Gurpinar Guler ⁶⁷, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut , K. Ozdemir ⁶⁸, A. Polatoz , B. Tali ⁶⁹, U.G. Tok , S. Turkapar , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac ⁷⁰

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya ⁷¹, O. Kaya ⁷², S. Tekten ⁷³

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak ^{66,74}, G.G. Dincer ⁶⁶, Y. Komurcu , S. Sen ⁷⁵

Istanbul University, Istanbul, Turkey

O. Aydilek ⁷⁶, B. Hacisahinoglu , I. Hos ⁷⁷, B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, TurkeyS. Cerci , B. Isildak , D. Sunar Cerci , T. Yetkin **Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine**A. Boyaryntsev , B. Grynyov **National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine**L. Levchuk **University of Bristol, Bristol, U.K.**D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , M. Glowacki, J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou , K. Walkingshaw Pass**Rutherford Appleton Laboratory, Didcot, U.K.**A.H. Ball, K.W. Bell , A. Belyaev ⁸⁰, C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams **Imperial College, London, U.K.**I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio, C.A. Carrillo Montoya , G.S. Chahal ⁸¹, D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi , A. Howard, G. Iles , C.R. Knight , J. Langford , J. León Holgado , L. Lyons , A.-M. Magnan , S. Mallios, M. Mieskolainen , J. Nash ⁸², M. Pesaresi , P.B. Pradeep, B.C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee ³², M. Vojinovic , N. Wardle , D. Winterbottom **Brunel University, Uxbridge, U.K.**K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid **Baylor University, Waco, Texas, U.S.A.**S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish ⁸³, J. Dittmann , K. Hatakeyama , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson **Catholic University of America, Washington, DC, U.S.A.**R. Bartek , A. Dominguez , C. Huerta Escamilla, A.E. Simsek , R. Uniya , A.M. Vargas Hernandez **The University of Alabama, Tuscaloosa, Alabama, U.S.A.**B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson, C.U. Perez , P. Rumerio ⁸⁴, E. Usai , R. Yi 

Boston University, Boston, Massachusetts, U.S.A.

A. Akpinar , C. Cosby , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier ,
C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain, I. Reed ,
J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli 

Brown University, Providence, Rhode Island, U.S.A.

G. Benelli , D. Cutts , L. Gouskos , M. Hadley , U. Heintz , J.M. Hogan ⁸⁵, T. Kwon ,
G. Landsberg , K.T. Lau , D. Li , J. Luo , S. Mondal , N. Pervan , T. Russell, S. Sagir ⁸⁶,
X. Shen , F. Simpson , M. Stamenkovic , N. Venkatasubramanian, X. Yan 

University of California, Davis, Davis, California, U.S.A.

S. Abbott , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez ,
M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral ,
G. Mocellin , M. Mulhearn , S. Ostrom , W. Wei , S. Yoo , F. Zhang 

University of California, Los Angeles, California, U.S.A.

M. Bachtis , R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko ,
M.A. Iqbal , T. Lam , E. Manca , A. Nunez Del Prado, D. Saltzberg , V. Valuev 

University of California, Riverside, Riverside, California, U.S.A.

R. Clare , J.W. Gary , M. Gordon, G. Hanson , W. Si 

University of California, San Diego, La Jolla, California, U.S.A.

A. Aportela, A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte ,
L. Giannini , Y. Gu, J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts ,
M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , M. Quinnan ,
B.V. Sathia Narayanan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang ,
A. Yagil 

University of California, Santa Barbara - Department of Physics, Santa Barbara, California, U.S.A.

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , J. Incandela ,
J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica ,
R. Schmitz , F. Setti , J. Sheplack , D. Stuart , T.Á. Vámi , S. Wang , D. Zhang

California Institute of Technology, Pasadena, California, U.S.A.

S. Bhattacharya , A. Bornheim , O. Cerri, A. Latorre, J. Mao , H.B. Newman ,
G. Reales Gutiérrez, M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 

Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.

J. Alison , S. An , P. Bryant , M. Cremonesi, V. Dutta , T. Ferguson ,
T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, C. Liu , T. Mudholkar , S. Murthy ,
P. Palit , K. Park, M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, U.S.A.

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganello ,
J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner ,
N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, U.S.A.

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , S. Hogan ,
P. Kotamnives, J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom ,
P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerick ,
D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati ,
H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira ,
Y. Feng , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer ,
S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig ,
J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma ,
B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid ,
K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn ,
J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro ,
C. Pena ⁸⁷, F. Ravera , A. Reinsvold Hall ⁸⁸, L. Ristori , M. Safdari , E. Sexton-Kennedy ,
N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk ,
N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 

University of Florida, Gainesville, Florida, U.S.A.

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh ,
E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez ,
G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig ,
Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, U.S.A.

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , V. Hagopian , R. Hashmi , R.S. Kim ,
S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay ,
J. Zhang

Florida Institute of Technology, Melbourne, Florida, U.S.A.

B. Alsufyani , M.M. Baarmand , S. Butalla , S. Das , T. Elkafrawy ⁸⁹, M. Hohlmann ,
E. Yanes

University of Illinois Chicago, Chicago, Illinois, U.S.A.

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov ,
C.E. Gerber , M. Hawsworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos ,
A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti,
T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

The University of Iowa, Iowa City, Iowa, U.S.A.

M. Alhusseini , D. Blend, K. Dilsiz ⁹⁰, L. Emediato , G. Karaman , O.K. Köseyan ,
J.-P. Merlo, A. Mestvirishvili ⁹¹, O. Neogi, H. Ogul ⁹², Y. Onel , A. Penzo , C. Snyder,
E. Tiras ⁹³

Johns Hopkins University, Baltimore, Maryland, U.S.A.

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou ,
P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

The University of Kansas, Lawrence, Kansas, U.S.A.

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean ,
Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu ,
J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹⁴, C. Rogan , C. Royon ,
R. Salvatico , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, U.S.A.

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli ,
D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, U.S.A.

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen ,
R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili ,
C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova ⁹⁵, A. Shevelev , L. Wang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.

J. Bendavid , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer ,
G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa ,
L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus ,
C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang ,
B. Wyslouch , T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, U.S.A.

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi ,
M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J.E. Siado ,
W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, U.S.A.

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava ,
M. Morris , D. Nguyen , J. Pekkanen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, U.S.A.

G. Alverson , E. Barberis , J. Bonilla , M. Campana , J. Dervan , Y. Haddad , Y. Han ,
A. Krishna , J. Li , M. Lu , G. Madigan , R. McCarthy , D.M. Morse , V. Nguyen ,
T. Orimoto , A. Parker , L. Skinnari , D. Wood 

Northwestern University, Evanston, Illinois, U.S.A.

J. Bueghly, S. Dittmer , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt ,
A. Taliencio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, U.S.A.

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth ,
K.W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence ,
N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady ,
C. Moore , Y. Musienko ²⁵, H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti ,
A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, U.S.A.

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas ,
K. Wei, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, U.S.A.

H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg ,
N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow ,
I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.

S. Malik 

Purdue University, West Lafayette, Indiana, U.S.A.

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung ,
A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer,
J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Virdi , F. Wang , A. Wildridge , W. Xie ,
Y. Yao 

Purdue University Northwest, Hammond, Indiana, U.S.A.

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, U.S.A.

D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner,
F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi,
J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, U.S.A.

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs ,
A. Khukhunaishvili , N. Parmar , P. Parygin ⁹⁵, R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis ,
M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath ,
R. Montalvo, K. Nash, J. Reichert , H. Routray , P. Saha , S. Salur , S. Schnetzer,
S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, U.S.A.

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti ,
N. Karunaratna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, U.S.A.

D. Aebi , M. Ahmad , T. Akhter , O. Bouhali ⁹⁶, R. Eusebi , J. Gilmore , T. Huang ,
T. Kamon ⁹⁷, H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, U.S.A.

N. Akchurin , J. Damgov , N. Gogate , V. Hegde , A. Hussain , Y. Kazhykarim,
K. Lamichhane , S.W. Lee , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, U.S.A.

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, U.S.A.

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovsky , C. Neu 

Wayne State University, Detroit, Michigan, U.S.A.

S. Bhattacharya , P.E. Karchin 

University of Wisconsin - Madison, Madison, Wisconsin, U.S.A.

A. Aravind , S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts ,
C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless ,
J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida ,
L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi ,
W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin [†], I. Gorbunov , V. Karjavine ,
V. Korenkov , A. Lanev , A. Malakhov , V. Matveev ⁹⁸, V. Palichik , V. Perelygin ,
M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin ,
B.S. Yuldashev⁹⁹, A. Zarubin , I. Zhizhin , Yu. Andreev , A. Dermenev , S. Gninenko ,
N. Golubev , A. Karneyeu , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova ,
A. Toropin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

G. Gavrilo , V. Golovtcov , Y. Ivanov , V. Kim ¹⁰⁰, P. Levchenko ¹⁰¹, V. Murzin ,
V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev[†], T. Aushev , V. Gavrilo ,
N. Lychkovskaya , A. Nikitenko ^{102,103}, V. Popov , A. Zhokin , M. Chadeeva ¹⁰⁰,
R. Chistov ¹⁰⁰, S. Polikarpov ¹⁰⁰, V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov ,
E. Boos , V. Bunichev , M. Dubinin ⁸⁷, L. Dudko , V. Klyukhin , O. Kodolova ¹⁰³,
O. Lukina , S. Obraztsov , M. Perfilov , V. Savrin , A. Snigirev , G. Vorotnikov 

V. Blinov¹⁰⁰, T. Dimova¹⁰⁰, A. Kozyrev¹⁰⁰, O. Radchenko¹⁰⁰, Y. Skovpen¹⁰⁰,
V. Kachanov¹⁰⁰, D. Konstantinov¹⁰⁰, S. Slabospitskii¹⁰⁰, A. Uzunian¹⁰⁰, A. Babaev¹⁰⁰, V. Borshch¹⁰⁰,
D. Druzhkin¹⁰⁴, V. Chekhovsky, V. Makarenko¹⁰⁰

[†] Deceased

¹ Also at Yerevan State University, Yerevan, Armenia

² Also at TU Wien, Vienna, Austria

³ Also at Ghent University, Ghent, Belgium

⁴ Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵ Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷ Also at UFMS, Nova Andradina, Brazil

⁸ Also at Nanjing Normal University, Nanjing, China

⁹ Now at The University of Iowa, Iowa City, Iowa, U.S.A.

¹⁰ Also at University of Chinese Academy of Sciences, Beijing, China

¹¹ Also at China Center of Advanced Science and Technology, Beijing, China

¹² Also at University of Chinese Academy of Sciences, Beijing, China

¹³ Also at China Spallation Neutron Source, Guangdong, China

¹⁴ Now at Henan Normal University, Xinxiang, China

¹⁵ Also at Université Libre de Bruxelles, Bruxelles, Belgium

¹⁶ Also at an institute formerly covered by a cooperation agreement with CERN

¹⁷ Also at Helwan University, Cairo, Egypt

¹⁸ Now at Zewail City of Science and Technology, Zewail, Egypt

¹⁹ Now at British University in Egypt, Cairo, Egypt

²⁰ Now at Cairo University, Cairo, Egypt

²¹ Also at Purdue University, West Lafayette, Indiana, U.S.A.

²² Also at Université de Haute Alsace, Mulhouse, France

²³ Also at Istinye University, Istanbul, Turkey

²⁴ Also at Tbilisi State University, Tbilisi, Georgia

²⁵ Also at an international laboratory covered by a cooperation agreement with CERN

²⁶ Also at The University of the State of Amazonas, Manaus, Brazil

²⁷ Also at University of Hamburg, Hamburg, Germany

²⁸ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁹ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

³⁰ Also at Brandenburg University of Technology, Cottbus, Germany

³¹ Also at Forschungszentrum Jülich, Jülich, Germany

³² Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

³³ Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

³⁴ Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

³⁵ Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³⁶ Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³⁷ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁸ Also at Punjab Agricultural University, Ludhiana, India

³⁹ Also at University of Visva-Bharati, Santiniketan, India

⁴⁰ Also at Indian Institute of Science (IISc), Bangalore, India

⁴¹ Also at IIT Bhubaneswar, Bhubaneswar, India

⁴² Also at Institute of Physics, Bhubaneswar, India

⁴³ Also at University of Hyderabad, Hyderabad, India

⁴⁴ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany

⁴⁵ Also at Isfahan University of Technology, Isfahan, Iran

⁴⁶ Also at Sharif University of Technology, Tehran, Iran

⁴⁷ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran

- ⁴⁸ Also at Department of Physics, Isfahan University of Technology, Isfahan, Iran
- ⁴⁹ Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁵⁰ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁵¹ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁵² Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁵³ Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁵⁴ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.
- ⁵⁵ Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵⁶ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁷ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- ⁵⁸ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁹ Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁶⁰ Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁶¹ Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁶² Also at University of Vienna, Vienna, Austria
- ⁶³ Also at Universität Zürich, Zurich, Switzerland
- ⁶⁴ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶⁵ Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- ⁶⁶ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶⁷ Also at Konya Technical University, Konya, Turkey
- ⁶⁸ Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁹ Also at Adiyaman University, Adiyaman, Turkey
- ⁷⁰ Also at Bozok Universitеси Rektörlüğü, Yozgat, Turkey
- ⁷¹ Also at Marmara University, Istanbul, Turkey
- ⁷² Also at Milli Savunma University, Istanbul, Turkey
- ⁷³ Also at Kafkas University, Kars, Turkey
- ⁷⁴ Now at Istanbul Okan University, Istanbul, Turkey
- ⁷⁵ Also at Hacettepe University, Ankara, Turkey
- ⁷⁶ Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- ⁷⁷ Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷⁸ Also at Yildiz Technical University, Istanbul, Turkey
- ⁷⁹ Also at Vrije Universiteit Brussel, Brussel, Belgium
- ⁸⁰ Also at School of Physics and Astronomy, University of Southampton, Southampton, U.K.
- ⁸¹ Also at IPPP Durham University, Durham, U.K.
- ⁸² Also at Monash University, Faculty of Science, Clayton, Australia
- ⁸³ Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt
- ⁸⁴ Also at Università di Torino, Torino, Italy
- ⁸⁵ Also at Bethel University, St. Paul, Minnesota, U.S.A.
- ⁸⁶ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁸⁷ Also at California Institute of Technology, Pasadena, California, U.S.A.
- ⁸⁸ Also at United States Naval Academy, Annapolis, Maryland, U.S.A.
- ⁸⁹ Also at Ain Shams University, Cairo, Egypt
- ⁹⁰ Also at Bingol University, Bingol, Turkey
- ⁹¹ Also at Georgian Technical University, Tbilisi, Georgia
- ⁹² Also at Sinop University, Sinop, Turkey
- ⁹³ Also at Erciyes University, Kayseri, Turkey
- ⁹⁴ Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁹⁵ Now at another institute formerly covered by a cooperation agreement with CERN
- ⁹⁶ Also at Texas A&M University at Qatar, Doha, Qatar

⁹⁷ Also at Kyungpook National University, Daegu, Korea

⁹⁸ Also at another international laboratory covered by a cooperation agreement with CERN

⁹⁹ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan

¹⁰⁰ Also at another institute formerly covered by a cooperation agreement with CERN

¹⁰¹ Also at Northeastern University, Boston, Massachusetts, U.S.A.

¹⁰² Also at Imperial College, London, U.K.

¹⁰³ Now at Yerevan Physics Institute, Yerevan, Armenia

¹⁰⁴ Also at Universiteit Antwerpen, Antwerpen, Belgium