

Electrochemical One-Step Synthesis of Alkyne Sulfonates and Sulfonamides: Building Blocks for Highly Substituted Alkenes and 1,3-Butadienes

Meysam Azizzade, Florian A. Breitschaft, Siegfried R. Waldvogel, and Till Opatz*



Cite This: *JACS Au* 2025, 5, 5493–5503




Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information



ABSTRACT: Alkyne motifs have a central place in organic synthesis due to their versatility as building blocks for complex molecule construction. The development of efficient and sustainable strategies for the synthesis of functionalized alkynes remains an utmost challenge in modern organic chemistry. Herein, we report a novel and efficient electrochemical multicomponent reaction (eMCR) that enables the one-step synthesis of valuable alkyne sulfonates and sulfonamides using SO₂ stock solutions and electric current as a clean oxidant. The generated alkynyl sulfonates and sulfonamides serve as powerful synthetic intermediates, undergoing diverse downstream transformations, including regio- and stereoselective copper-catalyzed dimerizations, hydroarylations, and cyclizations, to access 1,3-dienes, highly substituted alkenes, and heterocycles. Moreover, this method enables late-stage functionalization of complex molecules and pharmaceuticals, demonstrating its significant potential for natural product synthesis, medicinal chemistry, and materials science. The study underscores the promise of electrochemical MCRs as sustainable and versatile tools in modern synthetic methodology.

KEYWORDS: *alkynes, sulfonates, sulfur dioxide, electrochemistry, copper catalysis*


Alkynes serve as versatile building blocks in organic chemistry and therefore play a crucial role in the construction of complex molecules.¹ Their synthetic utility has resulted in countless methods, providing facile access to a wider array of alkynyl structures. Besides the direct formation of triple bonds through eliminations or methodologies like the Corey-Fuchs reaction² and the Seyferth-Gilbert homologation,³ modifying pre-existing triple bonds is a common approach.⁴ Nevertheless, many procedures for the synthesis of alkynyl motifs from simple and accessible alkynes heavily depend on the use of aggressive organometallic reagents,⁵ noble metal catalysts,⁶ iodine,⁷ or harsh reaction conditions. The metal-free synthesis of new functionalized alkynes is quite challenging and thus, developing new methods to generate diverse arrays has remained a topic of interest in the last few decades.⁸ Remarkable advances have been made in the realm of organic electrosynthesis recently, a metal-free approach, propelled by the facilitation of redox chemistry without

external oxidants or reducing agents.⁹ Although a few protocols have been developed for the synthesis of new alkynyl motifs from simpler alkynes under electrochemical conditions,¹⁰ metal-free, multicomponent reactions for this purpose are both still underexplored and desirable. In this scenario, multicomponent reactions (MCRs), which can generate multiple carbon–carbon or carbon–heteroatom bonds from simple starting materials in a single synthetic operation, represent an attractive strategy in organic synthesis and often allow to enhance reaction efficiency and productivity.¹¹ In this regard, Lei's group very recently reported electrochemical

Received: August 4, 2025
Revised: October 12, 2025
Accepted: October 14, 2025
Published: October 23, 2025



Scheme 1. Synthesis of Alkyne Derivatives; (A) Electrochemical MCRs for Alkyne Functionalization; (B) C(sp²) Functionalization; (C) Traditional Synthesis Strategy; (D) Present Work



MCRs for the functionalization of alkynes through a Sonogashira-type reaction using palladium catalysts (Scheme 1A).¹² As part of our efforts to valorize SO₂, recently gaining significant attention in electrochemical syntheses (Scheme 1B),¹³ traditionally considered a waste product, and in view of

the importance of sulfonates and sulfonamides in areas like the materials sciences,¹⁴ agrochemistry¹⁵ (serving e.g., as herbicides,¹⁶ insecticides,¹⁷ antifungals,¹⁸ or agents against plant pathogenic bacteria and viruses¹⁹), and pharmaceuticals,²⁰ we sought for a one-step synthesis of alkyne sulfonates and alkyne

Table 1. Reaction Optimization

A) Reaction optimization^a

B) Reproducibility challenges in the presence of non-nucleophilic organic bases

Entry	Electrolyte	Base	Anode cathode	Current / Q _{Faraday}	Yield ^b
1 ^c	Bu ₄ NBF ₄ (1 eq.)	DIPEA	C Pt	10mA / 3.73 F	10%
2	Bu ₄ NBF ₄ (1 eq.)	DIPEA	C Pt	10mA / 3.73 F	18%
3	Bu ₄ NBF ₄ (1 eq.)	DIPEA	C Pt	7mA / 3.73 F	< 8%
4	Bu ₄ NBF ₄ (1 eq.)	DIPEA	C Pt	15mA / 3.73 F	27%
5	Bu ₄ NBF ₄ (1 eq.)	DIPEA	C Pt	20mA / 3.73 F	23%
6	Bu ₄ NBF ₄ (1 eq.)	DIPEA	C Pt	15mA / 5.59 F	36%
7	Bu ₄ NBF ₄ (1 eq.)	DBU	C Pt	15mA / 5.59 F	55% - 96%
8	Bu ₄ NBF ₄ (1 eq.)	DMIPA	C Pt	15mA / 5.59 F	52 %
9	Bu ₄ NBF ₄ (1 eq.)	TEA	C Pt	15mA / 5.59 F	12 %
10 ^d	Bu ₄ NBF ₄ (1 eq.)	DMIPA	C Pt	15mA / 5.59 F	59 %
11 ^d	-	DMIPA	C Pt	15mA / 5.59 F	73% (71%) ^e
12 ^d	-	DMIPA	GC or RVC Cu	15mA / 5.59 F	< 8%
13 ^d	-	DMIPA	C SS or Ni or Cu	15mA / 5.59 F	73%
14 ^d	-	DMIPA	C Cu	15mA / 4.84 F	73% (71%) ^e
15 ^{d,f}	-	DMIPA	C Cu	15mA / 4.84 F	54%


^aReaction conditions: **1a** (0.6 mmol), **2h** (4.5 equiv), SO₂ in DMSO (0.8 mL, 6.0 M, 8 equiv), base (2 equiv), and DCM (6 mL) in an undivided cell. ^bNMR yield. ^cSO₂ in MeCN. ^dDMIPA (3.2 equiv). ^eIsolated yield. ^f4 × AA batteries as the power source.

sulfonamides, which are conventionally accessed via tedious and harsh thionyl chloride chemistry (Scheme 1C). Here, we present a novel and versatile MCR electrooxidation method for the generation of alkynyl sulfonates and sulfonamides (Scheme 1D). These compounds proved to be useful building blocks.

RESULTS AND DISCUSSION

Our study began with the investigation of the electrochemical decarboxylative coupling between 3-phenylpropionic acid **1a**, SO₂ stock solution (5 M in acetonitrile), and neopentyl alcohol **2h** (*N,N*-diisopropylethylamine (DIPEA) as base, Bu₄NBF₄ as electrolyte, constant current (10 mA, 4.63 mA/cm², 2.16 cm²), graphite (C) anode and platinum plate cathode, room temperature, undivided cell), see Table S1. Different solvents were examined (Table S1), showing dichloromethane (DCM) to be optimal (Table 1, entry 1). Since the stock solution can be conveniently prepared by simply bubbling SO₂ gas into the solvent, we also explored various SO₂ solutions and found that the reaction could generate a yield up to 18% when solution in DMSO was used (entry 2). The change of the current from 10 mA (4.63 mA/cm², 2.16 cm²) to 15 mA (6.94 mA/cm², 2.16 cm²) had a significant effect on the yield (entries 4,6). After an extensive evaluation of the reaction conditions (Table S3), it

turned out that using DBU as a base, along with its role in the formation of a Lewis adduct²¹ with SO₂, provided the highest efficiency compared to other amines (entry 7). However, poor reproducibility was encountered when DBU, DBN, 1,2-dimethyl-1,4,5,6-tetrahydropyrimidine, or 1,1,3,3-tetramethylguanidine were used (Table S3), most likely due to side reactions of the desired product (Table 1B). In contrast, the use of *N,N*-dimethylisopropylamine (DMIPA) provided a good and reproducible yield of 59% (entry 10). A systematic investigation of different supporting electrolytes, including control experiments performed in their absence (entry 11), revealed that the yield improved when no additional supporting electrolyte was used (71%). The replacement of graphite with other anode materials (glassy carbon (GC) and reticulated vitreous carbon (RVC)) showed a detrimental effect on the formation of **3ah** (entry 12), but gratifyingly, Ni, Cu, and stainless steel also proved to be suitable cathode materials for the reaction (entry 13).²² The desired product was also successfully synthesized using AA batteries as the power source (entry 15, 54%), showcasing the robustness and simplicity of our method. In addition, control experiments indicated that open-air conditions are crucial to the reaction. After systematically screening various conditions (see Support-

Scheme 2. Scope of the Electrochemical Synthesis of Alkyne Sulfonates and Sulfonamides; (A) Alcohol and Amine Scope; (B) Alkyne Scope^d



^awith (bromoethynyl)benzene. ^bwith 3-(4-chlorophenyl)propionic acid. ^c7 h ($Q_{\text{Faraday}} = 6.45 \text{ F}$). ^dReaction conditions: **1** (0.6 mmol), **2** (4.5 equiv), SO₂ in DMSO (0.8 mL, 6.0 M, 8 equiv), DMIPA (3.2 equiv), and DCM (6 mL) in an undivided cell with graphite (C) anode and copper plate cathode (each 2.7 cm × 0.8 cm), room temperature, 15 mA, 315 min ($Q_{\text{Applied}} = 280 \text{ C}$, $Q_{\text{Faraday}} = 4.84 \text{ F}$).

ing Information for details), the desired product **3ah** could be obtained in 71% yield through constant current electrolysis at 15 mA (6.94 mA/cm²), in the presence of 3.2 equiv of DMIPA in DCM at r.t. for 315 min ($Q_{\text{Applied}} = 280 \text{ C}$, $Q_{\text{Faraday}} = 4.84 \text{ F}$) under air (entry 14).

Reaction Scope of the Electrochemical Synthesis of Alkyne Sulfonates and Sulfonamides

We evaluated the scope of this electrochemical MCR first using various alcohols (Scheme 2). Generally, a wide range of primary alcohols bearing different functional groups, such as ether (**3al**, **3am**, **3an**), phenyl (**3ap**, **3aq**), halide (**3aj**, **3ak**, **3an**), and cyclic acetal (**3ao**), were all found to be compatible. Moreover, primary alcohols of different chain lengths and steric hindrance (**3aa**–**3ai**) were smoothly converted to the

corresponding alkenyl sulfonate products in 38–74% yields. Secondary alcohols gave slightly lower yields due to their increased hindrance (**3ar**–**3aza**). The natural products borneol, menthol, and coprostanol were also amenable to this two-bond-forming strategy, producing 59% (**3ay**), 54% (**3ay**), and 27% (**3aza**) yield. Gratifyingly, this electrosynthesis was also applicable to amines, enabling the one-pot generation of alkenyl sulfonamides. We explored its versatility with a range of cyclic and acyclic secondary amines, yielding products **3azb** to **3azg** in isolated yields of 29 to 57%. Following the successful development of a decarboxylative MCR on propionic acid derivatives, alkynyl bromides were also evaluated as coupling partners. Gratifyingly, the reaction exhibited very good compatibility with various functional groups in the bromoalkyne. Aliphatic alkynes with different chain lengths are

Scheme 3. Mechanistic Studies; (A) Proposed Mechanism; (B) Cyclic Voltammetry Studies; (C) Intermediate Trapping Experiments; (D) Mechanistic Experiments


also suitable substrates for this reaction (**3ka**, **3la**). A broad array of synthetic handles showed good functional group robustness, including halide (**3la**), nitrile (**3ma**), ether (**3pa**, **3ra**), ester (**3na**, **3qa**, **3ra**, **3sa**), sulfonate ester (**3oa**), and ketone (**3ra**). Promisingly, substrates based on the anti-inflammatory drugs isoxepac (**3ra**) and α -tocopherol (**3sa**) proved to be well tolerated. In the reaction of 1,4-bis(bromoethynyl)benzene, a monoaddition product was isolated (**3ta**) in 41% yield, and the unreacted bromoalkyne unit can subsequently be subjected to coupling reactions. Importantly, compound **3ta** also enables the construction of symmetrical or unsymmetrical bis(alkynyl) sulfonates (**3ua**). We also attempted to synthesize target compound **3ah** from phenylacetylene; however, the yield did not exceed 35% (Table S4).

Investigation of the Reaction Mechanism


After developing a general method for the eMCR assembly of alkyne sulfonates and sulfonamides, subsequent efforts were directed toward exploring the reaction mechanism (Scheme 3). Cyclic voltammetry (CV) measurements showed that the oxidation potentials increase in the following order: intermediate *O*-alkyl sulfite, 3-phenylpropiolate, and alkyne sulfonate **3ah**. Notably, **3ah** is not prone to overoxidation due to its significantly higher oxidation potential compared to the intermediate *O*-alkyl sulfite (Scheme 3B). Evidence for the involvement of radical species in the reaction mechanism was obtained through trapping experiments employing an excess of 2,6-di-*tert*-butyl-4-methylphenol (BHT) (Scheme 3C). Under these conditions, the formation of **3aa** was not observed; instead, **3va** was isolated in 37% yield, further confirming the presence of **Int II** as a key intermediate. Control experiments

revealed that both the electric charge and base are essential for the reaction (Scheme 3D). The radical pathway is evidenced by the behavior of radical probe substrate **3wa** (alkenyl radical addition), as shown in Scheme 3D. Based on these results and previous literature reports, we propose the following mechanism (Scheme 3A). As shown in Scheme 3A, sulfur dioxide and the alcohol form Lewis acid–base adducts, generating the intermediate *O*-alkyl sulfite (**Int I**) via deprotonation by DMIPA. Initial anodic oxidation of **Int I** produces the radical **Int II**. This radical then attacks either the acetylenic acid anion or the bromoalkyne, giving rise to radical intermediates **Int III** or **Int IV**. **Int III** undergoes further oxidation, leading to decarboxylation and formation of the desired product **3**, whereas **Int IV** may undergo cathodic reduction to yield **3**. As the cathodic reaction, most likely the SO₂ reduction or hydrogen evolution process occurs.

Synthesis of 1,3-Butadienes via Alkyne Dimerization and Arylation with RB(OH)₂

The straightforward one-pot synthesis of alkenyl sulfonates and sulfonamides offers the chance to develop modular synthetic methods based on C–C triple bonds through further manipulations of these functional groups. The combination of arylation and alkyne dimerization²³ provides a good preparative method for various conjugated 1,3-dienes, which constitute essential scaffolds in natural products and pharmaceuticals,²⁴ as well as functional materials.²⁵ Additionally, they serve as valuable intermediates in multifunctionalization²⁶ and the syntheses of carbocycles and heterocycles.²⁷ However, previous approaches for the synthesis of 1,3-dienes involve nonregioselective reactions utilizing costly catalysts such as Pd, Au, and Ru, which are sensitive to air and

Scheme 4. Copper-Catalyzed Dimerizations and Hydroarylations; (A) Synthesis of 1,3-Butadienes; (B) Synthesis of Highly Substituted Alkenes




^aReaction conditions A: **3ah** (0.1 mmol), RB(OH)_2 (0.3 mmol), Cu(OAc)_2 (10 mol %) and MeOH (0.5 mL). ^bReaction conditions B: **3** (0.1 mmol), RB(OH)_2 (0.3 mmol), Cu(OAc)_2 (10 mol %), and NMP (0.5 mL). ^cfrom boronic acid pinacol esters.

moisture.²⁸ In addition, the absence of effective methods for the synthesis of alkenyl sulfonates hampered the preparation of 1,3-dienes with sulfonate substituents. These challenges drove us to find a novel strategy to overcome these limitations. We first investigated the reactivity and regio- and *E/Z*-selectivity of alkenyl sulfonates in reaction with phenylboronic acid (Scheme 4A). As alkyne dimerization has historically involved a wide range of catalysts, established conditions from the literature²⁹ (entries 1–3, Table S5) were applied to this novel alkenyl motif, yet all of these conditions failed to yield the desired 1,3-butadiene. After careful optimization of different catalysts, it was found that a high yield can be achieved by carbocupration with 10 mol % Cu(OAc)_2 as an inexpensive catalyst in MeOH at room temperature (entry 14, Table S5). Contrary to previous reports, complete control over the regioselectivity of the triple-bond dimerization was achieved, owing to the polarization of the acetylene fragment by the electron-withdrawing sulfonate moiety. The stereoselective formation of highly substituted 1,3-butadiene can be rationalized as a migratory insertion of the alkyne into the Ar–Cu bond. With the optimized conditions at hand, the scope of this regio- and *E*-selective dimerization was explored. Arylboronic acids, regardless of their electronic nature, consistently afforded the products in synthetically useful yields. A broad range of functional groups could be employed, such as alkenes (**4g**), halogens (**4h–4m**), esters (**4n**), aldehydes (**4o**), and imides (**4p**), as well as heterocycles such as indole (**4q**, **4r**) and carbazole (**4s**). Notably, the alkenylation of **3ah** with 1-

phenylvinylboronic acid under similar conditions gave linear 1,3,5-triene **4t** in 68% yield.

Synthesis of Highly Substituted Alkenes via Hydroarylation of Alkenyl Sulfonates with RB(OH)_2

During the synthesis of various 1,3-diene derivatives, hydroarylated side products were consistently observed in 5–20% yield. The previously reported stereoselective synthesis of highly substituted alkene sulfonates and sulfonamides³⁰ involved a three-step process that required the use of precious metal: first, a photoredox-catalyzed fluorosulfonyl-borylation followed by a Pd-catalyzed Suzuki–Miyaura cross-coupling in the presence of SPhos, and finally a SuFEx click reaction to ligate sulfonyl fluoride with alcohols and amines. On the other hand, direct methods for the preparation of alkenyl sulfonates either suffer from poor *E/Z*-selectivity or are limited to the synthesis of (*E*)-2-arylethene-1-sulfonates.^{13c} Therefore, further optimization studies were undertaken to redirect the hydroarylation pathway from a byproduct to the major product. Increasing temperature and changing solvent were found to be beneficial to the reaction, and the desired coupling product (**5a**) was ultimately obtained in 82% yield when the reaction was conducted in NMP at 100 °C (entry 7, Table S5). To broaden the generality of this coupling reaction, a variety of arylboronic acids were coupled with **3ah** under standard conditions (Scheme 4B). Alkene sulfonates **5k–5t** were obtained in moderate to excellent yields regardless of whether the reaction was conducted in MeOH at room temperature

Scheme 5. Product Transformations; (A) Synthetic Applications; (B) Nucleophilic Modification^d

^awithout base in MeCN as solvent. ^bEt₃N (2 equiv) as base in MeCN as solvent. ^cEt₃N (2 equiv) as base in DMF as solvent. ^dSupporting Information for detailed experimental conditions.

(conditions A) or in NMP at 100 °C (conditions B); however, the yield was slightly higher under conditions B.

Notably, for these particular boronic acids, no dimerization products **4** were observed under either set of conditions. As shown in Scheme 4B, this protocol can be used not only for the hydroarylation but also for facile hydroallylation (**5o**), hydroallylation (**5p**), and hydroferrocenylation (**5q**) of the alkenyl sulfonates. The importance of the allyl and allenyl groups relates to their presence in biologically active natural products and their wide synthetic versatility, serving as key intermediates for derivatization and coupling reactions.³¹ The absence of 1,3-diene formation under conditions A for compounds **5r–5t** indicates that 1,3-diene formation is highly dependent on the steric hindrance of the substituents on the alkyne. To determine the origin of the hydrogen atom in copper-catalyzed reactions, a series of control experiments were carried out (Figures S4 and S5).

Other Product Transformations

To demonstrate the practicality of our electrochemical protocol in the synthesis of heterocycles (Scheme 5A), we first conducted a 1,3-dipolar reaction between an in situ generated Münchnone (from acetic anhydride and proline) and alkenyl sulfonate **3ah**. The corresponding 2,3-dihydro-1*H*-pyrrolizine **6a** was obtained in 72% yield.³² Pyrazolo[1,5-*a*]pyridine **6b** was synthesized via a desulfonylation reaction. Furthermore, the transformations in Scheme 5A represent a

nucleophilic phosphine-catalyzed [3 + 2] annulation between alkenyl sulfonate **3ah** and *N*-hydroxyphthalimide (NHPI), yielding the fused tricyclic product **6c**.³³ Hydrolysis of alkenyl sulfonate also proceeded smoothly, generating β -keto sulfonate **6d** in 87% yield, a synthetically valuable motif for further functionalization. The use of octasulfur in the synthesis of compound **6e** represents the first report of a [2 + 2 + 1] strategy for the synthesis of valuable dihydrothiophenes, which are important motifs in natural or non-natural product synthesis.³⁴ A catalyst-free desulfonylative C(sp)–P coupling reaction formed the corresponding alkynyl di(phenyl)-phosphine oxide **6f** in 58% yield. Moreover, fully regio- and stereoselective functionalization of the electrochemically generated **3ah** was conducted via the reaction with *N*, *O*, and *S*-nucleophiles (Scheme 5B). These transformations highlight the significant synthetic potential of this electrochemical MCR reaction. An efficient transformation of **3ah** into β -aminovinyl sulfonates takes place successfully in the presence of a range of *N*-nucleophiles, affording **6g–6j** in yields of up to 89%. Alkynyl motifs were found to be suitable substrates for hydrothiolation and, contrary to previous reports,^{30b} exclusively afforded the *Z*-isomers (**6k–6o**) in high yields with excellent stereoselectivity. Gram-scale reactions were also conducted, affording 0.800 g (99%) and 0.786 g (98%) of compounds **6m** and **6o**, respectively. With compound **6o** in hand, we attempted a metal-free desulfonylative

synthesis of *N*-fused benzimidazo[2,1-*b*]thiazoles, which have shown promising biological activities.³⁵ In the presence of Cs₂CO₃ as a base, *N*-fused heterocycle **7a** was synthesized in 80% yield. Interestingly, hydroalkoxylation involving 4-hydroxycoumarin yielded the *Z*-isomer **6p**, in contrast to the *syn*-addition products reported in previous studies.³⁶ The intermolecular addition of hydroxybenzotriazole to **3ah** at room temperature within 20 min, regio- and stereoselectively gives vinyl ether **6q** without requiring a gold catalyst.³⁷ Subsequently, access to sulfonate-substituted enol esters (**6r–6w**) was explored through the nucleophilic modification of **3ah**. The versatility of this hydrocarboxylation method was then demonstrated by applying it to the modification of pharmaceutical compounds (**6s–6w**). We have shown that employing nucleophilic addition on our electrochemical products would provide an exciting opportunity for linking complex molecules and synthesizing bifunctional compounds (**6v,6w**), thereby making this a very useful protocol for chemical biology applications. It is worth mentioning that the formation of *E*-isomers in compounds **6g**, **6h**, and **6q**, synthesized without base, suggests an intramolecular proton transfer mechanism (*syn*-addition),³⁸ in contrast to the *Z*-selectivity observed in other nucleophilic addition products (**6i–6p**, **6r–6w**) where Et₃N was used as the base, likely involving an intermolecular proton transfer (*anti*-addition).

CONCLUSION

In summary, here we have developed the first electrochemical multicomponent reaction (eMCR) approach to valuable alkyne sulfonates and sulfonamides in a one-step synthesis using SO₂ stock solutions as the sulfur source. This metal- and catalyst-free strategy utilizes electricity as a green oxidant without requiring any supporting electrolyte, offers excellent functional group tolerance and a broad substrate scope, and enables the efficient synthesis of 1,3-dienes, highly substituted alkenes, and various heterocycles. The practicality of the electro-oxidative reaction was confirmed through gram-scale reactions and numerous downstream functionalizations to indicate the potential of the electrochemical products to be valuable intermediates in natural product synthesis, pharmaceutical discovery, and materials construction. The current research is an indication of the strength of electrochemical MCRs to be potent tools in modern synthetic methodology.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacsau.5c00972>.

Experimental procedures, ¹H, ¹⁹F, ¹³C, 2D NMR spectra, characterization data of compounds, and crystal structures (PDF)

Accession Codes

Deposition Numbers 2464884–2464885 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via the joint Cambridge Crystallographic Data Center (CCDC) and Fachinformationszentrum Karlsruhe Access Structures service.

AUTHOR INFORMATION

Corresponding Author

Till Opatz – Department of Chemistry, Johannes Gutenberg-University Mainz, 55128 Mainz, Germany; orcid.org/0000-0002-3266-4050; Email: opatz@uni-mainz.de

Authors

Meysam Azizzade – Department of Chemistry, Johannes Gutenberg-University Mainz, 55128 Mainz, Germany
Florian A. Breitschaft – Department of Electrosynthesis, Max-Planck-Institute for Chemical Energy Conversion, 45470 Mülheim an der Ruhr, Germany
Siegfried R. Waldvogel – Department of Electrosynthesis, Max-Planck-Institute for Chemical Energy Conversion, 45470 Mülheim an der Ruhr, Germany; Karlsruhe Institute of Technology, Institute of Biological and Chemical Systems – Functional Molecular Systems (IBCS-FMS), 76131 Karlsruhe, Germany; orcid.org/0000-0002-7949-9638

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacsau.5c00972>

Author Contributions

CRediT: Meysam Azizzade data curation, formal analysis, investigation, methodology, writing - original draft, writing - review & editing; Florian A. Breitschaft formal analysis, investigation, methodology, validation, writing - review & editing; Siegfried R. Waldvogel conceptualization, project administration, resources, supervision, visualization, writing - review & editing; Till Opatz conceptualization, funding acquisition, project administration, resources, supervision, writing - original draft, writing - review & editing.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors acknowledge the financial support by the BMFT in the frame of the Cluster4Future ETOS by the project E-SONATA (FKZ 03ZU1205IA and FKZ 03ZU1205IB). Helpful discussions with the partners within the project E-SONATA are gratefully acknowledged. We thank Dr. Dieter Schollmeyer (Mainz) for X-ray crystallography, Dr. Christopher Kampf (Mainz) for HR mass spectrometry and Dr. Johannes C. Liermann (Mainz) for NMR spectroscopy.

REFERENCES

- (1) (a) Munslow, I. J. Alkyne Reductions. In *Modern Reduction Methods*; Andersson, P. G.; Munslow, I. J., Eds.; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2008; pp 363–385. (b) Harvey, D. F.; Sigano, D. M. Carbene–Alkyne–Alkene Cyclization Reactions. *Chem. Rev.* **1996**, *96*, 271–288. (c) Chinchilla, R.; Nájera, C. Chemicals from Alkynes with Palladium Catalysts. *Chem. Rev.* **2014**, *114*, 1783–1826. (d) Ohno, H.; Inuki, S. Nonbiomimetic Total Synthesis of Indole Alkaloids using Alkyne-Based Strategies. *Org. Biomol. Chem.* **2021**, *19*, 3551–3568. (e) Kushwaha, P.; Saxena, A.; von Münchow, T.; Dana, S.; Saha, B.; Ackermann, L. Metallalectro-Catalyzed Alkyne Annulations via C–H Activations for Sustainable Heterocycle Syntheses. *Chem. Commun.* **2024**, *60*, 12333–12364. (f) Karak, P.; Rana, S. S.; Choudhury, J. Cationic π -Extended Heteroaromatics via a Catalytic C–H Activation Annulative Alkyne-Insertion Sequence. *Chem. Commun.* **2021**, *58*, 133–154. (h) Gong, X.; Yang, M.; Liu, J.-B.; He, F.-S.; Fan, X.; Wu, J. A Metal Free Route to Alkynyl Sulfones

under Photoinduced Conditions with the Insertion of Sulfur Dioxide. *Green Chem.* **2020**, *22*, 1906–1910.

(2) Corey, E. J.; Fuchs, P. L. A Synthetic Method for Formyl → Ethynyl Conversion (RCHO → RC≡CH or RC≡CR'). *Tetrahedron Lett.* **1972**, *13*, 3769–3772.

(3) Seyferth, D.; Marmor, R. S.; Hilbert, P. Some Reactions of Dimethylphosphono-Substituted Diazoalkanes. (MeO)₂P(O)CR Transfer to Olefins and 1,3-Dipolar Additions of (MeO)₂P(O)C-(N₂)R¹. *J. Org. Chem.* **1971**, *36*, 1379–1386.

(4) (a) Siemsen, P.; Livingston, R. C.; Diederich, F. Acetylene Couplings: A Powerful Method for Building Molecules. *Angew. Chem.* **2000**, *112*, 2740–2767. Siemsen, A.; Livingston, R. C.; Diederich, F. Acetylenic Coupling: A Powerful Tool in Molecular Construction. *Angew. Chem., Int. Ed.* **2000**, *39*, 2632–2657. (b) Park, K.; Lee, S. Transition Metal-Catalyzed Decarboxylative Coupling Reactions of Alkynyl Carboxylic Acids. *RSC Adv.* **2013**, *3*, 14165–14182.

(5) (a) Smedley, C. J.; Li, G.; Barrow, A. S.; Gialelis, T. L.; Giel, M. C.; Ottonello, A.; Cheng, Y.; Kitamura, S.; Wolan, D. W.; Sharpless, K. B.; Moses, J. E. Diversity Oriented Clicking (DOC): Divergent Synthesis of SuFExable Pharmacophores from 2-Substituted-Alkynyl-1-Sulfonyl Fluoride (SASF) Hubs. *Angew. Chem., Int. Ed.* **2020**, *59*, 12460–12469. Diversity Oriented Clicking (DOC): Divergent Synthesis of SuFExable Pharmacophores from 2-Substituted-Alkynyl-1-Sulfonyl Fluoride (SASF) Hubs. *Angew. Chem.*, **2020** *132* 12560 12569. (b) Ye, M.; Hou, M.; Wang, Y.; Ma, X.; Yang, K.; Song, Q. Arylation of Terminal Alkynes: Transition-Metal-Free Sonogashira-Type Coupling for the Construction of C(sp)–C(sp²) Bonds. *Org. Lett.* **2023**, *25*, 1787–1792.

(6) (a) Chinchilla, R.; Nájera, C. Recent Advances in Sonogashira Reactions. *Chem. Soc. Rev.* **2011**, *40*, 5084–5121. (b) Sarala, A. S.; Bhowmick, S.; de Carvalho, R. L.; Al-Thabaiti, S. A.; Mokhtar, M.; da Silva Júnior, E. N.; Maiti, D. Transition-Metal-Catalyzed Selective Alkynylation of C–H Bonds. *Adv. Synth. Catal.* **2021**, *363*, 4994–5027.

(7) (a) Dai, C.; Wang, J.; Deng, S.; Zhou, C.; Zhang, W.; Zhu, Q.; Tang, X. Acid-Mediated Sulfonylation of Arylethynylene Bromides with Sodium Arylsulfonates: Synthesis of (*E*)-1,2-Bis(arylsulfonyl)-ethylenes and Arylacetylenic Sulfones. *RSC Adv.* **2017**, *7*, 36112–36116. (b) Meesin, J.; Katrun, P.; Paresee-charoen, C.; Pohmakotr, M.; Reutrakul, V.; Soorukram, D.; Kuhakarn, C. Iodine-Catalyzed Sulfonylation of Arylacetylenic Acids and Arylacetylenes with Sodium Sulfonates: Synthesis of Arylacetylenic Sulfones. *J. Org. Chem.* **2016**, *81*, 2744–2752. (c) Singh, R.; Allam, B. K.; Singh, N.; Kumari, K.; Singh, S. K.; Singh, K. N. A Direct Metal-Free Decarboxylative Sulfonyl Functionalization (DSF) of Cinnamic Acids to α,β -Unsaturated Phenylsulfones. *Org. Lett.* **2015**, *17*, 2656–2659.

(8) Le Vaillant, F.; Waser, J. Alkynylation of Radicals: Spotlight on the “Third Way” to Transfer Triple Bonds. *Chem. Sci.* **2019**, *10*, 8909–8923.

(9) (a) Ghosh, A.; Bera, A.; Bera, S.; Banerjee, D. *Handbook of CH-Functionalization*; Maiti, D., Ed.; Wiley–VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2022; pp 1–27. (b) Cembellin, S.; Batanero, B. Organic Electrosynthesis Towards Sustainability: Fundamentals and Greener Methodologies. *Chem. Rec.* **2021**, *21*, 2453–2471. (c) Dutta, S.; Kayser, J.; Waldvogel, S. R. Electrochemical Synthesis of Disubstituted Alkynes from Dihydrazones. *Org. Lett.* **2025**, *27*, 7053–7057.

(10) (a) Mo, Z.-Y.; Zhang, Y.-Z.; Huang, G.-B.; Wang, X.-Y.; Pan, Y.-M.; Tang, H.-T. Electrochemical Sulfonylation of Alkynes with Sulfonyl Hydrazides: A Metal- and Oxidant-Free Protocol for the Synthesis of Alkynyl Sulfones. *Adv. Synth. Catal.* **2020**, *362*, 2160–2165. (b) Zhong, Q.; Zhao, Y.; Sheng, S.; Chen, J. Electrochemical Decarboxylative Sulfonylation of Arylacetylenic Acids with Sodium Arylsulfonates: Access to Arylacetylenic Sulfones. *Synth. Commun.* **2020**, *50*, 161–167. (c) Meng, X.; Xu, H.; Cao, X.; Cai, X.-M.; Luo, J.; Wang, F.; Huang, S. Electrochemically Enabled Sulfonylation of Alkynes with Sodium Sulfonates. *Org. Lett.* **2020**, *22*, 6827–6831. (d) Gao, P.-S.; Weng, X.-J.; Wang, Z.-H.; Zheng, C.; Sun, B.; Chen, Z.-H.; You, S.-L.; Mei, T.-S. Cu^{II}/TEMPO-Catalyzed Enantioselective

C(sp³)–H Alkynylation of Tertiary Cyclic Amines through Shono-Type Oxidation. *Angew. Chem., Int. Ed.* **2020**, *59*, 15254–15259. (e) Ye, X.-H.; Zhao, P.-Y.; Zhang, S.-Y.; Zhang, Y.-B.; Wang, Q.-L.; Shan, C.; Wojtas, L.; Guo, H.; Chen, H.; Shi, X.-D. Facilitating Gold Redox Catalysis with Electrochemistry: An Efficient Chemical-Oxidant-Free Approach. *Angew. Chem., Int. Ed.* **2019**, *58*, 17226–17230. (f) Pollok, D.; Waldvogel, S. R. Electro-Organic Synthesis – A 21st Century Technique. *Chem. Sci.* **2020**, *11*, 12386–12400. (g) Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic Organic Electrochemical Methods Since 2000: On the Verge of a Renaissance. *Chem. Rev.* **2017**, *117*, 13230–13319. (h) Tay, N. E. S.; Lehnher, D.; Rovis, T. Photons or Electrons? A Critical Comparison of Electrochemistry and Photoredox Catalysis for Organic Synthesis. *Chem. Rev.* **2022**, *122*, 2487–2649. (i) Zhu, C.; Ang, N. W. J.; Meyer, T. H.; Qiu, Y.; Ackermann, L. Organic Electrochemistry: Molecular Syntheses with Potential. *ACS Cent. Sci.* **2021**, *7*, 415–431. (j) Xu, H.-C.; Moeller, K. D. Electrochemistry in Synthetic Organic Chemistry. *J. Org. Chem.* **2021**, *86*, 15845–15846. (k) Little, R. D. A Perspective on Organic Electrochemistry. *J. Org. Chem.* **2020**, *85*, 13375–13390.

(11) (a) Coppola, G. A.; Pillitteri, S.; Van der Eycken, E. V.; You, S.-L.; Sharma, U. K. Multicomponent Reactions and Photo/Electrochemistry Join Forces: Atom Economy Meets Energy Efficiency. *Chem. Soc. Rev.* **2022**, *51*, 2313–2382. (b) Imeni, S.; Makarem, A.; Javahershenas, R. Recent Advances in Multicomponent Electro-Organic (Electrochemical) Synthesis of Heterocycles. *Asian J. Org. Chem.* **2023**, *12*, No. e202300303. (c) Krumbiegel, C.; Breitschaft, F. A.; Waldvogel, S. R. Electrochemical Synthesis of Cyclic Sulfites Using Diols and Sulfur Dioxide. *Chem. Commun.* **2025**, *61*, 11267–11270. (d) Blum, S. P.; Hofman, K.; Manolikakes, G.; Waldvogel, S. R. Advances in Photochemical and Electrochemical Incorporation of Sulfur Dioxide for the Synthesis of Value-Added Compounds. *Chem. Commun.* **2021**, *57*, 8236–8249.

(12) (a) Zeng, L.; Li, H.; Hu, J.; Zhang, D.; Hu, J.; Peng, P.; Wang, S.; Shi, R.; Peng, J.; Pao, C.-W.; Chen, J.-L.; Lee, J.-F.; Zhang, H.; Chen, Y.-H.; Lei, A. Electrochemical Oxidative Aminocarbonylation of Terminal Alkynes. *Nat. Catal.* **2020**, *3*, 438–445. (b) Wu, Y.; Zeng, L.; Li, H.; Cao, Y.; Hu, J.; Xu, M.; Shi, R.; Yi, H.; Lei, A. Electrochemical Palladium-Catalyzed Oxidative Sonogashira Carbonylation of Arylhydrazines and Alkynes to Ynones. *J. Am. Chem. Soc.* **2021**, *143*, 12460–12466.

(13) (a) Blum, S. P.; Karakaya, T.; Schollmeyer, D.; Klapars, A.; Waldvogel, S. R. Metal-Free Electrochemical Synthesis of Sulfonamides Directly from (Hetero)Arenes, SO₂, and Amines. *Angew. Chem., Int. Ed.* **2021**, *60*, 5056–5062. (b) Blum, S. P.; Schollmeyer, D.; Turks, M.; Waldvogel, S. R. Metal- and Reagent-Free Electrochemical Synthesis of Alkyl Arylsulfonates in a Multi-Component Reaction. *Chem. - Eur. J.* **2020**, *26*, 8358–8362. (c) de A Bartolomeu, A.; Breitschaft, F. A.; Schollmeyer, D.; Pilli, R. A.; Waldvogel, S. R. Electrochemical Multicomponent Synthesis of Alkyl Alkenesulfonates Using Styrenes, SO₂ and Alcohols. *Chem. - Eur. J.* **2024**, *30*, No. e202400557. (d) Chien, P. C.; Breitschaft, F. A.; Kelm, H.; Waldvogel, S. R.; Manolikakes, G. Electrochemical Three-component Synthesis of Alkenesulfonates from Cinnamic Acids, SO₂, and Alcohols. *ChemSusChem* **2025**, *18*, No. e202500186.

(14) (a) Bříza, T.; Král, V.; Martásek, P.; Kapláněk, R. Electrophilic Polyfluoroalkylating Agents Based on Sulfonate Esters. *J. Fluorine Chem.* **2008**, *129*, 235–247. (b) Mori, H.; Kudo, E.; Saito, Y.; Onuma, A.; Morishima, M. *Macromolecules* **2010**, *43*, 7021–7032. (c) Kuwayama, A.; Yamamoto, S.; Sakagami, Y.; Yamagishi, M.; Okano, K.; Horie, M.; Funahashi, M.; Mori, A. Synthesis and Properties of Polythiophenes Bearing Alkylsulfonic Acid Esters at the Side Chain. *Polym. Chem.* **2024**, *15*, 1635–1641. (d) Zhang, M.; Moore, J. D.; Flynn, D. L.; Hanson, P. R. Development of High-Load, Soluble Oligomeric Sulfonate Esters via ROM Polymerization: Application to the Benzoylation of Amines. *Org. Lett.* **2004**, *6*, 2657–2660. (e) Li, X.; Jiang, Y.; Shuai, L.; Wang, L.; Meng, L.; Mu, X. Sulfonated Copolymers with SO₃H and COOH Groups for the Hydrolysis of Polysaccharides. *J. Mater. Chem.* **2012**, *22*, 1283–1289. (f) Wang, L.; Pan, X.; Wang, F.; Yang, L.; Liu, L. Structure-Properties

- Relationships Investigation on the Azo Dyes Derived from Benzene Sulfonamide Intermediates. *Dyes Pigm.* **2008**, *76*, 636–645. (g) Kang, S. I.; Bae, Y. H. pH-Induced Solubility Transition of Sulfonamide-Based Polymers. *J. Controlled Release* **2002**, *80*, 145–155. (h) Aelony, D. Some Sulfonamide Plasticizers and Waxes. *Ind. Eng. Chem. Res.* **1954**, *46*, 587–591. (i) de Groote, P.; Rouxhet, P. G.; Devaux, J.; Godard, P. Infrared Study of the Hydrogen Bonding Association in Polyamides Plasticized by Benzenesulfonamides. Part I: Self-Association in Amide and Sulfonamide Systems; Part II: Amide-Sulfonamide Interaction. *Appl. Spectrosc.* **2001**, *55*, 877–887.
- (15) (a) Scott, K. A.; Njardarson, J. T. Analysis of US FDA-Approved Drugs Containing Sulfur Atoms. *Top. Curr. Chem.* **2018**, *376*, No. 5. (b) Lamberth, C. Sulfur Chemistry in Crop Protection. *J. Sulfur Chem.* **2004**, *25*, 39–62.
- (16) Meng, F. F.; Sun, X. W.; Shang, M. H.; Zhang, J. S.; Niu, C. W.; Li, Y. H.; Wang, Z. W.; Wang, J. G.; Li, Z. M. Chemical Preparation, Degradation Analysis, Computational Docking and Biological Activities of Novel Sulfonylureas with 2,5-Disubstituted Groups. *Pestic. Biochem. Physiol.* **2022**, *188*, No. 105261.
- (17) (a) Yang, Y.; Liu, Y.; Song, H.; Li, Y.; Wang, Q. Design, Synthesis, and Insecticidal Activity of Novel Triazole Derivatives Containing Sulfonamide or Sulfonimide Moieties. *J. Agric. Food Chem.* **2021**, *69*, 10790–10796. (b) Yu, J.; Jiang, X. Synthesis and Perspective of Organosulfur Chemicals in Agrochemicals. *Adv. Agrochem* **2023**, *2*, 3–14.
- (18) (a) Wahid, S.; Hanif, M.; Jahangir, S.; Shafique, M.; Shahid, H. A.; Muhammad, H.; Shah, S. A. A.; Versiani, M. A.; Khan, K. M.; Tahiri, I. A. Secnidazole-Sulfonates: Synthesis, Physical, Electrochemical, Antibacterial & Antifungal Characteristics. *J. Mol. Struct.* **2019**, *1184*, 569–575. (b) Lin, J.; Zhou, S.; Xu, J. X.; Yao, W. Q.; Hao, G. F.; Li, Y. T. Design, Synthesis, and Structure–Activity Relationship of Economical Triazole Sulfonamide Aryl Derivatives with High Fungicidal Activity. *J. Agric. Food Chem.* **2020**, *68*, 6792–6801.
- (19) (a) Wu, R.; Liu, T.; Wu, S. K.; Li, H. D.; Song, R. J.; Song, B. A. Synthesis, Antibacterial Activity, and Action Mechanism of Novel Sulfonamides Containing Oxyacetal and Pyrimidine. *J. Agric. Food Chem.* **2022**, *70*, 9305–9318. (b) Yang, Y.; Song, R.; Yin, L.; Han, K.; Yan, F.; Song, B. Inactivating Activities and Mechanism of Imidazo[1,2-*c*]pyrimidin-5(6*H*)-one Nucleoside Derivatives Incorporating a Sulfonamide Scaffold. *J. Agric. Food Chem.* **2023**, *71*, 7977–7987.
- (20) (a) Drews, J. Drug Discovery: A Historical Perspective. *Science* **2000**, *287*, 1960–1964. (b) Tilby, M. J.; Willis, M. C. How Do We Address Neglected Sulfur Pharmacophores in Drug Discovery? Expert Opin. *Drug Discovery* **2021**, *16*, 1227–1231. (c) Feng, M.; Tang, B.; Liang, S. H.; Jiang, X. Sulfur Containing Scaffolds in Drugs: Synthesis and Application in Medicinal Chemistry. *Curr. Top. Med. Chem.* **2016**, *16*, 1200–1216. (d) Scott, K. A.; Njardarson, J. T. Analysis of US FDA-Approved Drugs Containing Sulfur Atoms. *Top. Curr. Chem.* **2018**, *376*, No. 5. (e) Zhao, C.; Rakesh, K. P.; Ravidar, L.; Fang, W.-Y.; Qin, H.-L. Pharmaceutical and Medicinal Significance of Sulfur (SVI)-Containing Motifs for Drug Discovery: A Critical Review. *Eur. J. Med. Chem.* **2019**, *162*, 679–734. (f) Boulard, E.; Zibulski, V.; Oertel, L.; Lienau, P.; Schäfer, M.; Ganzer, U.; Lücking, U. Increasing Complexity: A Practical Synthetic Approach to Three-Dimensional, Cyclic Sulfoximines and First Insights into Their in Vitro Properties. *Chem. - Eur. J.* **2020**, *26*, 4378–4388. (g) Scozzafava, A.; Owa, T.; Mastrolorenzo, A.; Supuran, C. T. Anticancer and Antiviral Sulfonamides. *Curr. Med. Chem.* **2003**, *10*, 925–953. (h) Elder, D.; Facchine, K. L.; Levy, J. N.; Parsons, R.; Ridge, D.; Semo, L.; Teasdale, A. An Approach to Control Strategies for Sulfonate Ester Formation in Pharmaceutical Manufacturing Based on Recent Scientific Understanding. *Org. Process Res. Dev.* **2012**, *16*, 1707–1710. (i) Başaran, E.; Tür, G.; Akkoc, S.; Taskin-Tok, T. Design, Synthesis, and in Silico and in Vitro Cytotoxic Activities of Novel Isoniazid–Hydrazone Analogues Linked to Fluorinated Sulfonate Esters. *ACS Omega* **2024**, *9* (15), 17551–17562. (j) Reddy, C. R.; Rathaur, A.; Sagar, B. K. Acid-Mediated Intramolecular Cyclizations of (N-Aryl)-Acetylenic Sulfonamides: Synthesis of Fused and Spirocyclic Sultams. *J. Org. Chem.* **2024**, *89*, 14120–14128.
- (21) (a) Hata, T.; Kinumaki, S. Reactions of Ammonia and Aliphatic Amines with Sulphur Dioxide. *Nature* **1964**, *203*, 1378–1379. (b) Douglas, J. E.; Kollman, P. A. Trimethylamine-SO₂, the Prototype Strong Charge-Transfer Complex. *J. Am. Chem. Soc.* **1978**, *100*, 5226–5227.
- (22) Klein, M.; Waldvogel, S. R. Counter Electrode Reactions—Important Stumbling Blocks on the Way to a Working Electro-Organic Synthesis. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202204140. Klein, M.; Waldvogel, S. R. Reaktionen an der Gegenelektrode – wichtige Stolpersteine auf dem Weg einer funktionierenden elektroorganischen Synthese. *Angew. Chem.* **2022**, *134* (47), No. e202204140.
- (23) (a) Chen, S.; Wang, Y.-N.; Xie, J.; Li, W.; Ye, M.; Ma, X.; Yang, K.; Li, S.; Lan, Y.; Song, Q. Chemo-, Regio- and Stereoselective Access to Polysubstituted 1,3-Dienes via Nickel-Catalyzed Four-Component Reactions. *Nat. Commun.* **2024**, *15*, No. 5479. (b) Liu, Y.; Zhang, G.; Huang, H. Ni-Catalyzed Dimerization and Arylation of Diarylacetylenes with Arylboronic Acids. *Org. Lett.* **2017**, *19*, 6674–6677. (c) Shirakawa, E.; Takahashi, G.; Tsuchimoto, T.; Kawakami, Y. Nickel-Catalyzed Hydroarylation of Alkynes Using Arylboron Compounds: Selective Synthesis of Multisubstituted Arylalkenes and Aryldienes. *Chem. Commun.* **2001**, 2688–2689.
- (24) (a) Staunton, J.; Weissman, K. J. Polyketide Biosynthesis: A Millennium Review. *Nat. Prod. Rep.* **2001**, *18*, 380–416. (b) Rychnovsky, S. D. Oxo Polyene Macrolide Antibiotics. *Chem. Rev.* **1995**, *95*, 2021–2040. (c) Negishi, E.-i.; Huang, Z.; Wang, G.; Mohan, S.; Wang, C.; Hattori, H. Recent Advances in Efficient and Selective Synthesis of Di-, Tri-, and Tetrasubstituted Alkenes via Pd-Catalyzed Alkenylation–Carbonyl Olefination Synergy. *Acc. Chem. Res.* **2008**, *41*, 1474–1485. (d) Ke, L.; Chen, Z. Dienedicyclic Acid as a Useful Diene Building Block via Directed Heck-Decarboxylate Coupling. *Commun. Chem.* **2020**, *3*, No. 48. (e) Keerthana, M. S.; Jeganmohan, M. Palladium-Catalyzed Aerobic α,β -Dehydrogenation of Aliphatic Amides. *J. Org. Chem.* **2022**, *87*, 4873–4882.
- (25) (a) Davis, R.; Kumar, N. S. S.; Abraham, S.; Suresh, C. H.; Rath, N. P.; Tamaoki, N.; Das, S. Molecular Packing and Solid-State Fluorescence of Alkoxy-Cyano Substituted Diphenylbutadienes: Structure of the Luminescent Aggregates. *J. Phys. Chem. C* **2008**, *112*, 2137–2146. (b) Gholami, M.; Tykwinski, R. R. Oligomeric and Polymeric Systems with a Cross-Conjugated π -Framework. *Chem. Rev.* **2006**, *106*, 4997–5027. (c) Rauschenbach, M.; Meier-Merziger, M.; Frey, H. Green Perspective Drives the Renaissance of Anionic Diene Polymerization. *Polym. Chem.* **2024**, *15*, 4297–4311.
- (26) (a) Thullen, S. M.; Rovis, T. A Mild Hydroaminoalkylation of Conjugated Dienes Using a Unified Cobalt and Photoredox Catalytic System. *J. Am. Chem. Soc.* **2017**, *139*, 15504–15508. (b) Chen, P.; Tian, L.; Ji, X.; Deng, G.-J.; Huang, H. Copper-Catalyzed 1,2-Sulfonyl etherification of 1,3-Dienes. *Org. Lett.* **2024**, *26*, 2939–2944. (c) Liu, X.-D.; Ye, A.-H.; Chen, Z.-M. Catalytic Enantioselective Intermolecular Three-Component Sulfonylative Difunctionalizations of 1,3-Dienes. *ACS Catal.* **2023**, *13*, 2715–2722. (d) Forster, D.; Guo, W.; Wang, Q.; Zhu, J. Fe-Catalyzed Three-Component Coupling Reaction of $\alpha,\beta,\gamma,\delta$ -Unsaturated Carbonyl Compounds and Conjugate Dienes with Alkylsilyl Peroxides and Nucleophiles. *ACS Catal.* **2021**, *11*, 10871–10877. (e) Liu, Z.-L.; Ye, Z.-P.; Liao, Z.-H.; Lu, W.-D.; Guan, J.-P.; Gao, Z.-Y.; Chen, K.; Chen, X.-Q.; Xiang, H.-Y.; Yang, H. Photoinduced, Palladium-Catalyzed Enantioselective 1,2-Alkylsulfonylation of 1,3-Dienes. *ACS Catal.* **2024**, *14*, 3725–3732. (f) Ren, S.-Y.; Zhou, Q.; Zhou, H.-Y.; Wang, L.-W.; Mulina, O. M.; Paveliev, S. A.; Tang, H.-T.; Terent'ev, A. O.; Pan, Y.-M.; Meng, X.-J. Three-Component Electrochemical Aminoselenation of 1,3-Dienes. *J. Org. Chem.* **2023**, *88*, 5760–5771. (g) Wang, H.; Liu, C.-F.; Tan, T.-D.; Khoo, K. R. B.; Koh, M. J. N-Heterocyclic Carbene–Nickel-Catalyzed Regioselective Diarylation of Aliphatic-1,3-Dienes. *ACS Catal.* **2022**, *12*, 724–732.
- (27) (a) Kennedy, C. R.; Joannou, M. V.; Steves, J. E.; Hoyt, J. M.; Kovel, C. B.; Chirik, P. J. Iron-Catalyzed Vinylsilane Dimerization and

- Cross-Cycloadditions with 1,3-Dienes: Probing the Origins of Chemo- and Regioselectivity. *ACS Catal.* **2021**, *11*, 1368–1379.
- (b) Eom, D.; Park, S.; Park, Y.; Ryu, T.; Lee, P. H. Synthesis of Indenes via Brønsted Acid Catalyzed Cyclization of Diaryl- and Alkyl Aryl-1,3-Dienes. *Org. Lett.* **2012**, *14*, 5392–5395.
- (c) Kastrati, A.; Jaquier, V.; Garbo, M.; Besnard, C.; Mazet, C. Pd-Catalyzed Regioselective Cyclopropanation of 2-Substituted 1,3-Dienes. *ACS Org. Inorg. Au* **2023**, *3*, 291–298.
- (d) Pezzati, B.; Chellat, M. F.; Murphy, J. J.; Besnard, C.; Reginato, G.; Stephens, J. C.; Alexakis, A. Organocatalytic Asymmetric Annulation of 1,3-Bis(alkoxycarbonyl)-buta-1,3-dienes and Aldehydes. *Org. Lett.* **2013**, *15*, 2950–2953.
- (e) Liang, S.-Y.; Zhang, T.-Y.; Chen, Z.-C.; Du, W.; Chen, Y.-C. Functional-Group-Directed Regiodivergent (3 + 2) Annulations of Electronically Distinct 1,3-Dienes and 2-Formyl Arylboronic Acids. *Org. Lett.* **2024**, *26*, 1483–1488.
- (f) Ebe, Y.; Nishimura, T. Iridium-Catalyzed Annulation of Salicylimines with 1,3-Dienes. *J. Am. Chem. Soc.* **2014**, *136*, 9284–9287.
- (g) Nishimura, T.; Ebe, Y.; Hayashi, T. Iridium-Catalyzed [3 + 2] Annulation of Cyclic N-Sulfonyl Ketimines with 1,3-Dienes via C–H Activation. *J. Am. Chem. Soc.* **2013**, *135*, 2092–2095.
- (h) Li, Y.-L.; Zhang, P.-C.; Wu, H.-H.; Zhang, J. Palladium-Catalyzed Asymmetric Tandem Denitrogenative Heck/Tsuji–Trost of Benzotriazoles with 1,3-Dienes. *J. Am. Chem. Soc.* **2021**, *143*, 13010–13015.
- (28) (a) Maikhuri, V. K.; Maity, J.; Srivastava, S.; Prasad, A. K. Transition Metal-Catalyzed Double Cvinyl–H Bond Activation: Synthesis of Conjugated Dienes. *Org. Biomol. Chem.* **2022**, *20*, 9522–9588.
- (b) Zhou, P.; Huang, L.; Jiang, H.; Wang, A.; Li, X. Highly Chemoselective Palladium-Catalyzed Cross-Trimerization between Alkyne and Alkenes Leading to 1,3,5-Trienes or 1,2,4,5-Tetrasubstituted Benzenes with Dioxxygen. *J. Org. Chem.* **2010**, *75*, 8279–8282.
- (c) Liu, J.; Yang, J.; Baumann, W.; Jackstell, R.; Beller, M. Stereoselective Synthesis of Highly Substituted Conjugated Dienes via Pd-Catalyzed Carbonylation of 1,3-Diynes. *Angew. Chem., Int. Ed.* **2019**, *58*, 10683–10687.
- Liu, J.; Yang, J.; Baumann, W.; et al. Stereoselective Synthesis of Highly Substituted Conjugated Dienes via Pd-Catalyzed Carbonylation of 1,3-Diynes. *Angew. Chem.* **2019**, *131*, 10793–10797.
- (d) Shiroodi, R. K.; Dudnik, A. S.; Gevorgyan, V. Stereoccontrolled 1,3-Phosphatylxy and 1,3-Halogen Migration Relay Toward Highly Functionalized 1,3-Dienes. *J. Am. Chem. Soc.* **2012**, *134*, 6928–6931.
- (29) (a) Horie, H.; Kurahashi, T.; Matsubara, S. Selective Synthesis of Trienes and Dienes via Nickel-Catalyzed Intermolecular Cotrimerization of Acrylates and Alkynes. *Chem. Commun.* **2010**, *46*, 7229–7231.
- (b) Nakao, Y.; Kanyiva, K. S.; Hiyama, T. A Strategy for C–H Activation of Pyridines: Direct C-2 Selective Alkenylation of Pyridines by Nickel/Lewis Acid Catalysis. *J. Am. Chem. Soc.* **2008**, *130*, 2448–2449.
- (c) Ogata, K.; Murayama, H.; Sugasawa, J.; Suzuki, N.; Fukuzawa, S.-i. Nickel-Catalyzed Highly Regio- and Stereoselective Cross-Trimerization Between Triisopropylsilylacetylene and Internal Alkynes Leading to 1,3-diene-5-ynes. *J. Am. Chem. Soc.* **2009**, *131*, 3176–3177.
- (d) Fukutani, T.; Hirano, K.; Satoh, T.; Miura, M. Synthesis of Highly Substituted Naphthalene and Anthracene Derivatives by Rhodium-Catalyzed Oxidative Coupling of Arylboronic Acids with Alkynes. *Org. Lett.* **2009**, *11*, 5198–5201.
- (30) (a) Li, H.; Huang, M.; Zou, Z.; Wang, Z.; Li, Y.; Sun, C.; Chen, W.; Pan, Y.; Zhang, W.; Wang, Y. Photoredox-Catalyzed Stereo- and Regioselective Vicinal Fluorosulfonyl-Borylation of Unsaturated Hydrocarbons. *Chem. Sci.* **2023**, *14*, 13893–13901.
- (b) Nie, X.; Xu, T.; Hong, Y.; Zhang, H.; Mao, C.; Liao, S. Introducing A New Class of Sulfonyl Fluoride Hubs via Radical Chloro-Fluorosulfonylation of Alkynes. *Angew. Chem., Int. Ed.* **2021**, *60*, 22035–22042.
- (31) (a) Blicke, R.; Taillefer, M.; Monnier, F. Metal-Catalyzed Intermolecular Hydrofunctionalization of Allenes: Easy Access to Allylic Structures via the Selective Formation of C–N, C–C, and C–O Bonds. *Chem. Rev.* **2020**, *120*, 13545–13598.
- (b) Yu, S.; Ma, S. How Easy Are the Syntheses of Allenes? *Chem. Commun.* **2011**, *47*, 5384–5418.
- (c) Yu, S.; Ma, S. Allenes in Catalytic Asymmetric Synthesis and Natural Product Syntheses. *Angew. Chem., Int. Ed.* **2012**, *51*, 3074–3112.
- Yu, S.; Ma, S. Allene in katalytischer asymmetrischer Synthese und Naturstoffsynthese. *Angew. Chem.* **2012**, *124*, 3128–3167.
- (d) López, F.; Mascareñas, J. L. [4 + 2] and [4 + 3] Catalytic Cycloadditions of Allenes. *Chem. Soc. Rev.* **2014**, *43*, 2904–2915.
- (e) Sahu, S. K.; Behera, P. K.; Choudhury, P.; Sethi, M.; Jena, S.; Rout, L. Recent Advances in [3 + 2] Cycloaddition of Allenes with 1,3-Carbonyl Ylides; Rh (II)-Catalyzed Access to Bridged Polyoxocarbocycles. *New J. Chem.* **2021**, *45*, 11018–11029.
- (f) Huo, H.-X.; Duvall, J. R.; Huang, M.-Y.; Hong, R. Catalytic Asymmetric Allylation of Carbonyl Compounds and Imines with Allylic Boronates. *Org. Chem. Front.* **2014**, *1*, 303–320.
- (g) Yus, M.; Gonzalez-Gomez, J. C.; Foubelo, F. Diastereoselective Allylation of Carbonyl Compounds and Imines: Application to the Synthesis of Natural Products. *Chem. Rev.* **2013**, *113*, 5595–5698.
- (32) Smedley, C. J.; Li, G.; Barrow, A. S.; Gialelis, T. L.; Giel, M.-C.; Ottonello, A.; Cheng, Y.; Kitamura, S.; Wolan, D. W.; Sharpless, K. B.; Moses, J. E. Diversity Oriented Clicking (DOC): Divergent Synthesis of SuFExable Pharmacophores from 2-Substituted-Alkynyl-1-Sulfonyl Fluoride (SASF) Hubs. *Angew. Chem., Int. Ed.* **2020**, *59*, 12460–12469.
- (33) Zhou, Q. F.; Chu, X. P.; Ge, F. F.; Wang, Y.; Lu, T. Phosphine-Catalyzed [3 + 2] Annulation of Electron-Deficient Alkynes with N-Hydroxyphthalimide: Synthesis of 3a-Hydroxyisoxazolo[3,2-a]-isoindol-8(3aH)-ones. *Adv. Synth. Catal.* **2013**, *355*, 2787–2792.
- (34) (a) Rogers, E.; Araki, H.; Batory, L. A.; McInnis, C. E.; Njardarson, J. T. Highly Selective Copper-Catalyzed Ring Expansion of Vinyl Thiiranes: Application to Synthesis of Biotin and the Heterocyclic Core of Plavix. *J. Am. Chem. Soc.* **2007**, *129*, 2768–2769.
- (b) Dang, Y. Z.; Chen, Y. Oxidation of 3,4-Diaryl-2,5-dihydrothiophenes to 3,4-Diarylthiophenes Using CuBr₂: Simple and Efficient Preparation of 3,4-Diarylthiophenes. *Eur. J. Org. Chem.* **2007**, *2007*, 5661–5664.
- (35) Kuang, J.; Xia, Y.; Yang, A.; Zhang, H.; Su, C.; Lee, D. Copper-Catalyzed Aminothiolation of Terminal Alkynes with Tunable Regioselectivity. *Chem. Commun.* **2019**, *55*, 1813–1816.
- (36) Plamont, R.; Graux, L. V.; Clavier, H. Highly Selective Syn Addition of 1,3-Diones to Internal Ynamides Catalyzed by Zinc Iodide. *Eur. J. Org. Chem.* **2018**, *2018*, 1372–1376.
- (37) Kumar, M.; Scobie, M.; Mashuta, M. S.; Hammond, G. B.; Xu, B. Gold-Catalyzed Addition of N-Hydroxy Heterocycles to Alkynes and Subsequent 3,3-Sigmatropic Rearrangement. *Org. Lett.* **2013**, *15*, 724–727.
- (38) Strozier, R. W.; Caramella, P.; Houk, K. N. Influence of Molecular Distortions upon Reactivity and Stereochemistry in Nucleophilic Additions to Acetylenes. *J. Am. Chem. Soc.* **1979**, *101*, 1340–1343.