

eScholarship

Combinatorial Theory

Title

Canonical theorems in geometric Ramsey theory

Permalink

<https://escholarship.org/uc/item/0d76t2c1>

Journal

Combinatorial Theory, 5(4)

ISSN

2766-1334

Authors

Gehér, Panna

Sagdeev, Arsenii

Tóth, Géza

Publication Date

2025

DOI

10.5070/C65465673

Supplemental Material

<https://escholarship.org/uc/item/0d76t2c1#supplemental>

Copyright Information

Copyright 2025 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at

<https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

CANONICAL THEOREMS IN GEOMETRIC RAMSEY THEORY

Panna Gehér^{*1}, Arsenii Sagdeev^{†2}, and Géza Tóth^{‡3}

^{1,2,3}*HUN-REN Alfréd Rényi Institute of Mathematics, Budapest, Hungary*

¹*Eötvös Loránd University, Budapest, Hungary*

geher.panna@ttk.elte.hu

²*KIT, Karlsruhe, Germany*

sagdeevarsenii@gmail.com

³*Department of Computer Science and Information Theory, Budapest University of Technology and Economics, Budapest, Hungary*

geza@renyi.hu

Submitted: Apr 17, 2024; Accepted: Jun 9, 2025; Published: Dec 20, 2025

© The authors. Released under the CC BY license (International 4.0).

Abstract. In Euclidean Ramsey Theory usually we are looking for monochromatic configurations in the Euclidean space, whose points are colored with a fixed number of colors. In the canonical version, the number of colors is arbitrary, and we are looking for an ‘unavoidable’ set of colorings of a finite configuration, that is, a set of colorings with the property that one of them always appears in any coloring of the space. This set definitely includes the monochromatic and the rainbow colorings. In the present paper, we prove the following two results of this type. First, for any acute triangle T , and any coloring of \mathbb{R}^3 , there is either a monochromatic or a rainbow copy of T . Second, for every m , there exists a sufficiently large n such that in any coloring of \mathbb{R}^n , there exists either a monochromatic or a rainbow m -dimensional unit hypercube. In the maximum norm, ℓ_∞ , we have a much stronger statement. For every finite M , there exists an n such that in any coloring of \mathbb{R}_∞^n , there is either a monochromatic or a rainbow isometric copy of M .

Keywords. Euclidean Ramsey theory, canonical Ramsey theorem, colorings of the space

Mathematics Subject Classifications. 05D10, 05C55

*Supported by ERC Advanced Grant ‘GeoScape’ No. 882971, the Lendület Programme of the Hungarian Academy of Sciences – grant number LP2021-1/2021 and by the Thematic Excellence Program TKP2021-NKTA-62 of the National Research, Development and Innovation Office.

†Supported by ERC Advanced Grant ‘GeoScape’ No. 882971.

‡Supported by ERC Advanced Grant ‘GeoScape’ No. 882971 and by the National Research, Development and Innovation Office, NKFIH, K-131529 and ADVANCED 152590.

1. Introduction

A typical problem in *Ramsey theory* [GRS91] is to show that every r -coloring of a sufficiently large ‘domain’ contains a monochromatic copy of some fixed ‘configuration’. A classical result in the field is the van der Waerden theorem [VdW27], which states that for all $k, r \in \mathbb{N}$, there exists n such that every r -coloring of $[n]$ contains a monochromatic k -term arithmetic progression. Each problem in Ramsey theory has its *canonical* counterpart, in which the goal is to list a minimal set S of colorings which is ‘unavoidable’ regardless of the number of colors in use. That is, in any coloring of the domain we have a configuration whose coloring belongs to S . A coloring of a configuration is called *rainbow* if all of its points are of different color. For example, the canonical van der Waerden [EG80] theorem states that for all $k \in \mathbb{N}$, there exists n such that every coloring of $[n]$ contains either a monochromatic or a *rainbow* k -term arithmetic progression. For modern exposition and further generalizations of this result, see [GKP12, Ga20] and the references therein. We note that the minimal sets of unavoidable colorings often contain *other* colorings along with the monochromatic and the rainbow. In fact, this is the case for canonical versions of both the Ramsey theorem due to Erdős and Rado [ER50], see also [JM00, AJ05, Wag06, JLL03], and the Hales–Jewett theorem [PV83, NPRV85], see Section 3.1 for the exact statement.

In their celebrated trilogy [EGM⁺73, EGM⁺75a, EGM⁺75b], Erdős, Graham, Montgomery, Rothschild, Spencer and Straus initiated a systematic study of the following general problem in geometric Ramsey theory. For $n, r \in \mathbb{N}$ and $A \subset \mathbb{R}^n$, does every r -coloring of \mathbb{R}^n contain a monochromatic isometric copy of A ? They also introduced the notation $\mathbb{R}^n \xrightarrow{r} A$ for an affirmative answer on the latter question. Note that only a few results in the field are tight. Even the innocent-looking famous special case, when $n = 2$ and $A = I$ is a fixed two-point set is open. It is equivalent to the notorious Hadwiger–Nelson problem which asks for the chromatic number of the plane. After 75 years of extensive study, there are still two values of r , namely $r = 5$ and $r = 6$, for which we do not know whether $\mathbb{R}^2 \xrightarrow{r} I$ or not, see [dG18, EI20] and the survey in [Soi09].

Another major open problem in the field is to describe all *Ramsey sets*, that is sets A with the property that for all $r \in \mathbb{N}$ there is a sufficiently large n with $\mathbb{R}^n \xrightarrow{r} A$. It is known that every Ramsey set is spherical [EGM⁺73, Theorem 13] and finite [EGM⁺75a, Theorem 19]. We refer the reader to [LRW12] for the discussion about if these two necessary conditions might also be sufficient. From the other direction, it is known [Kř1, Can07] that (the vertex sets of) all regular polytopes are Ramsey. The vertex set A of a box or a simplex in arbitrary dimension has an even stronger *exponentially Ramsey* property: there exists a positive constant $c(A)$ such that for all $r \in \mathbb{N}$ and $n \sim c(A) \log r$, we have $\mathbb{R}^n \xrightarrow{r} A$, see [FR90]. For the currently best bounds on $c(A)$, we refer the reader to [KSZ26] and the references therein.

To the best of our knowledge, the study of canonical results in Euclidean Ramsey theory was initiated only recently. In [MOW22] the authors proved that every r -coloring of \mathbb{R}^n contains either a monochromatic or a rainbow copy of a fixed rectangle for $n = \Theta(r)$. However, this result is far from optimal, because $n = \Theta(\log r)$ dimensions are already sufficient to ensure a monochromatic copy of a given rectangle as we mentioned above. In fact, it seems hard to avoid rainbow copies of a given set A whenever the number of colors in use is much larger than the cardinality $|A|$. Thus it feels only natural to expect that for most of the Ramsey sets A , if not for

all of them, there exists a finite n such that every coloring of \mathbb{R}^n contains either a monochromatic or a rainbow copy of A regardless of the number of colors in use. In the spirit of classic arrow notation, we denote the latter claim by $\mathbb{R}^n \xrightarrow{\text{MR}} A$ for short. Recently Cheng and Xu [CX25] found the first evidence supporting this intuition. Using our notation, their theorem can be stated as follows:

Theorem 1.1 (Cheng, Xu [CX25]).

- For every acute triangle T , there exists n such that $\mathbb{R}^n \xrightarrow{\text{MR}} T$.
- For every right triangle T , we have $\mathbb{R}^3 \xrightarrow{\text{MR}} T$.

In fact, they proved a stronger statement, which also covers simplices of larger dimension and obtuse triangles that are not ‘too flat’. In the present paper, we refine their technique and strengthen the above result.

Theorem 1.2. For every acute triangle T , we have $\mathbb{R}^3 \xrightarrow{\text{MR}} T$.

Another canonical result obtained by Cheng and Xu in [CX25] covers the case of a square I^2 .

Theorem 1.3 (Cheng, Xu [CX25]). There exists n such that $\mathbb{R}^n \xrightarrow{\text{MR}} I^2$.

Their method probably can be generalized to a cube I^3 but perhaps not further, because for $m \geq 4$, the diameter of the (vertex set of an) m -dimensional hypercube I^m becomes too large in comparison with its sidelength, which seems to be an obstacle. Here we present a completely different approach to bypass this limitation.

Theorem 1.4. For every $m \in \mathbb{N}$, there exists n such that $\mathbb{R}^n \xrightarrow{\text{MR}} I^m$.

Euclidean Ramsey theory questions (along with the corresponding notation) can be translated into non-Euclidean norms in a straightforward manner, with the only difference being the definition of an isometric copy¹, see [EFI21, Dav25, FGST24, Geh23, KS23, Kup11, Rai04, Vor23]. The case of an n -dimensional space \mathbb{R}_∞^n equipped with the max-norm² seems to stand out here due to the following simple classification of finite (exponentially) Ramsey sets. It is not hard to deduce from the Hales–Jewett theorem that for every $r \in \mathbb{N}$ and a finite $M \subset \mathbb{R}^d$, there exists n such that $\mathbb{R}_\infty^n \xrightarrow{r} M$. In fact, the latter claim holds already for $n = \Theta(\log r)$ as it was shown in [KS21, FKS24]. In the present paper, we deduce the following general canonical result in max-norm Ramsey theory from the canonical Hales–Jewett theorem.

Theorem 1.5. For every finite $M \subset \mathbb{R}^d$, there exists n such that $\mathbb{R}_\infty^n \xrightarrow{\text{MR}} M$.

Paper outline. In Section 2, we prove Theorems 1.2 and 1.4. In Section 3, we formally state the canonical Hales–Jewett theorem and use it to prove Theorem 1.5. Finally, in Section 4, we make some further comments and state more open problems.

¹For a norm N on \mathbb{R}^n , a set $A' \subset \mathbb{R}^n$ is an N -isometric copy of $A \subset \mathbb{R}^n$ if there exists a bijection $f : A \rightarrow A'$ such that $\|\mathbf{x} - \mathbf{y}\|_N = \|f(\mathbf{x}) - f(\mathbf{y})\|_N$ for all $\mathbf{x}, \mathbf{y} \in A$.

²Recall that for $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$, its max-norm is defined by $\|\mathbf{x}\|_\infty = \max_i |x_i|$

2. Euclidean norm

2.1. Triangles – proof of Theorem 1.2

Let T be an acute triangle with sides a, b, c . Consider an arbitrary coloring of \mathbb{R}^3 and assume for a contradiction that there is no monochromatic or rainbow copy of T . Then some two points, which we denote by Y_1 and Y_2 , are of different colors. Take a sequence $Y_1 = X_0, X_1, \dots, X_k = Y_2$ such that consecutive points are at distance c . Then, there are two consecutive points, $X_i = A$ and $X_{i+1} = B$, of different colors, say, A is red and B is blue.

We say that a pair of points (X, Y) is a *good pair* if they are at distance c , one of them is red while the other one is blue. In particular, (A, B) is a good pair. Let X, Y, U, V be points such that $|XY| = |UV| = c$, $|XU| = |YV| = b$, $|XV| = |YU| = a$. That is, $XYUV$ is a simplex, all of whose faces are congruent to T . Then we say that the pair (U, V) is an *extension* of the pair (X, Y) .

Claim 2.1. *If (X, Y) is a good pair, and (U, V) is its extension, then (U, V) is also a good pair.*

Proof. Suppose without loss of generality that X is red and Y is blue. Since both UXY and VXY are congruent to T , both U and V are red or blue. But UVX and UVY are also congruent to T , so one of U, V is red, and the other one is blue. \square

Claim 2.2. *Every good pair has an extension.*

Proof. Let (A, B) be a good pair. Take a ball of radius b (resp. a) about A (resp. B). Their intersection is the circle C_1 , its plane is perpendicular to AB , and its center is on the segment AB since T is an acute triangle. Similarly, take a ball of radius b (resp. a) about B (resp. A). Their intersection is the circle C_2 , its plane is perpendicular to AB , and its center is on the segment AB . Fix an arbitrary $C \in C_1$, and let X (resp. Y) be the point of C_2 closest (resp. farthest) from C . We have $|CX| < |AB|$. Indeed, if $a = b$, then it is trivial because $C = X$. Otherwise, it follows from the fact that CX and AB are parallel and the orthogonal projections of X and C to the line AB are inside the segment AB , see Figure 2.1. The point Y is the reflection of C through the midpoint O of the segment AB . Since $\angle ACB = \angle AYB < 90^\circ$, both C and Y are outside the ball of center O and radius $|AB|/2$. Consequently $|CY| > |AB|$. Therefore, there is a point D on C_2 with $|CD| = |AB|$. The pair (C, D) is the desired extension. \square

Lemma 2.3. *Let l_1 and l_2 be two skew lines in \mathbb{R}^3 . Suppose that the set $S \subset \mathbb{R}^3$ is invariant under rotations about l_1 and l_2 . Then $S = \emptyset$ or $S = \mathbb{R}^3$.*

Proof. We will use the following notation: E^+ is the group of all orientation-preserving isometries of \mathbb{R}^3 . For any point P , $SO(P)$ is the group of all orientation-preserving isometries that do not move the point P . For any line l , $R(l)$ is the group of all rotations about l . The subgroup of translations of \mathbb{R}^3 is denoted by T .

Suppose that l_1 and l_2 are two skew lines in \mathbb{R}^3 and S is invariant for the groups $R(l_1)$ and $R(l_2)$. Then it is also invariant for the generated group $G = \langle R(l_1), R(l_2) \rangle$. We claim that $G = E^+$, which is sufficient to show that either $S = \emptyset$ or $S = \mathbb{R}^3$. Indeed, if for some




Figure 2.1: Segment AB is longer than CX and shorter than CY .

points $X, Y \in \mathbb{R}^3$, we have $X \in S$ but $Y \notin S$, then S is not invariant for a translation by the vector \overrightarrow{XY} , a contradiction.

Rotate the line l_1 about l_2 . We get a hyperboloid H_1 which contains l_1 , and its axis is l_2 (if l_1 and l_2 are orthogonal, our hyperboloid is degenerate, it is a plane minus a disc, but the argument works also in this case). Similarly, rotate l_2 about l_1 . We get a hyperboloid H_2 which contains l_2 , and its axis is l_1 . It follows, that the two hyperboloids intersect. Therefore, there is a rotated copy l'_1 of l_1 , and a rotated copy l'_2 of l_2 , that intersect in a point P . It is not hard to see that a rotation about l'_1 is equal to a conjugate of the rotation about l_1 by the same angle with an element of $R(l_2)$ that transforms l_1 into l'_1 . Hence $R(l'_1) \subset G$. Similarly, $R(l'_2)$ is a subgroup of G , and thus so is $G' = \langle R(l'_1), R(l'_2) \rangle$.

We claim that $G' = SO(P)$. The inclusion $G' \subseteq SO(P)$ is obvious. Take a unit sphere \mathcal{S} of center P . The groups $R(l'_1), R(l'_2), SO(P)$ are orientation-preserving isometries of \mathcal{S} . It is known [Ced89, Theorem 3.20] that each element of $SO(P)$ is a rotation about a line through P . Let Q_1 (resp. Q_2) be an intersection of the line l'_1 (resp. l'_2) and \mathcal{S} . Consider the orbit $O(Q_1)$ of the point Q_1 under the action of G' . It contains at least one point Q'_1 different from Q_1 . As earlier, each element of $R(PQ'_1)$ is a conjugate of the corresponding element of $R(l'_1)$ with an element of G' that transforms Q_1 into Q'_1 . Hence $R(PQ'_1) \subset G'$. If $\angle Q_1PQ'_1 < 90^\circ$, then rotate Q_1 about the line PQ'_1 by 180° , let Q''_1 be the image. Then Q''_1 is also in $O(Q_1)$, and $\angle Q_1PQ''_1 = 2\angle Q_1PQ'_1$. By repeating this procedure if necessary, we get a point $V_1 \in O(Q_1)$ such that $\angle Q_1PV_1 \geq 90^\circ$. All elements G' are finite products of rotations about lines $l'_1 = Q_1P$ and $l'_2 = Q_2P$. So, we can decrease all rotation angles continuously to 0. This shows that G' is connected, and we can find $V'_1 \in O(Q_1)$ such that $\angle Q_1PV'_1 = 90^\circ$. But the rotations about two orthogonal axes, $R(PQ_1)$ and $R(PV'_1)$, act transitively on \mathcal{S} . Therefore, we can get all rotation axes, and, consequently, all rotations. This proves that $G' = SO(P)$.

Consider the lines l_1 and l'_1 . Line l_1 does not contain P , but l'_1 does. Applying an appropriate element of $SO(P)$, we can transform l'_1 into a line l''_1 which is parallel to l_1 , but they are not identical. Clearly, $R(l''_1) \subset G$. Take the composition of rotation about l_1 by angle α and a rotation about l''_1 by angle $-\alpha$. The result is a translation $T(\alpha)$. As α goes to 0, the translation distance

of $T(\alpha)$ also goes to 0 continuously. So, we can obtain a translation by any given distance, but the direction changes. Taking its conjugate with the elements of $\text{SO}(P)$, we can change the directions of the translations arbitrarily, therefore, we can obtain any translation, so $T \subset G$. Finally, $E^+ = T \cdot \text{SO}(P)$, therefore, $G = E^+$, as desired. \square

Return to the proof of Theorem 1.2. Let $S' \subset \mathbb{R}^3 \times \mathbb{R}^3$ be the following set of pairs of points. Let the pair (A, B) be in S' , and take its closure under the extension operation. In other words, the elements of S' are exactly those pairs that can be obtained by finitely many applications of the extension operation, starting with the pair (A, B) . By Claim 2.1, all pairs in S' are good pairs. Let $S \subset \mathbb{R}^3$ be the set of all points included in at least one pair from S' . Suppose that (C, D) is an extension of (A, B) , and let (C', D') be a rotation of (C, D) about the line AB , by an arbitrary angle. Then (C', D') is also an extension of (A, B) . Therefore, the set S is invariant under rotations about the lines AB and CD . So we can apply Lemma 2.3 to conclude that $S = \mathbb{R}^3$ or $S = \emptyset$. However, S is certainly not empty, as $A, B \in S$, therefore, $S = \mathbb{R}^3$. Since all points of S are either red or blue, we used only two colors for the whole space. Now the following result of Erdős et al. guarantees that some copy of T is monochromatic, which completes the proof of Theorem 1.2 by contradiction.

Theorem 2.4 (Theorem 8 in [EGM⁺73]). *For every triangle T , we have $\mathbb{R}^3 \xrightarrow{2} T$.*

2.2. Hypercubes – proof of Theorem 1.4

For any $A_1 \subset \mathbb{R}^{d_1}, \dots, A_n \subset \mathbb{R}^{d_n}$, their *Cartesian product* $A_1 \times \dots \times A_n$ is $\{(\mathbf{x}_1, \dots, \mathbf{x}_n) : \mathbf{x}_i \in A_i\} \subset \mathbb{R}^{d_1 + \dots + d_n}$. If $A_1 = \dots = A_n = A$, then $A_1 \times \dots \times A_n$ is the *Cartesian power* of A and it is denoted by A^n for short. Let $I^m(a)$ be an m -dimensional hypercube of sidelength \sqrt{a} and $S_m(a)$ be an m -vertex, that is, an $(m - 1)$ -dimensional regular simplex of sidelength \sqrt{a} . Considering the neighbors of a fixed vertex, it is easy to see that $I^m(1)$ contains $S_m(2)$ as a subset. We also need the following asymmetric generalization of the arrow notation: by $C \xrightarrow{\text{MR}}(A, B)$, we denote that claim that every coloring of C contains either a monochromatic copy of A , or a rainbow copy of B .

The main result of this section is the following statement.

Proposition 2.5. *For all $k, m \in \mathbb{N}$, there is $n = n_1(k, m)$ such that $I^n(1) \xrightarrow{\text{MR}}(I^k(2^k), I^m(2^k))$.*

Theorem 1.4 follows from the special case $k = m$, i.e., from the claim that $I^n(1) \xrightarrow{\text{MR}} I^m(2^m)$. We prove Proposition 2.5 by induction on k for any fixed m . First we prove the base case.

Proposition 2.6. *For all $m \in \mathbb{N}$, there exists $n = n_2(m)$ such that $I^n(1) \xrightarrow{\text{MR}}(I(2), I^m(2))$.*

Proof. We use the probabilistic method to prove the following technical statement.

Claim 2.7. *For all $m, d > 4^m$, the Cartesian power $S_d^m(2)$ satisfies $S_d^m(2) \xrightarrow{\text{MR}}(I(2), I^m(2))$.*

This implies Proposition 2.6 since $S_d^m(2) \subset I^{md}(1)$. Consider an arbitrary coloring of $S_d^m(2)$ and suppose that there is no monochromatic copy of $I(2)$.

Choose an ordered pair of points from each of the m simplices uniformly and independently at random. The Cartesian product C of these ordered 2-point sets is a copy of $I^m(2)$ with a

natural bijection between their points. If C is not rainbow, then some two of its vertices have the same color. For any pair of vertices of $I^m(2)$, the event that the corresponding points of C are of the same color, is called a *bad event*. The *type* of the bad event is the pair of vertices. Note that a non-rainbow C can belong to several types at the same time.

There are $2^m(2^m - 1)/2 < 4^m \leq d - 1$ pairs of vertices of $I^m(2)$, so there are fewer than $d - 1$ types of bad events. Consider now a fixed bad event. The corresponding points, p and q of $I^m(2)$ have different coordinates, assume without loss of generality that their first coordinates are different. We want to estimate the probability that the points of C corresponding to p and q are of the same color. Fix all pairs of points in each of the simplices $S_d(2)$, except in the first one. In the first one, fix only one of the points, say, a , so that the position of the point corresponding to p is already fixed. We still have to take the other point, b , from the first simplex. As we go over all vertices of the first simplex, except a , the possible locations of the point of C that corresponds to q go over the vertices of a simplex $S_{d-1}(2)$. Since no copy of $I(2)$ is monochromatic, this copy of $S_{d-1}(2)$ is rainbow. So we get that the bad event of the given type at most once. This implies that the probability of each bad event is at most $1/(d - 1)$. Now the union bound implies that a random copy of $I^m(2)$ is rainbow with a positive probability, which completes the proof. \square

Before the induction step, we need one more simple statement.

Proposition 2.8. *For all $r, m \in \mathbb{N}$, there exists $n = n_3(r, m)$ such that $I^n(1) \xrightarrow{r} I^m(2)$.*

Proof. We prove the following technical statement by induction.

Claim 2.9. *For all n_1, \dots, n_m such that $n_1 > r, n_2 > n_1^r, \dots, n_m > (n_1 \cdot \dots \cdot n_{m-1})^r$, we have $S_{n_1}(2) \times \dots \times S_{n_m}(2) \xrightarrow{r} I^m(2)$.*

This implies the original statement since $S_{n_1}(2) \times \dots \times S_{n_m}(2) \subset I^n(1)$ for $n = n_1 + \dots + n_m$. The $m = 1$ case is trivial, since among $n_1 > r$ vertices of $S_{n_1}(2)$ some two are of the same color by the pigeonhole principle. To prove the induction step, we consider $S_{n_1}(2) \times \dots \times S_{n_m}(2)$ as the union of n_m ‘layers’ of the form $S_{n_1}(2) \times \dots \times S_{n_{m-1}}(2) \times \{x\}$, $x \in S_{n_m}(2)$. Since $n_m > (n_1 \cdot \dots \cdot n_{m-1})^r$, some two layers are colored identically by the pigeonhole principle. Now the union of a monochromatic copy of $I^{m-1}(2)$ from one of these layers, which must exist by the induction hypothesis, with the corresponding set from the other layer form the desired monochromatic copy of $I^m(2)$. \square

Now we complete the proof of Proposition 2.5 for $k > 1$ by induction. We show that the desired statement holds for

$$n = n_1(k, m) := n' + n_3(r', m'), \text{ where } n' = n_2(m)2^{k-1}, r' = 2^{n'+k}, m' = n_1(k - 1, m).$$

Fix an arbitrary coloring of $I^n(1)$.

For each $\mathbf{x} \in I^{n-n'}(1)$, we consider a ‘layer’ $\{\mathbf{x}\} \times I^{n'}(1) \subset I^n(1)$. By partitioning the n' basic vectors of $I^{n'}(1)$ into $n_2(m)$ classes of size 2^{k-1} and applying Proposition 2.6, we conclude that $I^{n'}(1) \xrightarrow{\text{MR}} (I(2^k), I^m(2^k))$. If at least for one $\mathbf{x} \in I^{n-n'}(1)$, the second alternative holds, that is, there exists a rainbow copy of $I^m(2^k)$ in the layer $\{\mathbf{x}\} \times I^{n'}(1)$, then we are done. So we

can assume without loss of generality that for all $\mathbf{x} \in I^{n-n'}(1)$, there exists a monochromatic copy of $I(2^k)$ in the layer $\{\mathbf{x}\} \times I^{n'}(1)$.

Consider the auxiliary r' -coloring of $I^{n-n'}(1)$, where each vertex \mathbf{x} is colored according to which of the r' copies of $I(2^k)$ in the layer $\{\mathbf{x}\} \times I^{n'}(1)$ is monochromatic. If the layer contains several monochromatic copies of $I(2^k)$, we pick one of them arbitrarily. Now Proposition 2.8 implies that there exists a copy of $I^{m'}(2)$ that is monochromatic under this auxiliary coloring. In terms of the original coloring, this gives us two identically colored copies of $I^{m'}(2)$ that form a ‘prism’ $I^{m'}(2) \times I(2^k)$.

Applying the induction hypothesis to the base of this prism, we find either a monochromatic copy of $I^{k-1}(2 \cdot 2^{k-1})$, or a rainbow copy of $I^m(2 \cdot 2^{k-1})$ there. In the latter case, we are done immediately. In the former case, the union of this monochromatic copy of $I^{k-1}(2^k)$ in the first level of the prism with the corresponding identically colored set from the second level form the desired monochromatic copy of $I^k(2^k)$. This completes the proof of Proposition 2.5.

3. Maximum norm

3.1. Hales–Jewett theorem

In this subsection, we discuss the Hales–Jewett theorem, which is a central result in Ramsey theory and also the main tool of this section. Informally speaking, it states that a sufficiently high dimensional grid, colored with a given number of colors, contains a given dimensional, monochromatic subgrid of a certain type, described below.

For $k, n \in \mathbb{N}$, let τ be an element of $([k] \cup \{*\})^n$ with at least one $*$ -coordinate, where $*$ is an abstract symbol that we call a *variable*. We define a *combinatorial line* corresponding to τ as a subset $\{\tau(i) : i \in [k]\} \subset [k]^n$, where $\tau(i)$ is an element of $[k]^n$ that we obtain by replacing all the variable coordinates of τ by i . More generally, let τ be the element of $([k] \cup \{*_1, \dots, *_m\})^n$ with at least one variable coordinate of each of the m types. We define an *m -dimensional combinatorial subspace* corresponding to τ as a subset $\{\tau(i_1, \dots, i_m) : i_1, \dots, i_m \in [k]\} \subset [k]^n$, where $\tau(i_1, \dots, i_m)$ is an element of $[k]^n$ that we obtain by replacing all the variable coordinates of τ of the j -th type by i_j for each $j \in [m]$. In this notation, the multidimensional Hales–Jewett theorem is the following statement.

Theorem 3.1 (Multidimensional Hales–Jewett theorem). *For all $k, r, m \in \mathbb{N}$, there is a sufficiently large $n \in \mathbb{N}$ such that for every r -coloring of $[k]^n$, there exists a monochromatic m -dimensional combinatorial subspace.*

Hales and Jewett [HJ63] proved this theorem for $m = 1$ in 1963, while the multidimensional generalization is due to Graham and Rothschild [GR71]. Furstenberg and Katznelson [FK91] proved a density version of this theorem using ergodic theory, while in 2009, the Polymath Project [Pol12] developed a combinatorial proof.

For the canonical version, if $k > 2$, then it is not true that for every coloring of $[k]^n$, there exists either a monochromatic or a rainbow m -dimensional combinatorial subspace even if n is arbitrary large. Indeed, let \sim be an equivalence relation on $[k]$. Consider a coloring c_\sim of $[k]^n$

such that for all $\mathbf{x} = (x_1, \dots, x_n)$ and $\mathbf{y} = (y_1, \dots, y_n)$ in $[k]^n$, we have

$$c_{\sim}(\mathbf{x}) = c_{\sim}(\mathbf{y}) \text{ if and only if } x_1 \sim y_1, \dots, x_n \sim y_n.$$

Observe that the induced coloring on every m -dimensional combinatorial subspace of $[k]^n$ coincides with the coloring c_{\sim} on $[k]^m$ under a natural isomorphism. Therefore, no m -dimensional combinatorial subspace is monochromatic or rainbow, unless \sim is a *trivial* equivalence relation, that is, the elements of $[k]$ are either pairwise equivalent or pairwise non-equivalent. However, Prömel and Voigt [PV83, Theorem C.7] showed that one can always find a combinatorial subspace colored according to some c_{\sim} , see also [NPRV85].

Theorem 3.2 (Canonical Hales–Jewett theorem). *For all $k, m \in \mathbb{N}$, there is a sufficiently large $n \in \mathbb{N}$ such that for every coloring of $[k]^n$, there exists an m -dimensional combinatorial subspace such that the induced coloring on it coincides with c_{\sim} for some equivalence relation \sim on $[k]$.*

3.2. Proof of Theorem 1.5

In this section, we deduce Theorem 1.5 from the Canonical Hales–Jewett theorem. Observe that our finite $M \subset \mathbb{R}^d$ is a subset of a Cartesian power A^d for some finite $A \subset \mathbb{R}$. Indeed, let A be the union of the projections of M onto each of the d coordinate axes. Hence, it is sufficient to prove Theorem 1.5 only for A^d playing the role of M .

Label the elements of A by a_1, \dots, a_s in ascending order. By adding between every two of its consecutive elements sufficiently many equally spaced intermediate points, we obtain a set $B = \{b_1, \dots, b_k\} \supset A$ such that

$$\max_{1 \leq i < k} (b_{i+1} - b_i) \leq \min_{1 \leq j < s} (a_{j+1} - a_j), \tag{3.1}$$

where the elements of B are also labeled in ascending order, see Figure 3.1.

Let n be from the statement of Theorem 3.2 applied to our k and $m = d + \lceil d \log_2 s \rceil$. In what follows, we prove that every coloring of B^n contains either a monochromatic or a rainbow ℓ_{∞} -isometric copy of A^d , which implies that $\mathbb{R}_{\infty}^n \xrightarrow{\text{MR}} A^d$. Consider an arbitrary coloring of B^n . A natural bijection φ between $[k]^n$ and B^n defined by

$$\varphi(x_1, \dots, x_n) = (b_{x_1}, \dots, b_{x_n}) \text{ for all } (x_1, \dots, x_n) \in [k]^n$$

translates that coloring into a coloring of $[k]^n$. Now Theorem 3.2 implies that the m -dimensional combinatorial subspace corresponding to some $\tau \in ([k] \cup \{*_1, \dots, *_m\})^n$ is colored according to c_{\sim} for some equivalence relation \sim on $[k]$.

If all the elements of $[k]$ are pairwise equivalent with respect to \sim , i.e., if the combinatorial space is monochromatic, then it is easy to complete the proof. Indeed, consider the subset of our combinatorial space where the last $m - d$ variables stay fixed, say, $*_{d+1} = \dots = *_m = 1$, while the first d variables range over the indices of the elements of A inside B . It is not hard to see that the image of this subset under φ is the desired monochromatic ℓ_{∞} -isometric copy of A^d inside B^n . More precisely, let $1 = i_1 < \dots < i_s = k$ be a sequence such that $a_j = b_{i_j}$ for

all $j \in [s]$, and put $I = \{i_1, \dots, i_s\} \subset [k]$, see Figure 3.1. Let \mathcal{I}_1 be the set of all s^d points of $[k]^m$ such that their first d coordinates belong to I , while the remaining $m - d$ coordinates are equal to 1. Note that $\tau(\mathcal{I}_1) \subset [k]^n$ is a subset of our combinatorial subspace, and thus it is monochromatic. Moreover, $\varphi(\tau(\mathcal{I}_1)) \subset B^n$ is an ℓ_∞ -isometric copy of A^d , because the ℓ_∞ -norm is independent of the arguments' multiplicities, i.e., the number of $*_r$ -coordinates of τ for all $r \in [d]$, as desired.




Figure 3.1: For $A = \{0, 16, 20, 26\}$, we can take $B = \{0, 4, 8, 12, 16, 20, 23, 26\}$, $I = \{1, 5, 6, 8\}$.

Otherwise, there exists $1 \leq i' < k$ such that $i' \not\sim i' + 1$. Let σ be an arbitrary injection from I^d to $\{i', i' + 1\}^{m-d}$, where I is from the previous paragraph. Such an injection exists because $|I^d| = s^d \leq 2^{m-d} = |\{i', i' + 1\}^{m-d}|$ by our choice of m . Consider $\mathcal{I}_2 = \{(\mathbf{x}, \sigma(\mathbf{x})) : \mathbf{x} \in I^d\} \subset [k]^m$. Note that every two different elements of \mathcal{I}_2 have at least one non-equivalent coordinate, and thus the set $\tau(\mathcal{I}_2) \subset [k]^n$ is rainbow. Moreover, it is not hard to check that $\varphi(\tau(\mathcal{I}_2)) \subset B^n$ is an ℓ_∞ -isometric copy of A^d . Indeed, in the previous paragraph, we have already seen this when the last $m - d$ variable coordinates were fixed. Now they are not fixed, but vary between i' and $i' + 1$. However, (3.1) implies that $b_{i'+1} - b_{i'}$ is not larger than the smallest ℓ_∞ -distance between different elements of A^d . Therefore, these additional variable coordinates do not change any of the ℓ_∞ -distances. This completes the proof Theorem 1.5.

4. Concluding remarks

Canonical types. For any finite set A , let $\mathbb{R}^n \xrightarrow{\text{MR}}^r A$ denote the claim that every r -coloring of \mathbb{R}^n contains a monochromatic or a rainbow copy of A . Clearly, if $\mathbb{R}^n \xrightarrow{\text{MR}}^r A$ then also $\mathbb{R}^{n+1} \xrightarrow{\text{MR}}^r A$ and $\mathbb{R}^n \xrightarrow{\text{MR}}^{r-1} A$. Therefore, we can distinguish the following three types of sets, see Figure 4.1.

1. There is r such that $\mathbb{R}^n \xrightarrow{\text{MR}}^r A$ for every n .
2. There is n such that $\mathbb{R}^n \xrightarrow{\text{MR}}^r A$ for every r .
3. None of the above, i.e., for every r there is n with $\mathbb{R}^n \xrightarrow{\text{MR}}^r A$, and for every n there is r with $\mathbb{R}^n \not\xrightarrow{\text{MR}}^r A$.

It is clear that Ramsey sets cannot be of type 1, and it is natural to ask whether they are all of type 2. In fact, we cannot fully answer this question even if the Ramsey set A is a rectangle or an obtuse triangle. Our Theorem 1.4 implies that rectangles with sidelengths a and b are of type 2, provided that $(b/a)^2$ is rational, but we could not determine a type of the remaining rectangles. As we mentioned in the Introduction, Cheng and Xu [CX25] proved that the obtuse triangles that are not 'too flat' are of type 2. The type of 'flat' obtuse triangles remains uncertain. We do not have an example of a Ramsey set of type 3.




Figure 4.1: All three possible types of finite sets.

For non-Ramsey sets we cannot rule out that all three types are possible. As it was observed in [CX25], a classical spherical 3-coloring of \mathbb{R}^n without monochromatic copies of a 4-term unit arithmetic progression l_4 from [EGM⁺73] cannot contain a rainbow copy either, so l_4 is of type 1. We are not aware of any non-Ramsey set of other types. A 3-term unit arithmetic progression l_3 might be natural candidate for type 2 (with n being 2 or 3), since it was recently shown in [CMY24] that $\mathbb{R}^2 \xrightarrow{2} l_3$. A similar question was raised in [MP24, Section 4].

Tightness of Theorem 1.2. A classical 2-coloring of the plane with color-alternating strips of heights $\sqrt{3}/2$ does not contain monochromatic copies of a unit equilateral triangle \triangle . Hence, it is *not* true that $\mathbb{R}^2 \xrightarrow{MR} \triangle$. Are there other triangles T for which the dimension 3 in Theorem 1.2 is also tight? It was conjectured in [EGM⁺75b, Conjecture 3] that $\mathbb{R}^2 \xrightarrow{2} T$ for every non-equilateral triangle T . If true, this would be an evidence supporting that $\mathbb{R}^2 \xrightarrow{MR} T$ might hold for all non-equilateral triangles T , and thus our result is not tight. However, during the last 50 years, that conjecture was verified only for a few special families of triangles, see [NS24] and the references therein.

Quantitative bounds in Theorem 1.4. What is the minimum $n = n(m)$ such that $\mathbb{R}^n \xrightarrow{MR} I^m$? A careful analysis of our proof leads to the upper bound $n(m) = \exp_2^{2^{m-1}}(O(m))$, where $\exp_2^1(x) = 2^x$ and $\exp_2^{k+1}(x) = 2^{\exp_2^k(x)}$ for all $k > 1$. However, our argument is very wasteful in a sense that all the auxiliary propositions are stated for the hypercubes, but in fact we are working only with their small subsets: Cartesian product of simplices. Taking this into account, we can improve the upper bound to $n(m) = \exp_2^4(O(m))$, which is still a quadruple exponent. From the other direction, we know that $n(m) = \Omega(m \log m)$, since for smaller n , there exists a coloring of \mathbb{R}^n with no monochromatic isometric copies of I^m that uses less than 2^m colors, see [Pro18].

The case $m = 2$ is of a particular interest here. We can show that $K_{876} \square K_{123} \square K_4 \square K_2 \xrightarrow{MR} K_2 \square K_2$, where \square stands for the Cartesian product of these cliques. A straightforward translation of this purely graph theoretical statement into geometric language implies that $n(2) \leq 875 + 122 + 3 + 1 = 1001$. We do not know what is the best upper bound on $n(2)$ that can be derived from [CX25], but it might be of the same order. From the other direction, we do not even know if $n(2) > 3$.

Max-norm Ramsey theory. Though our Theorem 1.5 is rather general, the quantitative bounds on the minimum dimension $n = n(M)$ such that $\mathbb{R}_{\infty}^n \xrightarrow{\text{MR}} M$ it generates are probably extremely far from being tight. For instance, if M is a rectangle with sidelengths $a < b$, then the number of auxiliary points on Figure 3.1, as well as the resulting upper bound, will depend on the fraction b/a . We found an ad hoc argument for this case in the spirit of [CX25] which leads to the upper bound $O(b/a)$, i.e., this dependence is at most linear. Does there exist a universal constant n such that $\mathbb{R}_{\infty}^n \xrightarrow{\text{MR}} R$ for every rectangle R ?

Manhattan Ramsey theory. Let \mathbb{R}_1^n be the n -dimensional space equipped with the Manhattan ℓ_1 -norm. It is not hard to show that for every r and a finite $M \subset \mathbb{R}^d$, there exists n such that $\mathbb{R}_1^n \xrightarrow{r} M$, see [KSZ26]. Our Proposition 2.5 has the following canonical corollary: for every finite $M \subset \mathbb{Q}^d$, there exists n such that $\mathbb{R}_1^n \xrightarrow{\text{MR}} M$. Indeed, it is sufficient to note that a path of m pairwise orthogonal edges in the hypercube is an ℓ_1 -isometric copy of an arithmetic progression. Besides that, every ‘rational’ M can be embedded into a Cartesian power of this progression after a proper scaling. We wonder, if the same conclusion holds for ‘non-rational’ sets M as well, e.g. for all rectangles.

Acknowledgements

We are very grateful to Endre Szabó and Balázs Csikós for their help in the presentation of Lemma 1. Arsenii Sagdeev thanks Alexander Golovanov for the helpful discussion of Lemma 1.

References

- [AJ05] Maria Axenovich and Robert E. Jamison. Canonical pattern Ramsey numbers. *Graphs and Combinatorics*, 21:145–160, 2005. doi:10.1007/s00373-005-0603-6.
- [Can07] Kristal Cantwell. All regular polytopes are Ramsey. *Journal of Combinatorial Theory, Series A*, 114(3):555–562, 2007. doi:10.1016/j.jcta.2006.08.001.
- [Ced89] Judith N. Cederberg. *A Course in Modern Geometries*. Undergraduate Texts in Mathematics. Springer Science & Business Media, New York, 1989.
- [CMY24] Gabriel Currier, Kenneth Moore, and Chi Hoi Yip. Any two-coloring of the plane contains monochromatic 3-term arithmetic progressions. *Combinatorica*, 44(6):1367–1380, 2024. doi:10.1007/s00493-024-00122-2.
- [CX25] Xinbu Cheng and Zixiang Xu. Euclidean Gallai–Ramsey for various configurations. *Discrete & Computational Geometry*, 73:1037–1052, 2025. doi:10.1007/s00454-024-00693-3.
- [Dav25] James Davies. Chromatic number of spacetime. *Acta Arithmetica*, 218(1):65–76, 2025. doi:10.4064/aa240306-20-12.
- [dG18] Aubrey D. N. J. de Grey. The chromatic number of the plane is at least 5. *Geombinatorics*, 28(1):18–31, 2018. arXiv:1804.02385.

- [EFI21] Geoffrey Exoo, David Fisher, and Dan Ismailescu. The chromatic number of the Minkowski plane – the regular polygon case. *Geombinatorics*, 31:49–67, 2021. arXiv:2108.12861.
- [EG80] Paul Erdős and Ronald L. Graham. *Old and New Problems and Results in Combinatorial Number Theory*, volume 28. L'Enseignement Mathématique, Université de Genève, Geneva, Switzerland, 1980.
- [EGM⁺73] Paul Erdős, Ronald L. Graham, Peter Montgomery, Bruce L. Rothschild, Joel H. Spencer, and Ernst G. Straus. Euclidean Ramsey theorems i. *Journal of Combinatorial Theory, Series A*, 14(3):341–363, 1973. doi:10.1016/0097-3165(73)90011-3.
- [EGM⁺75a] Paul Erdős, Ronald L. Graham, Peter Montgomery, Bruce L. Rothschild, Joel H. Spencer, and Ernst G. Straus. Euclidean Ramsey theorems ii. In *Infinite and Finite Sets*, Colloquium Mathematicum Societatis János Bolyai, pages 520–557. North-Holland, 1975.
- [EGM⁺75b] Paul Erdős, Ronald L. Graham, Peter Montgomery, Bruce L. Rothschild, Joel H. Spencer, and Ernst G. Straus. Euclidean Ramsey theorems iii. In *Infinite and Finite Sets*, Colloquium Mathematicum Societatis János Bolyai, pages 559–583. North-Holland, 1975.
- [EI20] Geoffrey Exoo and Dan Ismailescu. The chromatic number of the plane is at least 5: A new proof. *Discrete & Computational Geometry*, 64(1):216–226, 2020. doi:10.1007/s00454-019-00058-1.
- [ER50] Paul Erdős and Richard Rado. A combinatorial theorem. *Journal of the London Mathematical Society*, 25(4):249–255, 1950. doi:10.1112/jlms/s1-25.4.249.
- [FGST24] Nóra Frankl, Panna Gehér, Arsenii Sagdeev, and Géza Tóth. Monochromatic infinite sets in Minkowski planes. *Discrete & Computational Geometry*, 74:1–15, 2024. doi:10.1007/s00454-024-00702-5.
- [FK91] Hillel Furstenberg and Yitzhak Katznelson. A density version of the Hales–Jewett theorem. *Journal d'Analyse Mathématiques*, 57(1):64–119, 1991. doi:10.1007/BF03041066.
- [FKS24] Nóra Frankl, Andrey Kupavskii, and Arsenii Sagdeev. Max-norm Ramsey theory. *European Journal of Combinatorics*, 118:103918, 2024. doi:10.1016/j.ejc.2024.103918.
- [FR90] Peter Frankl and Vojtěch Rödl. A partition property of simplices in Euclidean space. *Journal of the American Mathematical Society*, 3(1):1–7, 1990. doi:10.1090/S0894-0347-1990-1020148-2.
- [Ga20] António Girão. A canonical polynomial Van der Waerden's theorem. 2020. arXiv:2004.07766.

- [Geh23] Panna Gehér. Note on the chromatic number of Minkowski planes: the regular polygon case. *Studia Scientiarum Mathematicarum Hungarica*, 60(1):27–42, 2023. doi:10.1556/012.2023.01536.
- [GKP12] William Gasarch, Clyde Kruskal, and Andy Parrish. Van der Waerden’s theorem: Variants and applications. manuscript, 2012.
- [GR71] Ronald L. Graham and Bruce L. Rothschild. Ramsey’s theorem for n -parameter sets. *Transactions of the American Mathematical Society*, 159:257–292, 1971. doi:10.1090/S0002-9947-1971-0284352-8.
- [GRS91] Ronald L. Graham, Bruce L. Rothschild, and Joel H. Spencer. *Ramsey Theory*, volume 20 of *Wiley Series in Discrete Mathematics and Optimization*. John Wiley & Sons, New York, 1991.
- [HJ63] Alfred W. Hales and Robert I. Jewett. Regularity and positional games. *Transactions of the American Mathematical Society*, 106(2):222–229, 1963. doi:10.1090/S0002-9947-1963-0143712-1.
- [JL03] Robert E. Jamison, Tao Jiang, and Alan C. H. Ling. Constrained Ramsey numbers of graphs. *Journal of Graph Theory*, 42(1):1–16, 2003. doi:10.1002/jgt.10072.
- [JM00] Tao Jiang and Dhruv Mubayi. New upper bounds for a canonical Ramsey problem. *Combinatorica*, 20(1):141–146, 2000. doi:10.1007/s004930070037.
- [KS21] Andrey Kupavskii and Arsenii Sagdeev. All finite sets are Ramsey in the maximum norm. *Forum Mathematicum, Sigma*, 9:e55, 2021. doi:10.1017/fms.2021.50.
- [KS23] Valeriya Kirova and Arsenii Sagdeev. Two-colorings of normed spaces without long monochromatic unit arithmetic progressions. *SIAM Journal on Discrete Mathematics*, 37(2):718–732, 2023. doi:10.1137/22M1483700.
- [KSZ26] Andrey Kupavskii, Arsenii Sagdeev, and Dmitrii Zakharov. Cutting corners. *J. Combin. Theory Ser. B*, 177:273–292, 2026. doi:10.1016/j.jctb.2025.11.008.
- [Kup11] Andrey Kupavskiy. On the chromatic number of \mathbb{R}^n with an arbitrary norm. *Discrete Mathematics*, 311(6):437–440, 2011. doi:10.1016/j.disc.2010.12.005.
- [Kř1] Igor Kříž. Permutation groups in Euclidean Ramsey theory. *Proceedings of the American Mathematical Society*, 112(3):899–907, 1991. doi:10.2307/2048715.
- [LRW12] Imre Leader, Paul A. Russell, and Mark Walters. Transitive sets in Euclidean Ramsey theory. *Journal of Combinatorial Theory, Series A*, 119(2):382–396, 2012. doi:10.1016/j.jcta.2011.09.005.
- [MOW22] Yaping Mao, Kenta Oeki, and Zhao Wang. Euclidean Gallai–Ramsey theory. 2022. arXiv:2209.13247.
- [MP24] Isabel McGuigan and Katherine Pan. Bipartite and Euclidean Gallai–Ramsey theory. 2024. arXiv:2410.07634.

- [NPRV85] Jaroslav Nešetřil, Hans Jürgen Prömel, Vojtěch Rödl, and Bernd Voigt. Canonizing ordering theorems for Hales–Jewett structures. *Journal of Combinatorial Theory, Series A*, 40(2):394–408, 1985. doi:10.1016/0097-3165(85)90098-6.
- [NS24] Alexander Natalchenko and Arsenii Sagdeev. Monochromatic triangles in the max-norm plane. *European Journal of Combinatorics*, 120:103977, 2024. doi:10.1016/j.ejc.2024.103977.
- [Pol12] D. H. J. Polymath. A new proof of the density Hales–Jewett theorem. *Annals of Mathematics*, 175(3):1283–1327, 2012. doi:10.4007/annals.2012.175.3.6.
- [Pro18] Roman I. Prosanov. Upper bounds for the chromatic numbers of Euclidean spaces with forbidden Ramsey sets. *Mathematical Notes*, 103:243–250, 2018. doi:10.1134/S000143461801025X.
- [PV83] Hans Jürgen Prömel and Bernd Voigt. Canonical partition theorems for parameter sets. *Journal of Combinatorial Theory, Series A*, 35(3):309–327, 1983. doi:10.1016/0097-3165(83)90016-X.
- [Rai04] Andrey M. Raigorodskii. The chromatic number of a space with the metric ℓ_q . *Russian Mathematical Surveys*, 59(5):973–974, 2004. doi:10.1070/RM2004v059n05ABEH000784.
- [Soi09] Alexander Soifer. *The Mathematical Coloring Book*. Springer-Verlag, New York, 2009.
- [VdW27] B. L. Van der Waerden. Beweis einer baudetschen vermutung. *Nieuw Archief voor Wiskunde*, 15:212–216, 1927.
- [Vor23] Vsevolod Voronov. The chromatic number of the plane with an interval of forbidden distances is at least 7. 2023. arXiv:2304.10163.
- [Wag06] Peter Wagner. An upper bound for constrained Ramsey numbers. *Combinatorics, Probability and Computing*, 15(4):619–626, 2006. doi:10.1017/S0963548305007467.