

A GENERALIZED FRAMEWORK FOR HIGHER-ORDER LOCALIZED ORTHOGONAL DECOMPOSITION METHODS

MORITZ HAUCK¹, ALEXEI LOZINSKI^{2,*} AND ROLAND MAIER¹

Abstract. We revisit the higher-order Localized Orthogonal Decomposition variant by Maier [*SIAM J. Numer. Anal.* **59** (2021) 1067–1089] based on nonconforming constraints (discontinuous finite element spaces) and introduce a new variant based on conforming constraints (continuous finite elements), putting both approaches in a general unified framework. We propose a new localization strategy that is suitable for both approaches and offers a new perspective on the localization of LOD in general. We fully analyze the strategy for linear scalar elliptic problems and discuss extensions to the Helmholtz equation and the Gross–Pitaevskii eigenvalue problem. Numerical examples are presented that provide valuable comparisons between conforming and nonconforming constraints.

Mathematics Subject Classification. 65N12, 65N15, 65N30.

Received July 30, 2025. Accepted January 13, 2026.

1. INTRODUCTION

The numerical solution of partial differential equations (PDEs) with strongly heterogeneous and highly oscillatory coefficients is challenging. Classical finite element methods often fail to produce accurate results unless the fine-scale variations of the coefficients are fully resolved. However, globally resolving such fine-scale features can be prohibitively expensive. The associated computational cost and memory requirements often exceed the capabilities of available computing resources. To address this, numerous multiscale methods have been developed, particularly in the elliptic setting. These include the Heterogeneous Multiscale Method [2, 54, 55], (Generalized) Multiscale Finite Element Methods [6, 7, 10, 22, 38], Multiscale Spectral Generalized Finite Element Methods [6, 40], rough polyharmonic splines [50], the Localized Orthogonal Decomposition (LOD) [36, 44], and gamblets [48]. More recently, refined localization strategies within the LOD framework have been proposed; see, for instance, [26, 32]. Comprehensive overviews of numerical multiscale methods can be found in the textbooks [45, 49] and the review article [4].

While the techniques discussed above typically yield first-order convergence behavior, also higher-order multiscale methods have been developed. In the context of the Heterogeneous Multiscale Method, such extensions have been proposed in [1, 39]. For the Multiscale Finite Element Method, higher-order convergence rates are

Keywords and phrases. Multiscale method, localized orthogonal decomposition, higher-order, *a priori* error analysis, exponential decay.

¹ Institute for Applied and Numerical Mathematics, Karlsruhe Institute of Technology, Englerstr. 2, 76131 Karlsruhe, Germany.

² Université Marie et Louis Pasteur, CNRS, LmB (UMR 6623), F-25000 Besançon, France.

*Corresponding author: alexei.lozinski@univ-fcomte.fr

achieved in [3, 37]. Hybrid multiscale methods, which reduce global degrees of freedom to element boundaries, have also become quite popular to achieve higher-order rates; see, *e.g.*, [5, 18, 29]. However, all these higher-order approaches require certain smoothness assumptions on the domain, the coefficient, and/or the exact solution to obtain convergence rates beyond first order. For rough coefficients, as they arise in many applications (*e.g.*, in the context of composite materials, that do not have a smooth transition between materials properties), such conditions are typically not fulfilled and coefficients are only in L^∞ .

Here, we focus on higher-order variants of LOD methods, which are able to achieve optimal convergence assuming only sufficient smoothness of the right-hand side, without any smoothness or structural assumptions on the coefficients. The LOD method was first introduced in [44], where the authors defined the prototypical approximation space as the orthogonal complement of the fine-scale space, that is, the kernel of a suitable quasi-interpolation operator mapping onto the conforming piecewise linear finite element space. The resulting method was of first order and the construction could not easily be adapted to achieve higher orders of convergence. However, in some of the later contributions, for instance, the review article [4], the use of quasi-interpolation was abandoned in favor of a more general concept of *quantities of interest* (QOI), closely related to the construction of gamblets in [49]: the fine-scale space is now seen as the space of functions on which the QOI vanish. This concept turns out flexible enough to achieve higher-order convergence rates. In particular, the higher-order multiscale method in [42] (see also [41]) can be cast in this framework, taking the QOI as the L^2 -inner products with the basis functions of the nonconforming, discontinuous finite element space (polynomial functions of a given order on each cell of a given mesh). This leads to the natural question: can one instead use continuous finite elements to define the QOI, thereby reducing the number of degrees of freedom, while maintaining the convergence rates of the resulting method?

In the present article, we give a positive answer to this question by revisiting the construction of the LOD method based on QOI. We show that this framework naturally leads to variants of the LOD in which the fine-scale space is defined using either continuous or discontinuous finite elements of arbitrary order. We refer to these variants as CG-LOD and DG-LOD, respectively. Both achieve optimal higher-order convergence rates, in contrast to certain earlier approaches in the literature concerning CG-LOD-type methods. In particular, the method proposed in [35], which closely resembles the lowest-order CG-LOD considered here, has a convergence order of greater than one, but only under the restrictive assumption that the right-hand side vanishes on the boundary. We argue that achieving optimal rates without any additional restrictions on the right-hand side is only possible if the underlying finite element space used for defining the QOI is not restricted on the boundary. This space should remain nonconforming, *i.e.*, not a subspace of $H_0^1(\Omega)$ for a typical problem with Dirichlet boundary conditions, both for CG and DG variants of the LOD method. This is one of the key observations of our article. Incidentally, we apply it to the “classical” version of the LOD, cf. [45]. We prove that one can increase its convergence rate just by adding a few degrees of freedom associated to the boundary of the domain.

Moreover, we give a new perspective on localization strategies for both CG- and DG-LOD variants (and, in a wider scope, to other possible LOD-like methods). Historically, the first “naive” localization in [44] consisted in truncating the problems for each basis functions to a patch of ℓ layers of elements around the node to which the basis function is associated. This resulted in a deterioration of the error (both theoretically and experimentally) when refining the mesh for a fixed parameter ℓ . This drawback was removed by a modification in [36] at the expense of more costly problems to be solved to construct the new basis functions. The procedure of [36] is specific to the construction of the LOD based on a quasi-interpolation operator to a conforming finite element space, since it localizes the corrections of this space. Thus, it is not directly applicable to other variants of the LOD based on the concept of QOI. In the case of the DG-LOD, introduced in [42], an alternative improved localization strategy was proposed in [20]. However, this approach relies on an intricate construction of quasi-interpolation operators and bubble functions tailored to specific constraints. In contrast, we propose here a new, conceptually simple localization strategy that naturally accommodates both CG- and DG-LOD variants of arbitrary order. In principle, the strategy can be generalized to any LOD method based on an arbitrary set of finite element-based QOI. It requires only a readily identifiable quasi-interpolation operator mapping into a lowest-order conforming finite element space, consistent with the given set of QOI and satisfying standard

assumptions, without the need to explicitly construct any tailor-made bubble functions. We illustrate the concepts outlined above both on a theoretical and experimental level, primarily using a scalar multiscale diffusion problem as a model problem. Additionally, we indicate how the underlying principles extend to a broader class of PDEs, including heterogeneous Helmholtz problems and the Gross–Pitaevskii eigenvalue problem. We also provide numerical comparisons of different higher-order LOD variants, with particular focus on the exponential decay rates of the localized basis functions. Note that, in follow-up work to the present article, the proposed localization strategy has been successfully applied to heterogeneous elliptic mixed-dimensional problems [33] and to heterogeneous Stokes problems [30].

The paper is organized as follows. In Section 2, we introduce the elliptic model problem for which we define a prototypical multiscale method in Section 3 that is able to achieve higher-order convergence rates under minimal structural assumptions. We then establish the exponential decay of the basis functions in Section 4 and introduce a practical multiscale method using localized versions of the basis functions in Sections 5 and 6. We also discuss the applicability of the approach to the heterogeneous Helmholtz equation and the Gross–Pitaevskii eigenvalue problem in Section 7. An improved variant of the “classical” LOD of [45] is proposed in Section 8. Finally numerical examples supporting our theoretical results are presented in Section 9.

Notation

Throughout this work, we will write $a \lesssim b$ or $b \gtrsim a$ if it holds that $a \leq Cb$ or $a \geq Cb$, respectively, where $C > 0$ is a constant that is independent of the mesh size H , the oversampling parameter ℓ , and oscillations of the PDE solution u , but can depend on the dimension d , the bounds α and β on the coefficient, and the polynomial degree p . In particular, we do not explicitly track dependencies on p , as we are not considering an asymptotic behavior with respect to p .

2. MODEL PROBLEM

We consider the prototypical second-order elliptic PDE $-\operatorname{div}(A\nabla u) = f$ in weak form, with homogeneous Dirichlet boundary conditions, posed on a polygonal Lipschitz domain $\Omega \subset \mathbb{R}^d$, where $d \in \{2, 3\}$. The matrix-valued coefficient $A \in L^\infty(\Omega, \mathbb{R}^{d \times d})$ is symmetric and positive definite, satisfying for almost all $x \in \Omega$:

$$\alpha|\eta|^2 \leq (A(x)\eta) \cdot \eta \leq \beta|\eta|^2, \quad \forall \eta \in \mathbb{R}^d \tag{2.1}$$

with some constants $0 < \alpha \leq \beta < \infty$, where $|\cdot|$ denotes the Euclidean norm in \mathbb{R}^d . It is important to note that no regularity assumptions are imposed on the coefficient A . The scenario of particular interest here is when the coefficient A is rough, with oscillations across multiple length scales.

The solution space of the considered PDE is the Sobolev space $V := H_0^1(\Omega)$, and the associated bilinear form $a: V \times V \rightarrow \mathbb{R}$ is given by

$$a(u, v) := \int_{\Omega} (A\nabla u) \cdot \nabla v \, dx.$$

The symmetry of A and the ellipticity condition (2.1) ensure that the bilinear form a defines an inner product on the space V . The induced norm, referred to as the energy norm, is defined by $\|\cdot\|_a^2 := a(\cdot, \cdot)$, and it is equivalent to the standard $H^1(\Omega)$ -norm on V . Given a source term $f \in L^2(\Omega)$, the weak formulation of the problem seeks a solution $u \in V$ such that

$$a(u, v) = (f, v)_\Omega, \quad \forall v \in V, \tag{2.2}$$

where $(\cdot, \cdot)_\Omega$ denotes the $L^2(\Omega)$ -inner product. Denoting by $\|\cdot\|_\Omega^2 := (\cdot, \cdot)_\Omega$ the $L^2(\Omega)$ -norm, the Riesz representation theorem ensures the existence and uniqueness of the solution $u \in V$, along with the stability estimate $\|\nabla u\|_\Omega \leq \alpha^{-1}C_F\|f\|_\Omega$, where $C_F > 0$ is the constant in the Friedrichs inequality on the domain Ω .

3. PROTOTYPICAL MULTISCALE METHOD

As mentioned above, classical finite element methods often perform poorly for multiscale problems: they require resolving all microscopic details of the coefficient globally, which can be prohibitively expensive, and they typically suffer from slow convergence due to the low regularity of the underlying multiscale solution. In this section, we introduce a prototypical multiscale method that, under minimal structural assumptions on the coefficient A , yields accurate (optimal-order) approximations even on very coarse meshes. Following the standard construction of the LOD (cf. [44, 45]), we begin by introducing a subspace of V consisting of fine-scale functions with oscillations on length scales smaller than a prescribed parameter H , characterized by certain vanishing (possibly higher-order) weighted averages. To formalize this, we introduce a hierarchy of quasi-uniform, shape-regular, and geometrically conforming meshes $\{\mathcal{T}_H\}_H$, where each mesh \mathcal{T}_H is a finite subdivision of the closure of Ω into closed simplicial¹ elements T . The mesh size parameter $H > 0$ is defined as the maximum diameter of the elements in the mesh \mathcal{T}_H , *i.e.*, $H := \max_{T \in \mathcal{T}_H} \text{diam}(T)$. In what follows, we are primarily interested in the under-resolved regime where meshes are too coarse to capture the microscopic details of the coefficient. Given an arbitrary but fixed polynomial degree $p \in \mathbb{N}$, we denote by M_H the p -th order finite element space with respect to \mathcal{T}_H , associated with either the standard discontinuous Galerkin (DG) method or its continuous Galerkin (CG) counterpart. Specifically, in the DG case, M_H is given by $\mathcal{P}^p(\mathcal{T}_H)$, the space of \mathcal{T}_H -piecewise polynomials of total degree at most p , while in the CG case, the space is defined as $\mathcal{P}^p(\mathcal{T}_H) \cap H^1(\Omega)$. We define the fine-scale space of the LOD as

$$W := \{w \in V : (\mu, w)_\Omega = 0, \forall \mu \in M_H\}. \quad (3.1)$$

The problem-adapted approximation space of the LOD is then given by the orthogonal complement of W with respect to the energy inner product a , *i.e.*,

$$\tilde{V}_H := \{v \in V : a(v, w) = 0, \forall w \in W\}. \quad (3.2)$$

Since W has finite codimension in V , the space \tilde{V}_H is finite-dimensional. More precisely, we shall see in Lemma 3.3 that $\dim(\tilde{V}_H) = \dim(M_H)$. The use of tildes in the notation highlights that these spaces are specifically tailored to the problem at hand. The prototypical LOD method is defined as the Galerkin projection onto \tilde{V}_H , *i.e.*, it seeks $\tilde{u}_H \in \tilde{V}_H$ such that

$$a(\tilde{u}_H, \tilde{v}_H) = (f, \tilde{v}_H)_\Omega, \quad \forall \tilde{v}_H \in \tilde{V}_H. \quad (3.3)$$

Recall that M_H can be either a CG or a DG space, which leads to two distinct definitions of \tilde{V}_H and, consequently, two different (prototypical) LOD methods. We refer to these as the prototypical CG-LOD and DG-LOD methods, respectively. Before stating a convergence result for these methods, we introduce some notation for broken Sobolev spaces. For $s \in \mathbb{N}$, we denote by $H^s(\mathcal{T}_H)$ the space of \mathcal{T}_H -piecewise H^s -functions, with the corresponding broken seminorm $|f|_{s, \mathcal{T}_H}^2 := \sum_{T \in \mathcal{T}_H} |f|_{s, T}^2$, where $|\cdot|_{s, T}$ denotes the seminorm of order s of a function in $H^s(T)$.

Theorem 3.1 (Errors of the prototypical method). *The prototypical method (3.3) is well-posed and its solution is given by $\tilde{u}_H = \mathcal{R}u$, where $\mathcal{R}: V \rightarrow \tilde{V}_H$ denotes the a -orthogonal projection onto \tilde{V}_H . Moreover, for any $f \in H^s(\Omega)$ for the CG-LOD, and $f \in H^s(\mathcal{T}_H)$ for the DG-LOD, with $s \in \{0, 1, \dots, p+1\}$, we have*

$$\|u - \tilde{u}_H\|_\Omega \lesssim H^{s+2} |f|_{s, \mathcal{T}_H}, \quad \|\nabla(u - \tilde{u}_H)\|_\Omega \lesssim H^{s+1} |f|_{s, \mathcal{T}_H}. \quad (3.4)$$

¹This restriction to simplicial elements is not essential and is made for simplicity of presentation. In fact, quadrilateral/hexahedral elements may also be considered. The corresponding globally continuous finite element spaces can be constructed using polynomials on a reference element, which are then mapped to the mesh elements *via* multi-linear coordinate transformations.

Proof. First, note that the space \tilde{V}_H is a closed subspace of V , and hence, problem (3.3) is well-posed by the Riesz representation theorem. Comparing (3.3) with the weak formulation (2.2), we observe that $a(u - \tilde{u}_H, \tilde{v}_H) = 0$ for all $\tilde{v}_H \in \tilde{V}_H$, which implies that $\tilde{u}_H = \mathcal{R}u$, where $\mathcal{R}: V \rightarrow \tilde{V}_H$ is the a -orthogonal projection.

Since the prototypical DG-LOD is equivalent to the prototypical method presented in [42], the error estimates in (3.4) can be directly concluded from Theorem 3.1 of [42]. For the prototypical CG-LOD, the proof below follows along the same lines and we present it here for the sake of readability and completeness. We denote by $\Pi_H: L^2(\Omega) \rightarrow M_H$ the L^2 -orthogonal projection onto M_H and note that the error satisfies $e := u - \tilde{u}_H = u - \mathcal{R}u \in W$. This yields that

$$\alpha \|\nabla e\|_\Omega^2 \leq a(u - \mathcal{R}u, e) = (f, e)_\Omega = (f - \Pi_H f, e - \Pi_H e)_\Omega, \quad (3.5)$$

where we have subtracted the projection Π_H on both sides, using that $\Pi_H e = 0$ since $e \in W$ and due to the L^2 -orthogonality properties of Π_H . Using that

$$\|v - \Pi_H v\|_{L^2(\Omega)} \lesssim H^k |v|_{k, \Omega}$$

for $0 \leq k \leq p + 1$ and $v \in H^k(\Omega)$ (see, e.g., [53]), we get

$$\alpha \|\nabla e\|_\Omega^2 \lesssim H^s |f|_{s, \Omega} H \|\nabla e\|_\Omega,$$

which proves the error estimate in the H^1 -norm. The L^2 -error estimate follows by another application of the approximation properties of Π_H , i.e., $\|e\|_\Omega = \|e - \Pi_H e\|_\Omega \lesssim H \|\nabla e\|_\Omega$. \square

Remark 3.2 (Boundary conditions of M_H). In the CG-LOD variant, not enforcing boundary conditions for M_H is crucial to achieve higher-order convergence rates as stated in Theorem 3.1. For example, using the polynomial degree $p = 1$, Theorem 3.1 gives an H^1 -error of order $\mathcal{O}(H^3)$. In contrast, the more traditional approaches in [36, 44, 45], where the homogeneous Dirichlet boundary conditions are imposed on the analogue of M_H , yield an error which is only of order $\mathcal{O}(H)$. If $f \in H_0^1(\Omega)$, convergence rates beyond first order can also be achieved using $H_0^1(\Omega)$ -conforming spaces M_H ; see, e.g., Lemma 8.1 of [45]. This has been exploited, for instance, for the Gross–Pitaevskii problem in [35].

To better understand the structure of the space \tilde{V}_H , we characterize the a -orthogonal projection $\mathcal{R}: V \rightarrow \tilde{V}_H$ via a saddle point formulation using Lagrange multipliers in M_H . Specifically, for any $v \in V$, we can identify $\mathcal{R}v$ with the unique solution to the following problem: find $(\mathcal{R}v, \lambda) \in V \times M_H$ such that

$$a(\mathcal{R}v, w) + b(w, \lambda) = 0, \quad \forall w \in V, \quad (3.6a)$$

$$b(\mathcal{R}v, \mu) = b(v, \mu), \quad \forall \mu \in M_H, \quad (3.6b)$$

where the bilinear form b is given by $b(v, \mu) := (v, \mu)_\Omega$. Indeed, equation (3.6a) ensures that $\mathcal{R}v \in \tilde{V}_H$, while equation (3.6b) implies that $\mathcal{R}v - v \in W$. The well-posedness of (3.6) follows from the classical saddle point theory (cf. Corollary 4.2.1 of [13]) and, in particular, relies on the inf-sup condition

$$\inf_{\mu \in M_H} \sup_{v \in V} \frac{b(v, \mu)}{\|\nabla v\|_\Omega \|\mu\|_\Omega} \gtrsim H > 0, \quad (3.7)$$

which is a direct consequence of Lemma 3.4 from [42].

The operator \mathcal{R} and its representation given in (3.6), enables the construction of a basis for the space $\tilde{V}_H = \mathcal{R}V$. Before formalizing this idea, we briefly review two existing LOD constructions from the literature:

- (1) In [44, 45], the fine-scale space W is defined as the kernel of a quasi-interpolation operator $\mathcal{I}_H: V \rightarrow V_H$, where V_H is an underlying $H_0^1(\Omega)$ -conforming finite element space. A basis for \tilde{V}_H is obtained by correcting the basis functions of V_H with corrections in W . Specifically, the prototypical LOD space \tilde{V}_H is redefined as

$\tilde{V}_H = V_H - \mathcal{C}V_H$ with the a -orthogonal projection $\mathcal{C}: V \rightarrow W$ serving as a correction operator. This strategy is beneficial for localizing the space \tilde{V}_H by expressing \mathcal{C} as the sum of contributions from every element of the mesh. However, it is not applicable in the present work, as our underlying finite element spaces M_H are not $H_0^1(\Omega)$ -conforming, which is essential to obtain higher-order convergence, as discussed in Remark 3.2.

- (2) In [4], a more general framework is introduced in which the basis functions of the prototypical LOD space are associated with so-called quantities of interest (QOI), a concept that also plays an important role in the theory of gamblers; see, e.g., [49]. Such QOI are continuous functionals on V encoding information about the exact solution u to problem (2.2) that the method aims to preserve exactly. The fine-scale space W is then defined as the intersection of the kernels of all QOI. We adopt this general approach in the present work, choosing the QOI as the $L^2(\Omega)$ -inner products with basis functions of M_H ; see (3.8) below. Our DG-LOD corresponds to the choice of QOI in Example 3.1 (a) & (b) of [4]. The CG-LOD with $p = 1$ is closely related to ([4], Ex. 3.1 (d)) except for the treatment of boundary conditions of the space M_H , cf. Remark 3.2. Further note that the variant from Example 3.1 (c) of [4] is equivalent to the classical LOD as discussed in [45]. Modifying the boundary treatment, as in Remark 3.2 for the CG-LOD, yields a version of this method with an error estimate improved by one order in H compared to [45]; see (8).

In our setting, the basis functions of \tilde{V}_H are associated with QOI defined as

$$q_j(v) := \int_{\omega_j} v \Lambda_j \, dx, \quad j = 1, \dots, J, \quad (3.8)$$

where Λ_j are the basis functions of M_H , $J := \dim(M_H)$, and $\omega_j = \text{supp}(\Lambda_j)$. In case of the CG-LOD, $\{\Lambda_j\}_{j=1}^J$ can be chosen, for example, as the p -th order Lagrange nodal basis of M_H . For the DG-LOD, the basis functions can be taken as the $L^2(T)$ -orthonormal Legendre bases of $\mathcal{P}^p(T)$ on each element $T \in \mathcal{T}_H$. Henceforth, we denote by $\{\mu_j\}_{j=1}^J$ the coefficients of a function $\mu \in M_H$ with respect to the basis $\{\Lambda_j\}_{j=1}^J$, i.e.,

$$\mu = \sum_{j=1}^J \mu_j \Lambda_j. \quad (3.9)$$

The following lemma characterizes the basis functions of \tilde{V}_H as solutions to saddle point problems with a Kronecker-delta constraint on the QOI.

Lemma 3.3 (Prototypical basis). *A basis of the space \tilde{V}_H is given by $\{\tilde{\varphi}_j : j = 1, \dots, J\}$ with $\tilde{\varphi}_j$ defined for each $j \in \{1, \dots, J\}$ as the unique solution to the saddle point problem that seeks $(\tilde{\varphi}_j, \lambda_j) \in V \times M_H$ such that*

$$a(\tilde{\varphi}_j, v) + b(v, \lambda_j) = 0, \quad \forall v \in V, \quad (3.10a)$$

$$b(\tilde{\varphi}_j, \mu) = \mu_j, \quad \forall \mu \in M_H, \quad (3.10b)$$

where μ_j is the coordinate of Λ_j in the basis representation of μ , cf. (3.9). Furthermore, the projection operator \mathcal{R} can be represented as

$$\mathcal{R}v = \sum_{j=1}^J q_j(v) \tilde{\varphi}_j. \quad (3.11)$$

Proof. The well-posedness of problem (3.10) follows from classical saddle point theory, using the inf-sup condition (3.7); see, for instance, Corollary 4.2.1 of [13]. Taking $v \in W$ as test function in (3.10a) proves that $\tilde{\varphi}_j \in \tilde{V}_H$. To show (3.11), we define $e := \mathcal{R}v - \sum_{j=1}^J q_j(v) \tilde{\varphi}_j$ and observe that $e \in W$. Indeed, using (3.6b), (3.10b), and the identity $b(v, \mu) = \sum_{j=1}^J \mu_j q_j(v)$, cf. (3.9), we find that $q_i(e) = q_i(\mathcal{R}v) - q_i(v) = 0$ for all $i \in \{1, \dots, J\}$. Moreover, since $e \in \tilde{V}_H$, it follows that $e = 0$, proving (3.11) and that the functions $\{\tilde{\varphi}_j\}_{j=1}^J$ span the space \tilde{V}_H . These functions are also linear independent, since if a linear combination $\sum_{j=1}^J \alpha_j \tilde{\varphi}_j$ vanishes for some coefficients $\{\alpha_j\}_{j=1}^J$, then (3.10b) implies that $\sum_{j=1}^J \alpha_j \mu_j = 0$ for all $\mu \in M_H$, so that all coefficients α_j vanish. This means that the family $\{\tilde{\varphi}_j\}_{j=1}^J$ forms a basis of the space \tilde{V}_H . \square

4. EXPONENTIAL DECAY

We emphasize that the prototypical LOD basis functions defined in (3.10) are globally supported. Therefore, their computation would require the solution to global problems, which is infeasible for practical purposes. In this section, we show that the prototypical LOD basis functions decay exponentially, which motivates their approximation by locally computable counterparts in Section 5. A practical multiscale method based on such local approximations is presented in Section 6. To quantify the decay of the basis functions, and more generally of the solutions to saddle points problem of type (3.10), we introduce the notion of patches with respect to the mesh \mathcal{T}_H . Given an *oversampling parameter* $\ell \in \mathbb{N}$, we define the ℓ -th order patch of a union of elements $S \subset \mathcal{T}_H$ recursively for $\ell \geq 2$ by $\mathbf{N}^\ell(S) := \mathbf{N}^1(\mathbf{N}^{\ell-1}(S))$, where $\mathbf{N}^1(S) := \mathbf{N}(S)$ is the set of mesh elements sharing at least a node with the elements in S . Before proving the exponential decay result, we state a technical lemma used in the proof, which is a reformulation of Corollary 3.6 from [42].

Lemma 4.1 (Non-standard inverse inequality). *For all $T \in \mathcal{T}_H$ and all $\mu \in M_H$, we have*

$$\|\mu\|_T \lesssim H^{-1} \|\mu\|_{-1,T},$$

where the H^{-1} -norm is defined by $\|\mu\|_{-1,T} := \sup_{w \in H_0^1(T), w \neq 0} \frac{(\mu, w)_T}{\|\nabla w\|_T}$.

Proof. We denote by $b_T \in H_0^1(T)$ the usual bubble function (the product of the \mathcal{P}^1 -hat functions associated to the vertices of T) and use its well-known properties together with a standard inverse inequality for μ to conclude that

$$\|\mu\|_T^2 \lesssim (\mu, \mu b_T)_T \leq \frac{(\mu, \mu b_T)_T}{\|\nabla(\mu b_T)\|_T} (\|\nabla \mu\|_T \|b_T\|_{L^\infty(T)} + \|\mu\|_T \|\nabla b_T\|_{L^\infty(T)}) \lesssim \|\mu\|_{-1,T} \frac{1}{H} \|\mu\|_T.$$

□

The exponential decay will be proven in the following theorem in a rather general setting. Note that the exponential decay of the prototypical LOD basis functions can be recovered by setting $f_S(v) = 0$ for all $v \in V$ and $g_S(\mu) = \mu_j$ for all $\mu \in M_H$, where μ_j is the coordinate of Λ_j in the basis expansion of μ , cf. (3.9).

Theorem 4.2 (Exponential decay). *Let S be a union of mesh elements $S \subset \mathcal{T}_H$ and $\psi \in V$ be the solution to the problem: find $(\psi, \lambda) \in V \times M_H$ such that*

$$a(\psi, v) + b(v, \lambda) = f_S(v), \quad \forall v \in V, \quad (4.1a)$$

$$b(\psi, \mu) = g_S(\mu), \quad \forall \mu \in M_H, \quad (4.1b)$$

where f_S and g_S are bounded linear functionals on V and M_H , respectively, such that $f_S(v) = 0$ for all $v \in V$ with $\text{supp}(v) \subset \overline{\Omega \setminus S}$ and analogously for g_S . Then there exists $C_{\text{dec}} > 0$, independent of H , ℓ , and S , such that for all $\ell \in \mathbb{N}$

$$\|\nabla \psi\|_{\Omega \setminus \mathbf{N}^\ell(S)} \leq \exp(-C_{\text{dec}} \ell) \|\nabla \psi\|_\Omega. \quad (4.2)$$

Proof. The proof is based on ideas from [36, 44, 45]. For the DG-LOD, it follows directly from the arguments in Theorem 4.1 from [42]. In the CG-LOD case, some modifications are necessary, as outlined below. Fix an integer $\ell \geq 1$, and let $\eta \in W^{1,\infty}(\Omega)$ be the first-order finite element cut-off function characterized by

$$\eta = 0 \text{ in } \mathbf{N}^{\ell-1}(S), \quad \eta = 1 \text{ in } \Omega \setminus \mathbf{N}^\ell(S), \quad (4.3)$$

with the transition region $R := \mathbf{N}^\ell(S) \setminus \mathbf{N}^{\ell-1}(S)$, satisfying

$$\|\eta\|_{L^\infty(\Omega)} \leq 1, \quad \|\nabla \eta\|_{L^\infty(\Omega)} \lesssim H^{-1}. \quad (4.4)$$

Furthermore, let

$$a_\omega(v, w) := \int_\omega (A \nabla v) \cdot \nabla w \, dx \quad (4.5)$$

denote the restriction of the bilinear form a to a subdomain $\omega \subset \Omega$. Testing (4.1a) with $\eta\psi$ and decomposing the support of η into $\Omega \setminus \mathbf{N}^\ell(S)$ and R , we obtain that

$$a_{\Omega \setminus \mathbf{N}^\ell(S)}(\psi, \eta\psi) = f_S(\eta\psi) - a_R(\psi, \eta\psi) - b(\eta\psi, \lambda).$$

Using the lower bound from (2.1), we estimate this by

$$\alpha \|\nabla\psi\|_{\Omega \setminus \mathbf{N}^\ell(S)}^2 \leq |f_S(\eta\psi)| + |a_R(\psi, \eta\psi)| + |b(\eta\psi, \lambda)| =: \Xi_1 + \Xi_2 + \Xi_3. \quad (4.6)$$

We now consider terms Ξ_1 – Ξ_3 individually. We have $\Xi_1 = 0$ since $(\eta\psi)|_S = 0$. To estimate Ξ_2 , we apply (2.1) and (4.4), which yields

$$\Xi_2 \leq \beta \|\nabla\psi\|_R \|\nabla(\eta\psi)\|_R \lesssim \|\nabla\psi\|_R^2 + H^{-1} \|\nabla\psi\|_R \|\psi\|_R. \quad (4.7)$$

To estimate the right-hand side of the previous inequality, let Λ_z^1 denote the \mathcal{P}^1 -finite element hat function associated with the node z , and define $\omega_z := \text{supp}(\Lambda_z^1)$. Then, we have the Poincaré-type inequality

$$\|v\|_{\omega_z} \leq C_P H \|\nabla v\|_{\omega_z} \quad \forall v \in H^1(\omega_z) \quad \text{with} \quad \int_{\omega_z} \Lambda_z^1 v \, dx = 0, \quad (4.8)$$

which can be shown with the Peetre–Tartar lemma (see, e.g., [23], Lem. A.38), a scaling argument, and the maximization of C_P over all the patch configurations that are admissible by the mesh regularity. To be able to apply (4.8) for estimating the right-hand side of (4.7), we cover R by a collection of patches ω_z such that the union of the ω_z equals $R^2 := \text{int}(\mathbf{N}^{\ell+1}(S) \setminus \mathbf{N}^{\ell-1}(S))$. Testing (4.1b) with Λ_z^1 and noting that $\omega_z \cap S = \emptyset$ gives $\int_{\omega_z} \Lambda_z^1 \psi \, dx = 0$. Thus, applying (4.8) to ψ locally on each patch in the collection and summing up, we obtain that $\|\psi\|_R \lesssim H \|\nabla\psi\|_{R^2}$, using the finite overlap of the ω_z . Substituting this into (4.7), we get

$$\Xi_2 \lesssim \|\nabla\psi\|_{R^2}^2. \quad (4.9)$$

To estimate Ξ_3 , we localize it to the ring R^2 . To this end, we decompose λ as $\lambda = \lambda^{\text{in}} + \lambda^{\text{out}}$, where λ^{in} is the linear combination of basis functions of M_H supported only in $\mathbf{N}^{\ell+1}(S)$, and λ^{out} is the combination of the remaining basis functions, associated with Lagrange points not in the interior of $\mathbf{N}^{\ell+1}(S)$. This decomposition is well-defined and unique. Since $\eta\psi = \psi$ in $\Omega \setminus \mathbf{N}^\ell(S)$ and $\lambda^{\text{out}} = 0$ in $\mathbf{N}^\ell(S)$, it follows from (4.1b) that $b(\eta\psi, \lambda^{\text{out}}) = b(\psi, \lambda^{\text{out}}) = 0$. Thus,

$$\Xi_3 = b(\eta\psi, \lambda^{\text{in}}) = \int_{R^2} \eta\psi \lambda^{\text{in}} \, dx \leq \|\eta\psi\|_{R^2} \|\lambda^{\text{in}}\|_{R^2} \lesssim \|\psi\|_{R^2} \|\lambda\|_{R^2}, \quad (4.10)$$

noting that the integral above could be restricted to R^2 since $\eta = 0$ on $\mathbf{N}^{\ell-1}(S)$ and $\lambda^{\text{in}} = 0$ outside $\mathbf{N}^{\ell+1}(S)$. The last estimate in (4.10) follows from $\|\lambda^{\text{in}}\|_{R^2} \lesssim \|\lambda\|_{R^2}$, which can be proved by a scaling argument on each element, noting that λ^{in} is uniquely determined by λ . To further estimate (4.10), we consider an arbitrary element $T \subset \overline{R^2}$ and observe $\|\lambda\|_T \lesssim H^{-1} \|\lambda\|_{-1,T}$ by Lemma 4.1. Testing (4.1a) with any $v \in H_0^1(T)$ satisfying $\|\nabla v\|_T = 1$ and noting that $f_S(v) = 0$, yields

$$(\lambda, v)_T = b(v, \lambda) = -a(\psi, v) \leq \beta \|\nabla\psi\|_T \|\nabla v\|_T,$$

which implies that $\|\lambda\|_{-1,T} \leq \beta \|\nabla\psi\|_T$. Applying (4.8) as above, combining the previous estimates, and summing over all $T \subset \overline{R^2}$, we obtain that

$$\Xi_3 \lesssim H \|\nabla\psi\|_{R^2} \left(\sum_{T \subset R^2} H^{-2} \|\nabla\psi\|_T^2 \right)^{1/2} \lesssim \|\nabla\psi\|_{R^2}^2. \quad (4.11)$$

Substituting (4.9) and (4.11) into (4.6), and noting that the ring R^2 can be written as $R^2 = \text{int}((\Omega \setminus \mathbf{N}^{\ell-1}(S)) \setminus (\Omega \setminus \mathbf{N}^{\ell+1}(S)))$, we conclude, for a constant $C > 0$ independent of H and ℓ that

$$\|\nabla\psi\|_{\Omega \setminus \mathbf{N}^\ell(S)}^2 \leq C \|\nabla\psi\|_{R^2}^2 = C \|\nabla\psi\|_{\Omega \setminus \mathbf{N}^{\ell-1}(S)}^2 - C \|\nabla\psi\|_{\Omega \setminus \mathbf{N}^{\ell+1}(S)}^2,$$

which, after rearranging the terms, leads to

$$\|\nabla\psi\|_{\Omega \setminus \mathbf{N}^{\ell+1}(S)}^2 \leq \frac{C}{1+C} \|\nabla\psi\|_{\Omega \setminus \mathbf{N}^{\ell-1}(S)}^2. \quad (4.12)$$

Iterating the argument gives the assertion with $C_{\text{dec}} := \frac{1}{4} \log \frac{1+C}{C}$. \square

Remark 4.3 (Better decay rate for DG-LOD). As mentioned above, the proof of exponential decay for the DG-LOD is very similar. Inequality (4.8) can be replaced by the standard Poincaré inequality for mean-zero functions on each mesh element. Thus, estimate (4.9) can be localized to ring R of width one instead of R^2 , noting that piecewise constants lie in M_H . Likewise, in (4.10) the form b localizes to R , since contributions outside $\mathbf{N}^\ell(S)$ vanish. Therefore, (4.12) can be replaced by

$$\|\nabla\psi\|_{\Omega \setminus \mathbf{N}^\ell(S)}^2 \leq \frac{C}{1+C} \|\nabla\psi\|_{\Omega \setminus \mathbf{N}^{\ell-1}(S)}^2 \quad (4.13)$$

with a constant $C > 0$, and the resulting decay rate is $\frac{1}{2} \log \frac{1+C}{C}$. Although C might not be the same as in (4.12), the more local nature of estimate (4.13) gives some theoretical underpinning of the better localization properties of the DG-LOD compared to the CG-LOD. This is confirmed by numerical experiments in Section 9.

5. LOCALIZATION

The exponential decay properties established in the previous section motivate the localized computation of the prototypical LOD basis functions. However, the naive strategy of localizing these basis functions by restricting problem (3.10) to ℓ -th order patches has the drawback that the localization error increases when decreasing the mesh size if the parameter ℓ is kept fixed, cf. [42, 44]. One strategy to address this, described in Item 1 on Page 6, expresses the prototypical LOD space \tilde{V}_H as $\tilde{V}_H = V_H - \mathcal{C}V_H$, where V_H is the underlying $H_0^1(\Omega)$ -conforming first-order finite element space and $\mathcal{C}: V \rightarrow W$ denotes the a -orthogonal projection onto the fine-scale space W . The operator \mathcal{C} is then decomposed into a sum of element-wise contributions, which are subsequently localized. Note that the projection operator \mathcal{R} can then be written as $\mathcal{R} = \mathcal{I}_H - \mathcal{C}\mathcal{I}_H$ if $\mathcal{I}_H: V \rightarrow V_H$ is a projection. However, this strategy requires the fine-scale space to be defined as the kernel of a quasi-interpolation operator $\mathcal{I}_H: V \rightarrow V_H$. In the present setting, this construction is not directly applicable, since our underlying space M_H is not necessarily $H_0^1(\Omega)$ -conforming.

Our localization approach mimics the above construction by expressing \mathcal{R} as

$$\mathcal{R} = \mathcal{I}_H - \mathcal{K}, \quad (5.1)$$

where the operator $\mathcal{K}: V \rightarrow V$ will be characterized below, and $\mathcal{I}_H: V \rightarrow V_H$ is a quasi-interpolation operator onto the first-order $H_0^1(\Omega)$ -conforming finite element space V_H associated with the mesh \mathcal{T}_H . Note that V_H is fixed as the first-order finite element space regardless of the polynomial degree p . The quasi-interpolation operator \mathcal{I}_H (which is not necessarily a projection) is assumed to satisfy classical local approximation and stability estimates, *i.e.*,

$$H^{-1}\|v - \mathcal{I}_H v\|_T + \|\nabla \mathcal{I}_H v\|_T \lesssim \|\nabla v\|_{\mathbf{N}(T)}, \quad \forall T \in \mathcal{T}_H, \forall v \in V. \quad (5.2)$$

Moreover, the operator \mathcal{I}_H should depend on its argument only through its QOI defined in (3.8), *i.e.*, $\mathcal{I}_H w = 0$ for all $w \in W$. Such operators can be readily constructed for the two classes of methods considered. Indeed,

for the CG-LOD, a weighted Clément-type interpolation can be used. This interpolation is uniquely defined by assigning values at interior nodes $z \in \mathcal{N}_H^{\text{in}}$ as

$$(\mathcal{I}_H v)(z) := \left(\int_{\omega_z} \Lambda_z^1 dx \right)^{-1} \int_{\omega_z} v \Lambda_z^1 dx, \quad \forall v \in V, \quad (5.3)$$

where ω_z is the support of the first-order finite element hat function Λ_z^1 associated with node z . Nodal values at boundary nodes are set to zero. Such operators were, *e.g.*, introduced in [16]. Definition (5.3) suggests that it is convenient to include the first-order hat functions Λ_z^1 into the basis $\{\Lambda_j\}_{j=1}^J$ of the space M_H . In practice, we thus choose $\{\Lambda_j\}_{j=1}^J$ as a slight modification of the p -th order Lagrange nodal basis; namely we replace the standard Lagrange basis functions associated to the nodes with first-order hat functions on the same nodes, while keeping unchanged the other basis functions (associated to the interior points on edges/faces/elements). The QOI corresponding to the latter thus do not enter into the calculation of $\mathcal{I}_H v$.

For the DG-LOD method, a quasi-interpolation can be uniquely defined by specifying its nodal values at interior nodes $z \in \mathcal{N}_H^{\text{in}}$ as

$$(\mathcal{I}_H v)(z) := \sum_{T \in \omega_z} \frac{|T|}{|\omega_z|} \int_T v dx, \quad \forall v \in V, \quad (5.4)$$

where $|\cdot|$ denotes the d -dimensional volume of a subdomain, and values at boundary nodes are again set to zero; see, *e.g.*, [24] for an analysis in a more general setting. In practice, we choose a basis $\{\Lambda_j\}_{j=1}^J$ of M_H so that it contains the characteristic functions of all the mesh elements $T \in \mathcal{T}_H$. Any other function Λ_j is orthogonal to piecewise constants, so that the corresponding QOI do not enter into the calculation of $\mathcal{I}_H v$ according to (5.4).

The operator $\mathcal{K}: V \rightarrow V$, as introduced in (5.1), is characterized for each $v \in V$ as the unique solution $(\mathcal{K}v, \lambda) \in V \times M_H$ to the saddle point problem

$$a(\mathcal{K}v, w) + b(w, \lambda) = a(\mathcal{I}_H v, w), \quad \forall w \in V, \quad (5.5a)$$

$$b(\mathcal{K}v, \mu) = -c(v, \mu), \quad \forall \mu \in M_H, \quad (5.5b)$$

where we use the abbreviation $c(v, \mu) := b(v - \mathcal{I}_H v, \mu)$. The operator \mathcal{K} can now be expressed as the following sum of local element contributions, *i.e.*,

$$\mathcal{K} = \sum_{T \in \mathcal{T}_H} \mathcal{K}_T,$$

where, for all $T \in \mathcal{T}_H$, the operators $\mathcal{K}_T: V \rightarrow V$ are defined for $v \in V$ as the unique solution $(\mathcal{K}_T v, \lambda_T) \in V \times M_H$ to the modified saddle point problem

$$a(\mathcal{K}_T v, w) + b(w, \lambda_T) = a_T(\mathcal{I}_H v, w), \quad \forall w \in V, \quad (5.6a)$$

$$b(\mathcal{K}_T v, \mu) = -c_T(v, \mu), \quad \forall \mu \in M_H. \quad (5.6b)$$

Here, in contrast to (5.5), we use localized versions of the bilinear forms a and c on the right-hand side of the problem, denoted by a_T and c_T , respectively. The local form a_T is defined by restricting a to T , as in (4.5). To define the local form c_T , there are generally several meaningful options. We formulate the following four properties that the bilinear form c_T should satisfy:

- (1) *Summation property:* $\sum_{T \in \mathcal{T}_H} c_T(v, \mu) = c(v, \mu)$ for all $v \in V$ and $\mu \in M_H$;
- (2) *Vanishing on constants:* $c_T(v, \mu) = 0$ for all $\mu \in M_H$ if $v = 1$ on $\mathbf{N}(T)$ for elements T that do not contain any boundary nodes;
- (3) *Vanishing on fine scales:* $c_T(w, \mu) = 0$ for all $w \in W$ and $\mu \in M_H$;
- (4) *Locality:* $|c_T(v, \mu)| \lesssim \|v\|_{\mathbf{N}(T)} \|\mu\|_{\mathbf{N}(T)}$ for all $v \in V$ and $\mu \in M_H$.

For the CG-LOD, the form c_T can be constructed as

$$c_T(v, \mu) := \sum_{j=1}^J \frac{|T \cap \omega_j|}{|\omega_j|} \mu_j q_j(v) - \int_T \mu \mathcal{I}_H v \, dx, \tag{5.7}$$

where μ_j is the coordinates of Λ_j in the basis expansion of μ , cf. (3.9). To verify property (1) above, we note that

$$\sum_{T \in \mathcal{T}_H} c_T(v, \mu) = \sum_{j=1}^J \mu_j \int_{\Omega} \Lambda_j v \, dx - \int_{\Omega} \mu \mathcal{I}_H v \, dx = b(v - \mathcal{I}_H v, \mu).$$

For property (2), we assume $v|_{\mathbf{N}(T)} = 1$ and, therefore, $(\mathcal{I}_H v)|_T = 1$. We write

$$c_T(v, \mu) = \sum_{j:T \subset \bar{\omega}_j} \frac{|T|}{|\omega_j|} \mu_j \int_{\omega_j} \Lambda_j \, dx - \int_T \mu \, dx = \sum_{j:T \subset \bar{\omega}_j} \mu_j \int_T \Lambda_j \, dx - \int_T \mu \, dx = 0,$$

since $|T|^{-1} \int_T \Lambda_j \, dx = |T'|^{-1} \int_{T'} \Lambda_j \, dx$ for any two elements $T, T' \subset \bar{\omega}_j$ by the affine equivalence of finite element basis functions, which implies $\int_T \Lambda_j \, dx = \frac{|T|}{|\omega_j|} \int_{\omega_j} \Lambda_j \, dx$ for $T \subset \bar{\omega}_j$. Property (3) is evident since $q_j(w) = 0$ and $\mathcal{I}_H w = 0$ for $w \in W$. Finally, for property (4), we observe that

$$|c_T(v, \mu)| \leq \sum_{j:T \subset \bar{\omega}_j} \frac{|T|}{|\omega_j|} |\mu_j| \|\Lambda_j\|_{\omega_j} \|v\|_{\omega_j} + \|\mu\|_T \|\mathcal{I}_H v\|_T \lesssim \sum_{j:T \subset \bar{\omega}_j} |\mu_j| \|\Lambda_j\|_T \|v\|_{\mathbf{N}(T)} + \|v\|_{\mathbf{N}(T)} \|\mu\|_T,$$

since, similar to above, $\|\Lambda_j\|_{\omega_j}^2 = \frac{|\omega_j|}{|T|} \|\Lambda_j\|_T^2$. We also have $\sum_{j:T \subset \bar{\omega}_j} |\mu_j| \|\Lambda_j\|_T \lesssim \|\mu\|_T$ by the equivalence of norms on the finite dimensional space $\mathcal{P}^p(T)$ and scaling to a reference element. Hence, $|c_T(v, \mu)| \lesssim \|v\|_{\mathbf{N}(T)} \|\mu\|_T$.

For the DG-LOD method, (5.7) can be used as well and simplifies to

$$c_T(v, \mu) = \int_T (v - \mathcal{I}_H v) \mu \, dx,$$

since the supports ω_j of any Λ_j consists of only one mesh element in this case.

Due to the general setting considered in Theorem 4.2, the result applies to the function $\mathcal{K}_T v$ for any $v \in V$, and shows that it decays exponentially away from the element T . This motivates the localization of the operator \mathcal{K}_T to ℓ -th order patches around T . To this end, we introduce the corresponding localized spaces

$$\begin{aligned} V_T^\ell &:= \{v \in V : \text{supp}(v) \subset \mathbf{N}^\ell(T)\}, \\ M_T^\ell &:= \{\mu|_{\mathbf{N}^\ell(T)} : \mu \in M_H\}, \end{aligned}$$

where we implicitly extend functions in the space M_T^ℓ by zero. The localized operator $\mathcal{K}_T^\ell: V \rightarrow V_T^\ell$ can then be defined, for any $v \in V$, as the unique solution $(\mathcal{K}_T^\ell v, \lambda_T^\ell) \in V_T^\ell \times M_T^\ell$ to the local saddle point problem

$$a(\mathcal{K}_T^\ell v, w) + b(w, \lambda_T^\ell) = a_T(\mathcal{I}_H v, w), \quad \forall w \in V_T^\ell, \tag{5.8a}$$

$$b(\mathcal{K}_T^\ell v, \mu) = -c_T(v, \mu), \quad \forall \mu \in M_T^\ell. \tag{5.8b}$$

Finally, a localized version of the operator \mathcal{R} can be defined by

$$\mathcal{R}^\ell v := (\mathcal{I}_H - \mathcal{K}^\ell)v, \quad \mathcal{K}^\ell v := \sum_{T \in \mathcal{T}_H} \mathcal{K}_T^\ell v. \tag{5.9}$$

The following theorem shows that the localized operator \mathcal{K}^ℓ approximates \mathcal{K} exponentially well in the operator norm. Notably, it avoids the H^{-1} prefactor typically arising in naive localization strategies; see, e.g., [42, 44].

Theorem 5.1 (Localization error). *For all $v \in V$ and $\ell \in \mathbb{N}$, we have*

$$\|\nabla(\mathcal{R} - \mathcal{R}^\ell)v\|_\Omega \lesssim \ell^{(d-1)/2} \exp(-C_{\text{dec}}\ell) \|\nabla \mathcal{R}v\|_\Omega, \quad (5.10)$$

where C_{dec} is the constant from Theorem 4.2.

Proof. We suppose without loss of generality that $\ell \geq 2$, observing that the case $\ell = 1$ holds by scaling. We abbreviate the localization error by $e := (\mathcal{R} - \mathcal{R}^\ell)v$ and note that $e \in W$. Indeed, by (5.1), (5.9), (5.6b), and (5.8b) we have $b(e, \mu) = -\sum_{T \in \mathcal{T}_H} b(\mathcal{K}_T v - \mathcal{K}_T^\ell v, \mu) = 0$ for all $\mu \in M_H$. Thus,

$$\alpha \|\nabla(\mathcal{R} - \mathcal{R}^\ell)v\|_\Omega^2 \leq -a(\mathcal{R}^\ell v, e) = \sum_{T \in \mathcal{T}_H} \left(-a_T(\mathcal{I}_H v, e) + a(\mathcal{K}_T^\ell v, e) \right). \quad (5.11)$$

In the following, we consider each summand on the right-hand side separately. We use the cut-off function from (4.3) with $S = T$, now denoted by η_T . Note that $\eta_T = 0$ on T , so that $a_T(\mathcal{I}_H v, e) = a_T(\mathcal{I}_H v, (1 - \eta_T)e)$. Together with (5.8a) for the test function $w = (1 - \eta_T)e \in V_T^\ell$, this yields

$$\begin{aligned} -a_T(\mathcal{I}_H v, e) + a(\mathcal{K}_T^\ell v, e) &= -a_T(\mathcal{I}_H v, (1 - \eta_T)e) + a(\mathcal{K}_T^\ell v, (1 - \eta_T)e + \eta_T e) \\ &= -b((1 - \eta_T)e, \lambda_T^\ell) + a(\mathcal{K}_T^\ell v, \eta_T e) =: \Xi_1 + \Xi_2. \end{aligned}$$

To estimate the term Ξ_1 , we decompose $\lambda_T^\ell = \lambda_T^{\text{in}} + \lambda_T^{\text{out}}$, where λ_T^{in} is the linear combination of basis functions of M_H fully supported in $\mathbf{N}^{\ell-1}(T)$, and λ_T^{out} is the combination of the remaining basis functions. Specifically, for the CG-LOD, the basis functions forming λ_T^{out} are associated with Lagrange points that do not lie in $\text{int}(\mathbf{N}^{\ell-1}(T))$. For DG-LOD, we simply choose $\lambda_T^{\text{out}} = \lambda_T^\ell|_{\Omega \setminus \mathbf{N}^{\ell-1}(T)}$. Since $\eta_T = 0$ in $\mathbf{N}^{\ell-1}(S)$, it follows from (5.8b) that $b((1 - \eta_T)e, \lambda^{\text{in}}) = b(e, \lambda^{\text{in}}) = 0$. Thus, introducing $R_T^\ell := \text{int}(\mathbf{N}^\ell(T) \setminus \mathbf{N}^{\ell-2}(T))$, we obtain that

$$\Xi_1 = -b((1 - \eta_T)e, \lambda_T^{\text{out}}) \leq \int_{R_T^\ell} |e \lambda_T^{\text{out}}| \, dx \lesssim \|e\|_{R_T^\ell} \|\lambda_T^\ell\|_{R_T^\ell}.$$

To estimate the norm of λ_T^ℓ , we employ Lemma 4.1 on each mesh element $K \subset \overline{R_T^\ell}$, resulting in a negative power of H and the $H^{-1}(K)$ -norms of λ_T^ℓ . We now take any $w \in H_0^1(K)$ with $\|\nabla w\|_K = 1$ as a test function in (5.8a) and note that $a_T(\mathcal{I}_H v, w) = 0$ since $T \cap R_T^\ell = \emptyset$ for $\ell \geq 2$. Thus,

$$|(\lambda_T^\ell, w)_K| = |a(\mathcal{K}_T^\ell v, w)| \leq \beta \|\nabla \mathcal{K}_T^\ell v\|_K,$$

yielding $\|\lambda_T^\ell\|_K \lesssim \|\lambda_T^\ell\|_{-1, K} \leq \beta H^{-1} \|\nabla \mathcal{K}_T^\ell v\|_K$. Recalling that $\|e\|_{R_T^\ell} \lesssim H \|\nabla e\|_{R_T^\ell}$ by the Poincaré-type inequality (4.8) (or the usual Poincaré inequality for the DG-LOD), and substituting this into the estimate for Ξ_1 , we obtain that

$$\Xi_1 \lesssim \|\nabla \mathcal{K}_T^\ell v\|_{R_T^\ell} \|\nabla e\|_{R_T^\ell}. \quad (5.12)$$

Turning to Ξ_2 , we remark that the contributions to this term also vanish on all the mesh elements outside R_T^ℓ since $\ell \geq 2$. As above, we arrive at

$$\Xi_2 \lesssim \|\nabla \mathcal{K}_T^\ell v\|_{R_T^\ell} (\|\nabla e\|_{R_T^\ell} + \|e\|_{R_T^\ell}) \lesssim \|\nabla \mathcal{K}_T^\ell v\|_{R_T^\ell} \|\nabla e\|_{R_T^\ell}.$$

Putting the above estimates together, we obtain that

$$\|\nabla e\|_\Omega^2 \lesssim \sum_{T \in \mathcal{T}_H} \|\nabla \mathcal{K}_T^\ell v\|_{R_T^\ell} \|\nabla e\|_{R_T^\ell} \lesssim \exp(-C_{\text{dec}}\ell) \sum_{T \in \mathcal{T}_H} \|\nabla \mathcal{K}_T^\ell v\|_{\mathbf{N}^\ell(T)} \|\nabla e\|_{R_T^\ell}, \quad (5.13)$$

where we applied Theorem 4.2 to (5.8), treating the patch $\mathbf{N}^\ell(T)$ as the whole domain. In order to pass from the norm of \mathcal{K}_T^ℓ to that of v , we again use classical saddle point theory (see, e.g. [13], Cor. 4.2.1), recalling that

the inf-sup constant of b is of order H as outlined in (3.7). This results in

$$\begin{aligned} \|\nabla \mathcal{K}_T^\ell v\|_{\mathbf{N}^\ell(T)} &\lesssim \sup_{w \in V_T^\ell} \frac{a_T(\mathcal{I}_H v, w)}{\|\nabla w\|_\Omega} + H^{-1} \sup_{\mu \in M_T^\ell} \frac{c_T(v, \mu)}{\|\mu\|_\Omega} \\ &\lesssim \|\nabla v\|_{\mathbf{N}(T)} + H^{-1} \|v - \bar{v}_T\|_{\mathbf{N}(T)} \lesssim \|\nabla v\|_{\mathbf{N}(T)} \end{aligned} \quad (5.14)$$

where \bar{v}_T is the average of v over $\mathbf{N}(T)$ if T contains a boundary node, and $\bar{v}_T = 0$ otherwise. The justification of the last inequality in (5.14) requires to consider the cases where T is completely inside Ω and T is adjacent to the boundary separately. In the first case, we can subtract the average \bar{v}_T inside the bilinear form c_T due to Item 2 on Page 10, and then proceed by using Item 4 and the Poincaré inequality on $\mathbf{N}(T)$. Note that the constant in the Poincaré inequality is of order H , as seen from scaling and maximizing over all possible configurations of $\mathbf{N}(T)$ allowed by the mesh regularity. In the other case where T lies at the boundary $\partial\Omega$, at least one of the boundary faces of $\mathbf{N}(T)$ lies on $\partial\Omega$ and we can conclude by the Friedrichs-type inequality $\|v\|_{\mathbf{N}(T)} \lesssim H \|\nabla v\|_{\mathbf{N}(T)}$, which holds since v vanishes on $\partial\Omega$.

Returning to (5.13), we conclude that

$$\begin{aligned} \|\nabla e\|_\Omega^2 &\lesssim \exp(-C_{\text{dec}}\ell) \sqrt{\sum_{T \in \mathcal{T}_H} \|\nabla v\|_{\mathbf{N}(T)}^2} \sqrt{\sum_{T \in \mathcal{T}_H} \|\nabla e\|_{R_T^2}} \\ &\lesssim \ell^{(d-1)/2} \exp(-C_{\text{dec}}\ell) \|\nabla v\|_\Omega \|\nabla e\|_\Omega, \end{aligned}$$

where we have used the fact that each element $K \in \mathcal{T}_H$ belongs to at most $\mathcal{O}(\ell^{d-1})$ rings R_T^2 for different $T \in \mathcal{T}_H$. Dividing by $\|\nabla e\|_\Omega$ gives the assertion. \square

6. LOCALIZED MULTISCALE METHOD

In this section, we introduce a practical multiscale method with locally computable basis functions. This is reasonable due to the exponential decay behavior of the globally defined functions. We define the localized multiscale space as $\tilde{V}_H^\ell = \mathcal{R}^\ell V$. Using that the operator \mathcal{R}^ℓ depends on its argument only through the QOI introduced in (3.8), allows us to write

$$\tilde{V}_H^\ell := \text{span}\{\tilde{\varphi}_j^\ell : j = 1, \dots, J\}, \quad (6.1)$$

with appropriate basis functions associated with the QOI by the property that $q_i(\tilde{\varphi}_j^\ell) = \delta_{ij}$. Formally, these basis functions can be written as $\tilde{\varphi}_j^\ell = \mathcal{R}^\ell \tilde{\varphi}_j$. This property allows us to practically compute $\tilde{\varphi}_j^\ell$ as a linear combination of the solution to a few local problems as explained in the following. To this end, we compute the coefficients κ_{zj} for $z \in \mathcal{N}_H^{\text{in}}$ and $j = 1, \dots, J$ such that $\mathcal{I}_H \tilde{\varphi}_j = \sum_{z \in \mathcal{N}_H^{\text{in}}} \kappa_{zj} \Lambda_z^1$, where $\mathcal{N}_H^{\text{in}}$ denotes the set of interior nodes. These coefficients are easily deduced from the definitions in (5.3) and (5.4) for the CG-LOD and DG-LOD, respectively, and the known QOI of $\tilde{\varphi}_j$. We can then write the localized basis function $\tilde{\varphi}_j^\ell$ as

$$\tilde{\varphi}_j^\ell = \sum_{z \in \mathcal{N}_H^{\text{in}}} \kappa_{zj} \Lambda_z^1 - \sum_{T \in \mathcal{T}_H} \psi_{j,T}^\ell, \quad (6.2)$$

where the pair $(\psi_{j,T}^\ell, \lambda_{j,T}^\ell) \in V_T^\ell \times M_T^\ell$ solves the local saddle point problem

$$a(\psi_{j,T}^\ell, w) + b(w, \lambda_{j,T}^\ell) = \sum_{z \in \mathcal{N}_H^{\text{in}}} \kappa_{zj} a_T(\Lambda_z^1, w), \quad \forall w \in V_T^\ell, \quad (6.3a)$$

$$b(\psi_{j,T}^\ell, \mu) = -\frac{|T \cap \omega_j|}{|\omega_j|} \mu_j + \sum_{z \in \mathcal{N}_H^{\text{in}}} \kappa_{zj} (\mu, \Lambda_z^1)_T, \quad \forall \mu \in M_T^\ell. \quad (6.3b)$$

Note that only a small number of these problems have to be solved for each j . In the case of CG-LOD, the functions $\psi_{j,T}^\ell$ are non-zero only for mesh elements $T \subset \bar{\omega}_j$. For the DG-LOD, if the index j is associated with the characteristic function on an element K , then the functions $\psi_{j,T}^\ell$ are non-zero for elements $T \subset \mathbf{N}(K)$; for all other indices j corresponding to basis functions of M_H supported on K , we have that $\tilde{\varphi}_j^\ell = -\psi_{j,K}^\ell$. We emphasize that, compared to naive localization strategies as in [42, 44], the computational cost of the proposed stabilized localization strategy differs only for the basis functions associated with the lowest-order QOI. Specifically, these are the hat functions for the CG-LOD and the characteristic functions of elements for the DG-LOD.

The LOD method seeks the unique function $\tilde{u}_H \in \tilde{V}_H^\ell$ such that

$$a(\tilde{u}_H^\ell, \tilde{v}_H^\ell) = (f, \tilde{v}_H^\ell)_\Omega, \quad \forall \tilde{v}_H^\ell \in \tilde{V}_H^\ell. \quad (6.4)$$

Theorem 6.1 (Localized method). *The localized multiscale method (6.4) is well-posed. Moreover, for any right-hand side $f \in H^s(\Omega)$ in the case of the CG-LOD, and $f \in H^s(\mathcal{T}_H)$ for the DG-LOD, with $s \in \{0, 1, \dots, p+1\}$, we have the error estimates*

$$\|\nabla(u - \tilde{u}_H^\ell)\|_\Omega \lesssim H^{1+s}|f|_{s, \mathcal{T}_H} + \ell^{(d-1)/2} \exp(-C_{\text{dec}}\ell)\|f\|_\Omega, \quad (6.5)$$

$$\|u - \tilde{u}_H^\ell\|_\Omega \lesssim (H + \ell^{(d-1)/2} \exp(-C_{\text{dec}}\ell))\|\nabla(u - \tilde{u}_H^\ell)\|_\Omega, \quad (6.6)$$

where $C_{\text{dec}} > 0$ is the constant introduced in Theorem 4.2.

Proof. The H^1 -error estimate (6.5) follows from (3.4) and (5.10) using that

$$\|\nabla(u - \tilde{u}_H^\ell)\|_\Omega \lesssim \|\nabla(u - \mathcal{R}^\ell u)\|_\Omega \leq \|\nabla(u - \tilde{u}_H)\|_\Omega + \|\nabla(\mathcal{R}u - \mathcal{R}^\ell u)\|_\Omega,$$

where we have used Céa's lemma in the first step and the identity $\tilde{u}_H = \mathcal{R}u$ in the second step.

The L^2 -error estimate (6.6) is proven by the Aubin–Nitsche trick, introducing $w \in V$ such that

$$a(w, v) = (u - \tilde{u}_H^\ell, v)_\Omega$$

for all $v \in V$ and observing $\|u - \tilde{u}_H^\ell\|_\Omega^2 = a(u - \tilde{u}_H^\ell, w - \mathcal{R}^\ell w) \lesssim \|\nabla(u - \tilde{u}_H^\ell)\|_\Omega \|\nabla(w - \mathcal{R}^\ell w)\|_\Omega$. It remains to apply the H^1 -interpolation error estimate for w treated as the solution to the governing PDE with the right-hand side $u - \tilde{u}_H^\ell \in L^2(\Omega)$, similar to the already proven estimate for u with $s = 0$. A more detailed proof, in the DG-LOD case, can be found in Theorem 6.2 from [20]. \square

7. MORE GENERAL PROBLEMS

This section demonstrates that the general higher-order LOD framework developed for the elliptic model problem can be readily extended to more complex settings. We illustrate this with two examples: the heterogeneous Helmholtz problem and the Gross–Pitaevskii eigenvalue problem. Both examples showcase successful applications of LOD-type methods, which have also been extensively studied in the literature.

For the Helmholtz problem, the LOD method has the advantage of alleviating the severe mesh-size restrictions that arise for large wave numbers, a phenomenon commonly known as the pollution effect, which classical low-order finite element methods suffer from, provided suitable oversampling conditions are met. In addition, as a multiscale method, the LOD naturally handles heterogeneous coefficients, offering a further advantage. For the Gross–Pitaevskii eigenvalue problem, LOD-type methods are particularly appealing because they employ basis functions with superior approximation properties compared to classical finite elements of the same order. This effectively reduces the system size of the nonlinear eigenvalue problem.

7.1. Heterogeneous Helmholtz problem

In comparison with the second-order diffusion-type problems introduced in Section 2, the Helmholtz problem additionally features a zeroth-order term with a sign opposite to that of the second-order operator. The strength of this term is determined by the wavenumber $\kappa > 0$. For large κ , the problem becomes strongly indefinite and its solution is oscillatory, which typically results in significant numerical challenges, cf. [8]. The Helmholtz-type problem considered here seeks a complex-valued solution $u : \Omega \rightarrow \mathbb{C}$ satisfying

$$\begin{cases} -\operatorname{div}(A\nabla u) - \kappa^2 \mathcal{V}^2 u = f & \text{in } \Omega, \\ A\nabla u \cdot \nu - i\kappa \sigma u = 0 & \text{on } \partial\Omega, \end{cases} \quad (7.1)$$

where A is a matrix-valued coefficient as in Section 2, $\mathcal{V} \in L^\infty(\Omega; \mathbb{R})$ satisfies the uniform bounds $0 < \mathcal{V}_{\min} \leq \mathcal{V}(x) \leq \mathcal{V}_{\max} < \infty$ almost everywhere in Ω , and $\sigma \in L^\infty(\partial\Omega; \mathbb{R})$ is almost everywhere positive on $\partial\Omega$. The right-hand side f belongs to $L^2(\Omega; \mathbb{C})$. This problem describes the acoustic wave propagation in heterogeneous media and is a generalization of the classical (homogeneous) Helmholtz problem.

The weak formulation of (7.1) seeks a function $u \in V := H^1(\Omega; \mathbb{C})$ such that

$$a(u, v) = \int_{\Omega} f \bar{v} \, dx, \quad \forall v \in V, \quad (7.2)$$

where \bar{v} is the complex conjugate of v , and $a : V \times V \rightarrow \mathbb{C}$ is defined by

$$a(u, v) := \int_{\Omega} A\nabla u \cdot \nabla \bar{v} \, dx - \kappa^2 \int_{\Omega} \mathcal{V}^2 u \bar{v} \, dx - i\kappa \int_{\partial\Omega} \sigma u \bar{v} \, ds.$$

Note that a is sesquilinear. A natural norm for the Helmholtz problem is

$$\|v\|_{\kappa}^2 := \|A^{1/2} \nabla v\|_{\Omega}^2 + \kappa^2 \|\mathcal{V}v\|_{\Omega}^2. \quad (7.3)$$

Henceforth, we assume that the weak formulation (7.2) of the Helmholtz problem is well-posed and admits a unique solution that satisfies the stability estimate

$$\|u\|_{\kappa} \lesssim \kappa^n \|f\|_{\Omega}, \quad (7.4)$$

for some $n \geq 0$. For homogeneous coefficients and general Lipschitz domains, the well-posedness of (7.4) can be proved with $n = 5/2$; see [25]. In the case of variable coefficients, the analysis becomes substantially more involved; see, e.g., [17, 28]. Note that the polynomial-in- κ stability assumption (7.4) is classical for the numerical analysis of Helmholtz problems.

Several works use and analyze the LOD method for Helmholtz problems, including [14, 31, 43, 51, 52]; see also [27]. A key ingredient in the corresponding analysis is the coercivity of the sesquilinear form a when restricted to the fine-scale space W , defined similarly to (3.1). The following lemma establishes this property for both the proposed CG-LOD and DG-LOD method.

Lemma 7.1 (Coercivity on fine-scale space). *Assume the resolution condition*

$$H\kappa p^{-1} \leq \sqrt{\alpha}(\sqrt{2}V_{\max}C_{\text{ap}})^{-1} \quad (7.5)$$

with a constant $C_{\text{ap}} > 0$ depending solely on the shape-regularity of \mathcal{T}_H . Then the sesquilinear form a is coercive on $W \times W$, i.e.,

$$\Re a(w, w) \geq \frac{\alpha}{2} \|\nabla w\|_{\Omega}^2, \quad \forall w \in W.$$

Proof. We use the uniform bounds on the coefficients A and \mathcal{V} , and the approximation property of the L^2 -projection $\Pi_H: L^2(\Omega) \rightarrow M_H$, given by

$$\|v - \Pi_H v\|_\Omega \leq C_{\text{ap}} H p^{-1} \|\nabla v\|_\Omega,$$

for a constant $C_{\text{ap}} > 0$ independent of H and p . Recall that we choose $M_H = \mathcal{P}^p(\mathcal{T}_H)$ for the DG-LOD and $M_H = \mathcal{P}^p(\mathcal{T}_H) \cap H^1(\Omega)$ for the CG-LOD. Such a p -explicit approximation result can be found, for instance, in Section 3 from [9]. This approximation result then yields, for any $w \in W$, the following estimate:

$$\Re a(w, w) \geq \alpha \|\nabla w\|_\Omega^2 - \kappa^2 \mathcal{V}_{\max}^2 \|w\|_\Omega^2 \geq (\alpha - \kappa^2 \mathcal{V}_{\max}^2 C_{\text{ap}}^2 H^2 p^{-2}) \|\nabla w\|_\Omega^2.$$

The desired coercivity estimate then follows with the resolution condition. \square

Note that, unlike in the rest of the paper, we have now explicitly tracked the p -dependence to examine how the polynomial degree influences the resolution condition and to enable a comparison with other methods; see Remark 7.3.

To construct the basis functions of the LOD trial space, one solves local corrector problems analogous to (6.3), where the sesquilinear forms a and a_T now are those of the Helmholtz problem. The well-posedness of these problems follows directly from classical inf-sup theory (see, e.g., Corollary 4.2.1 from [13], which extends to the complex-valued setting), noting that the sesquilinear form a is coercive on the kernel of b as shown in Lemma 7.1 and the inf-sup condition (3.7). The global trial space of the LOD is then defined as the span of the resulting basis functions, in the spirit of (6.1) and (6.2). Since the Helmholtz problem is non-Hermitian, the LOD method employs different trial and test spaces. However, owing to the specific structure of the Helmholtz equation, the test space can be obtained as the complex conjugate of the trial space; see equation (4.6) from [51]. Therefore, no additional computations for the test space basis functions are required, and the LOD approximation to the Helmholtz problem is given by the solution $\tilde{u}_h^\ell \in \tilde{V}_H^\ell$ satisfying

$$a(\tilde{u}_H^\ell, \overline{\tilde{v}_H^\ell}) = \int_\Omega f \tilde{v}_H^\ell \, dx, \quad \forall \tilde{v}_H^\ell \in \tilde{V}_H^\ell.$$

The following theorem establishes the convergence of the CG-LOD and DG-LOD methods for the Helmholtz problem.

Theorem 7.2 (Localized method for Helmholtz). *Under the resolution condition from Lemma 7.1 and the oversampling condition $\ell \gtrsim \log(\kappa)$, the LOD method for (7.2) is well-posed. Moreover, with $f \in H^s(\Omega, \mathbb{C})$ for the CG-LOD, and $f \in H^s(\mathcal{T}_H; \mathbb{C})$ for the DG-LOD, and $s \in \{0, 1, \dots, p+1\}$, we have*

$$\|u - \tilde{u}_H^\ell\|_\kappa \lesssim H^{1+s} |f|_{s, \mathcal{T}_H} + \kappa^n \ell^{\frac{d-1}{2}} \exp(-C_{\text{dec}} \ell) \|f\|_\Omega \quad (7.6)$$

with n from (7.4) and the decay rate $C_{\text{dec}} > 0$ is independent of H , ℓ , and κ .

Proof. The proof combines arguments from the LOD analysis for the elliptic model problem (see, e.g., Thms. 4.2 and 5.1) with techniques specific to the Helmholtz setting (see, e.g., [31, 51]). For brevity, the details are omitted. \square

Remark 7.3 (Analogies to hp -FEM). The assumptions on the discretization parameters required for the stability and quasi-optimality of the hp -FEM for Helmholtz problems (see, e.g., [46]) are notably similar to those in the above theorem. In the case of homogeneous coefficients (for an extension to piecewise smooth coefficients, see [12]), it was shown that a stable and quasi-optimal finite element approximation can be achieved if the polynomial degree satisfies $p \approx \log(\kappa)$ and the resolution condition $H\kappa p^{-1} \lesssim 1$ holds, along with a suitable refinement strategy near geometric singularities. This resolution condition is the same as in Lemma 7.1 (up to a constant), and the role of the polynomial degree in the hp -FEM is similar to that of the localization parameter in the LOD; see Theorem 7.2.

7.2. Gross–Pitaevskii problem

As another example, we consider the Gross–Pitaevskii problem, which arises in quantum physics as a model for quantum states of so-called Bose–Einstein condensates. The problem is posed on a convex Lipschitz domain Ω , with homogeneous boundary conditions imposed on its boundary. This corresponds to the choice $V := H_0^1(\Omega)$. Note that due to the rapid decay of low-energy quantum states, the restriction to a sufficiently large domain, along with homogeneous Dirichlet boundary conditions, is a physically reasonable modeling assumption. Stationary quantum states correspond to the critical points of the Gross–Pitaevskii energy functional, defined as

$$\mathcal{E}(v) := \frac{1}{2}(\nabla v, \nabla v)_\Omega + \frac{1}{2}(\mathcal{V}v, v)_\Omega + \frac{\kappa}{4}(|v|^2 v, v)_\Omega, \quad v \in V, \tag{7.7}$$

subject to the L^2 -normalization constraint $\|v\|_\Omega = 1$. Here, $\mathcal{V} \in L^\infty(\Omega)$ is a non-negative (possibly rough) trapping potential and the parameter $\kappa > 0$ characterizes the strength of repulsive interactions between particles. Of particular physical interest is the ground state, which corresponds to the stationary quantum state of lowest energy, *i.e.*, it solves the constrained minimization problem

$$u \in \underset{v \in V : \|v\|_\Omega = 1}{\operatorname{arg\,min}} \mathcal{E}(v), \tag{7.8}$$

and the minimal energy is denoted by $E := \mathcal{E}(u)$. The corresponding Euler–Lagrange equations imply that the ground state u , together with an eigenvalue $\lambda \in \mathbb{R}$, solves the nonlinear eigenvalue problem

$$(\nabla u, \nabla v)_\Omega + (\mathcal{V}u, v)_\Omega + \kappa(|u|^2 u, v)_\Omega = \lambda(u, v)_\Omega, \quad \forall v \in V, \tag{7.9}$$

where λ is referred to as the ground state eigenvalue. In the above setting, it is a classical result that the ground state eigenvalue is the smallest eigenvalue among all eigenpairs of (7.9) and it is simple. Moreover, the ground state is unique up to sign and can be chosen to be strictly positive in the interior of Ω , cf. [15].

For the construction of the LOD method (we restrict ourselves to the prototypical method for simplicity; a localization can be performed analogously to Sects. 4 and 5), we consider only the terms on the left-hand side of (7.9) which are linear in u . The resulting bilinear form $a : V \times V \rightarrow \mathbb{R}$ is defined as

$$a(w, v) := (\nabla w, \nabla v)_\Omega + (\mathcal{V}w, v)_\Omega. \tag{7.10}$$

The approximation space \tilde{V}_H of the prototypical LOD is then defined as in (3.2), using the modified bilinear form from (7.10). Depending on whether the space M_H , used in the definition of the fine-scale space W in (3.1), consists of globally continuous or \mathcal{T}_H -piecewise polynomials, one obtains a CG- or DG-version of the LOD method, respectively. The prototypical LOD approximation is then defined as the solution to the finite-dimensional constrained minimization problem.

$$\tilde{u}_H \in \underset{\tilde{v}_H \in \tilde{V}_H : \|\tilde{v}_H\|_\Omega = 1}{\operatorname{arg\,min}} \mathcal{E}(\tilde{v}_H), \tag{7.11}$$

and the corresponding minimal energy is denoted by $E_H := \mathcal{E}(\tilde{u}_H)$. While the existence of such a minimizer follows from classical compactness arguments in the finite-dimensional setting, its uniqueness is generally not guaranteed. An overview of algorithms to practically solve (7.11) (after a localization of the basis functions) can be found in [34]. The Euler–Lagrange equations corresponding to (7.11) give rise to the following finite-dimensional nonlinear eigenvalue problem: seek $\tilde{u}_H \in \tilde{V}_H$ and an associated eigenvalue $\lambda_H \in \mathbb{R}$ such that

$$(\nabla \tilde{u}_H, \nabla \tilde{v}_H)_\Omega + (\mathcal{V}\tilde{u}_H, \tilde{v}_H)_\Omega + \kappa(|\tilde{u}_H|^2 \tilde{u}_H, \tilde{v}_H)_\Omega = \lambda_H(\tilde{u}_H, \tilde{v}_H)_\Omega, \quad \forall \tilde{v}_H \in \tilde{V}_H.$$

As a preliminary step towards proving the convergence of the prototypical LOD method for the Gross–Pitaevskii problem, we first analyze the approximation properties of the space \tilde{V}_H , as done in the following lemma.

Lemma 7.4 (Approximation properties of \tilde{V}_H). *Assume that $u \in H^s(\Omega)$ as well as $|u|^2u \in H^s(\Omega)$ with $s \in \{0, 1, \dots, p+1\}$, and let $\mathfrak{P}u \in \tilde{V}_H$ be the solution to*

$$a(\mathfrak{P}u, \tilde{v}_H) = -\kappa(|u|^2u, \tilde{v}_H)_\Omega - \lambda(u, \tilde{v}_H)_\Omega, \quad \forall \tilde{v}_H \in \tilde{V}_H. \quad (7.12)$$

Then, we have

$$\|\nabla(u - \mathfrak{P}u)\|_\Omega \lesssim H^{1+s}(|u|^2u|_{s,\Omega} + |u|_{s,\Omega}). \quad (7.13)$$

Proof. By coercivity and symmetry of a , and using (7.9) and (7.12), we obtain that

$$\|\nabla(u - \mathfrak{P}u)\|_\Omega^2 \leq a(u - \mathfrak{P}u, u) = -\kappa(|u|^2u, u - \mathfrak{P}u)_\Omega - \lambda(u, u - \mathfrak{P}u)_\Omega.$$

Since $u - \mathfrak{P}u \in W$ by construction, and using the approximation properties of the L^2 -projection $\Pi_H: L^2(\Omega) \rightarrow M_H$, it follows that

$$\begin{aligned} \|\nabla(u - \mathfrak{P}u)\|_\Omega^2 &\leq -\kappa(|u|^2u - \Pi_H(|u|^2u), (u - \mathfrak{P}u) - \Pi_H(u - \mathfrak{P}u))_\Omega \\ &\quad - \lambda(u - \Pi_H u, (u - \mathfrak{P}u) - \Pi_H(u - \mathfrak{P}u))_\Omega \\ &\lesssim H^{1+s}(|u|^2u|_{s,\Omega} + |u|_{s,\Omega}) \|\nabla(u - \mathfrak{P}u)\|_\Omega, \end{aligned}$$

where the hidden constant depends on λ . This yields (7.13). \square

The following theorem proves the convergence of the prototypical LOD method.

Theorem 7.5 (Prototypical method for the Gross–Pitaevskii problem).

Assume that $u \in H^s(\Omega)$ and $|u|^2u \in H^s(\Omega)$ with $s \in \{0, 1, \dots, p+1\}$. Then the discrete ground state and the corresponding energy fulfill

$$\|\nabla(u - \tilde{u}_H)\|_\Omega \lesssim H^{1+s}(|u|^2u|_{s,\Omega} + |u|_{s,\Omega}), \quad |E - E_H| \lesssim \|\nabla(u - \tilde{u}_H)\|_\Omega^2.$$

Proof. From the general convergence theory developed in Theorem 1 of [15], we derive under the stated assumptions that the solution to (7.2) fulfills a quasi-best approximation property in the $H^1(\Omega)$ -norm and the stated energy error holds. The result then follows directly invoking Lemma 7.4. \square

Remark 7.6 (L^2 - and eigenvalue approximation). Note that L^2 -error estimates for the ground state and the eigenvalue can be derived as well. Under suitable regularity assumptions on the dual problem defined in the proof of Theorem 3 from [15], we obtain an additional order of convergence in the $L^2(\Omega)$ -norm for the ground state approximation, compared to the H^1 -estimate in Theorem 7.5. Under similar assumptions, the eigenvalue approximation exhibits the same convergence rate as the energy approximation. A proof of these results is beyond the scope of this work; for details, see [35], which provides the proof for the CG-LOD with $p = 1$.

Remark 7.7 (Regularity of the solution). Under the assumptions made in this section, one can show that u and $|u|^2u$ belong to $H^2(\Omega)$, with corresponding norms bounded independently of the oscillations of \mathcal{V} ; see Lemma 2.2 from [35]. Using similar arguments and assuming more regular \mathcal{V} and a smooth boundary, even higher regularity of the solution can be shown, *e.g.*, $u \in H^{p+1}(\Omega)$, $|u|^2u \in H^{p+1}(\Omega)$ for some $p > 1$. These considerations justify the assumptions in Lemma 7.4 and Theorem 7.5. Note that, however, the seminorms of order greater than two may no longer be bounded independently of the oscillations of \mathcal{V} . The additional boundary regularity assumptions are not required if u is compactly supported in Ω . While exact compact support may not occur in practice, the ground state typically exhibits a rapid decay (see, *e.g.*, [11], Thm. 2.5), which allows to relax the boundary regularity assumptions. Overall, under the same regularity assumptions, the LOD solution is expected to be two orders more accurate than a classical higher-order finite element method; see also the numerical investigation in [19].

8. A VARIANT OF THE CLASSICAL LOD METHOD

We end the theoretical part of this work with some insight into an improvement of the classical LOD from [45] on the example of the model problem (2.2) from Section 2. This version is based on a suitable quasi-interpolation operator $\mathcal{I}_H^{\text{avg}}$ mapping to the continuous \mathcal{P}^1 -finite element space $V_H \subset V$. This operator is defined by $\mathcal{I}_H^{\text{avg}} = E_H \circ \Pi_H^{\text{dg}}$, where Π_H^{dg} is the $L^2(\Omega)$ -orthogonal projection onto the discontinuous \mathcal{P}_1 -finite element space V_H^{dg} , and $E_H: V_H^{\text{dg}} \rightarrow V_H$ is a classical averaging operator, defined at any interior node z by $E_H v(z) = \sum_{T \subset \omega_z} \frac{|T|}{|\omega_z|} v|_T(z)$ with ω_z denoting the patch of mesh elements around the node z . Note that E_H is also referred to as Oswald interpolation operator, see [47]. In [45], the nodal values of $\mathcal{I}_H^{\text{avg}} v$ on the boundary $\partial\Omega$ were set to zero. Here, we propose to apply the averaging E_H at all the nodes of the mesh, including the boundary ones, so that the target space of $\mathcal{I}_H^{\text{avg}}$ is no longer $H_0^1(\Omega)$ -conforming. With this choice in mind, we define a variant of the LOD method, referred to as CGA-LOD, by introducing the fine-scale space $W = \text{Ker}(\mathcal{I}_H^{\text{avg}}|_V) \subset V$, and defining the prototypical LOD space \tilde{V}_H by (3.2). We shall see that this slight modification in the treatment of the degrees of freedom at boundary nodes results in improving the error estimate by one order in H compared to the version in [45].

To accommodate the CGA-LOD within the framework of this article, we reformulate the fine-scale space W as the $L^2(\Omega)$ -orthogonal complement of a space of certain piecewise polynomials. To this end, consider any element $T \in \mathcal{T}_H$, which is a simplex with the vertices $\mathcal{N}(T)$, and introduce $\psi_{z,T} \in \mathcal{P}^1(T)$ for $z \in \mathcal{N}(T)$ such that $\int_T \Lambda_y^1 \psi_{z,T} dx = \delta_{yz}$ holds for all $y, z \in \mathcal{N}(T)$. Here, Λ_y^1 denotes the nodal basis function in \mathcal{P}^1 associated to the node y . Then, for any $v \in V$, we have locally on the element T , that $\Pi_H^{\text{dg}}(v) = \sum_{z \in \mathcal{N}(T)} \left(\int_T \psi_{z,T} v dx \right) \Lambda_z^1$. Combining this with the averaging operator E_H , we arrive at

$$\mathcal{I}_H^{\text{avg}}(v) = \sum_{z \in \mathcal{N}_H} \sum_{T \subset \omega_z} \frac{|T|}{|\omega_z|} \left(\int_T \psi_{z,T} v dx \right) \Lambda_z^1 = \sum_{z \in \mathcal{N}_H} \left(\int_{\omega_z} \psi_z v dx \right) \Lambda_z^1$$

with discontinuous piecewise linear functions ψ_z defined on $T \subset \omega_z$ by $\psi_z|_T = \frac{|T|}{|\omega_z|} \psi_{z,T}$. Outside ω_z , we extend ψ_z by zero. We can thus interpret the fine-scale space as in (3.1) with $M_H = \text{span}\{\psi_z, z \in \mathcal{N}_H\}$. This allows us to define the basis functions of the prototypical LOD space \tilde{V}_H as in (3.10) with $\Lambda_j = \psi_z$ (we identify here the numbering by j with the indexing by the nodes z). This leads to the prototypical CGA-LOD method, cf. (3.3). Its solution satisfies the following *a priori* error estimate.

Theorem 8.1 (Variant of classical LOD with different boundary treatment). *The CGA-LOD version of method (3.3) is well-posed, and its solution is given as $\tilde{u}_H = \mathcal{R}u$, where $\mathcal{R}: V \rightarrow \tilde{V}_H$ is the a -orthogonal projection onto \tilde{V}_H . Furthermore, if $f \in H^s(\Omega)$, with $s \in \{0, 1\}$, then*

$$\|\nabla(u - \tilde{u}_H)\|_{\Omega} \lesssim H^{1+s} |f|_{s,\Omega}. \tag{8.1}$$

Proof. The well-posedness of (3.3) and the identity $\tilde{u}_H = \mathcal{R}u$ are proved as in Theorem 3.1. With $e := u - \tilde{u}_H \in W$ and using the coercivity of a , we observe that

$$\alpha \|\nabla(u - \tilde{u}_H)\|_{\Omega}^2 \leq a(u - \mathcal{R}u, e) = (f, e - \mathcal{I}_H^{\text{avg}} e)_{\Omega}, \tag{8.2}$$

since $\mathcal{I}_H^{\text{avg}} e = 0$. Applying $\|e - \mathcal{I}_H^{\text{avg}} e\|_{\Omega} \lesssim H |e|_{1,\Omega}$ as proved in Lemma 3.2 of [45] gives (8.1) for $s = 0$.

To prove (8.1) in the case $s = 1$, we note that

$$(f, e - \mathcal{I}_H^{\text{avg}} e)_{\Omega} = (f - \mathcal{I}_H^{\psi} f, e - \mathcal{I}_H^{\text{avg}} e)_{\Omega}, \tag{8.3}$$

where $\mathcal{I}_H^{\psi}: L^2(\Omega) \rightarrow M_H$ is defined by

$$\mathcal{I}_H^{\psi} v = \sum_{z \in \mathcal{N}_H} \left(\int_{\omega_z} \Lambda_z^1 v \right) \psi_z.$$

Equality (8.3) holds since $e \in W$ is $L^2(\Omega)$ -orthogonal to all ψ_z . In order to bound $f - \mathcal{I}_H^\psi f$ in (8.3), we take any $T \in \mathcal{T}_H$ and observe that $(\mathcal{I}_H^\psi v)|_T = 1$ if $v|_{\mathbb{N}(T)} = 1$. Indeed, in this case we have

$$(\mathcal{I}_H^\psi v)|_T = \sum_{z \in \mathcal{N}(T)} \left(\int_{\omega_z} \Lambda_z^1 dx \right) \frac{|T|}{|\omega_z|} \psi_{z,T} = \sum_{z \in \mathcal{N}(T)} \left(\int_T \Lambda_z^1 dx \right) \psi_{z,T}.$$

Thus, $\mathcal{I}_H^\psi(v)|_T \in \mathcal{P}_1(T)$ and satisfies for all $y \in \mathcal{N}(T)$

$$\int_T (\mathcal{I}_H^\psi v) \Lambda_y^1 dx = \sum_{z \in \mathcal{N}(T)} \left(\int_T \Lambda_z^1 dx \right) \left(\int_T \psi_{z,T} \Lambda_y^1 dx \right) = \int_T \Lambda_y^1 dx.$$

Therefore, $(\mathcal{I}_H^\psi v)|_T = 1$ as claimed. For $f \in H^1(\Omega)$, this implies that

$$\|f - \mathcal{I}_H^\psi f\|_T = \inf_{p \in \mathbb{R}} \|f - p - \mathcal{I}_H^\psi(f - p)\|_T \lesssim \inf_{p \in \mathbb{R}} \|f - p\|_{\mathbb{N}(T)} \lesssim H \|\nabla f\|_{\mathbb{N}(T)}$$

by the Poincaré inequality. Using this in (8.2) and (8.3), together with the stated approximation properties of $\mathcal{I}_H^{\text{avg}}$, leads to estimate (8.1) for $s = 1$. \square

The order $\mathcal{O}(H^2)$ has also been proved for the original version of the LOD in Lemma 8.1 from [45] under the assumption that $f \in H_0^1(\Omega)$. The proof of Theorem 8.1 is inspired by that result. For a general $f \in H^1(\Omega)$, however, the original version only provides an error estimate of order $\mathcal{O}(H)$, see Theorem 3.4 from [45]. The CGA-LOD method, as presented in this section, can be localized in the same way as the other variants presented in this manuscript, following the strategy of Section 5. One needs to specify a quasi-interpolation operator $\mathcal{I}_H: V \rightarrow V_H$ onto the first-order $H_0^1(\Omega)$ -conforming finite element space V_H , compatible with the QOI of the current variant of LOD, and satisfying (5.2). In the context of CGA-LOD, \mathcal{I}_H can be readily constructed by reusing the expression for $\mathcal{I}_H^{\text{avg}}$ at the interior nodes, while setting it to 0 on the boundary. The resulting practical method satisfies the error estimates of Theorem 6.1 with $s \in \{0, 1\}$.

9. NUMERICAL EXPERIMENTS

In all numerical experiments, we consider the domain $\Omega = (0, 1)^2$, unless stated otherwise. For implementation purposes, we use uniform Cartesian meshes \mathcal{T}_H composed of square elements of side length H , and choose $M_H := \mathcal{Q}^p(\mathcal{T}_H)$ for the DG-LOD and $M_H := \mathcal{Q}^p(\mathcal{T}_H) \cap H^1(\Omega)$ for the CG-LOD, where $\mathcal{Q}^p(\mathcal{T}_H)$ denotes the space of \mathcal{T}_H -piecewise polynomials of coordinate degree at most p ; see also the footnote on Page 448. The local (infinite-dimensional) patch problems (5.8) are discretized using local submeshes of a fine Cartesian mesh \mathcal{T}_h , with mesh size $h < H$, fine enough to resolve all microscopic features of the coefficients. For this fine-scale discretization, we use the \mathcal{Q}^q -finite element method, with the polynomial degree $q \in \mathbb{N}$ specified individually in the subsections below. In the fully discrete convergence analysis, the space V is replaced by the fine-scale finite element space, and most arguments carry over directly; see, *e.g.*, Chapter 4.4 of [45]. This yields an *a priori* error estimate for the fully discrete LOD approximation with respect to the fine-scale finite element solution, analogous to Theorem 6.1. An estimate with respect to the weak solution of the original PDE then follows by applying the triangle inequality and standard finite element approximation results.

The numerical experiments presented below can be reproduced using the code available at https://github.com/moimmahauck/HO_LOD.

9.1. Heterogeneous elliptic problem

As a first numerical experiment, we consider the elliptic model problem (2.2) with the coefficient A_1 shown in Figure 1 (left). The coefficient is piecewise constant on a uniform Cartesian grid \mathcal{T}_ϵ with mesh size $\epsilon = 2^{-7}$. For all elements whose midpoints lie within a distance of 4ϵ from a given parabola, the coefficient is set to a




FIGURE 1. Coefficients A_1 and A_2 used in the first and second numerical experiments, respectively (*left* and *center*), and potential \mathcal{V} used in the third numerical experiment (*right*).

value of 2. In all other elements, the coefficient values are sampled independently from a uniform distribution on the interval $[0.1, 1]$. We further consider the two smooth source terms

$$f_1(x, y) = 2\pi^2 \sin(\pi x) \sin(\pi y), \quad f_2(x, y) = 1.$$

We emphasize that for the source term f_2 , as observed in Theorem 6.1, the first term on the right-hand side of error estimate (6.5) vanishes. This implies that, for f_2 , only the second summand, *i.e.*, the exponentially decaying localization error, remains. We abbreviate the relative errors with respect to the energy norm as

$$\text{err}_\alpha(H, \ell) := \frac{\|u_h - u_{H,h}^\ell\|_\alpha}{\|u_h\|_\alpha},$$

where $u_{H,h}^\ell$ denotes the fine-scale LOD approximation, and u_h is the fine-scale finite element solution as reference. The fine-scale discretization is performed using the Q^1 -finite element method on the fine mesh \mathcal{T}_h with mesh size $h = 2^{-9}$, which is sufficiently fine to resolve the microscopic details of the coefficient A . Note that, due to the expected low regularity of the analytical solution, higher-order finite elements do not offer an advantage in terms of convergence rates at the fine scale.

In Figure 2, the first row of plots illustrates the convergence behavior of the CG-LOD method with right-hand side f_1 . One can observe that the expected convergence rates can still be inferred from the first two data points, where all patches are global and the localization error is effectively zero. However, for mesh sizes H beyond 2^{-2} , the localization error completely dominates the convergence behavior. Although this observation is consistent with the theoretical results in Theorem 6.1, this numerical experiment clearly shows the unsatisfactory localization properties of the CG-LOD. The situation is quite different for the DG-LOD. With the same configuration as for the CG-LOD method, the second row of plots in Figure 2 clearly shows a convergence of order $p + 2$ for the LOD approximation as predicted by Theorem 6.1, provided that the oversampling parameter is sufficiently large. The noticeably larger gaps between the plateaus of the error curves for different oversampling parameters showcase the significantly improved localization properties of the DG-LOD compared to the CG-LOD.

In Figure 3 we compare the exponential decay of the localization error for the CG-LOD (first row) and the DG-LOD (second row). Recall that for the source term f_2 used in this experiment, the first term in the error estimate (6.5) vanishes, leaving only the exponentially decaying localization error. The numerical results in Figure 3 confirm the completely different localization behavior of the two methods. Specifically, as indicated by the dashed reference lines, the decay rate C_{dec} in Theorem 4.2 can be estimated to lie between 0.85 and 1.6 for the CG-LOD with polynomial orders $p \in \{1, 2, 3\}$. For the DG-LOD, the corresponding decay rates are notably higher, lying between 2 and 2.5. This showcases the faster exponential decay of the localization error and thus




FIGURE 2. Error plots for the CG-LOD and DG-LOD methods for polynomial degrees $p \in \{1, 2, 3\}$ for the source term f_1 . For fixed oversampling parameters ℓ , the relative energy norm error is plotted as a function of coarse mesh size H .

the superior localization properties for the DG-LOD compared to the CG-LOD. We believe that this difference is related to the design of the QOI, in particular the more local, discontinuous character of the QOI in the DG setting. Note that, in general, increasing the polynomial degree leads to better localization.

9.2. High frequency heterogeneous Helmholtz problem

In the second numerical experiment, we demonstrate the performance of the higher-order DG-LOD in the context of high-frequency heterogeneous Helmholtz problems. The coefficient A_2 used in this experiment is shown in Figure 1 (center), and the wave number is chosen as $\kappa = 2^8$. Furthermore, the coefficients \mathcal{V} and σ are set to one. As the source term f_3 , we use an approximate point source located at $(1/8, 1/8)$, which is supported within a circle of radius r , defined as

$$f_3(x, y) = \begin{cases} 10^4 \cdot \exp\left(\frac{-1}{1 - \frac{(x-1/8)^2 + (y-1/8)^2}{1/20^2}}\right) & \text{if } (x - 1/8)^2 + (y - 1/8)^2 < r^2, \\ 0 & \text{else.} \end{cases}$$

A high wavenumber, such as $\kappa = 2^8$, leads to a pronounced pollution effect when classical low-order finite elements are used; see [8]. As a result, obtaining a quasi-optimal approximation demands considerably more




FIGURE 3. Error plots for the CG-LOD and DG-LOD methods for polynomial degrees $p \in \{1, 2, 3\}$ for the source term f_2 . For fixed coarse mesh sizes H , the relative energy norm error is plotted as a function of the oversampling parameter ℓ .




FIGURE 4. Real part of the Helmholtz LOD solution (*left*), and LOD Gross-Pitaevskii ground state approximation (*right*).




FIGURE 5. Error plots for the DG-LOD method for polynomial degree $p = 1$ for the source term f_3 with $r = 1/4$ and wave numbers $\kappa \in \{8, 16, 32\}$. For fixed oversampling parameters ℓ , the relative error with respect to norm $\|\cdot\|_\kappa$ is plotted as a function of coarse mesh size H .




FIGURE 6. Error plots for the DG-LOD with $p = 2$ for the Gross-Pitaevskii problem. For fixed oversampling parameters ℓ , the L^2 - and H^1 -errors of the ground state approximation (left), as well as the errors in the energy and eigenvalue approximations (right), are plotted as functions of the coarse mesh size H .

restrictive conditions on the fine-scale mesh size than are required merely to capture the oscillatory nature of the solution. In this numerical experiment, we use the \mathcal{Q}^2 -finite element method on the mesh \mathcal{T}_h with $h = 2^{-10}$ for the fine-scale discretization. This choice aligns with the mesh size condition $h \sim \kappa^{-5/4}$ for quadratic finite elements to achieve a quasi-optimal approximation; see Theorem 5.1 from [21] for the constant coefficient case.

Figure 4 (left) shows the real part of the LOD approximation for $p = 3$, $H = 2^{-6}$, $\ell = 4$, and f_3 for $r = 1/20$. The corresponding relative error in the norm $\|\cdot\|_\kappa$, defined in (7.3), against the fine-scale solution is 7.2970×10^{-4} . This result demonstrates that higher polynomial degrees on the coarse scale can further relax the resolution requirements, which is consistent with the theoretical result presented in Lemma 7.1.

In Figure 5, we also illustrate the convergence behavior of the DG-LOD method for different wave numbers κ , here shown for the case $p = 1$. For $\kappa \in \{8, 16\}$ we observe optimal-order convergence. For $\kappa = 32$, the method produces large errors on coarse meshes that do not satisfy the resolution condition (7.5). Once the mesh is sufficiently refined, the optimal convergence behavior is recovered. These results are fully consistent with the theoretical prediction in Theorem 7.2.

9.3. Gross–Pitaevskii eigenvalue problem

In the third numerical experiment, we apply the higher-order DG-LOD method to approximate the Gross–Pitaevskii ground state. We consider the domain $\Omega = (-6, 6)^2$, the particle interaction parameter $\kappa = 100$, and choose the potential defined as

$$\mathcal{V}(x) := \frac{1}{2}|x|^2 + 40 \times \text{tent}(x)\text{tent}(y),$$

where tent denotes the periodized version of the tent function on the interval $[0, 1]$, which attains the value one at 0.5 and vanishes at 0 and 1. This potential is illustrated in Figure 1 (right) and a corresponding ground state approximation is shown in Figure 4 (right). Since $\mathcal{V} \in W^{1,\infty}(\Omega)$, the DG-LOD method with degree $p = 2$ achieves optimal order convergence. This is because the assumption in Theorem 7.5, that u and $|u|^2u$ belong to $H^3(\Omega)$, is satisfied.

In the convergence plot shown in Figure 6, one observes the expected optimal rates of convergence, for the ground state approximation in the L^2 - and H^1 -norms, as well as for the energy and eigenvalue approximations, provided the oversampling parameter is chosen sufficiently large. We emphasize that, initially, for very coarse mesh sizes H , a reduced convergence order (*e.g.*, around three for the H^1 -error of the ground state approximation) is observed. This behavior is based on the fact that seminorms of order greater than two for u and $|u|^2u$ are not bounded independently of the oscillations in \mathcal{V} . As soon as these oscillations are sufficiently resolved, the theoretically predicted optimal convergence rates are attained. A similar behavior is observed for the L^2 -error, as well as for the energy and eigenvalue errors. Note that machine precision effects can be seen in the eigenvalue errors for small mesh sizes.

FUNDING

M. Hauck and R. Maier acknowledge funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 258734477 – SFB 1173. A. Lozinski acknowledges funding from ANR (French National Research Agency), Project NQG (ANR-23-CE40-0005-01). Part of this research was performed while the authors were visiting the Institute for Mathematical and Statistical Innovation (IMSI), which is supported by the National Science Foundation (Grant No. DMS-1929348).

DATA AVAILABILITY STATEMENT

The research data associated with this article are provided in Matlab code to reproduce the numerical experiments of the paper is available from github.com/moimmahauck/HO_LOD.

REFERENCES

- [1] A. Abdulle and Y. Bai, Reduced basis finite element heterogeneous multiscale method for high-order discretizations of elliptic homogenization problems. *J. Comput. Phys.* **231** (2012) 7014–7036.
- [2] A. Abdulle, E. Weinan, B. Engquist and E. Vanden-Eijnden, The heterogeneous multiscale method. *Acta Numer.* **21** (2012) 1–87.
- [3] G. Allaire and R. Brizzi, A multiscale finite element method for numerical homogenization. *Multiscale Model. Simul.* **4** (2005) 790–812.
- [4] R. Altmann, P. Henning and D. Peterseim, Numerical homogenization beyond scale separation. *Acta Numer.* **30** (2021) 1–86.
- [5] R. Araya, C. Harder, D. Paredes and F. Valentin, Multiscale hybrid-mixed method. *SIAM J. Numer. Anal.* **51** (2013) 3505–3531.
- [6] I. Babuška and R. Lipton, Optimal local approximation spaces for generalized finite element methods with application to multiscale problems. *Multiscale Model. Simul.* **9** (2011) 373–406.
- [7] I. Babuška and J.E. Osborn, Generalized finite element methods: their performance and their relation to mixed methods. *SIAM J. Numer. Anal.* **20** (1983) 510–536.
- [8] I.M. Babuška and S.A. Sauter, Is the pollution effect of the FEM avoidable for the Helmholtz equation considering high wave numbers? *SIAM J. Numer. Anal.* **34** (1997) 2392–2423.
- [9] I. Babuška and M. Suri, The optimal convergence rate of the p -version of the finite element method. *SIAM J. Numer. Anal.* **24** (1987) 750–776.

- [10] I. Babuška, G. Caloz and J.E. Osborn, Special finite element methods for a class of second order elliptic problems with rough coefficients. *SIAM J. Numer. Anal.* **31** (1994) 945–981.
- [11] W. Bao and Y. Cai, Mathematical theory and numerical methods for Bose–Einstein condensation. *Kinet. Relat. Models* **6** (2013) 1–135.
- [12] M. Bernkopf, T. Chaumont-Frelet and J. Melenk, Wavenumber-explicit stability and convergence analysis of *hp* finite element discretizations of Helmholtz problems in piecewise smooth media. *Math. Comput.* **94** (2024) 73–122.
- [13] D. Boffi, F. Brezzi and M. Fortin, Mixed Finite Element Methods and Applications. Springer, Heidelberg (2013).
- [14] D.L. Brown, D. Gallistl and D. Peterseim, Multiscale Petrov–Galerkin method for high-frequency heterogeneous Helmholtz equations, in Meshfree Methods for Partial Differential Equations VIII. Springer International Publishing (2017) 85–115..
- [15] E. Cancès, R. Chakir and Y. Maday, Numerical analysis of nonlinear eigenvalue problems. *J. Sci. Comput.* **45** (2010) 90–117.
- [16] C. Carstensen, Quasi-interpolation and a posteriori error analysis in finite element methods. *M2AN Math. Model. Numer. Anal.* **33** (1999) 1187–1202.
- [17] T. Chaumont-Frelet and E.A. Spence, Scattering by finely layered obstacles: frequency-explicit bounds and homogenization. *SIAM J. Math. Anal.* **55** (2023) 1319–1363.
- [18] M. Cicuttin, A. Ern and S. Lemaire, A hybrid high-order method for highly oscillatory elliptic problems. *Comput. Methods Appl. Math.* **19** (2019) 723–748.
- [19] C. Döding, Localized orthogonal decomposition methods vs. classical FEM for the Gross–Pitaevskii equation, in Numerical Mathematics and Advanced Applications ENUMATH 2023. Vol. 153 of *Lect. Notes Comput. Sci. Eng.* Springer, Cham (2025) 268–277.
- [20] Z. Dong, M. Hauck and R. Maier, An improved high-order method for elliptic multiscale problems. *SIAM J. Numer. Anal.* **61** (2023) 1918–1937.
- [21] Y. Du and H. Wu, Preasymptotic error analysis of higher order FEM and CIP-FEM for Helmholtz equation with high wave number. *SIAM J. Numer. Anal.* **53** (2015) 782–804.
- [22] Y. Efendiev, J. Galvis and T.Y. Hou, Generalized multiscale finite element methods (GMsFEM). *J. Comput. Phys.* **251** (2013) 116–135.
- [23] A. Ern and J.-L. Guermond, Theory and Practice of Finite Elements. Vol. 159 of *Applied Mathematical Sciences*. Springer New York (2004).
- [24] A. Ern and J.-L. Guermond, Finite element quasi-interpolation and best approximation. *ESAIM Math. Model. Numer. Anal.* **51** (2017) 1367–1385.
- [25] S. Esterhazy and J.M. Melenk, On Stability of Discretizations of the Helmholtz Equation. Springer Berlin Heidelberg (2011) 285–324.
- [26] P. Freese, M. Hauck, T. Keil and D. Peterseim, A super-localized generalized finite element method. *Numer. Math.* **156** (2024) 205–235.
- [27] P. Freese, M. Hauck and D. Peterseim, Super-localized orthogonal decomposition for high-frequency Helmholtz problems. *SIAM J. Sci. Comput.* **46** (2024) A2377–A2397.
- [28] I.G. Graham and S.A. Sauter, Stability and finite element error analysis for the Helmholtz equation with variable coefficients. *Math. Comput.* **89** (2019) 105–138.
- [29] C. Harder, D. Paredes and F. Valentin, A family of multiscale hybrid-mixed finite element methods for the Darcy equation with rough coefficients. *J. Comput. Phys.* **245** (2013) 107–130.
- [30] M. Hauck and A. Lozinski, A high-order localized orthogonal decomposition method for heterogeneous Stokes problems. Preprint [arXiv:2511.22684](https://arxiv.org/abs/2511.22684) (2025).
- [31] M. Hauck and D. Peterseim, Multi-resolution localized orthogonal decomposition for Helmholtz problems. *Multiscale Model. Simul.* **20** (2022) 657–684.
- [32] M. Hauck and D. Peterseim, Super-localization of elliptic multiscale problems. *Math. Comput.* **92** (2023) 981–1003.
- [33] M. Hauck, A. Målqvist and M. Mosquera, A localized orthogonal decomposition method for heterogeneous mixed-dimensional problems. Preprint [arXiv:2510.09442](https://arxiv.org/abs/2510.09442) (2025).
- [34] P. Henning and E. Jarlebring, The Gross–Pitaevskii Equation and Eigenvector Nonlinearities: Numerical Methods and Algorithms. *SIAM Rev.* **67** (2025) 256–317.
- [35] P. Henning and A. Persson, On optimal convergence rates for discrete minimizers of the Gross–Pitaevskii energy in localized orthogonal decomposition spaces. *Multiscale Model. Simul.* **21** (2023) 993–1011.
- [36] P. Henning and D. Peterseim, Oversampling for the multiscale finite element method. *Multiscale Model. Simul.* **11** (2013) 1149–1175.

- [37] J.S. Hesthaven, S. Zhang and X. Zhu, High-order multiscale finite element method for elliptic problems. *Multiscale Model. Simul.* **12** (2014) 650–666.
- [38] T.Y. Hou and X.-H. Wu, A multiscale finite element method for elliptic problems in composite materials and porous media. *J. Comput. Phys.* **134** (1997) 169–189.
- [39] R. Li, P. Ming and F. Tang, An efficient high order heterogeneous multiscale method for elliptic problems. *Multiscale Model. Simul.* **10** (2012) 259–283.
- [40] C. Ma, R. Scheichl and T. Dodwell, Novel design and analysis of generalized finite element methods based on locally optimal spectral approximations. *SIAM J. Numer. Anal.* **60** (2022) 244–273.
- [41] R. Maier, *Computational multiscale methods in unstructured heterogeneous media*. Ph.D. thesis, University of Augsburg (2020).
- [42] R. Maier, A high-order approach to elliptic multiscale problems with general unstructured coefficients. *SIAM J. Numer. Anal.* **59** (2021) 1067–1089.
- [43] R. Maier and B. Verfürth, Multiscale scattering in nonlinear Kerr-type media. *Math. Comput.* **91** (2022) 1655–1685.
- [44] A. Målqvist and D. Peterseim, Localization of elliptic multiscale problems. *Math. Comput.* **83** (2014) 2583–2603.
- [45] A. Målqvist and D. Peterseim, Numerical Homogenization By Localized Orthogonal Decomposition. Vol. 5 of *SIAM Spotlights*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA (2020).
- [46] J.M. Melenk and S. Sauter, Wavenumber explicit convergence analysis for galerkin discretizations of the Helmholtz equation. *SIAM J. Numer. Anal.* **49** (2011) 1210–1243.
- [47] P. Oswald, On a BPX-preconditioner for P1 elements. *Computing (Wien. Print)* **51** (1993) 125–133.
- [48] H. Owhadi, Multigrid with rough coefficients and multiresolution operator decomposition from hierarchical information games. *SIAM Rev.* **59** (2017) 99–149.
- [49] H. Owhadi and C. Scovel, Operator-adapted Wavelets, Fast Solvers, and Numerical Homogenization. Vol. 35 of *Cambridge Monographs on Applied and Computational Mathematics*. Cambridge University Press, Cambridge (2019).
- [50] H. Owhadi, L. Zhang and L. Berlyand, Polyharmonic homogenization, rough polyharmonic splines and sparse super-localization. *ESAIM Math. Model. Numer. Anal.* **48** (2014) 517–552.
- [51] D. Peterseim, Eliminating the pollution effect in Helmholtz problems by local subscale correction. *Math. Comput.* **86** (2016) 1005–1036.
- [52] D. Peterseim and B. Verfürth, Computational high frequency scattering from high-contrast heterogeneous media. *Math. Comput.* **89** (2020) 2649–2674.
- [53] C. Schwab, Numerical mathematics and scientific computation, in *p*- and *hp*- Finite Element Methods. Clarendon Press, Oxford, England (1998).
- [54] E. Weinan and B. Engquist, The heterogeneous multiscale methods. *Commun. Math. Sci.* **1** (2003) 87–132.
- [55] E. Weinan and B. Engquist, The heterogeneous multi-scale method for homogenization problems, in *Multiscale Methods in Science and Engineering*. Vol. 44 of *Lect. Notes Comput. Sci. Eng.* Springer, Berlin, Heidelberg (2005) 89–110.



Please help to maintain this journal in open access!

This journal is currently published in open access under the Subscribe to Open model (S2O). We are thankful to our subscribers and supporters for making it possible to publish this journal in open access in the current year, free of charge for authors and readers.

Check with your library that it subscribes to the journal, or consider making a personal donation to the S2O programme by contacting subscribers@edpsciences.org.

More information, including a list of supporters and financial transparency reports, is available at <https://edpsciences.org/en/subscribe-to-open-s2o>.