



Measurement and effective field theory interpretation of the photon-fusion production cross section of a pair of W bosons in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration*

Abstract

This analysis presents an observation of the photon-fusion production of W boson pairs using the CMS detector at the LHC. The total cross section of the W^+W^- production in photon fusion is measured using proton-proton collision data with an integrated luminosity of 138 fb^{-1} collected with the CMS detector in 2016–2018 at a center-of-mass energy of $\sqrt{s} = 13$ TeV. Events are selected in the final state with one isolated electron and one isolated muon, and no additional tracks associated with the electron-muon production vertex. The total and fiducial production cross sections are $643^{+82}_{-78} \text{ fb}$ and $3.96^{+0.53}_{-0.51} \text{ fb}$, respectively, in agreement with the standard model predictions of $631 \pm 126 \text{ fb}$ and $3.87 \pm 0.77 \text{ fb}$. This agreement enables stringent constraints on anomalous quartic gauge couplings within a dimension-8 effective field theory framework.

Published in the Journal of High Energy Physics as doi:10.1007/JHEP06(2026)187.

1 Introduction

Photon-induced processes in proton-proton (pp) collisions at the LHC provide important tests of the standard model (SM) of particle physics, which are complementary to those accessible through the more common quark and gluon interactions. In the SM, the photon-fusion production of a pair of W bosons proceeds through interactions that involve trilinear and quartic gauge boson couplings. The measurement of this process can be used to constrain dimension-8 effective field theory (EFT) operators related to anomalous quartic gauge boson interactions.

The signal events considered in this analysis correspond to the process $pp(\gamma\gamma) \rightarrow p^{(*)}W^+W^-p^{(*)}$, where $p^{(*)}$ denotes a final-state proton that either stays intact or dissociates after the photon emission. When both protons coherently radiate a photon and both remain intact after their interaction, the process is called exclusive or elastic. When one or both incoming protons fragment after the photon emission, the process is either semi- or fully dissociative. Semi- and fully dissociative events can also give rise to an elastic-like signature when the proton remnants are not reconstructed because they fall outside of the detector acceptance.

The selection of signal events requires the absence of additional charged particles around the production vertex, as is characteristic of photon-fusion processes. Such a strategy was previously used by the CMS Collaboration to observe the $\gamma\gamma \rightarrow \tau\tau$ process in pp collisions [1] and in earlier searches and measurements at the LHC [2–7]. Evidence for the photon-fusion production of a pair of W bosons was reported by the CMS Collaboration using data collected at center-of-mass energies of 7 and 8 TeV [4], and this process was then observed by the ATLAS Collaboration with data collected at a center-of-mass energy of 13 TeV [7]. Limits were previously imposed on CP-even dimension-8 EFT operators using the $\gamma\gamma \rightarrow W^+W^-$ process by both the ATLAS [6] and CMS [4] Collaborations.

The present analysis relies on pp collision data collected at a center-of-mass energy of 13 TeV with the CMS detector in 2016–2018, corresponding to an integrated luminosity of 138 fb^{-1} . The $\gamma\gamma \rightarrow W^+W^-$ process is measured in the final state with an isolated electron, an isolated muon of opposite charge, and no additional tracks associated with the vertex formed by the electron and the muon. The dominant backgrounds are the inclusive production of W boson pairs from all sources except the signal $\gamma\gamma \rightarrow W^+W^-$, as well as $Z/\gamma^* \rightarrow \tau\tau$ events, which are constrained from data in selected events with a larger number of tracks. The cross section of the $\gamma\gamma \rightarrow W^+W^-$ process is measured both inclusively and in a fiducial phase space matching the analysis selection criteria. Potential beyond-the-SM (BSM) contributions are investigated using an EFT extension of the SM, from which stringent constraints on dimension-8 operators are derived, including for the first time limits on CP-odd dimension-8 operators.

The paper is structured as follows. Sections 2 and 3 describe the detector and the simulation of signal and background processes. The object reconstruction is detailed in Section 4, followed by the event selection in Section 5. The analysis strategy and methods to estimate signal and background processes are presented in Sections 6 and 7. After a discussion of the systematic uncertainties in Section 8, the results and the EFT interpretation are reported in Section 9. The paper is summarized in Section 10. Tabulated results are provided in the HEPData record for this analysis [8].

2 The CMS detector

The CMS apparatus [9, 10] is a multipurpose, nearly hermetic detector, designed to trigger on [11–13] and identify electrons, muons, photons, and (charged and neutral) hadrons [14–16].

Its central feature is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are reconstructed using gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. More detailed descriptions of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Refs. [9, 10].

Events of interest are selected using a two-tiered trigger system. The first level (L1), composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of 4 μ s [11]. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to a few kHz before data storage [12, 13].

3 Signal and background processes

Monte Carlo simulation samples corresponding to the elastic, semi-dissociative, and fully dissociative photon-fusion productions of pairs of τ leptons and W bosons are generated with the SUPERCHIC 4 generator [17, 18]. Elastic, semi-dissociative, and dissociative components are simulated independently and combined together according to the predictions from the SUPERCHIC 4 generator. Semi-dissociative and dissociative components amount to about five times the elastic contribution, for the selection criteria used in the analysis and detailed in Section 5. The $\gamma\gamma \rightarrow W^+W^-$ processes in which the observed electron or muon originates from a leptonic τ decay, with the τ lepton itself produced in a W boson decay, are also simulated and treated as a component of the signal.

The SUPERCHIC generator is used to model the SM prediction of the signal, and the GAMMA-UPC generator [19] is used to predict relative variations from nonzero dimension-8 EFT operators. The GAMMA-UPC generator, which is implemented in the MADGRAPH5_aMC@NLO v3.5.6 code [20, 21], derives photon fluxes from electric dipole and charge form factors for protons and ions, and includes hadronic survival probabilities. In this work, the photon fluxes used are those derived from the elastic charge form factor of the proton, including the effects of the photon virtualities [22], whereas the SUPERCHIC event generator applies form factors based on the ‘structure-function’ approach [23]. The EFT dimension-8 operators are simulated with MADGRAPH5_aMC@NLO at leading order (LO) accuracy using the reweighting technique [24]. We do not include the BSM effects on the background processes since they are expected to have negligible contributions.

The MADGRAPH5_aMC@NLO 2.6.5 event generator is used to generate background events originating from the Drell–Yan (DY) dilepton production process $pp \rightarrow Z/\gamma^*$, as well as the inclusive diboson production, $pp \rightarrow WW$, including the electroweak (EW) WW production in association with two jets, except for processes involving top quarks, which are simulated separately as described below. These processes are simulated at next-to-LO (NLO) accuracy in perturbative quantum chromodynamics (QCD) with up to two outgoing partons at Born level, with different jet multiplicities treated with the FxFx jet matching and merging procedure [21]. The MADGRAPH5_aMC@NLO generator is also used for the simulation of the minor contribution from other diboson production events, whereas POWHEG 2.0 [25–32] is used for $t\bar{t}$ and single top quark production. Contributions from EW WZ and ZZ production in association with two jets, as well as double parton scattering WW are at a percent level and are therefore

neglected.

The generators are interfaced with PYTHIA 8.240 [33] to model parton showering and hadronization, the underlying event activity, as well as the decay of the τ leptons. For the signal samples, the modeling of elastic and proton-dissociated contributions follows the procedure described in Ref. [17]. The PYTHIA generator with CP5 tune [34] is used for the description of the underlying event. The NNPDF3.1 parton distribution function (PDF) set [35] at next-to-NLO accuracy is used for background simulations. Additional pp interactions per bunch crossing, known as pileup (PU), are modeled with PYTHIA and added to the simulated samples with a distribution that is chosen to match that observed in the data. Generated events are processed through a GEANT4 [36] simulation of the CMS detector.

4 Object reconstruction

A particle-flow algorithm [37] is used to reconstruct and identify each individual particle in an event, with an optimized combination of information from the various elements of the CMS detector. The energy of photons is obtained from the ECAL measurement. The energy of electrons is determined from a combination of the electron momentum at the primary interaction vertex as determined by the tracker, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The energy of muons is obtained from the curvature of the corresponding track. The energy of charged hadrons is determined from a combination of their momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, the energy of neutral hadrons is obtained from the corresponding corrected ECAL and HCAL energies.

Electrons are measured in the pseudorapidity range $|\eta| < 2.5$. The momentum resolution for electrons with transverse momentum $p_T \approx 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.6 to 5%. It is generally better in the barrel region than in the endcaps, and also depends on the bremsstrahlung energy emitted by the electron as it traverses the material in front of the ECAL [14, 38]. Electrons are identified with a multivariate discriminant that includes isolation variables, as described in Ref. [14].

Muons are measured in the pseudorapidity range $|\eta| < 2.4$. The efficiency to reconstruct and identify muons is greater than 96%. Matching muons to tracks measured in the silicon tracker results in a relative transverse momentum resolution, for muons with p_T up to 100 GeV, of 1% in the barrel and 3% in the endcaps. The p_T resolution in the barrel is better than 7% for muons with p_T up to 1 TeV [15].

The relative isolation of electrons (muons) is calculated using the p_T of tracks in a cone of $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} < 0.3$ (0.4) centered on the lepton track, divided by the lepton p_T , corrected for charged and neutral pileup contributions [14] ([15]). A threshold of 0.20 is used, although this requirement is relaxed in certain cases to estimate specific background contributions, as detailed later.

The silicon tracker used in 2016 measured charged particles within the range $|\eta| < 2.5$. For nonisolated particles of $1 < p_T < 10$ GeV and $|\eta| < 1.4$, the track resolutions were typically 1.5% in p_T and 25–90 (45–150) μm in the transverse (longitudinal) impact parameter [16]. For tracks with p_T as low as 0.5 GeV, the transverse (longitudinal) impact parameter resolution was around 150 (300) μm . At the start of 2017, a new pixel detector was installed [39]; the upgraded tracker measured particles with typical resolutions of 1.5% in p_T and 20–75 μm in the transverse

impact parameter [40] for nonisolated particles of $1 < p_T < 10$ GeV. The transverse impact parameter resolution was up to about $105 \mu\text{m}$ for $p_T \simeq 0.5$ GeV.

During the 2016 and 2017 data taking, a gradual shift in the timing of the inputs of the ECAL L1 trigger in the region at $|\eta| > 2.0$ caused a specific trigger inefficiency. For events containing an electron (a jet) with p_T larger than ≈ 50 GeV (≈ 100 GeV), in the region $2.5 < |\eta| < 3.0$ the efficiency loss is ≈ 10 – 20% , depending on p_T , η , and time. Correction factors were computed from data and applied to the acceptance estimated with the simulation.

5 Event selection

The signal region (SR) and control region (CR) of the analysis feature an $e\mu$ final state. Events are selected by triggering on the presence of an electron, a muon, or both an electron and a muon. Offline, the leading (trailing) lepton p_T is required to be greater than 24 (15) GeV. The electron (muon) is required to have $|\eta| < 2.5$ (2.4), to pass the identification criteria [14] ([15]) and be isolated. The electron and muon must have opposite-sign (OS) electric charge, be separated by $\Delta R > 0.5$, and have an invariant mass above 20 GeV. The difference $|d_z|(e, \mu)$ between the longitudinal impact parameter of the electron and the muon is required to be less than 0.04 cm. The leptons are selected irrespectively of their $|d_z|$ with respect to the primary vertex. The $e\mu$ vertex is then built from the two leptons, and its z position is calculated as the average z intercepted of the tracks of the two objects. The event is vetoed if there is any additional identified and isolated electron or muon.

The track multiplicity associated with the $e\mu$ vertex, N_{tracks} , is used to separate the signal from the backgrounds. It is calculated considering tracks with $|d_z|$ less than 0.05 cm with respect to the electron-muon vertex, $p_T > 0.5$ GeV, and $|\eta| < 2.5$, and discarding the tracks used to build the selected electron and muon candidates. In the SR, N_{tracks} is required to be zero. As shown in Fig. 1 (left), this requirement keeps about half of the elastic photon-fusion signal events because of pileup track contamination, while reducing quark- and gluon-induced inclusive processes by more than two orders of magnitude. The N_{tracks} variable is poorly modelled by simulations, and data-driven corrections are determined and applied to signal and background samples, as detailed in Section 7.1. In the CR, N_{tracks} is required to be between 1 and 5, which provides a CR enriched in inclusive backgrounds. An upper requirement of $N_{\text{tracks}} = 5$ is imposed to select events with a similar topology as the SR and reduce extrapolation biases from the CR to the SR. The CR contains about 20 times more events than the SR.

In the $\gamma\gamma \rightarrow \tau\tau$ process, the τ leptons and their visible decay products are produced back-to-back [1], while the decay products of the W bosons from the $\gamma\gamma \rightarrow W^+W^-$ process are not. A minimum azimuth angle separation (acoplanarity) between the electron and the muon, $A = 1 - |\Delta\phi|/\pi > 0.015$, is applied to reduce the $\gamma\gamma \rightarrow \tau\tau$ contributions while keeping the signal efficiency higher than 95%, as shown in Fig. 1 (right). This selection criterion also removes a significant fraction of the DY background in the SR.

The $\mu\mu$ final state is used to derive corrections to the simulations based on data, as detailed in Section 7.1. Events in the $\mu\mu$ final state are selected with single-muon triggers. Offline, the leading (subleading) muon must have $p_T > 26$ – 29 (10) GeV depending on the data-taking year, and $|\eta| < 2.4$. The muons must have a transverse impact parameter with respect to the interaction vertex, d_{xy} , less than 0.05 cm, and be separated from each other by at most 0.1 cm in the z direction. The two muons must pass the identification and isolation criteria, and have OS charge and an invariant mass greater than 50 GeV.




Figure 1: Normalized N_{tracks} (left) and acoplanarity (right) distributions for simulated signal and background processes reconstructed in the detector with a PU distribution matching that of the data. The acoplanarity distribution is shown for events with $N_{\text{tracks}} = 0$.

The event selection criteria for the $e\mu$ SR and CR, as well as for the $\mu\mu$ final state, are summarized in Table 1.

Table 1: Selection criteria applied on the $e\mu$ SR [CR], and in the $\mu\mu$ final state. The p_T and pseudorapidity ranges correspond to different sets of triggers in different data-taking periods. The leading (subleading) lepton in the $e\mu$ SR [CR] must have $p_T > 24$ (15) GeV.

	$e\mu$ SR [CR]	$\mu\mu$
p_T^e (GeV)	>24(15)	—
$ \eta^e $	< 2.5	—
$ d_{xy}^e $ (cm)	<0.05 in the barrel <0.1 in the endcaps	—
p_T^μ (GeV)	>24(15)	>26–29(10)
$ \eta^\mu $	<2.4	<2.4
$ d_{xy}^\mu $ (cm)	<0.02	<0.05
$m_{\ell\ell}$ (GeV)	>20	>50
OS	Yes	Yes
$ dz(\ell, \ell) $ (cm)	<0.04	<0.1
ΔR	>0.5	>0.5
A	>0.015	—
N_{tracks}	0 [1–5]	—

6 Analysis strategy

The results of the analysis are extracted with a binned maximum likelihood fit of the $p_T^{e\mu}$ distribution in the SR and the CR, where $p_T^{e\mu}$ is defined as the magnitude of the vectorial p_T sum of the selected electron and muon. The observable $p_T^{e\mu}$ separates the processes with missing transverse momentum (p_T^{miss}), such as the signal and the inclusive WW production from processes with low p_T^{miss} , such as the DY background, which dominate the low- $p_T^{e\mu}$ bins, and the non-prompt background from multijet production. Systematic uncertainties are relatively larger for

p_T^{miss} than $p_T^{e\mu}$, which explains the choice of observable.

In the maximum likelihood fits to extract the total and fiducial cross sections, the parameter of interest is the signal strength, while three other parameters are left free to float:

1. the DY background normalization, correlated between the SR and the CR;
2. the inclusive WW normalization, correlated between the SR and the CR;
3. the event weight correction for $N_{\text{tracks}} = 0$ (described in Section 7.1), correlated between the DY and the inclusive WW backgrounds.

The DY and inclusive WW background normalizations are mostly constrained from the large CR sample, while the N_{tracks} correction gets constrained by the DY background in the low- $p_T^{e\mu}$ bins. In the likelihood scans to extract constraints on EFT operators, the SM $\gamma\gamma \rightarrow W^+W^-$ signal is constrained within uncertainties to its theoretical prediction and one EFT coefficient at a time, affecting the $\gamma\gamma \rightarrow W^+W^-$ process, is treated as a freely floating parameter of interest.

7 Background and signal estimation

7.1 Corrections to the simulated track multiplicity

The pileup multiplicity in simulation is reweighted to match that observed in the data. After this correction, the simulations still do not describe accurately the N_{tracks} distribution and need to be corrected further. Corrections to the multiplicity of tracks originating from the pileup ($N_{\text{tracks}}^{\text{PU}}$) and from the hard scattering ($N_{\text{tracks}}^{\text{HS}}$) vertices are derived as described in Ref. [1], using events in the $\mu\mu$ final state.

The $N_{\text{tracks}}^{\text{PU}}$ correction is obtained as an event weight by comparing the track multiplicity in data and in simulation inside 1-mm wide windows sampled along the z axis and at least 1 cm away from the dimuon vertex. It is applied to all simulated background and signal events. The $N_{\text{tracks}}^{\text{HS}}$ correction is applied as an event weight derived in a similar way, except the selected window is centered on the dimuon interaction vertex. This window includes both pileup and hard scattering tracks, and the pileup track multiplicity correction is applied first. This correction is measured for events with $A > 0.015$, and applied to both DY and inclusive diboson events, which are modeled with the same generator settings. The $N_{\text{tracks}}^{\text{HS}}$ correction reduces the event yields of the corresponding processes by about 40% for events with $N_{\text{tracks}} = 0$.

7.2 Estimation of backgrounds with prompt leptons

The DY background with prompt leptons is estimated from simulation, using the corrections described in Section 7.1. The inclusive quark-induced $pp \rightarrow WW (2\ell 2\nu)$ background has a similar $p_T^{e\mu}$ distribution as the signal, but, like the DY background, it is strongly suppressed by the N_{tracks} requirement. The contribution from this process is estimated from simulation with the corrections described in Section 7.1. The same $N_{\text{tracks}}^{\text{HS}}$ correction as that measured for the DY process is used, since the DY and diboson backgrounds are both simulated with the same generators and settings, and their $p_T(\tau\tau/WW)$ distributions are similar after selection. To make the analysis less dependent on the correction to $N_{\text{tracks}}^{\text{HS}}$ and its uncertainty, which is difficult to evaluate because of the generator dependence, the $N_{\text{tracks}}^{\text{HS}}$ correction to events with $N_{\text{tracks}} = 0$ is left as a free parameter in the maximum likelihood fit described in Section 6.

Other backgrounds, such as $\gamma\gamma \rightarrow \tau\tau$, minor diboson processes (WZ, ZZ), single top quark, and $t\bar{t}$ production, are estimated from simulation, with corrections to the simulated N_{tracks} .

For the $\gamma\gamma \rightarrow \tau\tau$ process, which was measured directly in Ref. [1], the simulated prediction (SUPERCHIC 4) reproduces the measured cross section and kinematic distributions within the experimental and modeling uncertainties. Corrections to $N_{\text{tracks}}^{\text{HS}}$ are not used for the single top quark and $t\bar{t}$ processes because these processes do not use the same generator. They are assigned large uncertainties (50%) to cover for $N_{\text{tracks}}^{\text{HS}}$ mismodelings, which have a negligible effect on the total background prediction.

7.3 Estimation of backgrounds with nonprompt leptons

The background with jets misidentified as electrons or muons, essentially coming from $W + \text{jets}$ and QCD multijet events, is estimated from observed events. Events where the electron and muon have same-sign (SS) electric charge are reweighted with an OS-to-SS scale factor (SF), denoted $\text{OSSS}(\text{id. e, iso. } \mu)$. Most events in the SS region correspond to the jet misidentification (mis-ID) background, but small contributions from other processes with genuine leptons are estimated from simulation and subtracted from the data.

The OS-to-SS SF depends on both the muon and electron p_T and is measured in a background-enriched region where the muon relative isolation is inverted (between 0.2 and 0.5 instead of <0.2 in the SR and CR) as:

$$\text{OSSS}(\text{id. e, anti-iso. } \mu) = \frac{N^{\text{OS}}(\text{id. e, anti-iso. } \mu)}{N^{\text{SS}}(\text{id. e, anti-iso. } \mu)} \quad (1)$$

where $N^{\text{OS}}(\text{id. e, anti-iso. } \mu)$ and $N^{\text{SS}}(\text{id. e, anti-iso. } \mu)$ are the number of OS and SS events with a well-identified electron and an anti-isolated muon. Inverting the muon isolation biases the OS-to-SS SFs, and a multiplicative correction to this SF is derived using events with an inverted electron identification. The $\text{OSSS}(\text{id. e, iso. } \mu)$ term is computed as:

$$\text{OSSS}(\text{id. e, iso. } \mu) = \text{OSSS}(\text{id. e, anti-iso. } \mu) \frac{\text{OSSS}(\text{anti-id. e, iso. } \mu)}{\text{OSSS}(\text{anti-id. e, anti-iso. } \mu)}. \quad (2)$$

In this expression, all the OSSS terms depend on the electron and muon p_T . They were verified not to depend on N_{tracks} .

To improve the statistical precision of the mis-ID background distribution, its $p_T^{e\mu}$ distribution is taken from events with $N_{\text{tracks}} < 10$, while the normalization is taken from the default estimation without relaxing the N_{tracks} selection. The compatibility of the $p_T^{e\mu}$ distributions in the SR with those in the region where events have $N_{\text{tracks}} < 10$ was confirmed.

8 Systematic uncertainties

Systematic uncertainties are considered as nuisance parameters in the statistical procedure to extract the signal cross section. They are treated with either Gaussian (shape uncertainties) or log-normal (normalization uncertainties) function priors included in the likelihood function [41]. Unless specified, systematic uncertainties are treated as shape uncertainties affecting the $p_T^{e\mu}$ distributions. Given that the cross section normalizations of the DY and diboson processes are both left floating in the maximum likelihood fit, overall normalization uncertainties are not considered for these processes. The sources of systematic uncertainties are summarized in Table 2.

The uncertainties in the electron and muon identification, isolation, and triggering efficiencies are up to 2% per lepton. A multiplicative correction to the identification and isolation SFs

Table 2: Systematic uncertainties and their relative effects on the inclusive cross section measurement. An uncertainty of 2% in the signal cross section is considered in the EFT interpretation.

Sources of uncertainties	Cross section uncertainty (%)
$N_{\text{tracks}}^{\text{HS}}$ correction	3.6
DY normalization	3.5
e selection	2.2
$N_{\text{tracks}}^{\text{PU}}$ correction	2.0
Integrated luminosity	1.6
$t\bar{t}$ normalization	1.2
Pileup modeling	1.2
Inclusive diboson normalization	1.0
μ selection	1.0
Jet mis-ID prediction	<1
$\mu_{\text{R}}, \mu_{\text{F}}, \text{PDF}$	<1

to account for the low- N_{tracks} environment is applied to electrons; its magnitude is 0.98. It is derived as the ratio of the SFs measured in $Z \rightarrow ee$ events with $N_{\text{tracks}} < 10$ and inclusively in N_{tracks} . Scale factors for muons do not need to be adjusted for the low- N_{tracks} environment.

The uncertainty in the $N_{\text{tracks}}^{\text{PU}}$ correction amounts to 2%, whereas no uncertainty in the $N_{\text{tracks}}^{\text{HS}}$ correction is provided since it is freely floating in the SR with $N_{\text{tracks}} = 0$. Theoretical uncertainties arising from the choice of renormalization and factorization scales (μ_{R} and μ_{F}) used in the calculations, as well as from the PDF choice, are also taken into account and propagated to the $p_{\text{T}}^{e\mu}$ distributions.

Statistical uncertainties in the two-dimensional p_{T} -dependent OS-to-SS scaling factors used to evaluate the jet mis-ID background are included. An additional 10% yield uncertainty is included to cover for a potential difference in the background composition in the SR and CR. It is determined by comparing OS-to-SS scaling factors measured in events with high or low $p_{\text{T}}^{\text{miss}}$.

The uncertainty in the cross section of the photon-fusion processes modeled with the SUPERCHIC 4 generator is obtained by varying the rescattering models of the proton survival factors. Following the recommendations of Ref. [19], conservative uncertainties of 1% for the elastic component, 10% for semi-dissociative, and 50% for fully dissociative production are assigned. Since the elastic, semi-dissociative, and fully dissociative components have similar $p_{\text{T}}^{e\mu}$ distributions, these cross section uncertainties are applied only in the determination of the constraints on EFT operators, presented in Section 9.2, and not in the cross section measurements.

Statistical uncertainties in the number of simulated background events selected in the SR and CR or observed event yields in the regions used to derive the jet mis-ID background are considered in all bins of the distributions using the approach from Ref. [42, 43]. The systematic uncertainty in the pileup modeling is included by varying by $\pm 4.6\%$ the total inelastic cross section used to calculate simulated pileup distributions [44]. The integrated luminosities for the 2016, 2017, and 2018 data-taking years have 1.2–2.5% individual uncertainties [45–47], whereas the overall uncertainty is 1.6%.




Figure 2: Observed and predicted $p_T^{e\mu}$ distributions for events with $N_{\text{tracks}} = 0$ (SR, left) or $1 \leq N_{\text{tracks}} \leq 5$ (CR, right), using 2016–2018 data. The distributions are shown after the maximum likelihood fit to the data (“postfit distributions”). The observed data and their associated Poissonian statistical uncertainty are shown with black markers with vertical error bars. The uncertainty band accounts for all sources of background and signal uncertainty, systematic as well as statistical, after the fit. The last bin includes the overflow. The lower panels show the ratio of data to sum of signal and background contributions, before (prefit, open red circles) and after (black full markers) the maximum likelihood fit.

9 Results

9.1 Cross section measurement

The inclusive and fiducial cross sections of the $\gamma\gamma \rightarrow W^+W^-$ process are extracted with the binned maximum likelihood fit described in Section 6, using the $p_T^{e\mu}$ distributions in the SR and CR shown in Fig. 2. In the maximum likelihood fit, the $N_{\text{tracks}}^{\text{HS}}$ correction is pulled down to 0.91 ± 0.06 times its initial value of 0.60. The experimental data agree well with the sum of signal and background contributions for all $p_T^{e\mu}$ regions. In the SR, the signal contribution dominates for $p_T^{e\mu} > 80$ GeV. The results presented hereafter have been determined from the measured $p_T^{e\mu}$ distributions using the CMS statistical analysis tool COMBINE [48], which is based on the ROOFIT [49] and ROOSTATS [50] frameworks. The sources of systematic uncertainties described in Section 8 are included in the statistical procedure as nuisance parameters.

The significance of the signal with respect to the background-only expectation is well above 5 standard deviations, and this process is observed for the first time using data collected with the CMS detector. The inclusive cross section for the $\gamma\gamma \rightarrow W^+W^-$ process is measured to be 643_{-78}^{+82} fb, whereas the value predicted by the SUPERCHIC 4 generator is 631 ± 126 fb [18]. Figure 3 shows how the predicted $\gamma\gamma \rightarrow W^+W^-$ signal matches the difference between the observed events and the simulated background events in the $p_T^{e\mu}$ distributions.

The cross section of the $\gamma\gamma \rightarrow W^+W^-$ process is also measured in a fiducial phase space matching the analysis selection criteria to the reconstructed event selection. All variables used in the definition of the fiducial region are calculated at the generator level after parton showering and hadronization, and the lepton momentum includes the momenta of photons radiated within a cone of $\Delta R < 0.1$ centered on the lepton. The fiducial region comprises events where the two W bosons decay leptonically in the final state of an electron and a muon of OS charge. Events




Figure 3: Distribution of $p_T^{e\mu}$ for the difference between the observed number of events and the expected number of background events (black markers) compared with the SM prediction $\gamma\gamma \rightarrow W^+W^-$ (red line), and $\gamma\gamma \rightarrow W^+W^-$ predictions for five benchmark points with nonzero EFT coefficients. The Poissonian statistical uncertainties associated with the observed data are represented by vertical error bars. The difference between the predictions for the SM and nonzero EFT coefficients is visible only in the overflow bin. The background uncertainty is shown with a gray band.

Table 3: Observed and predicted event yields in the SR and CR. The signal and background yields are the result of the global fit including all sources of uncertainties.

Process	$N_{\text{tracks}} = 0$ (SR)	$1 \leq N_{\text{tracks}} \leq 5$ (CR)
$Z/\gamma^* \rightarrow \tau\tau$	361 ± 22	$11\,260 \pm 170$
Jet mis-ID	119 ± 10	$2\,160 \pm 180$
Inclusive VV	117 ± 10	$3\,770 \pm 220$
$\gamma\gamma \rightarrow \tau\tau$	34.2 ± 1.5	37.3 ± 1.5
Top	6.6 ± 1.4	673 ± 98
EW WW +2 jets	3.6 ± 0.7	28.9 ± 4.9
Nonfiducial $\gamma\gamma \rightarrow WW$	11.4 ± 0.4	70.4 ± 2.4
Total bkg.	653 ± 30	$18\,000 \pm 160$
Fiducial $\gamma\gamma \rightarrow WW$	183 ± 22	121 ± 15
Total	835 ± 29	$18\,120 \pm 170$
Observed	829	18 112

in which the electron or muon is produced via a leptonic τ decay with the τ lepton originating from a W boson decay are also part of the signal within the fiducial region. The electron and muon at the generator level are required to have an acoplanarity $A > 0.015$ and an invariant mass > 20 GeV. There must not be any final-state charged particle with $c\tau > 1$ cm, $p_T > 0.5$ GeV, and $|\eta| < 2.5$, other than the two charged leptons themselves. The leading lepton must satisfy $p_T > 24$ GeV, whereas the requirement for the subleading lepton is $p_T > 15$ GeV. The electron (muon) must have $|\eta| < 2.5$ (2.4). Photon-induced diboson events not belonging to the fiducial region but entering the reconstruction-level categories are constrained to their expected normalizations and distributions, within uncertainties. In the SR, the nonfiducial fraction of the $\gamma\gamma \rightarrow W^+W^-$ process is about 6%. The measured fiducial cross section is $\sigma^{\text{fid}} = 3.96^{+0.53}_{-0.51}$ fb, whereas the value predicted by the SUPERCHIC 4 generator is 3.87 ± 0.77 fb [18]. The event yields after the maximum likelihood fit are shown in Table 3.

9.2 Effective field theory interpretation

In the SM, the photon-fusion production of a pair of W bosons arises from self-interactions of electroweak gauge bosons, in particular from the triple gauge coupling (TGC) γWW and the quartic gauge coupling (QGC) $\gamma\gamma WW$. Whereas the individual diagrams contributing to this process exhibit amplitudes that grow with energy, their combined SM contributions interfere destructively and preserve unitarity. This delicate balance between TGC and QGC terms can be easily disrupted by the presence of BSM physics.

Whereas constraints on γWW anomalous TGCs are typically derived from the inclusive production of $W\gamma$, WW , and electroweak production of $W + \text{jets}$ [51–53], the $\gamma\gamma \rightarrow W^+W^-$ process is primarily sensitive to anomalous QGCs and thus probes various dimension-8 operators within the effective field theory framework.

The results are interpreted with an EFT approach as constraints on the dimension-8 operators described in Ref. [54]. The selected operators are bi-quadratic in the Higgs and gauge field strengths (denoted f_M) or quartic in the gauge field strength (denoted f_T). The CP-odd operators are denoted $f_{\tilde{M}}$ and $f_{\tilde{T}}$. For CP-odd operators, the interference with the SM amplitude cancels since the analysis observables are not CP-sensitive. The results are therefore driven by the quadratic EFT contribution.

We use the elastic GAMMA-UPC simulations to compute, in each $p_T^{e\mu}$ bin, how each EFT opera-

Table 4: The 95% CL intervals on EFT operators varied one at a time while fixing the other ones to zero. Odd operators are denoted with a tilde sign.

	Expected 95% CL (TeV^{-4})	Observed 95% CL (TeV^{-4})
f_{M0}/Λ^4	[−4.03, 4.10]	[−3.24, 3.25]
f_{M1}/Λ^4	[−15.46, 14.93]	[−12.31, 12.03]
$f_{\widetilde{M1}}/\Lambda^4$	[−0.62, 0.62]	[−0.50, 0.49]
f_{M2}/Λ^4	[−0.61, 0.63]	[−0.49, 0.50]
$f_{\widetilde{M2}}/\Lambda^4$	[−2.24, 2.24]	[−1.80, 1.79]
f_{M3}/Λ^4	[−2.36, 2.28]	[−1.88, 1.84]
f_{M4}/Λ^4	[−2.22, 2.26]	[−1.79, 1.80]
$f_{\widetilde{M4}}/\Lambda^4$	[−4.49, 4.49]	[−3.59, 3.58]
f_{M5}/Λ^4	[−4.12, 4.27]	[−3.32, 3.40]
$f_{\widetilde{M5}}/\Lambda^4$	[−2.03, 2.03]	[−1.63, 1.62]
f_{M7}/Λ^4	[−29.86, 30.92]	[−24.06, 24.62]
f_{T0}/Λ^4	[−0.65, 0.53]	[−0.53, 0.42]
f_{T1}/Λ^4	[−2.13, 2.00]	[−1.71, 1.59]
f_{T2}/Λ^4	[−2.50, 2.01]	[−2.04, 1.56]
$f_{\widetilde{T2}}/\Lambda^4$	[−0.090, 0.090]	[−0.03, 0.03]
f_{T3}/Λ^4	[−3.28, 2.49]	[−2.69, 1.90]
$f_{\widetilde{T3}}/\Lambda^4$	[−0.089, 0.089]	[−0.071, 0.071]
f_{T4}/Λ^4	[−1.00, 0.76]	[−0.82, 0.58]
$f_{\widetilde{T4}}/\Lambda^4$	[−0.38, 0.38]	[−0.30, 0.30]
f_{T5}/Λ^4	[−0.20, 0.16]	[−0.16, 0.13]
$f_{\widetilde{T5}}/\Lambda^4$	[−0.21, 0.21]	[−0.17, 0.17]
f_{T6}/Λ^4	[−0.65, 0.61]	[−0.52, 0.48]
$f_{\widetilde{T6}}/\Lambda^4$	[−0.62, 0.62]	[−0.50, 0.50]
f_{T7}/Λ^4	[−0.76, 0.61]	[−0.62, 0.47]

tor changes the signal prediction. These relative shifts are applied as event weights to the elastic, semi-dissociative, and dissociative components of the signal modeled by the SUPERCHIC 4 generator, assuming the identical EFT effects for the nonelastic components. As a cross-check of this assumption, the EFT weights were compared using two different photon flux models: the elastic photon approximation and an inclusive photon PDF. Despite the different photon spectra predicted by these models, no significant difference in the EFT weights was observed as a function of $p_T^{e\mu}$. It should be noted that, when going beyond the collinear photon approximation, additional diagrams contribute to the semi- and fully dissociative productions, which do not strictly correspond to the $\gamma\gamma \rightarrow W^+W^-$ topology, as discussed in Ref. [18]. The quantitative impact of EFT corrections on these contributions has not been studied yet and, in the absence of such studies, the EFT effects are assumed in this analysis to be identical for the elastic and dissociative components. The impact is largest at high $p_T^{e\mu}$, as shown in Fig. 3.

The sensitivity of the measurement to the EFT operators, considered one at a time, is extracted from a likelihood scan performed by varying the corresponding EFT coefficient. The 95% confidence-level (CL) intervals on the EFT coefficients are extracted from the scan and reported in Table 4. Constraints on a series of operators are competitive with other measurements by the CMS Collaboration in different final states [55–60]. Figures 4 and 5 show a comparison of the observed limits with those from other measurements.




Figure 4: Summary of observed limits on the f_M Wilson coefficients for this analysis and previous results by the CMS Collaboration [55, 57–60]. The limits are arranged from most sensitive to least sensitive in descending order, with the current result highlighted in red. Horizontal error bars represent the 95% confidence level intervals. In some cases, limits on the Wilson coefficients are shown under the assumption of unitarity bounds, as indicated in the right-most column. The unitarity bound is defined as the scattering energy at which the aQGC coupling strength is set equal to the observed limit that would result in a scattering amplitude that violates unitarity [61].




Figure 5: Summary of observed limits on the f_T Wilson coefficients for this analysis and previous results by the CMS Collaboration [55–58, 60]. The limits are arranged from most sensitive to least sensitive in descending order, with the current result highlighted in red. Horizontal error bars represent the 95% confidence level intervals. In some cases, limits on the Wilson coefficients are shown under the assumption of unitarity bounds, as indicated in the right-most column. The unitarity bound is defined as the scattering energy at which the aQGC coupling strength is set equal to the observed limit that would result in a scattering amplitude that violates unitarity [61].

10 Summary

The observation of the photon-fusion production of W boson pairs using the CMS detector at the LHC is reported. The measurement utilizes proton-proton collision data collected in 2016–2018, corresponding to an integrated luminosity of 138 fb^{-1} at $\sqrt{s} = 13 \text{ TeV}$. The experimental strategy exploits the exclusive nature of the signal, requiring the absence of additional charged particles associated with the dilepton production vertex. Events are selected in the final state with one isolated electron and one isolated muon, and no additional tracks associated with the electron-muon production vertex. The analysis employs a maximum likelihood fit to the transverse momentum sum of the electron and muon to extract the cross section of the signal. The measured inclusive cross section for $\gamma\gamma \rightarrow W^+W^-$ production is $643_{-78}^{+82} \text{ fb}$, whereas the fiducial cross section, defined in a phase space matching the experimental selection criteria, is $3.96_{-0.51}^{+0.53} \text{ fb}$. These results show excellent agreement with the standard model predictions of $631 \pm 126 \text{ fb}$ and $3.87 \pm 0.77 \text{ fb}$ for the inclusive and fiducial cross sections, respectively. Beyond the cross section measurement, this analysis provides the most stringent constraints to date on several anomalous quartic gauge couplings using dimension-8 operators in an effective field theory approach.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMFTR, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES, NSC, and NAWA (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA) and Fonds voor Wetenschappelijk Onderzoek contract No. 1228724N (Belgium); the Beijing Municipal Science & Technology Commission, No.

Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, the Natural Science Foundation of China under Grant No. 12535004, and USTC Research Funds of the Double First-Class Initiative No. YD2030002017 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2025-1.1.5-NEMZ.KI-2025-00004 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing, FAIR – Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369, 2021/43/B/ST2/01552, 2023/49/B/ST2/03273, and the NAWA contract BPN/PPO/2021/1/00011 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR”, and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, the National Science, Research and Innovation Fund program IND_FF.68.369.2300.097, and the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

Data availability

Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the CMS data preservation, re-use and open access policy.

References

- [1] CMS Collaboration, “Observation of $\gamma\gamma \rightarrow \tau\tau$ in proton-proton collisions and limits on the anomalous electromagnetic moments of the τ lepton”, *Rept. Prog. Phys.* **87** (2024) 107801, doi:10.1088/1361-6633/ad6fcb, arXiv:2406.03975.
- [2] CMS Collaboration, “Exclusive $\gamma\gamma \rightarrow \mu^+\mu^-$ production in proton-proton collisions at $\sqrt{s} = 7$ TeV”, *JHEP* **01** (2012) 052, doi:10.1007/JHEP01(2012)052, arXiv:1111.5536.
- [3] CMS Collaboration, “Study of exclusive two-photon production of W^+W^- in pp collisions at $\sqrt{s} = 7$ TeV and constraints on anomalous quartic gauge couplings”, *JHEP* **07** (2013) 116, doi:10.1007/JHEP07(2013)116, arXiv:1305.5596.
- [4] CMS Collaboration, “Evidence for exclusive $\gamma\gamma \rightarrow W^+W^-$ production and constraints on anomalous quartic gauge couplings in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **08** (2016) 119, doi:10.1007/JHEP08(2016)119, arXiv:1604.04464.

- [5] CMS, TOTEM Collaboration, “Observation of proton-tagged, central (semi)exclusive production of high-mass lepton pairs in pp collisions at 13 TeV with the CMS-TOTEM precision proton spectrometer”, *JHEP* **07** (2018) 153, doi:10.1007/JHEP07(2018)153, arXiv:1803.04496.
- [6] ATLAS Collaboration, “Measurement of exclusive $\gamma\gamma \rightarrow W^+W^-$ production and search for exclusive Higgs boson production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, *Phys. Rev. D* **94** (2016) 032011, doi:10.1103/PhysRevD.94.032011, arXiv:1607.03745.
- [7] ATLAS Collaboration, “Observation of photon-induced W^+W^- production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Lett. B* **816** (2021) 136190, doi:10.1016/j.physletb.2021.136190, arXiv:2010.04019.
- [8] HEPData record for this analysis, 2025. doi:10.17182/hepdata.167352.
- [9] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [10] CMS Collaboration, “Development of the CMS detector for the CERN LHC Run 3”, *JINST* **19** (2024) P05064, doi:10.1088/1748-0221/19/05/P05064, arXiv:2309.05466.
- [11] CMS Collaboration, “Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **15** (2020) P10017, doi:10.1088/1748-0221/15/10/P10017, arXiv:2006.10165.
- [12] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [13] CMS Collaboration, “Performance of the CMS high-level trigger during LHC Run 2”, *JINST* **19** (2024) P11021, doi:10.1088/1748-0221/19/11/P11021, arXiv:2410.17038.
- [14] CMS Collaboration, “Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC”, *JINST* **16** (2021) P05014, doi:10.1088/1748-0221/16/05/P05014, arXiv:2012.06888.
- [15] CMS Collaboration, “Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JINST* **13** (2018) P06015, doi:10.1088/1748-0221/13/06/P06015, arXiv:1804.04528.
- [16] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014) P10009, doi:10.1088/1748-0221/9/10/P10009, arXiv:1405.6569.
- [17] L. A. Harland-Lang, M. Tasevsky, V. A. Khoze, and M. G. Ryskin, “A new approach to modelling elastic and inelastic photon-initiated production at the LHC: SuperChic 4”, *Eur. Phys. J. C* **80** (2020) 925, doi:10.1140/epjc/s10052-020-08455-0, arXiv:2007.12704.
- [18] S. Bailey and L. A. Harland-Lang, “Modeling W^+W^- production with rapidity gaps at the LHC”, *Phys. Rev. D* **105** (2022) 093010, doi:10.1103/PhysRevD.105.093010, arXiv:2201.08403.

-
- [19] H.-S. Shao and D. d’Enterra, “gamma-UPC: automated generation of exclusive photon-photon processes in ultraperipheral proton and nuclear collisions with varying form factors”, *JHEP* **09** (2022) 248, doi:10.1007/JHEP09(2022)248, arXiv:2207.03012.
- [20] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.
- [21] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO”, *JHEP* **12** (2012) 061, doi:10.1007/JHEP12(2012)061, arXiv:1209.6215.
- [22] H.-S. Shao and D. d’Enterra, “Dimuon and ditau production in photon-photon collisions at next-to-leading order in QED”, *JHEP* **02** (2025) 023, doi:10.1007/JHEP02(2025)023, arXiv:2407.13610.
- [23] T. Han, G. Valencia, and S. Willenbrock, “Structure function approach to vector boson scattering in p p collisions”, *Phys. Rev. Lett.* **69** (1992) 3274, doi:10.1103/PhysRevLett.69.3274, arXiv:hep-ph/9206246.
- [24] O. Mattelaer, “On the maximal use of Monte Carlo samples: re-weighting events at NLO accuracy”, *Eur. Phys. J. C* **76** (2016) 674, doi:10.1140/epjc/s10052-016-4533-7, arXiv:1607.00763.
- [25] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [26] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [27] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [28] S. Alioli et al., “Jet pair production in POWHEG”, *JHEP* **04** (2011) 081, doi:10.1007/JHEP04(2011)081, arXiv:1012.3380.
- [29] S. Alioli, P. Nason, C. Oleari, and E. Re, “NLO Higgs boson production via gluon fusion matched with shower in POWHEG”, *JHEP* **04** (2009) 002, doi:10.1088/1126-6708/2009/04/002, arXiv:0812.0578.
- [30] S. Frixione, G. Ridolfi, and P. Nason, “A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction”, *JHEP* **09** (2007) 126, doi:10.1088/1126-6708/2007/09/126, arXiv:0707.3088.
- [31] E. Re, “Single-top Wt-channel production matched with parton showers using the POWHEG method”, *Eur. Phys. J. C* **71** (2011) 1547, doi:10.1140/epjc/s10052-011-1547-z, arXiv:1009.2450.
- [32] R. Frederix, E. Re, and P. Torrielli, “Single-top t-channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO”, *JHEP* **09** (2012) 130, doi:10.1007/JHEP09(2012)130, arXiv:1207.5391.



- [33] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
- [34] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [35] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
- [36] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [37] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965. <https://cds.cern.ch/record/2270046>.
- [38] CMS Collaboration, “ECAL 2016 refined calibration and Run2 summary plots”, CMS Detector Performance Summary CMS-DP-2020-021, 2020. <https://cds.cern.ch/record/2717925>.
- [39] CMS Tracker Group Collaboration, “The CMS phase-1 pixel detector upgrade”, *JINST* **16** (2021) P02027, doi:10.1088/1748-0221/16/02/P02027, arXiv:2012.14304.
- [40] CMS Collaboration, “Track impact parameter resolution for the full pseudo rapidity coverage in the 2017 dataset with the CMS phase-1 pixel detector”, CMS Detector Performance Summary CMS-DP-2020-049, 2020. <https://cds.cern.ch/record/2743740>.
- [41] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].
- [42] R. Barlow and C. Beeston, “Fitting using finite Monte Carlo samples”, *Comput. Phys. Commun.* **77** (1993), no. 2, 219, doi:10.1016/0010-4655(93)90005-w. <https://cds.cern.ch/record/249779>.
- [43] J. S. Conway, “Incorporating nuisance parameters in likelihoods for multisource spectra”, in *PHYSTAT 2011*, p. 115. 2011. arXiv:1103.0354. <https://cds.cern.ch/record/1333496>. doi:10.5170/CERN-2011-006.115.
- [44] CMS Collaboration, “Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2018) 161, doi:10.1007/JHEP07(2018)161, arXiv:1802.02613.
- [45] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [46] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-17-004, 2018. <https://cds.cern.ch/record/2621960/>.

- [47] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV”, CMS Physics Analysis Summary CMS-PAS-LUM-18-002, 2019.
<https://cds.cern.ch/record/2676164/>.
- [48] CMS Collaboration, “The CMS statistical analysis and combination tool: Combine”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
- [49] W. Verkerke and D. P. Kirkby, “The RooFit toolkit for data modeling”, *eConf* (2003) MOLT007, doi:10.1142/9781860948985_0039, arXiv:physics/0306116. C0303241.
- [50] L. Moneta et al., “The RooStats project”, *PoS ACAT2010* (2010) 057, doi:10.22323/1.093.0057, arXiv:1009.1003.
- [51] CMS Collaboration, “Measurement of $W^{\pm}\gamma$ differential cross sections in proton-proton collisions at $\sqrt{s} = 13$ TeV and effective field theory constraints”, *Phys. Rev. D* **105** (2022) 052003, doi:10.1103/PhysRevD.105.052003, arXiv:2111.13948.
- [52] CMS Collaboration, “ W^+W^- boson pair production in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **102** (2020) 092001, doi:10.1103/PhysRevD.102.092001, arXiv:2009.00119.
- [53] CMS Collaboration, “Measurement of electroweak production of a W boson in association with two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Eur. Phys. J. C* **80** (2020) 43, doi:10.1140/epjc/s10052-019-7585-7, arXiv:1903.04040.
- [54] G. Durieux et al., “LHC EFT WG note: Basis for anomalous quartic gauge couplings”, *SciPost Phys. Comm. Rep.* (2025) 6, doi:10.21468/SciPostPhysCommRep.6, arXiv:2411.02483.
- [55] CMS Collaboration, “Measurements of production cross sections of WZ and same-sign WW boson pairs in association with two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Lett. B* **809** (2020) 135710, doi:10.1016/j.physletb.2020.135710, arXiv:2005.01173.
- [56] CMS Collaboration, “Measurements of the $pp \rightarrow W^{\pm}\gamma\gamma$ and $pp \rightarrow Z\gamma\gamma$ cross sections at $\sqrt{s} = 13$ TeV and limits on anomalous quartic gauge couplings”, *JHEP* **10** (2021) 174, doi:10.1007/JHEP10(2021)174, arXiv:2105.12780.
- [57] CMS Collaboration, “Measurement of the electroweak production of $Z\gamma$ and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV and constraints on anomalous quartic gauge couplings”, *Phys. Rev. D* **104** (2021) 072001, doi:10.1103/PhysRevD.104.072001, arXiv:2106.11082.
- [58] CMS Collaboration, “Measurement of the electroweak production of $W\gamma$ in association with two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *Phys. Rev. D* **108** (2023) 032017, doi:10.1103/PhysRevD.108.032017, arXiv:2212.12592.
- [59] CMS, TOTEM Collaboration, “Search for high-mass exclusive $\gamma\gamma \rightarrow WW$ and $\gamma\gamma \rightarrow ZZ$ production in proton-proton collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **07** (2023) 229, doi:10.1007/JHEP07(2023)229, arXiv:2211.16320.
















-
- [60] CMS Collaboration, “Vector boson scattering and anomalous quartic couplings in final states with $\ell\nu q\bar{q}$ or $\ell\ell q\bar{q}$ plus jets using proton-proton collisions at $\sqrt{s} = 13$ TeV”, 2025. [arXiv:2510.00118](#). Submitted to JHEP.
- [61] E. d. S. Almeida, O. J. P. Éboli, and M. C. Gonzalez–Garcia, “Unitarity constraints on anomalous quartic couplings”, *Phys. Rev. D* **101** (2020), no. 11, [doi:10.1103/physrevd.101.113003](#), [arXiv:2004.05174](#).

A The CMS Collaboration




Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, V. Makarenko , A. Tumasyan¹ 

















Institut für Hochenergiephysik, Vienna, Austria

W. Adam , L. Benato , T. Bergauer , M. Dragicevic , C. Giordano, P.S. Hussain , M. Jeitler² , N. Krammer , A. Li , D. Liko , M. Matthewman, J. Schieck² , R. Schöfbeck² , M. Shooshtari , M. Sonawane , W. Waltenberger , C.-E. Wulz² 
















Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , D. Ocampo Henao , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

J. Bierkens , N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , P. Kashko , S. Lowette , I. Makarenko , D. Müller , J. Song , S. Tavernier , M. Tytgat³ , G.P. Van Onsem , S. Van Putte , D. Vannerom 
















Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , A. Malara , M.A. Shahzad, A. Sharma , L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer , F. Zhang 












Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , D. Marckx , K. Skovpen , A.M. Tomaru, N. Van Den Bossche , J. van der Linden , J. Vandenbroeck , L. Wezenbeek 






















Université Catholique de Louvain, Louvain-la-Neuve, Belgium

H. Aarup Petersen , S. Bein , A. Benecke , A. Bethani , G. Bruno , A. Cappati , J. De Favereau De Jeneret , C. Delaere , F. Gameiro Casalinho , A. Giammanco , A.O. Guzel , V. Lemaître, J. Lidrych , P. Malek , P. Mastrapasqua , S. Turcpar 











Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

G.A. Alves , M. Barroso Ferreira Filho , E. Coelho , C. Hensel , D. Matos Figueiredo , T. Menezes De Oliveira , C. Mora Herrera , P. Rebello Teles , M. Soeiro , E.J. Tonelli Manganote⁴ , A. Vilela Pereira 







Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , H. Brandao Malbouisson , W. Carvalho , J. Chinellato⁵ , M. Costa Reis , E.M. Da Costa , G.G. Da Silveira⁶ , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza , S. S. Jesus , T. Laux Kuhn⁶ , M. Macedo , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel , F. Torres Da Silva De Araujo⁷ 






Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes⁶ , L. Calligaris , F. Damas , T.R. Fernandez Perez Tomei , E.M. Gregores , B. Lopes Da Costa , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , Sandra S. Padula , V. Scheurer




Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , P. Danev, R. Hadjiiska , P. Iaydjiev , M. Shopova , G. Sultanov 


University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Universidad Tecnica Federico Santa Maria, Valparaiso, Chile

W. Brooks 














Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Wang , L. Yuan 














Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, X. Wang , H. Yang

Institute of High Energy Physics, Beijing, China

G.M. Chen⁸ , H.S. Chen⁸ , M. Chen⁸ , Y. Chen , Q. Hou , X. Hou, F. Iemmi , C.H. Jiang, H. Liao , G. Liu , Z.-A. Liu⁹ , J.N. Song⁹, S. Song, J. Tao , C. Wang⁸, J. Wang , H. Zhang , J. Zhao 


State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, Q. Guo, Z. He, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, C. Quaranta , X. Sun , D. Wang , J. Wang, M. Zhang, Y. Zhao, C. Zhou 

State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 




Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer^{10,11}, Z. Cui¹¹, B. Li¹², H. Wang , K. Yi¹³ , J. Zhang 


Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Y. Li, Y. Zhou¹⁴





Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin , C. Lu , M. Xiao¹⁵ 




Universidad de Los Andes, Bogota, Colombia

C. Avila , D.A. Barbosa Trujillo , A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega




Universidad de Antioquia, Medellin, Colombia

C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

N. Godinovic , D. Lelas , A. Sculac 










University of Split, Faculty of Science, Split, Croatia

M. Kovac , A. Petkovic , T. Sculac 

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov , T. Susa 

University of Cyprus, Nicosia, Cyprus

A. Attikis , K. Christoforou , S. Konstantinou , C. Leonidou , L. Paizanos , F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

Charles University, Prague, Czech Republic

M. Finger[†] , M. Finger Jr. 

Escuela Politecnica Nacional, Quito, Ecuador

E. Ayala 



Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin 







Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

A.A. Abdelalim^{16,17} , B. El-mahdy¹⁸ 

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

A. Hussein , H. Mohammed 


National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , N. Seeba , L. Tani 

Department of Physics, University of Helsinki, Helsinki, Finland

E. Brücken , A. Milieva , K. Osterberg , M. Voutilainen 



















Helsinki Institute of Physics, Helsinki, Finland

F. Garcia , P. Inkaew , K.T.S. Kallonen , R. Kumar Verma , T. Lampén , K. Lassila-Perini , B. Lehtela , S. Lehti , T. Lindén , N.R. Mancilla Xinto , M. Myllymäki , M.m. Rantanen , S. Saariokari , N.T. Toikka , J. Tuominiemi 


























Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

N. Bin Norjoharuddeen , H. Kirschenmann , P. Luukka , H. Petrow 















IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, P. Devouge, J.L. Faure , F. Ferri , P. Gaigne, S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , Y. Maidannyk , J. Malcles , F. Orlandi , L. Portales , S. Ronchi , M.Ö. Sahin , A. Savoy-Navarro¹⁹ , P. Simkina , M. Titov , M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

R. Amella Ranz , F. Beaudette , G. Boldrini , P. Busson , C. Charlot , M. Chiusi , T.D. Cuisset , O. Davignon , A. De Wit , T. Debnath , I.T. Ehle , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , L. Kalipoliti , M. Manoni , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , A. Zabi , A. Zghiche 






















Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram²⁰ , J. Andrea , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , G. Coulon, S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , P. Vaucelle 




Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio , B. Orzari 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram , S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo , P. Depasse , H. El Mamouni , J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg , G. Grenier , B. Ille , E. Jourd'Huy , M. Lethuillier , B. Massoteau , L. Mirabito , A. Purohit , M. Vander Donckt , J. Xiao 





























Georgian Technical University, Tbilisi, Georgia

A. Khvedelidze²¹ , I. Lomidze , Z. Tsamalaidze²¹ 

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , P. Nattland , V. Oppenländer , A. Pauls , D. Pérez Adán , N. Röwert 






































































RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Daumann , S. Diekmann , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung , N. Kumar , M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , A. Pozdnyakov , W. Redjeb , H. Reithler , U. Sarkar , V. Sarkisovi , A. Schmidt , C. Seth , A. Sharma , J.L. Spah , V. Vaulin , S. Zaleski 










































RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

M.R. Beckers , C. Dziwok , G. Flügge , N. Hoeflich , T. Kress , A. Nowack , O. Pooth , A. Stahl , A. Zotz 




















































Deutsches Elektronen-Synchrotron, Hamburg, Germany

A. Abel , M. Aldaya Martin , J. Alimena , S. Amoroso , Y. An , I. Andreev , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman²² , K. Borras²³ , A. Campbell , S. Chatterjee , L.X. Coll Saravia , G. Eckerlin , D. Eckstein , E. Gallo²² , A. Geiser , M. Guthoff , A. Hinzmann , L. Jeppe , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange , D. Leyva Pernia , K.-Y. Lin , K. Lipka²⁴ , W. Lohmann²⁵ , J. Malvaso , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza , D. Rastorguev , L. Rygaard , M. Scham^{26,23} , S. Schnake²³ , P. Schütze , C. Schwanenberger²² , D. Schwarz , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford , M. Torkian , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann , L. Wiens²³ , C. Wissing , Y. Yang , S. Zakharov , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A.R. Alves Andrade , M. Antonello , S. Bollweg , M. Bonanomi , L. Ebeling , K. El Morabit , Y. Fischer , M. Frahm , E. Garutti , A. Grohsjean , A.A. Guvenli , J. Haller , D. Hundhausen , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer , C.c. Kuo , F. Labe , J. Lange , A. Lobanov , J. Matthiesen , L. Moureaux , K. Nikolopoulos , A. Paasch , K.J. Pena Rodriguez , N. Prouvost , B. Raciti , M. Rieger , D. Savoii , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , R. Ward , B. Wiederspan , M. Wolf , C. Yede 










Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , A. Brusamolino , E. Butz , Y.M. Chen , T. Chwalek , A. Dierlamm , G.G. Dincer , D. Druzhkin , U. Elicabuk , N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann²⁷ , M. Horzela , F. Hummer , U. Husemann , J. Kieseler , M. Klute , J. Knolle , R. Kunnilan Muhammed Rafeek , O. Lavoryk , J.M. Lawhorn , A. Lintuluoto , S. Maier , A.A. Monsch , M. Mormile , Th. Müller , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , R. Schmieder , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa , L. Stockmeier , K. Tauqeer , M. Toms , B. Topko , N. Trevisani , C. Verstege , T. Voigtländer , R.F. Von Cube , J. Von Den Driesch , M. Wassmer , C. Winter , R. Wolf , W.D. Zeuner , X. Zuo 






Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou , G. Daskalakis , A. Kyriakis 









National and Kapodistrian University of Athens, Athens, Greece

G. Melachroinos , Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , E. Tzovara , K. Vellidis , I. Zisopoulos 







National Technical University of Athens, Athens, Greece

T. Chatzistavrou , G. Karapostoli , K. Kousouris , E. Siamarkou , G. Tsipolitis 

University of Ioánnina, Ioánnina, Greece

I. Bestintzanos , I. Evangelou , C. Foudas , P. Katsoulis , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath^{28,29} , K. Márton , A.J. Rádl³⁰ , F. Sikler , V. Veszpremi 





MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkuti³¹ , M.M.A. Gadallah³² , Á. Kadlecik , M. León Coello , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 




HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

G. Bencze , S. Czellar , J. Molnar , Z. Szillasi 



Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

T. Csorgo³¹ , F. Nemes³¹ , T. Novak , I. Szanyi³³ 

IIT Bhubaneswar, Bhubaneswar, India

S. Bahinipati , S. Nayak , R. Raturi 

Panjab University, Chandigarh, India




S. Bansal , S.B. Beri , V. Bhatnagar , S. Chauhan , N. Dhingra³⁴ , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand , J.B. Singh , A. Singla 

University of Delhi, Delhi, India





































































A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , S. Phor , K. Ranjan , M.K. Saini 















Indian Institute of Technology Mandi (IIT-Mandi), Himachal Pradesh, India

P. Palni 




University of Hyderabad, Hyderabad, IndiaS. Acharya³⁵ , B. Gomber **Indian Institute of Technology Kanpur, Kanpur, India**S. Mukherjee **Saha Institute of Nuclear Physics, HBNI, Kolkata, India**S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar**Indian Institute of Technology Madras, Madras, India**M.M. Ameen , P.K. Behera , S. Chatterjee , G. Dash , A. Dattamunsi, P. Jana , P. Kalbhor , S. Kamble , J.R. Komaragiri³⁶ , T. Mishra , P.R. Pujahari , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay **IISER Mohali, India, Mohali, India**

B.K. Sirasva
















Tata Institute of Fundamental Research-A, Mumbai, IndiaL. Bhatt, S. Dugad , G.B. Mohanty , M. Shelake , P. Suryadevara**Tata Institute of Fundamental Research-B, Mumbai, India**A. Bala , S. Banerjee , S. Barman³⁷ , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar , M. Maity³⁷, G. Majumder , K. Mazumdar , S. Parolia , R. Saxena , A. Thachayath **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**D. Maity³⁸ , P. Mal , K. Naskar³⁸ , A. Nayak³⁸ , K. Pal , P. Sadangi, S.K. Swain , S. Varghese³⁸ , D. Vats³⁸ **Indian Institute of Science Education and Research (IISER), Pune, India**S. Dube , P. Hazarika , B. Kansal , A. Laha , R. Sharma , S. Sharma , K.Y. Vaish **Indian Institute of Technology Hyderabad, Telangana, India**S. Ghosh **Isfahan University of Technology, Isfahan, Iran**H. Bakhshiansohi³⁹ , A. Jafari⁴⁰ , V. Sedighzadeh Dalavi , M. Zeinali⁴¹ **Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**S. Bashiri , S. Chenarani⁴² , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang⁴³ **University College Dublin, Dublin, Ireland**M. Felcini , M. Grunewald **INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy**M. Abbrescia^{a,b} , M. Barbieri^{a,b}, M. Buonsante^{a,b} , A. Colaleo^{a,b} , D. Creanza^{a,c} , N. De Filippis^{a,c} , M. De Palma^{a,b} , W. Elmetenawee^{a,b,16} , N. Ferrara^{a,c} , L. Fiore^a , L. Generoso^{a,b}, L. Longo^a , M. Louka^{a,b} , G. Maggi^{a,c} , M. Maggi^a , I. Margjeka^a , V. Mastrapasqua^{a,b} , S. My^{a,b} , F. Nenna^{a,b} , S. Nuzzo^{a,b} , A. Pellecchia^{a,b} , A. Pompili^{a,b} , G. Pugliese^{a,c} , R. Radogna^{a,b} , D. Ramos^a , A. Ranieri^a , L. Silvestris^a , F.M. Simone^{a,c} , Ü. Sözbilir^a , A. Stamerra^{a,b} , D. Troiano^{a,b} , R. Venditti^{a,b} , P. Verwilligen^a , A. Zaza^{a,b} **INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy**

G. Abbiendi^a , C. Battilana^{a,b} , D. Bonacorsi^{a,b} , P. Capiluppi^{a,b} , F.R. Cavallo^a , T. Diotalevi^{a,b} , F. Fabbri^a , A. Fanfani^{a,b} , R. Farinelli^a , D. Fasanella^a , P. Giacomelli^a , C. Grandi^a , L. Guiducci^{a,b} , S. Lo Meo^{a,44} , M. Lorusso^{a,b} , L. Lunerti^a , S. Marcellini^a , G. Masetti^a , F.L. Navarra^{a,b} , G. Paggi^{a,b} , A. Perrotta^a , A.M. Rossi^{a,b} , S. Rossi Tisbeni^{a,b} , T. Rovelli^{a,b} , G.P. Siroli^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,45} , A. Di Mattia^a , A. Lapertosa^a , R. Potenza^{a,b} , A. Tricomi^{a,b,45} 





INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

J. Altork^{a,b} , P. Assiouras^a , G. Barbaglia^a , G. Bardelli^a , M. Bartolini^{a,b} , A. Calandri^{a,b} , B. Camaiani^{a,b} , A. Cassese^a , R. Ceccarelli^a , V. Ciulli^{a,b} , C. Civinini^a , R. D'Alessandro^{a,b} , L. Damenti^{a,b} , E. Focardi^{a,b} , T. Kello^a , G. Latino^{a,b} , P. Lenzi^{a,b} , M. Lizzo^a , M. Meschini^a , S. Paoletti^a , A. Papanastassiou^{a,b} , G. Sguazzoni^a , L. Viliani^a



















INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi , S. Bianco , S. Meola⁴⁶ , D. Piccolo 





INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

M. Alves Gallo Pereira^a , F. Ferro^a , E. Robutti^a , S. Tosi^{a,b} 



















INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a , F. Brivio^a , V. Camagni^{a,b} , F. Cetorelli^{a,b} , F. De Guio^{a,b} , M.E. Dinardo^{a,b} , P. Dini^a , S. Gennai^a , R. Gerosa^{a,b} , A. Ghezzi^{a,b} , P. Govoni^{a,b} , L. Guzzi^a , M.R. Kim^a , G. Lavizzari^{a,b} , M.T. Lucchini^{a,b} , M. Malberti^a , S. Malvezzi^a , A. Massironi^a , D. Menasce^a , L. Moroni^a , M. Paganoni^{a,b} , S. Palluotto^{a,b} , D. Pedrini^a , A. Perego^{a,b} , G. Pizzati^{a,b} , T. Tabarelli de Fatis^{a,b}











INFN Sezione di Napoli^a, Università di Napoli 'Federico II'^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a , C. Di Fraia^{a,b} , F. Fabozzi^{a,c} , L. Favilla^{a,d} , A.O.M. Iorio^{a,b} , L. Lista^{a,b,47} , P. Paolucci^{a,27} , B. Rossi^a 















INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università degli Studi di Cagliari^c, Cagliari, Italy

P. Azzi^a , N. Bacchetta^{a,48} , D. Bisello^{a,b} , L. Borella^a , P. Bortignon^{a,c} , G. Bortolato^{a,b} , A.C.M. Bulla^{a,c} , P. Checchia^a , T. Dorigo^{a,49} , F. Gasparini^{a,b} , U. Gasparini^{a,b} , S. Giorgetti^a , E. Lusiani^a , M. Margoni^{a,b} , A.T. Meneguzzo^{a,b} , M. Michelotto^a , J. Pazzini^{a,b} , F. Primavera^{a,b} , P. Ronchese^{a,b} , R. Rossin^{a,b} , F. Simonetto^{a,b} , M. Tosi^{a,b} , A. Triossi^{a,b} , S. Ventura^a , M. Zanetti^{a,b} , P. Zotto^{a,b} , A. Zucchetta^{a,b}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy


































A. Braghieri^a , M. Brunoldi^{a,b} , S. Calzaferri^{a,b} , P. Montagna^{a,b} , M. Pelliccioni^{a,b} , V. Re^a , C. Riccardi^{a,b} , P. Salvini^a , I. Vai^{a,b} , P. Vitulo^{a,b} 

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy


























S. Ajmal^{a,b} , M.E. Ascioti^{a,b} , G.M. Bilei^a , C. Carrivale^{a,b} , D. Ciangottini^{a,b} , L. Della Penna^{a,b} , L. Fanò^{a,b} , V. Mariani^{a,b} , M. Menichelli^a , F. Moscatelli^{a,50} , A. Rossi^{a,b} , A. Santocchia^{a,b} , D. Spiga^a , T. Tedeschi^{a,b} 

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy





























C. Aimè^{a,b} , C.A. Alexe^{a,c} , P. Asenov^{a,b} , P. Azzurri^a , G. Bagliesi^a , L. Bianchini^{a,b} 

T. Boccali^a , E. Bossini^a , D. Bruschini^{a,c} , R. Castaldi^a , F. Cattafesta^{a,c} , M.A. Ciocci^{a,d} , M. Cipriani^{a,b} , R. Dell'Orso^a , S. Donato^{a,b} , R. Forti^{a,b} , A. Giassi^a , F. Ligabue^{a,c} , A.C. Marini^{a,b} , A. Messineo^{a,b} , S. Mishra^a , V.K. Muraleedharan Nair Bindhu^{a,b} , S. Nandan^a , F. Palla^a , M. Riggirello^{a,c} , A. Rizzi^{a,b} , G. Rolandi^{a,c} , S. Roy Chowdhury^{a,51} , T. Sarkar^a , A. Scribano^a , P. Solanki^{a,b} , P. Spagnolo^a , F. Tenchini^{a,b} , R. Tenchini^a , G. Tonelli^{a,b} , N. Turini^{a,d} , F. Vaselli^{a,c} , A. Venturi^a , P.G. Verdini^a 






INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

P. Akrap^{a,b} , C. Basile^{a,b} , S.C. Behera^a , F. Cavallari^a , L. Cunqueiro Mendez^{a,b} , F. De Ruggi^{a,b} , D. Del Re^{a,b} , E. Di Marco^a , M. Diemoz^a , F. Errico^a , L. Frosina^{a,b} , R. Gargiulo^{a,b} , B. Harikrishnan^{a,b} , F. Lombardi^{a,b} , E. Longo^{a,b} , L. Martikainen^{a,b} , J. Mijuskovic^{a,b} , G. Organtini^{a,b} , N. Palmeri^{a,b} , R. Paramatti^{a,b} , S. Rahatlou^{a,b} , C. Rovelli^a , F. Santanastasio^{a,b} , L. Soffi^a , V. Vladimirov^{a,b} 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane^{a,b} , R. Arcidiacono^{a,c} , S. Argiro^{a,b} , M. Arneodo^{a,c} , N. Bartosik^{a,c} , R. Bellan^{a,b} , A. Bellora^{a,b} , C. Biino^a , C. Borca^{a,b} , N. Cartiglia^a , M. Costa^{a,b} , R. Covarelli^{a,b} , N. Demaria^a , L. Finco^a , M. Grippo^{a,b} , B. Kiani^{a,b} , L. Lanteri^{a,b} , F. Legger^a , F. Luongo^{a,b} , C. Mariotti^a , S. Maselli^a , A. Mecca^{a,b} , L. Menzio^{a,b} , P. Meridiani^a , E. Migliore^{a,b} , M. Monteno^a , M.M. Obertino^{a,b} , G. Ortona^a , L. Pacher^{a,b} , N. Pastrone^a , M. Ruspa^{a,c} , F. Siviero^{a,b} , V. Sola^{a,b} , A. Solano^{a,b} , A. Staiano^a , C. Tarricone^{a,b} , D. Trocino^a , G. Umoret^{a,b} , E. Vlasov^{a,b} , R. White^{a,b}

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar^{a,b,51} , S. Belforte^a , V. Candelise^{a,b} , M. Casarsa^a , F. Cossutti^a , K. De Leo^a , G. Della Ricca^{a,b} , R. Delli Gatti^{a,b} 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , J. Kim, T. Kim , D. Lee , H. Lee , J. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , S. Sekmen , B. Tae, Y.C. Yang 


Department of Mathematics and Physics - GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon , J. Seo 




Hanyang University, Seoul, Korea

E. Asilar , F. Carnevali , J. Choi⁵² , T.J. Kim , Y. Ryou 

Korea University, Seoul, Korea

S. Ha , S. Han, B. Hong , J. Kim , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, Korea

J. Goh , J. Shin , S. Yang 

Sejong University, Seoul, Korea

Y. Kang , H. S. Kim , Y. Kim , B. Ko, S. Lee 

Seoul National University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , H. Kim , J. Kim , T. Kim, Y. Kim , Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , J. Shin , U.K. Yang, I. Yoon

University of Seoul, Seoul, Korea

W. Jang , D.Y. Kang, D. Kim , S. Kim , J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

Yonsei University, Department of Physics, Seoul, Korea

G. Cho, K. Hwang , B. Kim , S. Kim, K. Lee , H.D. Yoo 


Sungkyunkwan University, Suwon, Korea

Y. Lee , I. Yu 

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

T. Beyrouthy , Y. Gharbia 


Kuwait University - College of Science - Department of Physics, Safat, Kuwait

F. Alazemi 

Riga Technical University, Riga, Latvia

K. Dreimanis , O.M. Eberlins , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs , R. Plese , A. Potrebko , M. Seidel , D. Sidiropoulos Kontos 

University of Latvia (LU), Riga, Latvia

N.R. Strautnieks 

Vilnius University, Vilnius, Lithuania

M. Ambrozas , A. Juodagalvis , S. Nargelas , A. Rinkevicius , G. Tamulaitis 








National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

I. Yusuff⁵³ , Z. Zolkapli



Universidad de Sonora (UNISON), Hermosillo, Mexico

J.F. Benitez , A. Castaneda Hernandez , A. Cota Rodriguez , L.E. Cuevas Picos, H.A. Encinas Acosta, L.G. Gallegos Maríñez, J.A. Murillo Quijada , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma , R. Lopez-Fernandez , J. Mejia Guisao , R. Reyes-Almanza , A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 





University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 





National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar , A. Awais , M.I.M. Awan, W.A. Khan 







AGH University of Krakow, Krakow, Poland

V. Avati, L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski 



National Centre for Nuclear Research, Swierk, Poland

M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

















Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,
A. Muhammad 





Warsaw University of Technology, Warsaw, Poland

P. Fokow , K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo ,
T. Camporesi , G. Da Molin , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo ,
A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 
































Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , L. Markovic , P. Milenovic , V. Milosevic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak , M. Dordevic , J. Milosevic , L. Nadderd , V. Rekoivic, M. Stojanovic 
















Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

M. Alcalde Martinez , J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes ,
Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz ,
A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos ,
J. Flix , M.C. Fouz , M. Gonzalez Hernandez , O. Gonzalez Lopez , S. Goy Lopez ,
J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasil-
las , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , R. Paz Herrera , A. Pérez-
Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Vazquez Escobar 




















Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 



Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Ayllon Torresano , A. Cardini , J. Cuevas , J. Del Riego Badas ,
D. Estrada Acevedo , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Ca-
ballero , P. Leguina , M. Obeso Menendez , E. Palencia Cortezon , J. Prado Pico ,
A. Soto Rodríguez , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fer-
nandez , G. Gomez , C. Lasosa García , R. Lopez Ruiz , C. Martinez Rivero ,
P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos ,
J. Piedra Gomez , C. Quintana San Emeterio , L. Scodellaro , I. Vila , R. Vi-
lar Cortabitarte , J.M. Vizan Garcia 


















University of Colombo, Colombo, Sri Lanka

B. Kailasapathy⁵⁴ , D.D.C. Wickramarathna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka














W.G.D. Dharmaratna⁵⁵ , K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland



























D. Abbaneo , C. Amendola , R. Ardino , E. Auffray , J. Baechler, D. Barney ,
J. Bendavid , M. Bianco , A. Bocci , L. Borgonovi , C. Botta , A. Bragagnolo ,
C.E. Brown , C. Caillol , G. Cerminara , P. Connor , K. Cormier , D. d'Enterria 

A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , P.J. Fernández Manteca , B.A. Fontana Santos Alves , E. Fontanesi , W. Funk , A. Gaddi, S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, A. Gruber , J. Hegeman , J.K. Heikkilä , R. Hofsaess , B. Huber , T. James , P. Janot , O. Kaluzinska , O. Karacheban²⁵ , G. Karathanasis , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , A.-M. Lyon , M. Magherini , L. Malgeri , M. Mannelli , A. Mehta , F. Meijers , J.A. Merlin, S. Mersi , E. Meschi , M. Migliorini , F. Monti , F. Moortgat , M. Mulders , M. Musich , I. Neutelings , S. Orfanelli, F. Pantaleo , M. Pari , G. Petrucciani , A. Pfeiffer , M. Pierini , M. Pitt , H. Qu , D. Rabadý , A. Reimers , B. Ribeiro Lopes , F. Riti , P. Rosado , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , M. Selvaggi , K. Shchelina , P. Silva , P. Sphicas⁵⁶ , A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea , E. Vernazza , J. Wanczyk⁵⁷ , S. Wuchterl , M. Zarucki , P. Zehetner , P. Zejdl , G. Zevi Della Porta 




















PSI Center for Neutron and Muon Sciences, Villigen, Switzerland

T. Bevilacqua⁵⁸ , L. Caminada⁵⁸ , W. Erdmann , R. Horisberger , Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , U. Langenegger , A. Nigamova , L. Noehte⁵⁸ , T. Rohe , A. Samalan 

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , M. Backhaus , G. Bonomelli , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'Archiacchiac⁵⁷ , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Glessgen , C. Grab , N. Härringer , T.G. Harte , W. Lustermann , M. Malucchi , R.A. Manzoni , L. Marchese , A. Mascellani⁵⁷ , F. Nessi-Tedaldi , F. Pauss , B. Ristic , R. Seidita , J. Steggemann⁵⁷ , A. Tarabini , D. Valsecchi , R. Wallny 













Universität Zürich, Zurich, Switzerland

C. Amsler⁵⁹ , P. Bärtschi , F. Bilandzija , M.F. Canelli , G. Celotto , V. Guglielmi , A. Jofrehei , B. Kilminster , T.H. Kwok , S. Leontsinis , V. Lukashenko , A. Macchiolo , F. Meng , M. Missiroli , J. Motta , P. Robmann, E. Shokr , F. Stäger , R. Tramontano , P. Viscone 

National Central University, Chung-Li, Taiwan

D. Bhowmik, C.M. Kuo, P.K. Rout , S. Taj , P.C. Tiwari³⁶ 


National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu , E. Yazgan 
















High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas 

Tunis El Manar University, Tunis, Tunisia

Y. Maghrbi 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Dolek , I. Dumanoglu⁶⁰ , Y. Guler⁶¹ , E. Gurpinar Guler⁶¹ , C. Isik , O. Kara⁶² , A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir⁶³ , B. Tali⁶⁴ , U.G. Tok , E. Uslan , I.S. Zorbakir 

Hacettepe University, Ankara, Turkey

S. Sen 




Middle East Technical University, Physics Department, Ankara, Turkey

M. Yalvac⁶⁵ 









Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi⁶⁶ , E. Gülmez , M. Kaya⁶⁷ , O. Kaya⁶⁸ , M.A. Sarkisla⁶⁹,
S. Tekten⁷⁰ 







Istanbul Technical University, Istanbul, Turkey

D. Boncukcu , A. Cakir , K. Cankocak^{60,71} 

Istanbul University, Istanbul, Turkey

B. Hacisahinoglu , I. Hos⁷² , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert ,
C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci , C. Dozen⁷³ , B. Isildak , E. Simsek , D. Sunar Cerci , T. Yetkin⁷³ 














Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , O. Dadazhanova, B. Grynyov 






















National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 








































University of Bristol, Bristol, United Kingdom

J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , D. Dharmender,
H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Parames-
varan , L. Robertshaw , M.S. Sanjrani³⁹, J. Segal, V.J. Smith 




Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev⁷⁴ , C. Brew , R.M. Brown , D.J.A. Cockerill ,
A. Elliot , K.V. Ellis, J. Gajownik , K. Harder , S. Harper , J. Linacre , K. Manolopoulos,
M. Moallemi , D.M. Newbold , E. Olaiya , D. Petyt , T. Reis , A.R. Sahasransu ,
G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen ,
T. Williams 














Imperial College, London, United Kingdom





















































































































I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller, C.A. Carrillo Montoya ,
D. Colling , I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi ,
G. Hall , H.R. Hoorani , A. Howard, G. Iles , C.R. Knight , P. Krueper , J. Langford ,
K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios,
A. Mastronikolis , M. Mieskolainen , J. Nash⁷⁵ , M. Pesaresi , P.B. Pradeep ,
B.C. Radburn-Smith , A. Richards, A. Rose , L. Russell , K. Savva , C. Seez ,
R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee²⁷ , M. Vojinovic ,
N. Wardle , D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, USA

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann ,
K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant ,
C. Sutantawibul , J. Wilson 














Bethel University, St. Paul, Minnesota, USAJ.M. Hogan **Catholic University of America, Washington, DC, USA**R. Bartek , A. Dominguez , S. Raj , B. Sahu , A.E. Simsek , S.S. Yu **The University of Alabama, Tuscaloosa, Alabama, USA**B. Bam , A. Buchot Perraguin , S. Campbell, R. Chudasama , S.I. Cooper , C. Crovella , G. Fidalgo , S.V. Gleyzer , A. Khukhunaishvili , K. Matchev , E. Pearson, P. Rumerio⁷⁶ , E. Usai , R. Yi **Boston University, Boston, Massachusetts, USA**S. Cholak , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , J. Fulcher , F. Golf , S. Jeon , J. O'Cain, I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , E. Wurtz, A.G. Zecchinelli **Brown University, Providence, Rhode Island, USA**G. Barone , G. Benelli , D. Cutts , S. Ellis , L. Gouskos , M. Hadley , U. Heintz , K.W. Ho , T. Kwon , L. Lambrecht , G. Landsberg , K.T. Lau , J. Luo , S. Mondal , J. Roloff, T. Russell , S. Sagir⁷⁷ , X. Shen , M. Stamenkovic , N. Venkatasubramanian **University of California, Davis, Davis, California, USA**S. Abbott , S. Baradia , B. Barton , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , E. Cannaert, M. Chertok , M. Citron , J. Conway , P.T. Cox , F. Eble , R. Erbacher , O. Kukral , G. Mocellin , S. Ostrom , I. Salazar Segovia, J.S. Tafoya Vargas , W. Wei , S. Yoo **University of California, Los Angeles, California, USA**K. Adamidis, M. Bachtis , D. Campos, R. Cousins , S. Crossley , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca , A. Nunez Del Prado , D. Saltzberg , V. Valuev **University of California, Riverside, Riverside, California, USA**R. Clare , J.W. Gary , G. Hanson **University of California, San Diego, La Jolla, California, USA**A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , B. D'Anzi , D. Diaz , J. Duarte , L. Giannini , Y. Gu, J. Guiang , V. Krutelyov , R. Lee , J. Letts , H. Li, M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch, M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , A. Yagil , Z. Zhao **University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA**A. Barzdukas , L. Brennan , C. Campagnari , S. Carron Montero⁷⁸ , K. Downham , C. Grieco , M.M. Hussain, J. Incandela , M.W.K. Lai, A.J. Li , P. Masterson , J. Richman , S.N. Santpur , R. Schmitz , D. Stuart , T.Á. Vámi , X. Yan , D. Zhang **California Institute of Technology, Pasadena, California, USA**A. Albert , S. Bhattacharya , A. Bornheim , O. Cerri, R. Kansal , J. Mao , H.B. Newman , G. Reales Gutiérrez, T. Sievert, M. Spiropulu , J.R. Vlimant , R.A. Wynne , S. Xie **Carnegie Mellon University, Pittsburgh, Pennsylvania, USA**J. Alison , S. An , M. Cremonesi, V. Dutta , E.Y. Ertorer , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil, M. Kanemura, C. Liu , M. Marchegiani 

P. Meiring , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 









































































University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , S. Kwan , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 



















Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , J. Dickinson , A. Duquette, J. Fan , X. Fan , J. Grassi , S. Hogan , P. Kotamnives , J. Monroy , G. Niendorf , M. Oshiro , J.R. Patterson , A. Ryd , J. Thom , P. Wittich , R. Zou , L. Zygala 













Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , C. Cosby , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green, A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , J. Hirschauer , V. Innocente , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , B. Klima , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg, V. Papadimitriou , N. Pastika , K. Pedro , C. Pena⁷⁹ , C.E. Perez Lara , V. Perovic , F. Ravera , A. Reinsvold Hall⁸⁰ , L. Ristori , M. Safdari , E. Sexton-Kennedy , E. Smith , N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , C. Wang , I. Zoi 

University of Florida, Gainesville, Florida, USA

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , V. Sulimov , Y. Takahashi , J. Wang 























Florida State University, Tallahassee, Florida, USA

T. Adams , A. Al Kadhim , A. Askew , S. Bower , R. Goff, R. Hashmi , A. Hassani , R.S. Kim , T. Kolberg , G. Martinez , M. Mazza , H. Prosper , P.R. Prova, R. Yohay 










Florida Institute of Technology, Melbourne, Florida, USA

B. Alsufyani , S. Butalla , S. Das , M. Hohlmann , M. Lavinsky, E. Yanes











University of Illinois Chicago, Chicago, Illinois, USA

M.R. Adams , N. Barnett, A. Baty , C. Bennett , R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , H. Gupta , M. Hawksworth, A. Hingrajiya, D.J. Hofman , Z. Huang , J.h. Lee , C. Mills , S. Nanda , G. Nigmatkulov , B. Ozek , T. Phan, D. Pilipovic , R. Pradhan , E. Prifti, P. Roy, T. Roy , D. Shekar, N. Singh, A. Thielen, M.B. Tonjes , N. Varelas , M.A. Wadud , J. Yoo 
























The University of Iowa, Iowa City, Iowa, USA

M. Alhusseini , D. Blend , K. Dilsiz⁸¹ , O.K. Köseyan , A. Mestvirishvili⁸² , O. Neogi, H. Ogul⁸³ , Y. Onel , A. Penzo , C. Snyder, E. Tiras⁸⁴ 










Johns Hopkins University, Baltimore, Maryland, USA

B. Blumenfeld , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , S. Sekhar , M.V. Srivastav , M. Swartz 



















The University of Kansas, Lawrence, Kansas, USA

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , R. Bhattacharya , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , S. Popescu⁸⁵ , C. Rogan , C. Royon , S. Rudrabhatla , S. Sanders , C. Smith , G. Wilson 





































Kansas State University, Manhattan, Kansas, USA

B. Allmond , N. Islam , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , G.G. Reddy , D. Roy , G. Sorrentino 













University of Maryland, College Park, Maryland, USA

A. Baden , A. Belloni , J. Bistany-riebman , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim , S. Lascio , P. Major , A.C. Mignerey , C. Palmer , C. Papageorgakis , M.M. Paranjpe , E. Popova⁸⁶ , A. Shevelev , L. Zhang 













Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

C. Baldenegro Barrera , H. Bossi , S. Bright-Thonney , I.A. Cali , Y.c. Chen , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov , G. Grosso , P. Harris , D. Hoang , G.M. Innocenti , K. Ivanov , D. Kovalskiy , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. Mcginn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , T.a. Sheng , G.S.F. Stephans , D. Walter , J. Wang , Z. Wang , B. Wyslouch , T. J. Yang 














University of Minnesota, Minneapolis, Minnesota, USA

A. Alpana , B. Crossman , W.J. Jackson , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi , R. Rusack , O. Sancar , R. Saradhy , N. Strobbe 





















University of Nebraska-Lincoln, Lincoln, Nebraska, USA

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , K.H.M. Kwok , A. Rohilla , J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan 










State University of New York at Buffalo, Buffalo, New York, USA

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos , A. Kharchilava , A. Mandal , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar , A. Williams , D. Yu 
































Northeastern University, Boston, Massachusetts, USA

A. Aarif , G. Alverson , E. Barberis , J. Bonilla , B. Bylsma , M. Campana , J. Dervan , Y. Haddad , Y. Han , I. Israr , A. Krishna , M. Lu , N. Manganelli , R. Mccarthy , D.M. Morse , T. Orimoto , L. Skinnari , C.S. Thoreson , E. Tsai , D. Wood 

Northwestern University, Evanston, Illinois, USA

S. Dittmer , K.A. Hahn , M. Mcginnis , Y. Miao , D.G. Monk , M.H. Schmitt , A. Taliercio , M. Velasco , J. Wang 


















University of Notre Dame, Notre Dame, Indiana, USA

G. Agarwal , R. Band , R. Bucci , S. Castells , A. Das , A. Datta , A. Ehnis , R. Goldouzian , M. Hildreth , K. Hurtado Anampa , T. Ivanov , C. Jessop , A. Karneyeu , K. Lannon , J. Lawrence , N. Loukas , L. Lutton , J. Mariano , N. Marinelli , T. McCauley , C. Mcgrady , C. Moore , Y. Musienko²¹ , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend , Y. Wan , M. Wayne , H. Yockey 

The Ohio State University, Columbus, Ohio, USA

M. Carrigan , R. De Los Santos , L.S. Durkin , C. Hill , M. Joyce , D.A. Wenzl, B.L. Winer , B. R. Yates 



















Princeton University, Princeton, New Jersey, USA

H. Bouchamaoui , G. Dezoort , P. Elmer , A. Frankenthal , M. Galli , B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , Y. Lai , D. Lange , A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, USA

S. Malik , R. Sharma 




















Purdue University, West Lafayette, Indiana, USA

S. Chandra , A. Gu , L. Gutay, M. Huwiler , M. Jones , A.W. Jung , D. Kondratyev , J. Li , M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , N.R. Saha , J.F. Schulte , F. Wang , A. Wildridge , W. Xie , Y. Yao , Y. Zhong 











Purdue University Northwest, Hammond, Indiana, USA

N. Parashar , A. Pathak , E. Shumka 





















Rice University, Houston, Texas, USA

D. Acosta , A. Agrawal , C. Arbour , T. Carnahan , P. Das , K.M. Ecklund , F.J.M. Geurts , T. Huang , I. Krommydas , N. Lewis, W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi , J. Rotter , C. Vico Villalba , M. Wulansatiti , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, USA

O. Bessidskaia Bylund, A. Bodek , P. de Barbaro[†] , R. Demina , A. Garcia-Bellido , H.S. Hare , O. Hindrichs , N. Parmar , P. Parygin⁸⁶ , H. Seo , R. Taus 
















Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA

B. Chiarito, J.P. Chou , S.V. Clark , S. Donnelly, D. Gadkari , Y. Gershtein , E. Halkiadakis , C. Houghton , D. Jaroslawski , A. Kobert , I. Laflotte , A. Lath , J. Martins , M. Perez Prada , B. Rand , J. Reichert , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

University of Tennessee, Knoxville, Tennessee, USA

D. Ally , A.G. Delannoy , S. Fiorendi , J. Harris, T. Holmes , A.R. Kanuganti , N. Karunarathna , J. Lawless, L. Lee , E. Nibigira , B. Skipworth, S. Spanier 













Texas A&M University, College Station, Texas, USA

D. Aebi , M. Ahmad , T. Akhter , K. Androsov , A. Basnet , A. Bolshov, O. Bouhali⁸⁷ , A. Cagnotta , V. D'Amante , R. Eusebi , P. Flanagan , J. Gilmore , Y. Guo, T. Kamon , S. Luo , R. Mueller , A. Safonov 

Texas Tech University, Lubbock, Texas, USA

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , W. Jin , Y. Kazhykarim, K. Lamichhane , S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, USA

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen , J. Zhang

University of Virginia, Charlottesville, Virginia, USA

























B. Cardwell , H. Chung , B. Cox , J. Hakala , G. Hamilton Ilha Machado, R. Hirosky 

M. Jose, A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez , Z. Wu


Wayne State University, Detroit, Michigan, USA

S. Bhattacharya , P.E. Karchin 











University of Wisconsin - Madison, Madison, Wisconsin, USA

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , S. Lomte , R. Loveless , A. Mallampalli , A. Mohammadi , S. Mondal, T. Nelson, G. Parida , L. Pétré , D. Pinna , A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

Authors affiliated with an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , Yu. Andreev , T. Aushev , D. Budkouski , R. Chistov , M. Danilov , T. Dimova , A. Ershov , S. Gninenko , I. Gorbunov , A. Gribushin , A. Kamenev , V. Karjavine , M. Kirsanov , V. Klyukhin , O. Kodolova⁸⁸ , V. Koronkov , I. Korsakov, A. Kozyrev , N. Krasnikov , A. Lanev , A. Malakhov , V. Matveev , A. Nikitenko^{89,88} , V. Palichik , V. Perelygin , S. Petrushanko , O. Radchenko , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , Y. Skovpen , K. Slizhevskiy, V. Smirnov , O. Teryaev , I. Tlisova , A. Toropin , N. Voytishin , A. Zarubin , I. Zhizhin 

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

E. Boos , V. Bunichev , M. Dubinin⁷⁹ , V. Savrin , A. Snigirev , L. Dudko , V. Kim²¹ , V. Murzin , V. Oreshkin , D. Sosnov 

†: Deceased

¹Also at Yerevan State University, Yerevan, Armenia

²Also at TU Wien, Vienna, Austria

³Also at Ghent University, Ghent, Belgium

⁴Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

⁵Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷Also at The University of the State of Amazonas, Manaus, Brazil

⁸Also at University of Chinese Academy of Sciences, Beijing, China

⁹Also at University of Chinese Academy of Sciences, Beijing, China

¹⁰Also at School of Physics, Zhengzhou University, Zhengzhou, China

¹¹Now at Henan Normal University, Xinxiang, China

¹²Also at University of Shanghai for Science and Technology, Shanghai, China

¹³Also at The University of Iowa, Iowa City, Iowa, USA

¹⁴Also at Nanjing Normal University, Nanjing, China

¹⁵Also at Center for High Energy Physics, Peking University, Beijing, China

¹⁶Also at Helwan University, Cairo, Egypt

¹⁷Now at Zewail City of Science and Technology, Zewail, Egypt

¹⁸Also at Cairo University, Cairo, Egypt

¹⁹Also at Purdue University, West Lafayette, Indiana, USA

²⁰Also at Université de Haute Alsace, Mulhouse, France

²¹Also at an institute formerly covered by a cooperation agreement with CERN

²²Also at University of Hamburg, Hamburg, Germany

²³Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁴Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

- ²⁵Also at Brandenburg University of Technology, Cottbus, Germany
- ²⁶Also at Forschungszentrum Jülich, Juelich, Germany
- ²⁷Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ²⁸Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary
- ²⁹Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania
- ³⁰Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary
- ³¹Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary
- ³²Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt
- ³³Also at The University of Kansas, Lawrence, Kansas, USA
- ³⁴Also at Punjab Agricultural University, Ludhiana, India
- ³⁵Also at University of Hyderabad, Hyderabad, India
- ³⁶Also at Indian Institute of Science (IISc), Bangalore, India
- ³⁷Also at University of Visva-Bharati, Santiniketan, India
- ³⁸Also at Institute of Physics, Bhubaneswar, India
- ³⁹Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- ⁴⁰Also at Isfahan University of Technology, Isfahan, Iran
- ⁴¹Also at Sharif University of Technology, Tehran, Iran
- ⁴²Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- ⁴³Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- ⁴⁴Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- ⁴⁵Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- ⁴⁶Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- ⁴⁷Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- ⁴⁸Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- ⁴⁹Also at Lulea University of Technology, Lulea, Sweden
- ⁵⁰Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- ⁵¹Also at UPES - University of Petroleum and Energy Studies, Dehradun, India
- ⁵²Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France
- ⁵³Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- ⁵⁴Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- ⁵⁵Also at Saegis Campus, Nugegoda, Sri Lanka
- ⁵⁶Also at National and Kapodistrian University of Athens, Athens, Greece
- ⁵⁷Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- ⁵⁸Also at Universität Zürich, Zurich, Switzerland
- ⁵⁹Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- ⁶⁰Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- ⁶¹Also at Konya Technical University, Konya, Turkey
- ⁶²Also at Istanbul Topkapi University, Istanbul, Turkey
- ⁶³Also at Izmir Bakircay University, Izmir, Turkey
- ⁶⁴Also at Adiyaman University, Adiyaman, Turkey
- ⁶⁵Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- ⁶⁶Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey
- ⁶⁷Also at Marmara University, Istanbul, Turkey
- ⁶⁸Also at Milli Savunma University, Istanbul, Turkey

-
- ⁶⁹ Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey
- ⁷⁰ Also at Kafkas University, Kars, Turkey
- ⁷¹ Now at Istanbul Okan University, Istanbul, Turkey
- ⁷² Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- ⁷³ Also at Istinye University, Istanbul, Turkey
- ⁷⁴ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- ⁷⁵ Also at Monash University, Faculty of Science, Clayton, Australia
- ⁷⁶ Also at Università di Torino, Torino, Italy
- ⁷⁷ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- ⁷⁸ Also at California Lutheran University,, Thousand Oaks, California, USA
- ⁷⁹ Also at California Institute of Technology, Pasadena, California, USA
- ⁸⁰ Also at United States Naval Academy, Annapolis, Maryland, USA
- ⁸¹ Also at Bingol University, Bingol, Turkey
- ⁸² Also at Georgian Technical University, Tbilisi, Georgia
- ⁸³ Also at Sinop University, Sinop, Turkey
- ⁸⁴ Also at Erciyes University, Kayseri, Turkey
- ⁸⁵ Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania
- ⁸⁶ Now at another institute formerly covered by a cooperation agreement with CERN
- ⁸⁷ Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar
- ⁸⁸ Also at Yerevan Physics Institute, Yerevan, Armenia
- ⁸⁹ Also at Imperial College, London, United Kingdom