

On uniqueness of coarse median structures

ELIA FIORAVANTI AND ALESSANDRO SISTO

We show that any product of bushy hyperbolic spaces has a unique coarse median structure, and that having a unique coarse median structure is a property closed under relative hyperbolicity. As a consequence, in contrast with the case of mapping class groups, there are nonhyperbolic pants graphs that have unique coarse median structures.

1 Introduction

Coarse median structures were introduced by Bowditch in [2] as a coarsening of the classical notion of median algebra. Roughly, a coarse median is a notion of centre for triangles in a given metric space. Bowditch’s motivation was to study the coarse geometry of mapping class groups and Teichmüller spaces, in particular quasi-isometric rigidity [8; 5]. Other examples of coarse median groups include cocompactly cubulated groups and hierarchically hyperbolic groups [8].

Since a coarse median structure on a given metric space is an additional structure, it is natural to wonder how canonical it is. A coarse median on the metric space X is in particular a map $\mu: X^3 \rightarrow X$, and therefore it is natural to define two coarse medians to be *equivalent* if they are bounded distance from each other. If all coarse medians on a given space are within bounded distance of each other, then we say that the space admits a *unique coarse median structure*.

The problem of uniqueness of coarse median structures, besides being a natural question, is further motivated by the theory of coarse-median preserving automorphisms, which are better behaved than general automorphisms [19; 18; 20]. For instance, fixed subgroups are finitely generated for coarse-median preserving automorphisms of cocompactly cubulated groups, but this is not true for general automorphisms. Clearly, if a group admits a unique coarse median structure, then all automorphisms preserve it.

The only groups that are so far known to have a unique coarse median structure are Gromov-hyperbolic groups [26]. There are further examples having a coarse median structure that is unique among those coming from cocompact cubulations [21], and in fact very recently this property has been shown to be generic in certain random models for right-angled Coxeter groups [1]. Surprisingly to the authors, the mapping class group of the five-holed sphere does not have a unique coarse median structure [25], and in fact it has uncountably many. The same also holds for a lot of right-angled Artin groups, including the one on the pentagon (which however has a unique coarse median structure coming from cocompact cubulations).

MSC2020: 20F65, 20F67, 20F69.

© 2026 The Authors, under license to MSP (Mathematical Sciences Publishers). Distributed under the [Creative Commons Attribution License 4.0 \(CC BY\)](#). Open Access made possible by subscribing institutions via [Subscribe to Open](#).

Our goal here is to provide examples of groups and spaces that do have unique coarse median structures. First, we consider products of bushy hyperbolic spaces. A hyperbolic space X is *bushy* if there exists a constant $\lambda \geq 0$ such that, for all points $x \in X$, there exist three ideal points $\xi_1, \xi_2, \xi_3 \in \partial_\infty X$ with pairwise Gromov products based at x bounded above by λ .

Theorem A *Let X be a finite product of bushy geodesic hyperbolic spaces. Then X admits a unique coarse median structure.*

Note that we do need the hyperbolic spaces to be bushy (Remark 4.8), and in fact this is the phenomenon exploited in [25] to construct different coarse median structures on a mapping class group.

Next, we show that having a unique coarse median structure is a property closed under relative hyperbolicity. Existence of coarse median structures in this case is due to Bowditch [3]; we deal with uniqueness only.

Theorem B *Let $(X, \mathcal{P})$ be a geodesic relatively hyperbolic space. Suppose that each $P \in \mathcal{P}$ admits a unique coarse median structure, and that there are only finitely many isometry types in $\mathcal{P}$. Then X admits a unique coarse median structure.*

The two theorems imply for instance that the right-angled Artin groups $F_2 \times F_2$ and $\mathbb{Z} * (F_2 \times F_2)$ admit unique coarse median structures.

Combining the two results we also obtain uniqueness of coarse median structures for pants graphs in low complexity, in contrast with the mapping class group case. The pants graph of a finite-type surface was introduced by Hatcher and Thurston [22]. Brock showed that it is quasi-isometric to Teichmüller space with the Weil–Petersson metric [12].

Corollary C *Let S be a connected finite-type surface of genus g with p punctures, and suppose that $3g + p \leq 6$. Then the pants graph of S has a unique coarse median structure.*

Proof Given the constraints on g and p , the pants graph is (either hyperbolic or) hyperbolic relative to products of two Farey graphs; see [13, Theorem 1.1; 14, Theorem 1]. Theorems A and B therefore apply. $\square$

Mangioni’s arguments in [25] suggest that mapping class groups of higher-complexity surfaces should have uncountably many coarse median structures, but there is a chance that the answer to the following is affirmative:

Question Do all pants graphs of finite-type surfaces have a unique coarse median structure?

The main challenge in answering the question above is that we have much less control on quasigeodesics in pants graphs that are not relatively hyperbolic.

There are several other natural problems that arise regarding the classification of groups with unique coarse median structures, including of particular interest:

Problem Classify right-angled Artin groups with unique coarse median structures.

Outline and proof ideas

In [Section 2](#) we cover preliminaries, the only novel result being [Corollary 2.6](#), which gives a criterion for a quasiflat in a coarse median space to be quasiconvex.

In [Section 3](#), we prove [Theorem B \(Theorem 3.5\)](#). Here we exploit the fact that intervals in coarse median spaces contain uniform quasigeodesics ([Lemmas 3.1](#) and [3.3](#)) and that quasigeodesic triangles in relatively hyperbolic spaces are well-understood.

In [Section 4](#), we prove [Theorem A \(Theorem 4.7\)](#). We exploit uniqueness of median operators in products of furry trees (which follows from work of Bowditch) and tricks with asymptotic cones to show that flats in products of bushy hyperbolic spaces are quasiconvex with respect to any coarse median structure. From this it is not hard to deduce uniqueness of the coarse median structure. (Uniqueness of the median operators on asymptotic cones does not seem to directly pull back to uniqueness of the coarse median structure on the space, in general.)

2 Preliminaries

2.1 Median algebras and median spaces

In this subsection we recall basic notions related to median algebras; see, e.g., [\[15; 3; 10\]](#) for an introduction.

Let Ω be a set and $m: \Omega^3 \rightarrow \Omega$ a map. The pair (Ω, m) is a *median algebra* if the following identities are satisfied for all elements $x, y, z, p \in \Omega$:

(M0) Symmetry and localisation: $m(x, y, z) = m(y, x, z) = m(x, z, y)$ and $m(x, x, y) = x$.

(M1) Associativity: $m(m(x, p, y), p, z) = m(x, p, m(y, p, z))$.

We refer to any map m satisfying these identities as a *median operator* on the set Ω .

The *n -dimensional discrete cube*, for some $n \geq 0$, is the finite median algebra consisting of the set $\{0, 1\}^n$ with the median operator computing majority votes on coordinates. The *rank* of a median algebra (Ω, m) is the largest integer r such that Ω contains an r -dimensional discrete cube as a median subalgebra; if such a largest integer does not exist, then the rank is infinite. We also define an *n -dimensional Euclidean cube* to be a product $\prod_{1 \leq i \leq n} [a_i, b_i]$ of nontrivial compact intervals $[a_i, b_i] \subseteq \mathbb{R}$, equipped with the median operator that chooses middle coordinates.

An *interval* in a median algebra (Ω, m) is a set of the form $[x, y] = \{m(x, y, z) \mid z \in \Omega\}$ for some $x, y \in \Omega$. A subset $A \subseteq \Omega$ is *convex* if we have $[a_1, a_2] \subseteq A$ for all $a_1, a_2 \in A$. One can show that intervals are convex.

A *topological median algebra* is the data of a median algebra (Ω, m) and a Hausdorff topology on Ω with respect to which the map m is continuous. If Ω is a metric space, we say that the median operator m is *Lipschitz* if there exists a constant $L \geq 1$ such that the map $x \mapsto m(x, y, z)$ is L -Lipschitz for all $y, z \in \Omega$.

A *median space* is a metric space (Ω, ρ) with the property that, for any three points $x_1, x_2, x_3 \in \Omega$, there exists a unique point $m = m(x_1, x_2, x_3) \in \Omega$ such that

$$\rho(x_i, x_j) = \rho(x_i, m) + \rho(m, x_j)$$

for all indices $1 \leq i < j \leq 3$. If (Ω, ρ) is a median space and $m: \Omega^3 \rightarrow \Omega$ is the resulting ternary map, then the pair (Ω, m) is a median algebra; we refer to m as the *induced* median operator. Conversely, if (Ω, m) is a median algebra and ρ is a metric on Ω with the property that $\rho(x, y) = \rho(x, m(x, y, z)) + \rho(m(x, y, z), y)$ for all $x, y, z \in \Omega$, then the pair (Ω, ρ) is a median space.

The following result is due to Zeidler [29, Proposition 3.3 and Lemma 3.5] (and to Bowditch for colourable median operators [4, Theorem 1.1]).

Proposition 2.1 *Let (Ω, d) be a geodesic metric space, and let $m: \Omega^3 \rightarrow \Omega$ be a Lipschitz median operator with finite rank. Then Ω admits a metric ρ bi-Lipschitz to d such that (Ω, ρ) is a median space inducing the median operator m .*

2.2 Coarse median spaces

In this section we recall the definition of coarse median space and related notions, and establish basic properties that will be useful later on.

Let X be a metric space. For $x, y \in X$ and $D \geq 0$, we write “ $x \approx_D y$ ” with the meaning of “ $d(x, y) \leq D$ ”. A *C-coarse median* on X , for some constant $C \geq 1$, is a map $\mu: X^3 \rightarrow X$ satisfying the following identities for all points $x, y, z, p \in X$:

- (M0) Symmetry and localisation: $\mu(x, y, z) = \mu(y, x, z) = \mu(x, z, y)$ and $\mu(x, x, y) = x$.
- (CM1) Coarse associativity: $\mu(\mu(x, p, y), p, z) \approx_C \mu(x, p, \mu(y, p, z))$.
- (CM2) Coarse Lipschitz property: $d(\mu(x, y, z), \mu(p, y, z)) \leq Cd(x, p) + C$.

We say that μ is a *coarse median* if it is a C -coarse median for some $C \geq 1$. The pair (X, μ) is then a *coarse median space*, or a C -coarse median space if we wish to specify the constant. (See [26] for the equivalence between the above definition and Bowditch’s original one in [2].)

Two coarse medians $\mu, \nu: X^3 \rightarrow X$ are *D-close*, for a constant $D \geq 0$, if we have $\mu(x, y, z) \approx_D \nu(x, y, z)$ for all points $x, y, z \in X$. We simply say that μ and ν are *close* if we do not wish to specify the constant, and note that closeness is an equivalence relation. We refer to each equivalence class of coarse medians on X , denoted $[\mu]$, as a *coarse median structure*.

There is a notion of *rank* for coarse median spaces extending the analogous notion for median algebras. Roughly, (X, μ) has $\text{rank} \leq r$ if all its finite subsets can be approximated by subsets of median algebras with $\text{rank} \leq r$. The exact definition is a bit more involved and will never be needed explicitly, so we refer to [2, Section 8].

In a coarse median space (X, μ) , the *interval* between two points $x, y \in X$ is the set

$$[x, y]_\mu := \{\mu(x, y, z) \mid z \in X\}.$$

We simply denote this by $[x, y]$ when there is no ambiguity on the coarse median under consideration. There is a natural map $\pi_{x,y}: X \rightarrow [x, y]$ given by $\pi_{x,y}(z) = \mu(x, y, z)$, and we refer to it as the *projection* to the interval. A subset $A \subseteq X$ is *D-quasiconvex*, for some $D \geq 0$, if the interval $[a_1, a_2]$ is contained in the D -neighbourhood of A for all points $a_1, a_2 \in A$. Intervals are uniformly quasiconvex; see Lemma 2.3(4).

Remark 2.2 Whenever we use the term *quasiconvex*, this will always be meant with respect to a fixed coarse median structure. We will never consider quasiconvexity in a metric sense, which would amount to requiring geodesics (or quasigeodesics) with endpoints in the set to remain close to it. Note that the only general connection between coarse median intervals and quasigeodesics is the rather weak one provided by Lemma 3.1 below. We refer to [26, Theorem 5.1] for an example where geodesics stray arbitrarily far from the interval between their endpoints.

We emphasise that, on the other hand, the term *quasigeodesic* is always meant in its classical metric sense. That is, given constants $K \geq 1$ and $L \geq 0$, a (K, L) -*quasigeodesic* in a metric space (Z, d) is a map $\gamma: I \rightarrow Z$, for a closed connected subset $I \subseteq \mathbb{R}$, such that we have

$$\frac{1}{K}|s - t| - L \leq d(\gamma(s), \gamma(t)) \leq K|s - t| + L \quad \text{for all } s, t \in I.$$

We will typically also refer to the image $\gamma(I) \subseteq Z$ as a quasigeodesic.

For a subset $A \subseteq X$ and $R \geq 0$, we denote by $\mathcal{N}_R(A)$ the (closed) R -neighbourhood of A in X . The following lemma collects various basic facts and identities that will be useful in the sequel. All of these are consequences of the general principle that every identity satisfied by median operators must hold for all C -coarse median operators up to bounded distance, and the error only depends on C and the specific identity. This is discussed in [8, Section 6] (see also [26, Section 3]).

Lemma 2.3 *Let (X, μ) be a C -coarse median space. There is $K = K(C) \geq 1$ such that the following hold.*

- (1) *For all $a, b, c, d, e \in X$, we have $\mu(\mu(a, b, c), d, e) \approx_K \mu(\mu(a, d, e), \mu(b, d, e), c)$.*
- (2) *Given $x, y \in X$, $p \in [x, y]$ and points $x' \in [x, p]$, $y' \in [p, y]$, we have $\mu(x', y', p) \approx_K p$.*
- (3) *For any $x, y, z \in X$, we have $\mu(x, y, z) \in [x, y] \cap [y, z] \cap [z, x]$. For any $R \geq 0$, the intersection*

$$\mathcal{N}_R([x, y]) \cap \mathcal{N}_R([y, z]) \cap \mathcal{N}_R([z, x])$$

has diameter at most $KR + K$.

- (4) *If $x, y \in X$ and $a, b \in \mathcal{N}_R([x, y])$ for some $R \geq 0$, then the interval $[a, b]$ is contained in the neighbourhood of $[x, y]$ of radius $KR + K$.*
- (5) *For $x, y, z \in X$ and $R \geq 0$, consider the points $a \in \mathcal{N}_R([x, y]) \cap \mathcal{N}_R([x, z])$, $b \in \mathcal{N}_R([y, z]) \cap \mathcal{N}_R([y, x])$ and $c \in \mathcal{N}_R([z, x]) \cap \mathcal{N}_R([z, y])$. Then $\mu(a, b, c)$ and $\mu(x, y, z)$ are at distance $\leq KR + K$.*
- (6) *If $A \subseteq X$ is a D -quasiconvex subset and $B \subseteq X$ is a set with $d_{\text{Haus}}(A, B) \leq D$, for some $D \geq 0$, then B is $2(CD + C + D)$ -quasiconvex.*

Proof Part (1) follows from the general principle explained above, since for $K = 0$ the stated approximate identity becomes the so-called *5-point identity* for median algebras. The fact that identities (M0) and (M1) imply the 5-point identity is a surprisingly difficult result originally proven by Kolibiar and Marcisová [24, Theorem 2] using an argument of Sholander [28]; see [10, Section 26.4] for further discussion. (Another way to prove this would be to go through the proof of the 5-point identity and use (CM1) and (CM2) to check that at every step the identities hold up to bounded distance.)

Parts (2) and (3) were shown in [27]: they follow, respectively, from a double application of Lemma 3.6 and from Proposition 3.2(I3) in that article.

We now prove part (4). Suppose first that $R = 0$. In this case we have $a, b \in [x, y]$, and hence $\mu(x, y, a) \approx_C a$ and $\mu(x, y, b) \approx_C b$, by identity (CM1). Given $p \in [a, b]$, we can write $p = \mu(a, b, p')$ for some $p' \in X$ and, using part (1) and identity (CM2), we obtain

$$\mu(\mu(a, b, p'), x, y) \approx_K \mu(\mu(a, x, y), \mu(b, x, y), p') \approx_{2C^2+2C} \mu(a, b, p') = p.$$

This shows that $[a, b]$ is contained in the neighbourhood of $[x, y]$ of radius $K + 2C^2 + 2C$, as required. For the general case, it suffices to consider points $a', b' \in [x, y]$ at distance $\leq R$ from a and b , respectively. The interval $[a, b]$ is then contained in the $2(CR + R)$ -neighbourhood of $[a', b']$, by a double application of identity (CM2), and $[a', b']$ is contained in the $(K + 2C^2 + 2C)$ -neighbourhood of $[x, y]$ by the previous discussion.

Regarding part (5), the point $\mu(a, b, c)$ lies in the intersection $[a, b] \cap [b, c] \cap [c, a]$. By part (4), the latter is contained in $\mathcal{N}_{R'}([x, y]) \cap \mathcal{N}_{R'}([y, z]) \cap \mathcal{N}_{R'}([z, x])$ for $R' = KR + K$. By part (3), the latter intersection has diameter $\leq KR' + K$ and also contains the point $\mu(x, y, z)$. In conclusion, the distance of $\mu(a, b, c)$ and $\mu(x, y, z)$ is at most $K^2R + K^2 + K$, as desired.

Finally, we address part (6). Consider two points $b_1, b_2 \in B$ and let $a_1, a_2 \in A$ be points with $d(a_i, b_i) \leq D$. By identity (CM2), the interval $[b_1, b_2]$ is contained in the $2(CD + C)$ -neighbourhood of the interval $[a_1, a_2]$. The latter is contained in $\mathcal{N}_D(A) \subseteq \mathcal{N}_{2D}(B)$; hence we obtain the desired conclusion. $\square$

2.3 Ultralimits

Let ω be a nonprincipal ultrafilter on $\mathbb{N}$. Given a sequence of metric spaces (X_n, d_n) and basepoints $p_n \in X$, we can consider the set of all sequences $x_n \in X_n$ such that $\lim_{\omega} d_n(x_n, p_n) < +\infty$, and endow this set with the pseudometric

$$d_{\omega}((x_n), (x'_n)) := \lim_{\omega} d_n(x_n, x'_n).$$

We denote by (X_{ω}, d_{ω}) the quotient metric space of this pseudometric space, which is known as the ω -limit (or *ultralimit*) of the sequence of pointed metric spaces (X_n, p_n) . We think of a sequence of points $x_n \in X_n$ as converging to the point $(x_n) \in X_{\omega}$. We refer to [16, Chapter 10] for generalities on ultralimits and ultrafilters.

To any sequence of subsets $A_n \subseteq X_n$, we can associate the set $A_{\omega} \subseteq X_{\omega}$ formed by all points of the form (a_n) with $a_n \in A_n$ for all n (or, equivalently, only those with $a_n \in A_n$ for ω -all n). Following [9, Section 10],

we say that A_ω is the *strong limit* of the A_n and write $A_n \rightarrow A_\omega$. If $\mathcal{A}_n$ are families of subsets of X_n , we also say that A_ω is a *strong limit* of the $\mathcal{A}_n$ if there exist sets $A_n \in \mathcal{A}_n$ with $A_n \rightarrow A_\omega$. Finally, a set $A \subseteq X_\omega$ is a *weak limit* of the $\mathcal{A}_n$ if each of its bounded subsets is contained in a strong limit of the $\mathcal{A}_n$.

If $\mu_n: X_n^3 \rightarrow X_n$ is a C_n -coarse median and $C_\omega := \lim_\omega C_n < +\infty$, then we obtain a C_ω -coarse median $\mu_\omega: X_\omega^3 \rightarrow X_\omega$, defined (with an abuse) by the formula

$$\mu_\omega((x_n), (y_n), (z_n)) = (\mu_n(x_n, y_n, z_n)).$$

To be precise, one should choose specific sequences x'_n, y'_n, z'_n representing the points $(x_n), (y_n), (z_n)$ and define the coarse median as the point $(\mu_n(x'_n, y'_n, z'_n))$. Choosing different sequences x'_n, y'_n, z'_n can alter the definition of the map μ_ω , because of the additive error in identity (CM2). However, independently of the choices, we obtain a C_ω -coarse median on X_ω , and different choices will yield a C_ω -coarse median that is $3C_\omega$ -close to it. Thus, the corresponding coarse median *structure* on X_ω is unaffected by all choices.

We have a particular kind of ultralimit when the metric spaces (X_n, d_n) are rescaled copies of a fixed metric space (X, d) , namely $(X, \lambda_n d)$ for a sequence $\lambda_n \rightarrow 0$. In this case, the ultralimit is known as an *asymptotic cone* of X . Note that asymptotic cones still depend on the choice of basepoints and ultrafilter, in general. It is important to observe that asymptotic cones of coarse median spaces actually inherit a uniquely defined structure of median algebra, with a Lipschitz median operator. Indeed, since $\lambda_n \rightarrow 0$, the additive errors in identities (CM1) and (CM2) disappear in the limit.

Ultralimits are well-behaved with respect to the notion of rank. An ultralimit of coarse median spaces of rank $\leq r$ is again a coarse median space of rank $\leq r$. All asymptotic cones of a coarse median space of rank $\leq r$ are median algebras of rank $\leq r$; see [2, Proposition 9.3].

We will also need a result of Bowditch that can be used to compare two families of subsets of a coarse median space, based on the behaviour of their strong limits in all asymptotic cones (Lemma 2.4 below). In order to state this, we need to introduce some of Bowditch's notation. (In the original application, roughly, $\mathcal{E}$ is the collection of all quasiflats in X , $\mathcal{F}$ are certain coarsely cubulated subspaces, $\mathcal{E}^\infty(X^\infty)$ are bilipschitz flats in the asymptotic cone, and $\mathcal{F}^\infty(X^\infty)$ are certain cubulated subspaces.)

Let (X, μ) be a coarse median space and let $\mathcal{E}, \mathcal{F}$ be two families of subsets of X . For each asymptotic cone X^∞ of X , let $\mathcal{E}^\infty(X^\infty)$ and $\mathcal{F}^\infty(X^\infty)$ be two families of subsets of X^∞ . We assume that the following conditions are satisfied for all asymptotic cones X^∞ of X (these are a slightly simplified version of Bowditch's conditions in [9, Section 10], which suffices for our purposes).

- (E1) All strong limits of $\mathcal{E}$ lie in $\mathcal{E}^\infty(X^\infty)$, and all strong limits of $\mathcal{F}$ lie in $\mathcal{F}^\infty(X^\infty)$. All elements of $\mathcal{F}^\infty(X^\infty)$ are weak limits of $\mathcal{F}$.
- (E2) Each element of $\mathcal{E}^\infty(X^\infty)$ is a subset of an element of $\mathcal{F}^\infty(X^\infty)$.
- (E3) If some $E \in \mathcal{E}^\infty(X^\infty)$ is contained in a bounded neighbourhood of some $F \in \mathcal{F}^\infty(X^\infty)$, then $E \subseteq F$.

We will use the following result.

Lemma 2.4 [9, Lemma 10.5] *For every $C \geq 1$, there exists a constant $R = R(C)$ such that the following holds. Let (X, μ) be a C -coarse median space, with X geodesic. Let $\mathcal{E}, \mathcal{F}, \mathcal{E}^\infty(\cdot), \mathcal{F}^\infty(\cdot)$ be families of subsets satisfying conditions (E1)–(E3) above. Then each bounded subset of each element of $\mathcal{E}$ is contained in the R -neighbourhood of an element of $\mathcal{F}$.*

2.4 Quasiflats in coarse median spaces

In this subsection we establish a criterion for a quasiflat in a coarse median space to be quasiconvex.

For simplicity, given a constant $C \geq 1$, we refer to (C, C) -quasi-isometric embeddings and (C, C) -quasi-isometries simply as C -quasi-isometric embeddings and C -quasi-isometries. We always equip the spaces $\mathbb{R}^p$ with their ℓ^1 -metric (or with any other metric bi-Lipschitz equivalent to it, e.g., the Euclidean metric). We will need the following classical result; see, e.g., [23, Corollary 2.6] for a proof.

Proposition 2.5 *There exists a constant $D = D(C, p)$ such that, for any C -quasi-isometric embedding $f: \mathbb{R}^p \rightarrow \mathbb{R}^p$, the image of f is D -dense in the codomain.*

We will use the proposition in the following form. To avoid confusion, we specify that we equip subsets of metric spaces with the restricted distance function (rather than with the induced path metric). Also recall that quasiconvexity is always meant in the coarse median sense, and not in a metric sense.

Corollary 2.6 *For all $C \geq 0$ and $p \in \mathbb{N}$, there exists a constant $C_1 = C_1(C, p)$ such that the following holds. Let (X, μ) be a C -coarse median space and let $E \subseteq X$ be a subset C -quasi-isometric to $\mathbb{R}^p$. Suppose that, for every bounded subset $B \subseteq E$, we have $B \subseteq \mathcal{N}_C(Y_B)$ for a C -quasiconvex subset $Y_B \subseteq X$ such that Y_B embeds C -quasi-isometrically in $\mathbb{R}^p$. Then E is C_1 -quasiconvex.*

Proof To begin with, suppose that we actually have a single subset $Y \subseteq X$ such that $E \subseteq \mathcal{N}_C(Y)$ and such that Y is C -quasiconvex and admits a C -quasi-isometric embedding $f: Y \rightarrow \mathbb{R}^p$. Mapping each point of E to a C -close point of Y and then composing with f , we obtain a quasi-isometric embedding $E \rightarrow \mathbb{R}^p$. Since E is itself C -quasi-isometric to $\mathbb{R}^p$, Proposition 2.5 implies that E and Y are at Hausdorff distance at most C' , for a constant C' depending only on C and p . Finally, the fact that Y is C -quasiconvex implies that E is C'' -quasiconvex for some constant $C'' = C''(C, C')$, using Lemma 2.3(6).

Now, we reduce the general case to the previous one by taking an ultralimit (without rescaling!). For this, let C'' be the constant obtained in the previous paragraph, and suppose for the sake of contradiction that there exist X, μ, E , as in the statement of the corollary, for which E is not $(C''+1)$ -quasiconvex. Thus, there are points $x, y \in E$ and $z \in [x, y]$ such that $d(z, E) \geq C'' + 1$. For each $n \in \mathbb{N}$, there exists a C -quasiconvex subset $Y_n \subseteq X$ such that Y_n admits a C -quasi-isometric embedding in $\mathbb{R}^p$ and such that the neighbourhood $\mathcal{N}_C(Y_n)$ contains the intersection $E \cap \mathcal{N}_n(x)$.

Choose a nonprincipal ultrafilter ω on $\mathbb{N}$ and define X_ω as the ω -limit of the constant sequence consisting of copies of the metric space X , each based at the point x . Let $E_\omega \subseteq X_\omega$ and $Y_\omega \subseteq X_\omega$ be the subspaces obtained as the strong limits, respectively, of the constant sequence E and of the sequence Y_n . Also let μ_ω be the ω -limit of the coarse medians μ_n , as defined above. We have that μ_ω is again a

C -coarse median, that E_ω is again C -quasi-isometric to $\mathbb{R}^p$, and that Y_ω is again C -quasiconvex and C -quasi-isometrically embeddable in $\mathbb{R}^p$. In addition, since the neighbourhoods $\mathcal{N}_C(Y_n)$ contain larger and larger balls in the set E , we have gained in the ultralimit that $E_\omega \subseteq \mathcal{N}_C(Y_\omega)$. We are thus in the situation discussed at the beginning of the proof, so E_ω must be C'' -quasiconvex. At the same time, the constant sequences $x_n = x, y_n = y, z_n = z$ converge to points $x_\omega, y_\omega \in E_\omega$ and $z_\omega \in [x_\omega, y_\omega]$ with $d(z_\omega, E_\omega) \geq C'' + 1$, a contradiction. $\square$

3 Coarse medians and (relative) hyperbolicity

3.1 Hyperbolic spaces and the Morse property

In this subsection, we give a short proof of the fact that Gromov-hyperbolic metric spaces admit a unique coarse median operator, a fact originally shown in [26, Theorem 4.2]. We then go on to deduce that Morse subsets of metric spaces are quasiconvex with respect to all coarse median operators. A subset of a metric space $A \subseteq X$ is said to be N -Morse, for a function $N: \mathbb{R}_{\geq 1} \rightarrow \mathbb{R}_{\geq 0}$, if every (L, L) -quasigeodesic in X with endpoints on A is contained in the $N(L)$ -neighbourhood of A . Recall that we always use the term “quasiconvex” with its median meaning, and the term “quasigeodesic” in the metric sense (Remark 2.2).

For both results mentioned above, the key observation is the following.

Lemma 3.1 *Let (X, μ) be a C -coarse median space, with X a geodesic metric space. For any $x, y \in X$, the interval $[x, y]$ contains a $(2C, 4C)$ -quasigeodesic from x to y .*

Proof We will construct a sequence of points $u_0, \dots, u_m \in [x, y]$ with $u_0 = x, u_m = y, d(u_i, u_{i+1}) \leq 2C$ and $d(u_i, u_j) \geq |i - j| - 3$ for all $0 \leq i < j \leq m$. In order to get the required quasigeodesic $\gamma: [0, m] \rightarrow [x, y]$ from x to y , one can then simply define $\gamma(t) := u_{\lfloor t \rfloor}$.

Thus, let $u_0 := x, u_1, \dots, u_m := y$ be a sequence of points in $[x, y]$ such that $d(u_i, u_{i+1}) \leq 2C$ for all i , and such that m is the smallest possible. (The following argument also shows that such a sequence exists.) For some indices i, j , let $\alpha \subseteq X$ be a geodesic from u_i to u_j and consider, for each integer $0 \leq k \leq d(u_i, u_j)$, the point $y_k \in \alpha$ with $d(u_i, y_k) = k$. Let $y'_k := \mu(x, y, y_k)$ be the projection of y_k to the interval $[x, y]$. By identity (CM2), we have $d(y'_k, y'_{k+1}) \leq 2C$ for all k . Moreover, by identity (CM1), the point $y'_0 = \mu(x, y, u_i)$ is at distance at most C from u_i , as we have $u_i \in [x, y]$. Similarly, the point $\mu(x, y, u_j)$ is within distance C of u_j , and within distance $2C$ of the last of the y'_k .

In conclusion, the points y'_k (together with the point $\mu(x, y, u_j)$) give a sequence of at most $d(u_i, u_j) + 3$ steps of length at most $2C$ leading from u_i and u_j within the interval $[x, y]$. By minimality of the integer m in our choice of the points u_i , it follows that $|i - j| \leq d(u_i, u_j) + 3$, as desired. $\square$

Corollary 3.2 *If X is a geodesic hyperbolic space, then X has a unique coarse median structure. More precisely, if X is δ -hyperbolic, there is a constant $D_1 = D_1(\delta, C)$ such that any two C -coarse medians on X are D -close to each other.*

Proof By hyperbolicity, there exist a constant $D' = D'(\delta, C)$ and a map $m: X^3 \rightarrow X$ such that, for any three points $x, y, z \in X$ and any $(2C, 4C)$ -quasigeodesic triangle T with these vertices, the point

$m(x, y, z)$ is at distance at most D' from each of the three sides of T . If μ is any C -coarse median on X , then each of the three intervals $[x, y]_\mu$, $[y, z]_\mu$, $[z, x]_\mu$ contains a $(2C, 4C)$ -quasigeodesic joining its endpoints, by Lemma 3.1. It follows that the point $m(x, y, z)$ is at distance at most D' from each of the intervals $[x, y]_\mu$, $[y, z]_\mu$, $[z, x]_\mu$. Now, Lemma 2.3(3) implies that the points $m(x, y, z)$ and $\mu(x, y, z)$ are at distance at most $\frac{1}{2}D := KD' + K$, for a constant K depending only on C . Thus, any two C -coarse medians on X are D -close. $\square$

In order to handle Morse subsets, we will require the following strengthening of Lemma 3.1.

Lemma 3.3 *Let (X, μ) be a C -coarse median space, with X a geodesic metric space.*

- (1) *Consider points $x, y \in X$ and $p \in [x, y]$. For some $L \geq 1$, let $\alpha \subseteq [x, p]$ and $\beta \subseteq [p, y]$ be (L, L) -quasigeodesics, respectively, from x to p and from p to y . Then, the union $\alpha \cup \beta$ is a (L', L') -quasigeodesic from x to y , for some $L' = L'(C, L)$.*
- (2) *For all $x, y \in X$, each point $p \in [x, y]$ lies on a (C_2, C_2) -quasigeodesic from x to y contained in $[x, y]$, for some constant $C_2 = C_2(C)$.*

Proof We begin with the proof of part (1). Let $K = K(C)$ be the constant provided by Lemma 3.1. For any $x' \in \alpha$ and $y' \in \beta$, we have $\mu(x', p, y') \approx_K p$ by Lemma 3.1(2). It then follows that

$$d(x', p) \approx_K d(x', \mu(x', p, y')) = d(\mu(x', p, x'), \mu(x', p, y')) \leq Cd(x', y') + C,$$

where the last inequality is given by identity (CM2), after using symmetry of μ . As an analogous chain of inequalities applies to the distance $d(p, y')$, we conclude that

$$d(x', p) + d(p, y') \leq 2Cd(x', y') + 2(C + K).$$

Now, setting $\eta := \alpha \cup \beta$ and letting t, s be the times at which $\eta(t) = x'$ and $\eta(s) = y'$, and using the fact that α and β are (L, L) -quasigeodesics, we obtain

$$|t - s| \leq Ld(x', p) + Ld(p, y') + 2L^2 \leq 2CLd(x', y') + 2L(C + K + L).$$

Since the points $x' \in \alpha$ and $y' \in \beta$ were arbitrary, this shows that η is a quasigeodesic with parameters depending only on L and C , proving part (1).

Regarding part (2), consider $x, y \in X$ and a point $p \in [x, y]$. By Lemma 3.1, there exist $(2C, 4C)$ -quasigeodesics $\alpha \subseteq [x, p]$ and $\beta \subseteq [p, y]$ connecting the endpoints of these two intervals. By part (1), the union $\eta := \alpha \cup \beta$ is a quasigeodesic from x to y , with quality only depending on C . The quasigeodesic η is contained in the union $[x, p] \cup [p, y]$, which is contained in the K -neighbourhood of the interval $[x, y]$ by Lemma 2.3(4). Projecting η onto $[x, y]$ yields the required quasigeodesic from x to y . $\square$

Thus, part (2) of Lemma 3.3 immediately implies the following:

Corollary 3.4 *Let (X, μ) be a C -coarse median space, with X a geodesic metric space. Let $A \subseteq X$ be an N -Morse subset. Then A is $N(C_2)$ -quasiconvex, for the constant $C_2 = C_2(C)$ from Lemma 3.3.*

3.2 Relatively hyperbolic spaces

Bowditch showed that the existence of a coarse median structure is invariant under relative hyperbolicity. That is, if a geodesic metric space X is hyperbolic relative to a collection of subspaces $\mathcal{P}$, and if there exists a constant $C \geq 1$ such that each $P \in \mathcal{P}$ admits a C -coarse median, then X itself admits a coarse median [3, Theorem 2.1].

Here, we address the question of the uniqueness of such coarse medians, extending Corollary 3.2 above.

Theorem 3.5 *Let a geodesic metric space X be hyperbolic relative to a collection of subspaces $\mathcal{P}$. Suppose that for all constants $C \geq 1$ there exists a constant $D \geq 1$ such that, for all $P \in \mathcal{P}$, all C -coarse medians on P are D -close to each other. Then X admits at most one coarse median structure.*

Proof Let us suppose that μ and ν are two C -coarse medians on X , for some $C \geq 1$, and show that μ and ν must be close to each other. Each subspace $P \in \mathcal{P}$ is N -Morse in X , for a fixed function N , by [17, Lemma 4.15]. Appealing to Corollary 3.4, it follows that there exists a constant $D' \geq 0$ such that all $P \in \mathcal{P}$ are D' -quasiconvex with respect to both μ and ν . In particular, the restrictions of μ and ν to each $P \in \mathcal{P}$ are coarse medians of bounded quality and, by our hypothesis, there exists a constant $D'' \geq 0$ such that $\mu(p_1, p_2, p_3) \approx_{D''} \nu(p_1, p_2, p_3)$ for all $P \in \mathcal{P}$ and all $p_1, p_2, p_3 \in P$.

Now, consider three arbitrary points $x, y, z \in X$. By [17, Lemma 8.19], these three points have a “barycentre”, which is either a point $b \in X$ or a subspace $B \in \mathcal{P}$. This barycentre is characterised by the fact that there exists a constant $\Delta \geq 0$ with the following property: Whenever α, β, γ are $(2C, 4C)$ -quasigeodesics in X forming a triangle with vertices x, y, z (respectively, opposite to α, β, γ):

- (1) in the former case, α, β, γ all come within distance Δ of the point b ;
- (2) in the latter, there exist points $a, b, c \in B$ such that the quasigeodesic α comes within distance Δ of the points b and c , the quasigeodesic β does of a and c , and finally γ does of a and b .

By Lemma 3.1, we can choose the above quasigeodesics so that $\alpha \subseteq [y, z]_\mu$, $\beta \subseteq [x, z]_\mu$, $\gamma \subseteq [x, y]_\mu$. If the barycentre is a point b , this shows that b is Δ -close to all three intervals $[x, y]_\mu, [y, z]_\mu, [z, x]_\mu$, and hence b is (uniformly) close to $\mu(x, y, z)$, as in the proof of Corollary 3.2. Since the same argument applies to ν , it follows that the point $\nu(x, y, z)$ is also near b , and hence near $\mu(x, y, z)$.

If instead the barycentre is a subspace $B \in \mathcal{P}$, we have that $\mu(x, y, z)$ is within bounded distance of the point $\mu(a, b, c)$ by Lemma 2.3(5), and similarly $\nu(x, y, z)$ is within bounded distance of $\nu(a, b, c)$. Finally, $\mu(a, b, c)$ and $\nu(a, b, c)$ are uniformly close, because $\mu|_B$ and $\nu|_B$ are close, as observed above.

In conclusion, in all cases, we have $\mu(x, y, z) \approx_D \nu(x, y, z)$ for a constant $D \geq 0$ independent of the points $x, y, z \in X$, proving the theorem. $\square$

4 Medians on products

Following [6], we say that a geodesic hyperbolic space Z is *bushy* if there exists a constant $\lambda \geq 0$ such that, for any point $z \in Z$, there exist three ideal points $\xi_1, \xi_2, \xi_3 \in \partial_\infty Z$ with all pairwise Gromov

products based at x bounded above by λ (see, e.g., [11, Definition III.H.3.15] for the definition of Gromov products of ideal points). An $\mathbb{R}$ -tree T is *furry* if removing any point of T results in at least 3 connected components. Note that all asymptotic cones of a bushy hyperbolic space are furry.

The goal of this section is to show that products of furry trees admit a unique structure as a (Lipschitz) median algebra, and deduce from this that products of bushy hyperbolic spaces admit a unique coarse median structure.

4.1 Products of trees

The following result can be deduced by combining various results of Bowditch.

Proposition 4.1 *Let $P := \prod_{1 \leq i \leq p} \mathcal{T}_i$ be a finite product of furry $\mathbb{R}$ -trees. Then P admits a unique Lipschitz median operator.*

Proof Let $n: P^3 \rightarrow P$ be the standard median operator on P , obtained by computing medians coordinatewise. Let ν be some other Lipschitz median operator on P . Let d be the metric on P that is the ℓ^1 product of the metrics on the factors $\mathcal{T}_i$.

Since P has covering dimension equal to p , the median algebra (P, ν) has rank equal to p : one inequality follows from [2, Theorem 2.2 and Lemma 7.6] and the other, e.g., from [7, Proposition 5.6]. By Proposition 2.1, it follows that there exists a metric ρ on P such that (P, ρ) is a median space with induced median operator ν , and such that ρ is bi-Lipschitz equivalent to d . In particular, (P, ρ) is a complete metric space and it has the same induced topology as (P, d) .

Now, [8, Proposition 4.8] implies that the identity map $(P, n) \rightarrow (P, \nu)$ is an isomorphism of median algebras. In other words, $\nu = n$ as desired. $\square$

4.2 Products of hyperbolic spaces

Let $X = \prod_{1 \leq i \leq p} X_i$ be a product of bushy geodesic hyperbolic spaces. By Corollary 3.2, each space X_i admits a unique coarse median structure $[m_i]$. Let $m: X^3 \rightarrow X$ denote any coarse median obtained as the coordinatewise product of the m_i . We refer to $[m]$ as the *standard* coarse median structure on X . Our goal is now to show that this is the only possible coarse median structure on the product X .

Thus, consider an arbitrary coarse median μ on X . We wish to analyse the structure of μ by appealing to Lemma 2.4 above. For this, we need suitable families of subsets $\mathcal{E}, \mathcal{F}, \mathcal{E}^\infty(\cdot), \mathcal{F}^\infty(\cdot)$ of X and its asymptotic cones. The families $\mathcal{E}, \mathcal{E}^\infty(\cdot)$ and $\mathcal{F}^\infty(\cdot)$ will be defined purely in terms of the metric on X and its asymptotic cones, whereas the family $\mathcal{F}$ will only depend on the chosen coarse median μ .

The family $\mathcal{E}$ consists of *maximal flats* in X , namely products $\prod_{1 \leq i \leq p} \gamma_i$ with each $\gamma_i \subseteq X_i$ a geodesic line. Each asymptotic cone X^∞ of X is a product of $\mathbb{R}$ -trees $\prod_{1 \leq i \leq p} T_i$. We similarly define $\mathcal{E}^\infty(X^\infty)$ to be the family of maximal flats in X^∞ , that is, products $\prod_{1 \leq i \leq p} \gamma_i$ with each $\gamma_i \subseteq T_i$ a geodesic line. Finally, $\mathcal{F}^\infty(X^\infty)$ is the family of *metric panels* in X^∞ : products $\prod_{1 \leq i \leq p} \alpha_i$ of geodesics $\alpha_i \subseteq T_i$. Each α_i might be a single point, a geodesic segment, a ray, or a line; thus, elements of $\mathcal{F}^\infty(X^\infty)$ need not be p -dimensional and they might be compact. Note that we have $\mathcal{E}^\infty(X^\infty) \subseteq \mathcal{F}^\infty(X^\infty)$.

Before describing the family $\mathcal{F}$, we need a definition and a lemma.

Definition 4.2 Let (Z, ν) be a coarse median space and $h \geq 1$ a constant.

- (1) A map $f: M \rightarrow Z$, where (M, m) is a median algebra, is an h -quasimorphism if, for all points $x, y, z \in M$, we have $\nu(f(x), f(y), f(z)) \approx_h f(m(x, y, z))$.
- (2) If M is also equipped with a metric, a map $f: M \rightarrow Z$ is an h -strong quasimorphism if it is both an h -quasimorphism and an h -quasi-isometric embedding.
- (3) An n -dimensional h -cuboid in (Z, ν) is an h -quasiconvex subset of Z that is also the image of an h -strong quasimorphism $f: \mathfrak{c} \rightarrow Z$, where $\mathfrak{c}$ is an n -dimensional Euclidean cube.

Recall that p is the number of factors of the product X . Weak and strong limits were defined in [Section 2.3](#).

Lemma 4.3 *There exists a constant $h = h(X, \mu)$ such that the following holds. For every asymptotic cone X^∞ of X , each element of the family $\mathcal{F}^\infty(X^\infty)$ is a weak limit of p -dimensional h -cuboids in X .*

Proof Recall that each asymptotic cone X^∞ is a product $\prod_{1 \leq i \leq p} T_i$ of $\mathbb{R}$ -trees. We need to show that, for every $F \in \mathcal{F}^\infty(X^\infty)$, every bounded subset of F is contained in a strong limit of p -dimensional uniform-quality cuboids of X . Each element of $\mathcal{F}^\infty(X^\infty)$ is exhausted by products $\prod_i \alpha_i$ of geodesic segments $\alpha_i \subseteq T_i$ and, up to enlarging the product, we can assume that none of the α_i is a single point. Thus, it suffices to consider the case when $F = \prod_{1 \leq i \leq p} \alpha_i$ for nontrivial segments $\alpha_i \subseteq T_i$.

The coarse median μ induces a Lipschitz median operator μ^∞ on X^∞ . Since the X_i are bushy by hypothesis, the trees T_i are furry, and thus [Proposition 4.1](#) implies that μ^∞ is the standard median operator on the product of trees. As a consequence, the set $F \subseteq X^\infty$ is convex for μ^∞ , and its vertex set (i.e., the finite set of points all of whose coordinates are endpoints of the α_i) is a median subalgebra of (X^∞, μ^∞) isomorphic to a p -dimensional discrete cube.

Let $f: \{0, 1\}^p \rightarrow X^\infty$ be an injective 0-quasimorphism (i.e., a median morphism) with image the vertex set of F . By [\[9, Lemma 11.2\]](#), there exists a constant $h' = h'(X, \mu)$ such that there is a sequence of h' -quasimorphisms $f_n: \{0, 1\}^p \rightarrow (X, \mu)$ converging strongly to f ; moreover, the images of the f_n are uniformly “straight” in Bowditch’s sense. The latter implies that the f_n can be extended to h -strong quasimorphisms $f_n: \mathfrak{c}_n \rightarrow X$ with h -quasiconvex image, where the $\mathfrak{c}_n$ are p -dimensional Euclidean cubes and $h = h(X, \mu)$; this follows from [Lemmas 9.2 and 9.3 in \[9\]](#), as well as the discussion between them.

In conclusion, the sets $f_n(\mathfrak{c}_n)$ are h -cuboids in (X, μ) and they converge strongly to a subset $C^\infty \subseteq X^\infty$ containing the vertices of the set F . In the limit, the set C^∞ is 0-quasiconvex, that is, convex. Since F is the convex hull of its vertex set (again because μ^∞ is standard), this shows that $F \subseteq C^\infty$. In conclusion, F is a weak limit of h -cuboids, as we wanted. $\square$

We thus define $\mathcal{F}$ as the collection of all p -dimensional h -cuboids in X , for the constant $h = h(X, \mu)$ provided by [Lemma 4.3](#). We can now check that our families satisfy Bowditch’s conditions.

Lemma 4.4 *The families $\mathcal{E}, \mathcal{F}, \mathcal{E}^\infty(\cdot), \mathcal{F}^\infty(\cdot)$ defined above satisfy conditions (E1)–(E3) of [Section 2.3](#).*

Proof Conditions (E2) and (E3) are immediate from the definitions. Regarding Condition (E1), it is also clear that all strong limits of the family $\mathcal{E}$ lie in $\mathcal{E}^\infty(X^\infty)$. Strong limits of the family $\mathcal{F}$ lie in $\mathcal{F}^\infty(X^\infty)$ because the median operator μ^∞ on X^∞ is standard by [Proposition 4.1](#). Finally, every set in $\mathcal{F}^\infty(X^\infty)$ is a weak limit of sets in $\mathcal{F}$ by [Lemma 4.3](#). $\square$

We can finally deduce that maximal flats in X are uniformly quasiconvex:

Proposition 4.5 *There exists a constant $k = k(X, \mu)$ such that all elements of $\mathcal{E}$ are k -quasiconvex.*

Proof By [Lemma 4.4](#), we can apply [Lemma 2.4](#) to the families $\mathcal{E}, \mathcal{F}, \mathcal{E}^\infty(\cdot), \mathcal{F}^\infty(\cdot)$. As a result, there exists a constant $R = R(X, \mu)$ such that every bounded subset of an element of $\mathcal{E}$ is contained in the R -neighbourhood of an element of $\mathcal{F}$.

By construction, all elements of $\mathcal{E}$ are bilipschitz equivalent to $\mathbb{R}^p$ and all elements of $\mathcal{F}$ embed in $\mathbb{R}^p$ uniformly quasi-isometrically. Moreover, the elements of $\mathcal{F}$ are uniformly quasiconvex. Thus, [Corollary 2.6](#) implies that the elements of $\mathcal{E}$ are uniformly quasiconvex, as claimed. $\square$

Corollary 4.6 *There exists a constant $k' = k'(X, \mu)$ such that all products of p geodesic segments in the X_i are k' -quasiconvex.*

Proof Since the spaces X_i are bushy, there exists a constant λ such that for any geodesic segment $\alpha_i \subseteq X_i$, there exist two (λ, λ) -quasigeodesic lines $\gamma_i, \gamma'_i \subseteq X_i$ with the following properties:

- $\alpha_i \subseteq \mathcal{N}_\lambda(\gamma_i) \cap \mathcal{N}_\lambda(\gamma'_i)$;
- for any $D \geq 0$, if $x \in X_i$ satisfies $d(x, \gamma_i), d(x, \gamma'_i) \leq D$, then $d(x, \alpha_i) \leq D + \lambda$.

Now, consider a product $\prod_{1 \leq i \leq p} \alpha_i$ of geodesic segments $\alpha_i \subseteq X_i$, and let $\gamma_i, \gamma'_i \subseteq X_i$ be quasigeodesic lines as above. The products $\prod_{1 \leq i \leq p} \mathcal{N}_\lambda(\gamma_i)$ and $\prod_{1 \leq i \leq p} \mathcal{N}_\lambda(\gamma'_i)$ are k'' -quasiconvex for some $k'' = k''(X, \mu)$, by [Proposition 4.5](#) and [Lemma 2.3\(6\)](#).

Using the two properties above it is then easy to show that $\prod_i \alpha_i$ is uniformly quasiconvex (roughly, given $x, y \in \prod_i \alpha_i$ and z in the interval between them, z is close to intervals with endpoints on $\prod_{1 \leq i \leq p} \gamma_i$ and $\prod_{1 \leq i \leq p} \gamma'_i$ by [Lemma 2.3\(4\)](#); therefore it is close to both these two products by quasiconvexity, and therefore z is close to $\prod_i \alpha_i$). $\square$

Theorem 4.7 *Let $X = \prod_{1 \leq i \leq p} X_i$ for some bushy geodesic hyperbolic spaces X_i . Then there is a unique coarse median structure on X .*

Proof As in the above discussion, let $[m_i]$ be the unique coarse median structures on the X_i , and let $[m]$ be the standard coarse median structure on X . Let μ an arbitrary coarse median on X . [Corollary 4.6](#) yields a constant $k' \geq 0$ such that all products of geodesics $\prod_{1 \leq i \leq p} \alpha_i$ are k' -quasiconvex with respect to μ . In particular, for $x, y \in X$, the intervals $[x, y]_\mu$ are contained in uniformly bounded neighbourhoods of the intervals $[x, y]_m$ (which are at uniformly bounded Hausdorff distance from products of geodesics). An application of [Lemma 2.3\(3\)](#) then yields that m and μ are close, and hence $[m] = [\mu]$ as required. $\square$

Remark 4.8 A product $Y = X \times \mathbb{R}$, where X is an unbounded coarse median space, cannot have a unique coarse median structure. This is essentially because the product structure on Y is not canonical, as

we can replace the map to the $\mathbb{R}$ -factor with any map of the form $(x, y) \mapsto y + f(x)$ for f an unbounded Lipschitz function on X (e.g., the distance function from a basepoint). Different ways of describing Y as a product yield different coarse median structures.

Acknowledgements

We are grateful to the referee for their many useful comments. Fioravanti is supported by Emmy Noether grant 515507199 and grant 541703614 of the Deutsche Forschungsgemeinschaft (DFG).

References

- [1] **J Behrstock, R A Ciceksiz, V Falgas-Ravry**, *Connectivity for square percolation and coarse cubical rigidity in random right-angled Coxeter groups* (2025) [arXiv 2502.18165](#)
- [2] **B H Bowditch**, *Coarse median spaces and groups*, Pacific J. Math. 261:1 (2013) 53–93 [MR](#)
- [3] **B H Bowditch**, *Invariance of coarse median spaces under relative hyperbolicity*, Math. Proc. Cambridge Philos. Soc. 154:1 (2013) 85–95 [MR](#)
- [4] **B H Bowditch**, *Embedding median algebras in products of trees*, Geom. Dedicata 170 (2014) 157–176 [MR](#)
- [5] **B H Bowditch**, *Large-scale rank and rigidity of the Teichmüller metric*, J. Topol. 9:4 (2016) 985–1020 [MR](#)
- [6] **B H Bowditch**, *Quasi-isometric maps between direct products of hyperbolic spaces*, Internat. J. Algebra Comput. 26:4 (2016) 619–633 [MR](#)
- [7] **B H Bowditch**, *Some properties of median metric spaces*, Groups Geom. Dyn. 10:1 (2016) 279–317 [MR](#)
- [8] **B H Bowditch**, *Large-scale rigidity properties of the mapping class groups*, Pacific J. Math. 293:1 (2018) 1–73 [MR](#)
- [9] **B H Bowditch**, *Quasiflats in coarse median spaces* (2022) Available at <https://bhbowditch.com/papers/quasiflats.pdf>
- [10] **B H Bowditch**, *Median algebras* (2024) Available at <https://bhbowditch.com/papers/median-algebras.pdf>
- [11] **M R Bridson, A Haefliger**, *Metric spaces of non-positive curvature*, Grundle Math. Wissen. 319, Springer (1999) [MR](#)
- [12] **J F Brock**, *The Weil–Petersson metric and volumes of 3-dimensional hyperbolic convex cores*, J. Amer. Math. Soc. 16:3 (2003) 495–535 [MR](#)
- [13] **J Brock, B Farb**, *Curvature and rank of Teichmüller space*, Amer. J. Math. 128:1 (2006) 1–22 [MR](#)
- [14] **J Brock, H Masur**, *Coarse and synthetic Weil–Petersson geometry: quasi-flats, geodesics and relative hyperbolicity*, Geom. Topol. 12:4 (2008) 2453–2495 [MR](#)
- [15] **I Chatterji, C Druţu, F Haglund**, *Kazhdan and Haagerup properties from the median viewpoint*, Adv. Math. 225:2 (2010) 882–921 [MR](#)
- [16] **C Druţu, M Kapovich**, *Geometric group theory*, American Mathematical Society Colloquium Publications 63, Amer. Math. Soc., Providence, RI (2018) [MR](#)
- [17] **C Druţu, M Sapir**, *Tree-graded spaces and asymptotic cones of groups*, Topology 44:5 (2005) 959–1058 [MR](#)
- [18] **E Fioravanti**, *On automorphisms and splittings of special groups*, Compos. Math. 159:2 (2023) 232–305 [MR](#)
- [19] **E Fioravanti**, *Coarse-median preserving automorphisms*, Geom. Topol. 28:1 (2024) 161–266 [MR](#)
- [20] **E Fioravanti**, *Growth of automorphisms of virtually special groups* (2025) [arXiv 2501.12321](#)
- [21] **E Fioravanti, I Levcovitz, M Sageev**, *Coarse cubical rigidity*, J. Topol. 17:3 (2024) art. id. e12353 [MR](#)
- [22] **A Hatcher, W Thurston**, *A presentation for the mapping class group of a closed orientable surface*, Topology 19:3 (1980) 221–237 [MR](#)
- [23] **M Kapovich, B Leeb**, *Quasi-isometries preserve the geometric decomposition of Haken manifolds*, Invent. Math. 128:2 (1997) 393–416 [MR](#)
- [24] **M Kolibiar, T Marcisová**, *On a question of J. Hashimoto*, Mat. Časopis Sloven. Akad. Vied 24 (1974) 179–185 [MR](#)

- [25] **G Mangioni**, *Short hierarchically hyperbolic groups, I: Uncountably many coarse median structures* (2024) [arXiv 2410.09232](#)
- [26] **G A Niblo, N Wright, J Zhang**, *A four point characterisation for coarse median spaces*, Groups Geom. Dyn. 13:3 (2019) 939–980 [MR](#)
- [27] **G A Niblo, N Wright, J Zhang**, *Coarse median algebras: the intrinsic geometry of coarse median spaces and their intervals*, Selecta Math. 27:2 (2021) art. id. 20 [MR](#)
- [28] **M Sholander**, *Medians and betweenness*, Proc. Amer. Math. Soc. 5 (1954) 801–807 [MR](#)
- [29] **R Zeidler**, *Coarse median structures and homomorphisms from Kazhdan groups*, Geom. Dedicata 180 (2016) 49–68 [MR](#)

ELIA FIORAVANTI elia.fioravanti@kit.edu

Institute for Algebra and Geometry, Karlsruhe Institute of Technology, Karlsruhe, Germany

ALESSANDRO SISTO a.sisto@hw.ac.uk

Department of Mathematics, Heriot-Watt University, Edinburgh, United Kingdom

Received: February 27, 2025 Revised: May 27, 2025

ALGEBRAIC & GEOMETRIC TOPOLOGY

msp.org/agt

EDITORS

PRINCIPAL ACADEMIC EDITORS

John Etnyre
etnyre@math.gatech.edu
Georgia Institute of Technology

Vesna Stojanoska
vesna@illinois.edu
University of Illinois at Urbana-Champaign

BOARD OF EDITORS

Julie Bergner	University of Virginia jeb2md@eservices.virginia.edu	Daniel Isaksen	Wayne State University isaksen@math.wayne.edu
Steven Boyer	Université du Québec à Montréal boyer.steven@uqam.ca	Thomas Koberda	University of Virginia thomas.koberda@virginia.edu
Tara E Brendle	University of Glasgow tara.brendle@glasgow.ac.uk	Markus Land	JGU Mainz mland@uni-mainz.de
Indira Chatterji	CNRS & Univ. Côte d'Azur (Nice) indira.chatterji@math.cnrs.fr	Christine Lescop	Université Joseph Fourier lescop@ujf-grenoble.fr
Octav Cornea	Université de Montreal cornea@dms.umontreal.ca	Norihiko Minami	OCAMI (Osaka Central Adv. Math. Inst.) norihikominami@gmail.com
Alexander Dranishnikov	University of Florida dranish@math.ufl.edu	Andrés Navas	Universidad de Santiago de Chile andres.navas@usach.cl
Tobias Ekholm	Uppsala University, Sweden tobias.ekholm@math.uu.se	Jessica S Purcell	Monash University jessica.purcell@monash.edu
Mario Eudave-Muñoz	Univ. Nacional Autónoma de México mario@matem.unam.mx	Birgit Richter	Universität Hamburg birgit.richter@uni-hamburg.de
David Futер	Temple University dfuter@temple.edu	Jérôme Scherer	École Polytech. Féd. de Lausanne jerome.scherer@epfl.ch
John Greenlees	University of Warwick john.greenlees@warwick.ac.uk	Zoltán Szabó	Princeton University szabo@math.princeton.edu
Matthew Hedden	Michigan State University mhedden@math.msu.edu	Maggy Tomova	University of Iowa maggy-tomova@uiowa.edu
Kristen Hendricks	Rutgers University kristen.hendricks@rutgers.edu	Daniel T Wise	McGill University, Canada daniel.wise@mcgill.ca
Hans-Werner Henn	Université Louis Pasteur henn@math.u-strasbg.fr	Lior Yanovski	Hebrew University of Jerusalem lior.yanovski@gmail.com
Kathryn Hess	École Polytechnique Féd. de Lausanne kathryn.hess@epfl.ch		

See inside back cover or msp.org/agt for submission instructions.

The subscription price for 2026 is US \$795/year for the electronic version, and \$1170/year (+\$80, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP. Algebraic & Geometric Topology is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

Algebraic & Geometric Topology (ISSN 1472-2747 printed, 1472-2739 electronic) is published 9 times per year and continuously online, by Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004. Periodical rate postage paid at Oakland, CA 94615-9651, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

AGT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY



mathematical sciences publishers

nonprofit scientific publishing

<https://msp.org/>

© 2026 Mathematical Sciences Publishers

Coxeter quotients of the automorphism group of a Coxeter group	2353
OLGA VARGHESE	
An elementary proof of Novikov’s theorem	2363
SAMUEL RANZ and LAURAN TOUSSAINT	
Magnitude homology of geodesic spaces	2377
KIYONORI GOMI	
(Non)vanishing results for extensions between simple outer functors on free groups	2421
LOUIS HAINAUT	
Whitehead filtrations for computations in topological Hochschild homology	2443
LOGAN HYSLOP	
Around subgroups of Artin groups: derived subgroups and acylindrical hyperbolicity in the even FC-case	2475
JONE LOPEZ DE GAMIZ ZEARRA and CONCHITA MARTÍNEZ-PÉREZ	
Reversible maps in groups of interval exchange transformations	2497
NANCY GUELMAN and ISABELLE LIOUSSE	
On stratifications and poset-stratified spaces	2519
LUKAS WAAS, JON WOOLF and SHOJI YOKURA	
On the girth of groups acting on CAT(0) cube complexes	2539
ARKA BANERJEE, DANIEL L GULBRANDSEN, PRATYUSH MISHRA and PRAYAGDEEP PARIJA	
Commutator-stable representations of hyperbolic groups	2559
ULYSSE REMFORT-AURAT	
Spaces of nonresultant systems of bounded multiplicity with real coefficients	2589
ANDRZEJ KOZŁOWSKI and KOHHEI YAMAGUCHI	
Crossed modules and cohomology of algebras over an operad	2635
JOHAN LERAY, SALIM RIVIÈRE and FRIEDRICH WAGEMANN	
On lens spaces bounding smooth 4-manifolds with $b_2 = 1$	2659
WOOHYEOK JO, JONGIL PARK and KYUNGBAE PARK	
On uniqueness of coarse median structures	2689
ELIA FIORAVANTI and ALESSANDRO SISTO	
A note on a nontrivial connected component of the space of symplectic structures	2705
LUYA WANG	