ChemComm

CORRECTION

Published on 26 October 2026
Licensed under CC-BY 4.0

Check for updates

Cite this: DOI: 10.1039/d6cc90351g

Correction: Modular synthesis and facile network formation of catechol functionalized triblock copolymers

Fahed Albreiki,[a] Tobias Göckler[ab] and Samanvaya Srivastava*[acde]

DOI: 10.1039/d6cc90351g

rsc.li/chemcomm


Correction for 'Modular synthesis and facile network formation of catechol functionalized triblock copolymers' by Fahed Albreiki *et al.*, *Chem. Commun.*, 2024, **60**, 14952–14955, https://doi.org/10.1039/d4cc04503c.

The authors regret that the chemical structure of the triblock polycatechol (compound **5**) is missing a carbon–carbon (C–C) bond in some of the images in the original article. Specifically, an additional C–C bond is missing before the carbon–sulfur (C–S) in the end-block chemical structure. The error appears in the graphical abstract, Scheme 1, and Fig. 2C. The corrected images are provided here.


There was also a missing C–C bond in the ^{1}H NMR peak assignment on page S6, Section 2.3 of the original version of the supplementary information. This has been corrected in an updated supplementary information file.

Please note that the polymers that were synthesized and characterized are as described in the experimental section; only a minor error in the chemical structure and a missing ^{1}H NMR peak assignment were present. This correction does not change any of the findings or conclusions of the paper.

Graphical abstract



[a] Department of Chemical and Biomolecular Engineering, University of California, Los Angeles, Los Angeles, CA 90095, USA. E-mail: samsri@ucla.edu
[b] Institute of Functional Interfaces, Karlsruhe Institute of Technology (KIT), 76344 Eggenstein-Leopoldshafen, Germany
[c] California NanoSystems Institute, University of California, Los Angeles, Los Angeles, CA 90095, USA
[d] BioPACIFIC MIP, University of California, Los Angeles, Los Angeles, CA 90095, USA
[e] Institute for Carbon Management, University of California, Los Angeles, Los Angeles, CA 90095, USA



Scheme 1 Synthesis of catechol-functionalized triblock copolymers.




Fig. 2 ^{1}H NMR spectra of (A) PEG **3**, (B) triblock polymer (P(AGE-*b*-EG-*b*-AGE)) **4**, and (C) tbPC polymer **5**. (D) GPC chromatograms of PEG **3** and triblock polymer (P(AGE-*b*-EG-*b*-AGE)) **4**. (E) DOSY ^{1}H NMR spectra of tbPC polymer dissolved in D_2O at 298 K.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

 This journal is © The Royal Society of Chemistry 2026