

# Topic 6: Extreme events - How will the magnitude and frequency of extreme weather events change in a future climate?

www.reklim.de

## An ensemble assessment of the climate change impact on flood hazard for three small to medium sized catchments in Germany

D. Duethmann<sup>1</sup>, I. Ott<sup>2</sup>; J. Liebert<sup>3</sup>; P. Berg<sup>4</sup>; H. Feldmann<sup>4</sup>; J. Ihringer<sup>3</sup>; H. Kunstmann<sup>2</sup>; B. Merz<sup>1</sup>; G. Schaedler<sup>4</sup>; S. Wagner<sup>2</sup>  
(1) GFZ, Potsdam, Germany (2) IMK-IFU, KIT, Garmisch-Partenkirchen, Germany (3) IWG, KIT, Karlsruhe, Germany (4) IMK-TRO, KIT, Karlsruhe, Germany.



### Introduction

- Aim: quantification of the climate change impact on changes in flood discharges for the near future.
  - ⇒ Important for the adaptation of flood management.
- Scenario assessments based on only one combination of global climate model (GCM), regional climate model (RCM) and hydrological model (HM) might be misleading.
  - ⇒ Ensemble approach: based on 2 GCMs, 2 high-resolution RCMs (7km) and 3 HMs.

Figure 1: Locations of the three selected catchments.





Figure 2: The ensemble approach.

### Methods: study area, climate input, hydrological models

**Study area:** Selection of three catchments representing different flood regimes in Germany: Ruhr (4485 km<sup>2</sup>, winter floods), Mulde (6171 km<sup>2</sup>, floods in winter/spring, but also in summer), Ammer (710 km<sup>2</sup>, summer floods).

#### Ensemble:

- Emission scenario: A1B; only one scenario, as other projects already showed that the emission scenario has only a smaller impact for the near future (2021-2050).
- GCMs: The climate projections are based on three realizations of ECHAM5 and one realization of CCCma3.
- RCMs: All GCM simulations are downscaled by the RCM CLM, and one realization of ECHAM5 is downscaled also with the RCM WRF.
- HMs: In each of the catchments two of three selected HMs (PRMS, SWIM and WaSiMETH) are applied.

### Results

#### Projected changes in meteorological input:

- Temperature: shows an increase (on average +1.1 C, range 0.8-1.5 C) in all catchments and for all ensemble members.
- Precipitation: Only in the Ruhr catchment all ensemble members project a change in the same direction. In the other two catchments, the ensemble mean indicates an increase, but some ensemble members show decreases.

#### Hydrological modeling results control period:

- Despite bias correction of temperature and precipitation, there are still some deviations between simulations with observed climate input and climate input from the RCMs (Fig. 3, left column).
- Caused by: - differences in the precipitation data sets used as model input and for bias correction  
- deviations in climate variables which are not bias corrected (radiation and humidity).




Figure 4: Changes in discharge for given return periods. Values are based on the change of the 2\*30 maximum annual, summer (SH) or winter (WH) values for the period 2021-2050 versus 1971-2000. R1-3: Realizations of ECHAM5. The gray shaded areas mark the maximum range of the single ensemble members. One color implies both HMs.

#### Hydrological modeling results scenario period:

- Large spread of changes in mean monthly maximum discharges and return values in all three catchments (Fig. 3+4, gray area).
- Ammer: the ensemble mean indicates a change in the seasonality with higher flood discharges in winter and lower flood discharges in summer (Fig. 3+4, top row).
- Mulde: no overall trend towards increasing or decreasing flood discharges can be observed (Fig. 3+4, middle row).
- Ruhr: the changes in mean monthly maximum discharge and return values, suggest increasing flood probabilities (Fig. 3+4, bottom row).




Figure 3 Left: Simulated MHQ for the control period 1971-2000 with climate input from observations (SimObs) and the RCMs. Right: Percentage change of MHQ between the scenario (2021-2050) and control (1971-2000) period. The gray shaded areas mark the range of the individual model projections.

### Conclusions

- The ten member ensemble based on 2 GCMs, 2 high-resolution RCMs and 2 HMs demonstrates large uncertainties for the possible impact of climate change on flood hazard in the near future.
- Largest contributions to the overall uncertainty are from the different RCMs in the summer half year, and the different GCMs and their realizations in the winter half year.
- Implications: water infrastructure needs to be planned and designed with these uncertainties, for example by planning water infrastructure in an adaptable way.