

Solubility and spectroscopic study of An^{III}/Ln^{III} in dilute to concentrated Na–Mg–Ca–Cl–NO₃ solutions

M. Herm^{1,*}, X. Gaona¹, Th. Rabung¹, D. Fellhauer¹, C. Crepin², K. Dardenne¹, M. Altmaier¹, H. Geckeis¹

¹ Karlsruhe Institute of Technology, Institute for Nuclear Waste Disposal, P.O. Box 3640, 76021 Karlsruhe, Germany

² Électricité de France SA (EDF), 22–30 Avenue de Wagram, 75008 Paris, France

* michel.herm@kit.edu

Introduction

- Long-term performance assessment of deep geological nuclear waste repositories
→ prediction of chemical behavior of An and long lived FP in aqueous solutions needed.
- Waste disposal in rock-salt formations: WIPP in USA, under consideration in Germany
→ high [Na⁺], [Mg²⁺] and [Cl⁻] expected in water intrusion scenarios.
- Nitrate can be found in high concentrations (≥ 1 M) as part of certain waste forms
→ i.e. waste originated from reprocessing facilities.

- Previous complexation studies with nitrate focused on acidic conditions; no MgCl₂ systems considered.

Objectives of this work

- Assessment of NO₃⁻ effect on Ln(III)/An(III) solubility under repository relevant conditions.
- Development of chemical, thermodynamic and activity models for the system Ln(III)/An(III) in NaCl–NaNO₃, MgCl₂–Mg(NO₃)₂ and CaCl₂–Ca(NO₃)₂ solutions.

Experimental

Solubility experiments

- Batch experiments in Ar atmosphere ($22 \pm 2^\circ\text{C}$)
- Undersaturation approach in 0.1–6.02 m NaCl–NaNO₃, 0.25–5.2 m MgCl₂–Mg(NO₃)₂ and 2.91–4.02 m CaCl₂–Ca(NO₃)₂ mixtures → up to 8.0 m NO₃⁻
- pH range: $7.5 \leq \text{pH}_m \leq 13.2$
- 6–12 mg Nd(OH)₃(am) solid phase in each experiment
- Equilibration time: $t \leq 500$ days

- pH measurements: $\text{pH}_m = -\log m_{\text{H}^+} = \text{pH}_{\text{exp}} + A_m$ [1]; A_m for Cl⁻–NO₃⁻ mixtures determined in this study
- $m_{\text{Nd(III)}}$ measured by ICP-MS after 10 kD (2–3 nm) ultrafiltration
- Solid phase characterization: XRD, SEM-EDX

Nd–L_{III} EXAFS @ ANKA (Karlsruhe, Germany)


- 4.06 m MgCl₂–Mg(NO₃)₂ with 5.81 m NO₃⁻; pH_m = 8.15
- $m_{\text{Nd(III)}} = 1.49 \times 10^{-3}$ m after 10 kD (2–3 nm) ultrafiltration

Cm(III)–TRLFS

- Sample preparation in Ar atmosphere ($22 \pm 2^\circ\text{C}$)
- TRLFS studies in 5.61/6.02 m NaCl–NaNO₃, 0.25/4.1 m MgCl₂–Mg(NO₃)₂ and 4.02 m CaCl₂–Ca(NO₃)₂ mixtures → up to 8.0 m NO₃⁻
- pH range: $1 \leq \text{pH}_m \leq 9$
- [Cm(III)] $\sim 1 \times 10^{-7}$ M per sample


Results and discussion

Solubility of Nd(III) in 5.61 m NaCl–NaNO₃



- Very good agreement with nitrate-free solubility data reported in [2].
- No effect of NO₃⁻ on Nd(OH)₃(am) solubility in NaCl–NaNO₃ systems.

Solubility of Nd(III) in 4.06 m MgCl₂–Mg(NO₃)₂ & 4.02 m CaCl₂–Ca(NO₃)₂



- No effect of NO₃⁻ on Nd(OH)₃(am) solubility in CaCl₂–Ca(NO₃)₂ systems.
- Significant effect of m_{NO3}⁻ on Nd(OH)₃(am) solubility in MgCl₂–Mg(NO₃)₂ systems.
- Slope analysis indicates the formation of aqueous species Mg–Nd–OH–NO₃ with stoichiometries Nd:OH 1:1 (pH_m ≤ 8.3) and Nd:OH 1:2 (pH_m > 8.3).

Chemical and thermodynamic model for the system Ln³⁺/An³⁺–H⁺–Na⁺–Mg²⁺–Ca²⁺–OH⁻–Cl⁻–NO₃⁻

- Solid phase controlling solubility: Nd(OH)₃(am) (XRD, SEM-EDX).
- Ternary/quaternary species with slope -1 and -2 in MgCl₂–Mg(NO₃)₂ systems.
- Formation of Mg[An^{III}/Ln^{III}OHNO₃]³⁺ and Mg[An^{III}/Ln^{III}(OH)₂NO₃]²⁺ indicated by Cm(III)–TRLFS and Nd–L_{III} EXAFS.



$$\log *K^{\circ}_{\text{s},(1,1,1,1)} = 10.80; \beta^{(0)} = 0.93; \beta^{(1)} = 4.30; C^{\phi} = 0$$




$$\log *K^{\circ}_{\text{s},(1,1,2,1)} = 1.56; \beta^{(0)} = 0.18; \beta^{(1)} = 1.60; C^{\phi} = 0$$

Ternary Pitzer parameter generally set to 0.


Interaction of the cation with the anion is assumed to be almost the same for chloride and nitrate.

Nd–L_{III} EXAFS in 4.06 m MgCl₂–Mg(NO₃)₂




- Peak at ~ 4 Å is properly fitted with Mg as backscatterer.
- EXAFS evaluation shows no Nd–Nd interaction.

TRLFS of Cm(III) in Na–Mg–Ca–Cl–NO₃



- No ternary/quaternary species needed to explain TRLFS observations in NaCl–NaNO₃ and CaCl₂–Ca(NO₃)₂ (data not shown).
- Ternary/quaternary species do form in MgCl₂–Mg(NO₃)₂ with pH_m ≥ 4.94 .

Pure component spectra of 4.06 m MgCl₂–Mg(NO₃)₂



- Nitrate effect → genuine complexation reaction!
- Very complex Cm(III) speciation found in MgCl₂–Mg(NO₃)₂ mixtures.

Conclusion

- Nitrate significantly influences solubility of Nd(OH)₃(am) in concentrated and weakly alkaline MgCl₂–Mg(NO₃)₂ solutions at m_{Mg2+} ≥ 2.83 m, m_{NO3}⁻ ≥ 1.13 m and pH_m 8–9.
- No effect of nitrate in NaCl–NaNO₃ and CaCl₂–Ca(NO₃)₂ solutions hinting towards the participation of Mg²⁺ in the complex formation reaction.
- Cm(III)–TRLFS and Nd–L_{III} EXAFS further confirm the participation of Mg²⁺.
- The chemical model proposed includes the formation of the quaternary aqueous species Mg[An^{III}/Ln^{III}OHNO₃]³⁺ and Mg[An^{III}/Ln^{III}(OH)₂NO₃]²⁺ in equilibrium with solid Nd(OH)₃(am).
- Thermodynamic and activity models (Pitzer) derived for the system Nd³⁺/Cm³⁺–H⁺–Mg²⁺–OH⁻–Cl⁻–NO₃⁻.