

Introduction

In the long term safety assessment of repositories for radioactive waste disposal, the event of water intrusion and consequent formation of aqueous systems needs to be evaluated. In the case of repositories in rock salt formations, brine solutions are dominated by NaCl and $MgCl_2$. The use of brucite-based backfill material in such repository concepts is responsible for the buffering of pH_m to ~ 9 . Gluconic acid ($C_6H_{12}O_7$) is a poly-hydroxycarboxylic acid expected in repositories for low and intermediate-level radioactive waste as a component in the formulation of cement.

Formation of stable An(III)/Ln(III)-gluconate/GLU binary complexes has been reported in the literature [1]. The presence of Ca(II) enhances complex stability through formation of ternary species with An(III)/Ln(III) in the hyperalkaline pH range [2]. Despite the relevance of Mg(II) in several repository concepts for radioactive waste disposal, no studies assessing the role of Mg(II) in An(III)/Ln(III)-gluconate complexation have been conducted so far. In $MgCl_2$ -rich brines, the formation of highly stable Mg(II)-gluconate complexes may outcompete the formation of An(III)-gluconate complexes.

Objectives

- Evaluation of the binary Mg(II)-gluconate (GLU) system, deriving the corresponding thermodynamic and activity models
- Assessing the solubility and aqueous speciation of Ln(III)/An(III) in the presence of gluconate and dilute to concentrated $MgCl_2$ solutions in alkaline media
- Deriving comprehensive thermodynamic and activity models for Mg(II)-Ln(III)/An(III)-gluconate systems under repository-relevant conditions
- Providing a sound basis to interpret An(III) behavior in saline systems in the presence of gluconate

Experimental conditions

NMR experiments

- 1H and ^{13}C NMR spectra recorded at 25.0 ± 0.2 °C
- Bruker Avance DRX-500 spectrometer operating at 499.9 MHz for 1H and 125.7 MHz for ^{13}C
- $[GLU]_{tot} = 0.20$ M, $pH_c = 8$ (TRIS)
- NaCl and $MgCl_2$ as background electrolytes:
 - 0.03 M $\leq [MgCl_2]_{tot} \leq 0.27$ M
 - $I = 1.00, 2.00$ M (NaCl-MgCl₂-NaGLU)


Nd(III) – solubility experiments

- Samples prepared and stored at 22 ± 2 °C in Ar-gloveboxes (O_2 content < 2 ppm)
- Undersaturation solubility experiments with Nd(OH)₃(am) (7–15 mg per batch experiment)
- $pH_m = 8$ (TRIS/MES), 9 (Mg(OH)₂ or Mg₂(OH)₃Cl·4H₂O(cr)) (with $pH_m = pH_{exp} + A_m$)
- BGE: 0.25 M – 4.50 M $MgCl_2$
- 10^{-4} M $\leq [GLU]_{tot} \leq 0.10$ M
- Equilibration time: ≤ 360 days
- $[Nd(III)]_{tot}$ determined by ICP-MS after 10 kD ultrafiltration
- Solid phase characterization: XRD

²⁴⁸Cm(III)-TRLFS measurements

- Excitation at $\lambda = 396.6$ nm
- $[Cm(III)]_{tot} = 1 \cdot 10^{-7}$ M
- 0.25 M, 4.50 M $MgCl_2$ and 0.10 M, 4.50 M NaCl as background electrolyte
- $pH_m = 6$ (MES), 8, 8.7 (MOPS/TRIS),
- 10^{-4} M $\leq [GLU]_{tot} \leq 0.10$ M, or
- $5 \leq pH_m \leq 8.7$ ($pH_m = pH_{exp} + A_m$)

Mg(II)-GLU



- Mg(II)-GLU interaction is weaker than for the Ca(II)-GLU system [4]
- rigid coordination structure (octahedral) of Mg(II) aq
- Formation of MgGLU⁺ proposed
- Apparent stability constant in $I = 1.00$ M in NaCl-MgCl₂-NaGLU-system:


M²⁺ + GLU ⇌ MGLU

$$M = Mg \rightarrow \log K' (I = 1 M) = 0.18 \pm 0.41 \text{ (} ^1H, ^{13}C \text{ NMR)}$$

$$\log K' (I = 2 M) = 0.49 \pm 0.06 \text{ (} ^1H, ^{13}C \text{ NMR)}$$

$$M = Ca \rightarrow \log K' (I = 1 M) = 1.02 \pm 0.05 [4]$$

XRD




- [1] TITS, J. et al. "The effect of isosaccharinic acid and gluconic acid on the retention of Eu(III), Am(III) and Th(IV) by calcite" *Applied Geochemistry*, 20, 2082-2096 (2005).
- [2] GAONA, X. et al. "Complexation of An(III) and Ln(III) with gluconate under hyperalkaline pH conditions: solubility and TRLFS studies" (2014). ISSP Conference, Karlsruhe.
- [3] NECK, V. et al. "Thermodynamics of trivalent actinides and neodymium in NaCl, MgCl₂, and CaCl₂ solutions: Solubility, hydrolysis, and ternary Ca-M(III)-OH complexes" *Pure Appl. Chem.*, 81, 1555–1568 (2009)
- [4] PALLAGI, A. et al. "Multinuclear NMR and molecular modelling investigations on the structure and equilibria of complexes that form in aqueous solutions of Ca²⁺ and gluconate" *Carbohydrate Research*, 345, 1856–1864 (2010)
- [5] Joint Committee on Powder Diffraction Standards, 2001. JCPDS-Powder diffraction files. Swarthmore, USA.

References


Mg(II)-An(III)/Ln(III)-GLU

Solubility-Nd(III)



- Series in NaCl and 0.25 M $MgCl_2$ at $pH_c = 8.7$:
 - large increase of intensity and red-shift in wavelength with increasing $[GLU]_{tot}$
 - single isobestic point observed → spectra deconvolution indicates one new Cm(III)-GLU species at ~ 606.2 nm
 - Stoichiometry of the complex: 1:1 or 1:2 (see also [2])
- Spectra series from 4.50 M $MgCl_2$ at $pH_c = 8.7$:
 - less pronounced changes in the speciation
 - no direct evidence for the presence of ternary species
- Spectra collected at $pH_c = 6$ (0.10 M NaCl and 0.25 M $MgCl_2$):
 - complicated speciation scheme
 - continuous shift of peak maxima
 - Step-wise formation of binary Cm(III)-GLU species (?)

TRLFS-Cm(III)



(* peak positions for Cm(III)-Cl-OH species as reported in [3])

Summary and conclusions

- GLU forms weak complexes with Mg^{2+} at $pH_c = 8$. Binary Mg(II)-gluconate species may become relevant in concentrated $MgCl_2$ brines
- GLU importantly increases the solubility of Nd(III) in diluted $MgCl_2$ solutions at pH_{max} , in good agreement with the binary Cm(III)-GLU complex observed by TRLFS
- Solubility and aqueous speciation of Ln(III)/An(III) are less impacted by GLU in concentrated $MgCl_2$ systems:
 - Likely competition between Mg(II) and Ln(III)/An(III) for GLU complexation
 - No exp. evidence on the formation of ternary Mg(II)-Nd(III)-GLU species
- Further exp. studies needed to gain insight on the Mg(II)-Nd(III)-GLU speciation and derive the corresponding thermodynamic and activity models