

A G^2 -subdivision algorithm

Hartmut Prautzsch¹, Georg Umlauf²

Fakultät für Informatik, Universität Karlsruhe, D-76128 Karlsruhe, Germany

Abstract

In this paper we present a method to optimize the smoothness order of subdivision algorithms generating surfaces of arbitrary topology. In particular we construct a subdivision algorithm which produces G^2 -surfaces consisting of bicubic patches. The underlying ideas can also be used to improve the smoothness order of subdivision algorithms for surfaces of higher degree or triangular nets.

Keywords: Catmull/Clark algorithm, subdivision, extraordinary points.

1 Introduction

Subdivision algorithms were first introduced to CAGD by Chaikin [1974] and Lane & Riesenfeld [1980] for the generation of curves and tensor product surfaces. Efforts to understand and analyze these and other subdivision algorithms lead to the general class of stationary subdivision algorithms [Micchelli & Prautzsch '87, Prautzsch '91, Dyn & Levin '92]. These algorithms which operate on regular control nets are a well-understood and rife tool today [Cavaretta et al. '91].

However, most types of surfaces cannot be generated from regular control nets as for example closed surfaces of genus 0. Therefore stationary subdivision algorithms were modified to be applicable to arbitrary control nets with irregular vertices or irregular meshes. [Catmull & Clark '78, Doo & Sabin '78, Loop '87, Dyn et al. '90, Qu '90]. All these algorithms base on a simple common concept: given a control net a new net is computed by simple affine combinations of the given vertices. Iterating this process leads to a sequence of ever denser nets which converge to a continuous surface. The main

¹e-mail: prau@ira.uka.de

²e-mail: umlauf@ira.uka.de

problem for these algorithms is to analyze the smoothness of this limiting surface. First attempts to solve these questions were already made by Doo & Sabin [1978]. Loop [1987] and Ball & Storry [1988] refined their ideas. But in 1993 Reif [1993] showed that all former proofs were incomplete. Recently Reif [1995] and Prautzsch [1996] could construct conditions that guarantee the convergence of subdivision algorithms over irregular meshes to C^1 - and C^k -surfaces respectively. These conditions are the key to analyze existing subdivision algorithms and to improve the smoothness order of their limiting surface.

2 The Catmull/Clark Algorithm

We will show how the general conditions can be applied for the example of the Catmull/Clark algorithm. We investigate its smoothness order and modify the algorithm so as to increase its smoothness order.

First we describe the Catmull/Clark algorithm. Given a quadrilateral net \mathcal{N}_0 it produces a new quadrilateral net \mathcal{N}_1 whose vertices are classified as M-, E-, and V-vertices. Averaging the four vertices of each mesh in \mathcal{N}_0 gives the M-vertices. Averaging the midpoint of each edge common to two adjacent meshes in \mathcal{N}_0 with the M-vertices of the meshes gives the E-Vertices. Finally, computing a weighted average of the $2n + 1$ vertices of all meshes in \mathcal{N}_0 with a common interior vertex gives the V-vertices of \mathcal{N}_1 . The weights are given by the masks in Figure 1.

Connecting each E-vertex with the M- and V-vertices corresponding to the two pairs of meshes and vertices defining both the underlying edge gives the new net \mathcal{N}_1 .

By the same procedure a new net \mathcal{N}_2 is then obtained from \mathcal{N}_1 and so on.

Note that the new nets $\mathcal{N}_i, i \geq 1$, contain only four-sided meshes. Therefore the only extraordinary vertices of a net $\mathcal{N}_i, i \geq 2$, are V-vertices which correspond to an extraordinary vertex of \mathcal{N}_{i-1} . Consequently, all nets $\mathcal{N}_i, i \geq 1$ have the same number of extraordinary vertices and these vertices are the more isolated the higher n is.

If \mathcal{N}_0 is a regular quadrilateral net, then the Catmull/Clark algorithm is simply the subdivision algorithm for the bicubic box splines over a rectangular grid. Thus each regular part of 4×4 vertices of any control net \mathcal{N}_i




Figure 1: The masks of the Catmull/Clark algorithm.

defines a bicubic patch of the limiting surface where the 4×4 vertices are the usual spline control points.

The only interesting parts of the limiting surface are those which are determined by (sub)nets consisting of one extraordinary vertex surrounded by 3 rings of quadrilateral meshes as shown in Figure 2. Let \mathcal{N}_0 denote such a (sub)net and let $\mathbf{p}_1, \dots, \mathbf{p}_m$ be its vertices. Refining \mathcal{N}_0 gives a net \mathcal{N}_1 with 6 rings of quadrilateral meshes around an extraordinary vertex. Discarding the 3 outer rings of control points results in a net \mathcal{N}_1 with the same number of vertices and connectedness as the initial one. Let $\mathbf{q}_1, \dots, \mathbf{q}_m$ be its vertices. Then there is an $m \times m$ matrix A such that

$$[\mathbf{q}_1 \dots \mathbf{q}_m]^t = A[\mathbf{p}_1 \dots \mathbf{p}_m]^t.$$

For later reference consider the surface \mathbf{s}_0 defined by the net \mathcal{N}_0 (and the Catmull/Clark algorithm). It is also given by \mathcal{N}_1 , whereas the subnet $\mathbf{q}_1, \dots, \mathbf{q}_m$ defines only a part \mathbf{s}_1 of \mathbf{s}_0 . The surface ring obtained from \mathbf{s}_0 by taking \mathbf{s}_1 away will be called the *first surface ring associated with \mathcal{N}_0* .

Next consider the class of all subdivision algorithms obtained from the Catmull/Clark algorithm by changing the matrix A . The spectral properties




Figure 2: A control net with an extraordinary vertex of valence 6.

of A determine whether such an algorithm generates regular C^k -manifolds. Specializing the results in [Prautzsch '96] we have

Theorem 2.1 *Let $1, \lambda, \lambda, \mu, \dots, \zeta$ be the m (possibly complex) eigenvalues of A where $1 > |\lambda| > |\mu| \geq \dots \geq |\zeta|$ and assume two eigenvectors \mathbf{c} and \mathbf{d} associated with λ . If the first surface ring of the net given by $[\mathbf{c}_1 \dots \mathbf{c}_m]^t = [\mathbf{c} \mathbf{d}]^t$ is regular without self-intersections and*

$$|\lambda|^r > |\mu|, \quad r = 1, 2,$$

then the limiting surface is a C^r -manifold for almost all initial nets \mathcal{N}_0 . Further, the condition $|\lambda|^r > |\mu|$ is also necessary.

Remark 2.2 *If the limiting surface is a C^2 -manifold, then it has a flat point corresponding to the extraordinary point of \mathcal{N}_0 . In general it is not possible to generate C^k -manifolds by subdivision algorithms consisting of polynomial patches of total bidegree $k + 1$ without flat points, see [Reif '94, Reif & Prautzsch '96].*

Remark 2.3 *Reif was the first to find out the importance of the first surface ring of $[\mathbf{c} \mathbf{d}]$. He called it a characteristic map of A [Reif '95].*

3 A modification of the Catmull–Clark–Algorithm

The matrix A for different n satisfy the C^1 -conditions, but not the C^2 -conditions of the Theorem above [Catmull & Clark '78, Ball & Storry '88]. Moreover, all matrices obtained from a Catmull/Clark matrix A by changing only the weights of the V-masks in Figure 1 do not satisfy the C^2 -conditions [Umlauf '96, Ball & Storry '88].

Still in order to obtain a C^2 -algorithm one can just diagonalize the matrices $A = V\Lambda V^{-1}$, where $\Lambda = \text{diag}(1, \lambda, \lambda, \mu, \dots, \zeta)$, change the modal matrix Λ into $\Lambda' = \text{diag}(1, \lambda, \lambda, \mu', \dots, \zeta')$ and compute the modified subdivision matrix $A' = V\Lambda'V^{-1}$.

Lemma 3.1 *The matrices A and A' have the same characteristic maps.*

Proof Let \mathcal{N}_0 denote the control net given by the eigenvectors of A and also A' associated with λ . Subdividing \mathcal{N}_0 by the Catmull/Clark algorithm and also by the modification results both times in the same net \mathcal{N}_1 . Namely the two outer rings of control points in \mathcal{N}_1 are produced by the subdivision rules for regular nets and the remaining inner control rings are a scaled version of \mathcal{N}_0 with λ being the scaling factor. \square

The symmetry of the Catmull/Clark algorithm masks with respect to the extraordinary point corresponds to a block circulant structure of A [Doo & Sabin '78]. Changing Λ does not affect the block circulance which means that the masks given by A' are again symmetric with respect to the extraordinary point.

Because of the block circulance one can use a block Fourier transformation to bring A into block-diagonal form. Then it suffices to further diagonalize only the blocks belonging to the eigenvalues with modulus in $(\lambda, \lambda^2]$. Changing all eigenvalues of A with modulus in $(\lambda, \lambda^2]$ to 0.3 which is always less than λ^2 one obtains the masks of Figure 3 and Table 4. We listed only the most interesting masks for extraordinary vertices of valence $n = 5, 6, 7, 8, 9$ and 10. For $n = 3$ the M- and E-masks are not changed.

Figure 6 shows an example. The surface on the left was produced by the Catmull/Clark algorithm and the surface on the right by the above modification from the control net shown in Figure 5. The surfaces are shown with a visualization of their Gaussian curvature.




Figure 3: The modified masks for $n = 5$. The weights are listed in Table 4.

Remark 3.2 *Similar to the above modification, one can modify the subdivision algorithms by Loop and Qu so as to obtain C^2 - and C^3 -algorithms, respectively. For details see [Umlauf '96].*

Acknowledgements

We wish to thank Max Ebbers for the implementation of the modified algorithms and for the generation of Figure 6.




Figure 5: The control net used for Figure 6.




Figure 6: The surfaces produced by the Catmull/Clark algorithm (left) and its modification (right).

		$n = 5$		$n = 6$		$n = 7$	
		$\alpha_i =$	$\beta_i =$	$\alpha_i =$	$\beta_i =$	$\alpha_i =$	$\beta_i =$
0 ↓ i ↓ $2n$		0.375000	0.250000	0.375000	0.250000	0.375000	0.250000
		0.365736	0.245100	0.352200	0.232198	0.344994	0.222472
		0.061275	0.243203	0.058049	0.236100	0.055618	0.233758
		0.069980	0.245100	0.073899	0.232198	0.069176	0.222472
		0.003206	0.005498	0.008900	0.006949	0.009944	0.003614
		-0.002857	0.012826	0.011399	0.035603	0.027033	0.039778
		-0.003963	-0.002100	-0.004450	0.006949	-0.018707	0.009824
		-0.002857	-0.015854	-0.022799	-0.017801	-0.011037	-0.010126
		0.003206	-0.002100	-0.004450	-0.013899	-0.018707	-0.044151
		0.069980	0.012826	0.011399	-0.017801	0.002456	-0.010126
	0.061275	0.005498	0.008900	0.006949	0.027033	0.009824	
			0.073899	0.035603	0.009944	0.014633	
			0.058049	0.006949	0.069176	0.039778	
					0.055618	0.003614	
		$n = 8$		$n = 9$		$n = 10$	
		$\alpha_i =$	$\beta_i =$	$\alpha_i =$	$\beta_i =$	$\alpha_i =$	$\beta_i =$
0 ↓ i ↓ $2n$		0.375000	0.250000	0.375000	0.250000	0.375000	0.250000
		0.341666	0.216666	0.325340	0.201818	0.321865	0.195996
		0.054166	0.233333	0.050454	0.224431	0.048999	0.223867
		0.062500	0.216666	0.064116	0.201818	0.057675	0.195996
		0.008333	0.000000	0.011859	0.001802	0.010257	-0.001581
		0.033333	0.033333	0.039981	0.047437	0.042986	0.041030
		0.008333	0.016666	0.008169	0.019952	0.011587	0.021141
		0.000000	0.033333	0.002029	0.032677	0.012630	0.046349
		-0.008333	0.000000	-0.005024	-0.001114	-0.003096	0.004139
		-0.033333	-0.033333	-0.018797	-0.020099	-0.01641	-0.012384
		-0.008333	-0.016666	-0.005917	-0.007855	-0.005247	-0.008075
		0.000000	-0.033333	-0.018797	-0.023668	-0.015611	-0.020991
		0.008333	0.000000	-0.005024	-0.007855	-0.005247	-0.005116
		0.033333	0.033333	0.002029	-0.020099	-0.016419	-0.020991
		0.008333	0.016666	0.008169	-0.001114	-0.003096	-0.008075
	0.062500	0.033333	0.039981	0.032677	0.012630	-0.012384	
	0.054166	0.000000	0.011859	0.019952	0.011587	0.004139	
			0.064116	0.047437	0.042986	0.046349	
			0.050454	0.001802	0.010257	0.021141	
					0.057675	0.041030	
					0.048999	-0.001581	

$$\gamma_0 = \frac{5}{12}, \quad \gamma_{2i-1} = \frac{1}{6}, \quad \gamma_{2i} = \frac{1}{36}, \quad i = 1, \dots, n, \text{ for } n = 3.$$

$$\gamma_0 = \frac{1}{4}, \quad \gamma_{2i-1} = \frac{1}{2n}, \quad \gamma_{2i} = \frac{1}{4n}, \quad i = 1, \dots, n, \text{ for } n \geq 5.$$

Table 4: The weights of the modified masks for $n = 5, 6, 7, 8, 9$ and 10 .

References

- A.A. BALL AND D.J.T. STORRY. Conditions for Tangent Plane Continuity over Recursively Generated B-spline Surfaces. *ACM Transactions on Graphics*, 7(2):83–102, 1988.
- E. CATMULL AND J. CLARK. Recursive generated B-spline surfaces on arbitrary topological meshes. *Computer Aided-Design*, 10(6):350–355, 1978.
- ALFRED S. CAVARETTA, WOLFGANG DAHMEN, AND CHARLES A. MICCHELLI. Stationary Subdivision. Memoirs of the American Mathematical Society, AMS, Providence, Rhode Island, September 1991.
- G.M. CHAIKIN. An Algorithm for High-Speed Curve Generation. *Computer Graphics and Image Processing*, 3:346–349, 1974.
- D.W.H. DOO AND M. SABIN. Behaviour of recursive division surfaces near extraordinary points. *Computer Aided-Design*, 10(6):356–360, 1978.
- NIRA DYN AND DAVID LEVIN. Stationary and Non-Stationary Subdivision Schemes. In T. Lyche and L.L. Schumaker, editors, *Mathematical Methods in CAGD II*, pages 209–216. Academic Press, Inc., Boston, San Diego, New York, Berkeley, Sydney, Tokyo, Toronto, 1992.
- NIRA DYN, DAVID LEVIN, AND J.D. GREGORY. A Butterfly Subdivision Scheme for Surface Interpolation with Tension Control. *ACM Transactions on Graphics*, 9(2):160–169, 1990.
- J.M. LANE AND R.F. RIESENFELD. A Theoretical Development for the Computer Generation and Display of Piecewise Polynomial Surfaces. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2(1):35–46, 1980.
- CHARLES TEORELL LOOP. Smooth Subdivision Surfaces Based on Triangles. Master's thesis, Department of Mathematics, University of Utah, August 1987.
- CHARLES A. MICCHELLI AND HARTMUT PRAUTZSCH. Computing surfaces invariant under subdivision. *Computer Aided Geometric Design*, 4(4):321–328, 1987.

- HARTMUT PRAUTZSCH. Linear Subdivision. *Linear Algebra and its Applications*, 143:223–230, 1991.
- HARTMUT PRAUTZSCH. Analysis of C^k -subdivision surfaces at extraordinary points. Submitted, 1996.
- RUIBIN QU. *Recursive subdivision algorithms for curve and surface design*. PhD thesis, Department of Mathematics and Statistics, Burnel University, Uxbridge, Middlesex, England, August 1990.
- ULRICH REIF. *Neue Aspekte in der Theorie der Freiformflächen beliebiger Topologie*. PhD thesis, Mathematisches Institut A, Universität Stuttgart, Stuttgart, August 1993.
- ULRICH REIF. A degree estimate for subdivision surfaces of higher regularity. To appear in *Proceedings of the AMS*, 1994.
- ULRICH REIF. A unified approach to subdivision algorithms near extraordinary vertices. *Computer Aided Geometric Design*, 12:153–174, 1995.
- ULRICH REIF AND HARTMUT PRAUTZSCH. Necessary conditions for subdivision surfaces. Submitted to *ACM Transactions on Graphics*, 1996.
- GEORG UMLAUF. Verbesserung der Glattheitsordnung von Unterteilungsalgorithmen für Flächen beliebiger Topologie. Master's thesis, IBDS, Universität Karlsruhe, April 1996.